






Defining sepsis in small animals

Stefano Cortellini DVM, MVetMed, DACVECC, DECVECC¹ | Amy E. DeClue DVM, MSc, DACVIM² | Massimo Giunti DVM, DECVECC³ | Robert Goggs BVSc, PhD, DACVECC, DECVECC⁴  | Kate Hopper DVM, PhD, DACVECC⁵ | Julie M. Menard DVM DACVECC⁶ | Rodrigo C. Rabelo DVM, DSc⁷  | Elizabeth A. Rozanski DVM, DACVECC⁸  | Claire R. Sharp DVM, DACVECC⁹  | Deborah C. Silverstein DVM, DACVECC¹⁰ | Virginia Sinnott-Stutzman DVM, DACVECC¹¹  | Giacomo Stanzani DVM, MVetMed, DACVECC, DECVECC¹²

¹Department of Clinical Science and Services, The Royal Veterinary College, University of London, Hatfield, UK

²Fetch Specialty and Emergency Veterinary Center, Greenville, South Carolina, USA

³Department of Veterinary Medical Sciences, Alma Mater Studiorum, University of Bologna, Bologna, Italy

⁴Department of Clinical Sciences, Cornell University College of Veterinary Medicine, Ithaca, New York, USA

⁵Department of Veterinary Surgical and Radiological Sciences, School of Veterinary Medicine, University of California, Davis, Davis, California, USA

⁶Department of Veterinary Clinical and Diagnostic Sciences, Faculty of Veterinary Medicine, University of Calgary, Calgary, Alberta, Canada

⁷Intensivet Veterinary Consulting, Brasília, Brazil

⁸Department of Clinical Sciences, Cummings School of Veterinary Medicine, Tufts University, North Grafton, Massachusetts, USA

⁹School of Veterinary Medicine, Murdoch University, Perth, Western Australia, Australia

¹⁰Department of Clinical Studies and Advanced Medicine, School of Veterinary Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, USA

¹¹BluePearl, Tampa, Florida, USA

¹²Dick White Referrals, Six Mile Bottom, UK

Correspondence

Robert Goggs, Department of Clinical Sciences, Cornell University College of Veterinary Medicine, S1-068 Schurman Hall, 240 Farrier Road, Ithaca, NY 14853, USA.
Email: r.goggs@cornell.edu

Robert Goggs presented some of the material contained within this review at the 2021 European Veterinary Emergency and Critical Care Online Congress.

Funding information

American College of Veterinary Emergency and Critical Care

Abstract

Objective: To discuss the definitions of sepsis in human and veterinary medicine.

Design: International, multicenter position statement on the need for consensus definitions of sepsis in veterinary medicine.

Setting: Veterinary private practice and university teaching hospitals.

Animals: Dogs and cats.

Interventions: None.

Measurements and Main Results: Sepsis is a life-threatening condition associated with the body's response to an infection. In human medicine, sepsis has been defined by consensus on 3 occasions, most recently in 2016. In veterinary medicine, there is

Abbreviations: DAMP, damage-associated molecular pattern; PAMP, pathogen-associated molecular pattern; PIRO, predisposition, infection, response, organ dysfunction; PRR, pattern recognition receptor; qSOFA, quick sequential organ failure assessment; SIRS, systemic inflammatory response syndrome; SOFA, sequential organ failure assessment.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2024 The Authors. *Journal of Veterinary Emergency and Critical Care* published by Wiley Periodicals LLC on behalf of Veterinary Emergency and Critical Care Society.



little uniformity in how sepsis is defined and no consensus on how to identify it clinically. Most publications rely on modified criteria derived from the 1991 and 2001 human consensus definitions. There is a divergence between the human and veterinary descriptions of sepsis and no consensus on how to diagnose the syndrome. This impedes research, hampers the translation of pathophysiology insights to the clinic, and limits our abilities to optimize patient care. It may be time to formally define sepsis in veterinary medicine to help the field move forward. In this narrative review, we present a synopsis of prior attempts to define sepsis in human and veterinary medicine, discuss developments in our understanding, and highlight some criticisms and shortcomings of existing schemes.

Conclusions: This review is intended to serve as the foundation of current efforts to establish a consensus definition for sepsis in small animals and ultimately generate evidence-based criteria for its recognition in veterinary clinical practice.

KEYWORDS

canine, feline, immune response, infection, systemic inflammatory response syndrome

1 | INTRODUCTION

The word “sepsis” and the disorder caused by severe infection have ancient origins,¹ but our understanding of the syndrome has greatly evolved over the last 120 years.² Minor infections that incite a localized host immune response and elicit circumscribed tissue reaction are commonplace and do not constitute sepsis. By contrast, sepsis represents a life-threatening host response to an infection that is overwhelming in its severity or widespread in its extent.^{3,4} Sepsis is a major public health concern and the leading cause of death in human non-cardiac ICUs.⁵ It is estimated that in 2017, almost 49 million adults developed sepsis worldwide resulting in 11 million sepsis-related deaths, accounting for nearly 20% of global mortality.⁶ In the United States alone, the Centers for Disease Control and Prevention estimate that over 1.7 million adults develop sepsis annually, resulting in roughly 350,000 fatalities.⁷ While mortality rates have improved in high-income countries, the global burden of sepsis remains substantial, particularly in low- and middle-income countries, with noticeable heterogeneity in worldwide survival rates.^{8,9}

In veterinary medicine, there are no accurate estimates of the incidence of sepsis, but mortality rates of between 20% and 68% are frequently reported.^{10–14} Mortality rates are higher in patients with greater degrees of physiologic parameter disturbance¹⁵ and in those with organ dysfunction.^{16,17} In people, early diagnosis and subsequent prompt therapeutic intervention are crucial in the management of sepsis.^{18–20} Delays in recognition and initiation of definitive care worsen outcomes.^{21,22} Equivalent evidence is lacking but delayed diagnosis likely contributes to worsened outcomes in animals also. Consequently, the definition of sepsis and its implications for patient care are of great importance.^{23–25} The distinction between a definition and the criteria that facilitate clinical recognition is critical and not simply a question of syntax. A sepsis definition should describe what

sepsis “is.”⁴ In contrast, the syndrome cannot presently be diagnosed using any standardized, validated test and hence sepsis definitions cannot readily be applied to the clinical setting. Consequently, there is a need to establish objective parameters that can be measured in individual animals, and which relate to the pathophysiology and sequelae of the sepsis. To enable early recognition, these parameters should be easy and inexpensive to measure in diverse settings, without requiring costly equipment or specialized laboratories.^{26,27} Rapid identification of sepsis is complicated by the clinical heterogeneity of the syndrome. The PIRO concept (predisposition, infection, response, organ dysfunction) introduced by the human 2001 definition process proposed considering factors that influence this heterogeneity to stage sepsis akin to cancer staging.²⁸ Surprisingly, despite offering insights into sepsis, the PIRO approach has not been widely adopted.^{29,30}

A major challenge facing any attempt to define sepsis is reconciling the distinct needs of clinicians and researchers. In the emergency room and ICU, clinicians require a sensitive definition of sepsis with clinical correlates that facilitate early, accurate diagnoses to ensure all patients with sepsis are identified. An increased false positive rate (overdiagnosis) is preferable to overlooking patients with a potentially life-threatening condition. In contrast, researchers evaluating novel therapies require a more specific sepsis definition to avoid inadvertently dismissing promising therapies as ineffective by testing them in individuals with no likelihood of response. Clinical research can accommodate a higher false negative rate that minimizes enrollment of patients without definitive sepsis. Higher specificity, perhaps achieved by including biomarkers, might enhance study comparability between distinct locations, improve assessment of illness severity, and enable patient stratification during analysis.³¹ Consequently, clinicians might find it easier to allocate resources and prognosticate. Potential drawbacks of more stringent criteria include false negatives, delays in diagnosis, and

TABLE 1 Concepts and terminology from the 1991 human consensus conference.

Term or concept	Definition
Infection	Microbial phenomenon characterized by an inflammatory response to the presence of microorganisms or the invasion of normally sterile host tissue by those organisms
Bacteremia	The presence of viable bacteria in the blood
Sepsis-induced hypotension	A systolic arterial pressure (SAP) <90 mm Hg or a reduction of ≥ 40 mm Hg from baseline in the absence of other causes for hypotension
Multiple Organ Dysfunction Syndrome (MODS)	Presence of altered organ function in an acutely ill patient such that homeostasis cannot be maintained without intervention
Systemic Inflammatory Response Syndrome (SIRS)	The systemic inflammatory response to a variety of severe clinical insults. The response is manifested by 2 or more of the following: <ul style="list-style-type: none"> - Hyperthermia ($>38^{\circ}\text{C}$, $>100.4^{\circ}\text{F}$) or hypothermia ($<36^{\circ}\text{C}$, $<96.8^{\circ}\text{F}$) - Tachycardia ($>90/\text{min}$) - Tachypnea ($>20/\text{min}$) or hyperventilation ($\text{PaCO}_2 < 32$ mm Hg) - Leukocytosis ($>12 \times 10^9/\text{L}$, $>12,000/\mu\text{L}$) or leukopenia ($<4 \times 10^9/\text{L}$, $<4000/\mu\text{L}$) or left shift ($>10\%$)
Sepsis	The systemic response to infection manifested by SIRS resulting from infection
Severe sepsis	Sepsis associated with organ dysfunction, hypoperfusion, or hypotension. Hypoperfusion includes, but is not limited to, lactic acidosis, oliguria, and acute alteration in mental status.
Septic shock	Sepsis-induced hypotension despite adequate fluid resuscitation with hypoperfusion including, but not limited to, lactic acidosis, oliguria, and acute alteration in mental status

increased costs, which may impede access to care or hinder dissemination of research findings. Below, we present the historical context of prior attempts to define and codify sepsis to serve as the foundation of our efforts to establish a consensus definition for sepsis and evidence-based criteria for its recognition in veterinary clinical practice.

2 | SEPSIS-1 (AMERICAN COLLEGE OF CHEST PHYSICIANS/SOCIETY OF CRITICAL CARE MEDICINE CONSENSUS 1991)

Before 1991, it was well-established that sepsis was a common cause of morbidity and mortality and that its incidence was increasing in parallel with changing patient demographics, developments in therapeutic strategies, and the use of new procedures. It was also recognized that the multitude of terms associated with the syndrome affecting these patients was impeding the interpretation and application of clinical trial data. To address these issues, the 1991 consensus conference defined various terms and concepts relevant to sepsis, many of which remain in use (Table 1).³² Three levels of sepsis severity (sepsis, severe sepsis, and septic shock) were defined, and these terms remained in use until 2016.⁴ Central to the 1991 definitions was the concept of the systemic inflammatory response syndrome (SIRS) wherein sepsis was the presence of SIRS due to an infection (Figure 1).³³ This definition derived from the then-prevailing premise that sepsis resulted from a hyperinflammatory response to infection. Inflammation is an important component of the immune response to injury or infection that is usually localized, constrained, controlled, and protective. In contrast, when an inflammatory response to a profound or sustained insult is no longer localized, but generalized, it is termed systemic inflamma-

tion or SIRS.³³ Crucially, the identification of SIRS is highly context dependent. For instance, an exercising dog with physiologic increases in heart rate, respiratory rate, and body temperature meets the established criteria for SIRS, which is nonsensical.¹² Assessments for SIRS criteria are valid only in visibly unwell patients, where clinical signs cannot be adequately explained by pain, anxiety, and medication use, and when the physiologic parameters are altered at rest. Only patients with compatible history, clinical signs, and physical examination findings where there is genuine concern for sepsis should be assessed. The SIRS criteria cease to be informative when applied without this context specificity.

Unfortunately for clinicians, SIRS due to tissue injury resulting from trauma, pancreatitis, thrombosis, or ischemia, and chemical peritonitis from bile or urine leakage, is clinically indistinguishable from SIRS provoked by an infection caused by bacteria, viruses, or fungi.³³ Development of SIRS results from similar biochemical responses to distinct stimuli due to the limited response repertoire of the innate immune system. Pathogens and tissue injury elicit similar responses because they activate the same pattern recognition receptors (PRRs) via pathogen-associated molecular patterns (PAMPs) and damage-associated molecular patterns (DAMPs), respectively.³⁴ The chemical mediators (eg, cytokines and chemokines) released in response to PRR activation generate the host response we recognize clinically as SIRS.³⁵ Microbial molecules including bacterial cell wall components, exotoxins, bacterial and viral DNA, and RNA function as PAMPs, while host molecules such as extracellular histones, neutrophil elastase, heat shock proteins, DNA and RNA, and high mobility group box-1, are DAMPs. Stimulation of membrane-bound PRRs such as Toll-like receptors by either PAMPs or DAMPs activates common intracellular signaling pathways resulting in the clinical manifestations we recognize




FIGURE 1 A schematic representation of sepsis as defined by the 1991 Consensus Conference. Per the 1991 (Sepsis-1) consensus definitions, sepsis is the result of the systemic inflammatory response to infection. The systemic inflammatory response syndrome (SIRS) can result from both infectious or noninfectious insults (eg, pancreatitis, trauma, or, in veterinary patients, syndromes such as gastric dilation volvulus). The combination of infection (bacterial, viral, or fungal) with systemic inflammation is termed sepsis. Severe sepsis is characterized by the development of organ dysfunction, while septic shock represents a subset of these patients in which cardiovascular compromise has occurred. In this scheme, there is a progression in severity from infection to sepsis, from sepsis to severe sepsis, and from severe sepsis to septic shock. Modified from Delano and Ward (2016)¹³¹ after Bone et al (1992).³²

as SIRS. The lack of response specificity poses diagnostic challenges for clinicians attempting to differentiate noninfectious causes of SIRS from sepsis (eg, acute pancreatitis from septic peritonitis).

Despite subsequent criticisms, the 1991 sepsis consensus conference represents a milestone in our understanding and recognition of sepsis. The definitions it established have had a lasting impact on clinical practice and research in both human and veterinary medicine as evidenced by the >8200 citations of the 1992 publication to date. In particular, the 1991 consensus definitions enabled a structured approach to diagnosis, a framework for the evaluation of novel therapies, and the foundation for future guidelines. After the 1991 consensus conference, an epidemiologic survey across 8 academic medical centers defined sepsis as hyper- or hypothermia, tachypnea or the need for mechanical ventilation, tachycardia, and clinical evidence of infection or a positive blood culture.³⁶ The study also used clinical criteria to define severe sepsis and septic shock based on evidence of organ dysfunction or hypotension despite a fluid challenge. The Rivers trial of early goal-directed therapy in sepsis³⁷ combined the 1991 consensus definitions and criteria from the Sands study to define the patient population for their trial. The Rivers study enrolled patients with 2/4 SIRS criteria and hypotension (systolic blood pressure <90 mm Hg) despite a fluid challenge or a plasma lactate concentration >4 mmol/L. The subsequent influence of the Rivers trial on clinical practice helped to embed these cutoffs in the literature.

3 | SEPSIS-2 (INTERNATIONAL SEPSIS CONSENSUS DEFINITIONS 2001; SOCIETY OF CRITICAL CARE MEDICINE, EUROPEAN SOCIETY OF INTENSIVE CARE MEDICINE, AMERICAN COLLEGE OF CHEST PHYSICIANS, AMERICAN THORACIC SOCIETY, SURGICAL INFECTION SOCIETY)

The use of SIRS to define sepsis per the 1991 consensus conference was not universally accepted.³⁸ In particular, the SIRS criteria were felt to be overly sensitive, nonspecific, and poorly reflective of illness severity.³⁹ A second sepsis definitions conference was held in 2001, but concluded that there was insufficient evidence to warrant a change to the definition.³² The diagnostic criteria upon which suspicion of sepsis could be based were expanded to better reflect clinical responses to infection (Table 2),²⁸ but it was recognized that these were inherently arbitrary in the absence of a gold standard against which to calibrate them. Additionally, the 2001 definitions allowed for the use of various organ dysfunction definitions and scores. Use of the multiple organ dysfunction score⁴⁰ or the sequential organ failure assessment (SOFA) score⁴¹ was suggested for adults, while 4 different systems were permitted for use in pediatrics.^{42–45} While this approach offered flexibility in patient assessment and organ dysfunction scoring, it simultaneously reduced the ability of clinicians and investigators to compare studies or trials employing distinct criteria and hence limited research

TABLE 2 Diagnostic criteria for sepsis proposed in 2001 including human cutoff values.

Infection	Documented, or suspected, and some of the following
General variables	Fever (core temperature $>38.3^{\circ}\text{C}$, $>100.9^{\circ}\text{F}$) Hypothermia (core temperature $<36^{\circ}\text{C}$, $<96.8^{\circ}\text{F}$) Heart rate $>90/\text{min}$ or >2 standard deviations (SD) above the normal value for age Tachypnea Altered mental status Significant edema or positive fluid balance (>20 mL/kg over 24 h) Hyperglycemia (plasma glucose >7.7 mmol/L, >120 mg/dL) in the absence of diabetes
Inflammatory variables	Leukocytosis (WBC $>12 \times 10^9/\text{L}$, $>12,000/\mu\text{L}$) Leukopenia (WBC $<4 \times 10^9/\text{L}$, $<4000/\mu\text{L}$) Normal WBC with $>10\%$ immature forms Plasma C-reactive protein (CRP) >2 SD above the normal value Plasma procalcitonin (PCT) >2 SD above the normal value
Hemodynamic variables	Arterial hypotension (SAP <90 mm Hg, MAP <70 mm Hg, or SAP decrease >40 mm Hg in adults or <2 SD below normal for age) $\text{SvO}_2 >70\%$ (adults only) Cardiac index >3.5 L/min
Organ dysfunction variables	Arterial hypoxemia ($\text{PaO}_2/\text{FiO}_2 <300$) Acute oliguria (urine output [UOP] <0.5 mL/kg/h) Creatinine increase (>44.2 $\mu\text{mol/L}$, >0.5 mg/dL) Coagulation abnormalities (INR >1.5 or aPTT >60 s) Ileus (absent bowel sounds) Thrombocytopenia (platelet count $<100 \times 10^9/\text{L}$, $<100,000/\mu\text{L}$) Hyperbilirubinemia (total bilirubin >70 $\mu\text{mol/L}$, >4 mg/dL)
Tissue perfusion variables	Hyperlactatemia (>3 mmol/L or >27 mg/dL) Decreased capillary refill time or mottling

Abbreviations: aPTT, activated partial thromboplastin time; FiO_2 , fraction of inspired oxygen; INR, international normalized ratio; MAP, mean arterial pressure; PaO_2 , partial pressure of oxygen in the arterial blood; SvO_2 , mixed venous oxygen saturation, SAP, systolic blood pressure.

reproducibility and generalizability. This potentially unintended consequence warrants consideration in any future effort to codify a complex syndrome like sepsis.

4 | PREDISPOSITION, INFECTION, RESPONSE, ORGAN DYSFUNCTION (PIRO)

A key issue for research definition and clinical diagnosis of sepsis is disease heterogeneity. A puppy with parvoviral enteritis is markedly different from an adult dog with septic peritonitis due to an intestinal perforation, which is markedly different again from an elderly dog with sepsis secondary to pyelonephritis. Yet, all these patients could be termed “septic.” To address these disparate disease presentations, the human 2001 consensus definitions introduced the PIRO concept (predisposing factors, infection factors, host response, and organ dysfunction) for the staging of sepsis in people (Figure 2).^{28,46} The PIRO model was intended as a concept upon which to base future research,⁴⁷

but was not validated at the time it was proposed,^{48,49} which likely limited its adoption.

4.1 | Predisposing factors

Predisposing factors including age, comorbidities, and concurrent therapies influence host susceptibility to infection and alter the likelihood of sepsis development. Genetic polymorphisms likely play important roles in determining which patients develop sepsis and influence the severity of the resulting syndrome.⁵⁰ In people, epidemiological data and candidate gene investigations suggest that genetic risk factors increase susceptibility to, and severity of, sepsis.^{51,52} Breed predispositions to parvoviral enteritis and pyometra suggest that genetic factors are relevant to sepsis development in small animals also.^{53,54} Veterinary researchers are increasingly applying high-throughput screening technologies to study sepsis,^{55–57} but our understanding of these factors in veterinary medicine is rudimentary.^{58–61}



FIGURE 2 The PIRO concept (predisposition, infection, response, organ dysfunction). The 2001 (Sepsis-2) consensus definitions developed the PIRO concept to better understand the heterogeneity of sepsis and provide a means to “stage” the disease. In this scheme, predisposing factors might include genetic susceptibilities, comorbidities, physiologic reserve, or the colonization by microorganisms expressing antimicrobial drug resistance. The nature of the infection would likely vary with each patient in terms of pathogen type, organism species, virulence factor expression, pathogen load, infection location, and the host immunocompetence. The host response to infection can vary between patients and within an individual over time and could be assessed through measurement of biomarkers of the disease or of the host response. The patient’s physiologic responses to infection manifest as the clinical signs detectable in the emergency room or ICU. Ultimately, in sepsis, the combination of these predisposing factors, the infection, and the resulting host response give rise to organ dysfunction, necessitating intensification of care through organ system support and correspondingly increasing the risk of mortality.

4.2 | Infection factors

The specific type, source, location, organism, virulence, and resistance profile of the pathogen strongly influence the nature, severity, and prognosis of the resulting sepsis syndrome. Most veterinary sepsis studies to date have focused on bacterial^{62–64} or viral infections,^{30,65} with fungal forms of sepsis more rarely recognized.^{66,67} Few studies differentiate the infecting organism or the location of the infection, which likely impacts the utility of study findings and their applicability to other patient populations. Without better delineation of the type of infection, we may be unable to identify the patients who would benefit the most from a particular intervention. For example, in the PROWESS (Protein C Worldwide Evaluation in Severe Sepsis) trial,⁶⁸ patients with urosepsis had significantly lower 28-day mortality compared with patients with a pulmonary source of sepsis. Although only indirectly related to the infection, the timing of sepsis development might also be relevant to outcome. In 1 human study, those who developed septic shock within 24 hours of ICU admission were more severely ill but had better outcomes than those who deteriorated later in their ICU stay.⁶⁹

4.3 | Host response

The host response to infection can vary between patients with the same pathogen, between distinct infections in the same patient, as well as in the same patient over time.² The presence of comorbidities, gastrointestinal microbiome composition,⁷⁰ genetic and epigenetic variation, and the influence of therapeutic interventions all contribute to this variation. Evaluation of host gene expression (transcriptomics) involves the measurement of messenger RNA in blood or circulating leukocytes.⁷¹ This approach has been widely studied in human sepsis, but there are few publications in veterinary medicine to date.^{72–74} The

host response can be assessed with biomarkers to gain insights into the disease process, the patient’s immune response, and their organ function.^{31,75,76} Such assessments may provide a deeper understanding of the host response and enable future treatments to be truly individualized.^{77–81}

4.4 | Organ dysfunction

Conceptually in PIRO, organ dysfunction derives from the interplay of predisposing factors, the infection itself, and the resulting host response, and the model suggests that organ dysfunction in sepsis is preventable. Certainly, the severity of organ dysfunction is an important determinant of prognosis in sepsis in both people and small animals.^{16,17,40} Composite organ failure assessment scores such as the SOFA⁴¹ or the logistic organ dysfunction score⁸² were proposed to aid the quantification of the degree of organ dysfunction while recognizing that measures capable of detecting early cell and tissue damage might be preferable in the future.

5 | PEDIATRIC SEPSIS DEFINITIONS

The 2001 Consensus Conference report included pediatric-specific diagnostic criteria for sepsis, namely, evidence of infection combined with signs and symptoms of inflammation such as hyper- or hypothermia, tachycardia, and evidence of organ dysfunction including altered mental status, hypoxemia, and hyperlactatemia.²⁸ It was recommended that organ dysfunction in children should be scored using previously reported criteria or established systems such as the Pediatric Logistic Organ Dysfunction (PELOD) score.⁴⁵ The 2001 report also noted that owing to their higher basal vascular tone, children

with sepsis exhibit hypotension only when shock is decompensated and blood pressure criteria should not be used to identify cardiovascular dysfunction. Instead, tachycardia, poor peripheral pulses, altered mentation, delayed capillary refill time, and mottled or cool extremities are preferred indicators of shock in childhood sepsis.

The limited pediatric-specific recommendations within the 2001 Consensus Conference report warranted the generation of guidelines focused on children. A pediatric sepsis consensus conference held in 2002 defined sepsis as systemic inflammation secondary to infection,⁸³ but despite this definition being contingent on SIRS criteria, specific cutpoints for children were not set. Instead, variations in physiologic parameters that exceeded 2 standard deviations above or were in the lowest 10th percentile of what is considered normal for the age group were preferred to account for the substantial age-related variations between infants, toddlers, school-age children, and adolescents. Additionally, persistent unexplained heart rate alterations were also considered to provide sufficient evidence of cardiovascular instability. Similar peer-group-referenced, population-indexed parameters were also used to define organ dysfunction. For instance, blood pressure in the lowest 5th percentile or exceeding 2 standard deviations below normal for age indicates cardiovascular dysfunction.

Following the 2016 redefinition of sepsis in adults, it was recognized that variables specific to children for sepsis identification and the prediction of patient-centered outcomes had not been systematically reviewed. To address this gap, the Pediatric Sepsis Definition Taskforce posed 2 questions: (i) In children with infections, what factors are associated with the development of sepsis? (ii) In children with sepsis, what factors help predict mortality?^{84,85} Answering the first question identified clinical predictors of sepsis, while the second enabled the identification of sepsis severity criteria and codified severe forms of the syndrome including septic shock. The systematic review determined that in children with infections, decreased level of consciousness and higher Pediatric Risk of Mortality (PRISM) scores⁸⁶ are associated with sepsis. In children diagnosed with sepsis, comorbidities including cancer; clinicopathologic parameters including plasma lactate, platelet count, fibrinogen, and procalcitonin; and the presence of organ dysfunction are all associated with mortality.⁸⁵ This approach to establishing the evidence basis for clinical sepsis diagnosis and prognostication is well suited for application in veterinary medicine.

6 | SEPSIS-3 (SOCIETY OF CRITICAL CARE MEDICINE, EUROPEAN SOCIETY OF INTENSIVE CARE MEDICINE 2016)

The SIRS criteria provide rapid, simple, and objective means to identify potential systemic inflammation. Nevertheless, it is widely recognized that in people, the SIRS criteria lack both discriminant and convergent validity resulting in excessive false positives and false negatives.^{39,87,88} In 2016, sepsis was redefined using a data-driven approach to enhance the specificity of the associated clinical criteria,^{4,89,90} and with a focus on the host response rather than the pathogen trigger. These changes were also intended to lessen the focus on inflammation as




FIGURE 3 A schematic representation of sepsis as defined by the 2016 Sepsis-3 process. In 2016, sepsis in humans was redefined as life-threatening organ dysfunction caused by a dysregulated host response to infection. The term severe sepsis was eliminated, and septic shock was formally defined as a subset of sepsis in which underlying circulatory and cellular/metabolic abnormalities are profound enough to substantially increase mortality. Clinical criteria derived from the review of large medical record databases were used to “operationalize” the definitions. Recognition of organ dysfunction replaced the identification of the systemic inflammatory response syndrome as the means to identify a dysfunctional host response to infection. Two systems for the identification of organ dysfunction were recommended: (i) the sequential organ failure assessment score (SOFA) primarily for the evaluation of patients in ICUs, and (ii) quick SOFA (qSOFA) consisting of hypotension, altered mentation, and tachypnea for the rapid identification of at-risk patients in non-ICU settings. Modified from Codina and Zeitlinger (2022).¹³²

the driving pathological process in sepsis because it is now known that sepsis involves activation of both pro- and anti-inflammatory responses⁹¹ and is accompanied by dysfunction in the cardiovascular, neuronal, autonomic, hormonal, bioenergetic, metabolic, and coagulation systems.^{92,93} Per the Sepsis-3 consensus, sepsis is the “life-threatening organ dysfunction caused by a dysregulated host response to infection” (Figure 3). The Sepsis-3 definitions removed the term “severe sepsis” to correct the misapprehension that sepsis involves a stepwise progressive deterioration in status and reframe all sepsis as life-threatening. The Sepsis-3 definitions also included a lay description of sepsis: “sepsis is a life-threatening condition that arises when the body’s response to an infection injures its own tissues and organs,” which is a potentially valuable means to convey this complex concept to clients.

To identify organ dysfunction, Sepsis-3 definitions recommend a ≥ 2 -point increase in the well-established SOFA score (Table 3),^{41,94} for which a veterinary equivalent has been evaluated.^{16,95} This represented a substantial shift from the 2001 definitions that were permissive in their use of various published organ function assessment tools. An alternative rapid assessment termed quick sequential organ failure assessment (qSOFA) was also proposed as a means to enable the identification of potential sepsis patients in out-of-hospital, emergency department, or general hospital ward settings.^{89,96} To identify organ dysfunction, the qSOFA score requires 2 or more of altered mentation

TABLE 3 Human sequential organ failure assessment (SOFA) score.

Organ system	Score				
	0	1	2	3	4
Respiratory					
PaO ₂ /FiO ₂	>400	<400	<300	<200	<100
Coagulation					
Platelet count, ×10 ⁹ /L (×10 ³ /μL)	>150	<150	<100	<50	<20
Liver					
Bilirubin, μmol/L	20	20–32	33–101	102–204	204
Bilirubin, mg/dL	<1.2	1.2–1.9	2.0–5.9	9.0–11.9	>12
Cardiovascular					
Blood pressure or catecholamine usage (μg/kg/min for at least 1 h)	MAP > 70 mm Hg	MAP < 70 mm Hg	Dopamine <5 or any dobutamine dose	Dopamine 5.1–15 or epinephrine <0.1 or norepinephrine <0.1	Dopamine >15 or epinephrine >0.1 or norepinephrine >0.1
Central nervous system					
Glasgow coma scale score	15	13–14	10–12	6–9	<6
Kidney					
Creatinine, μmol/L	<110	110–170	171–299	300–440	>440
Creatinine, mg/dL	<1.2	1.2–1.9	2.0–3.4	3.5–4.9	>5.0
Or urine output, mL/h				<500	<200

Abbreviations: FiO₂, fraction of inspired oxygen; MAP, mean arterial pressure; PaO₂, partial pressure of oxygen in the arterial blood.

(a diminished Glasgow Coma Scale score), hypotension (systolic blood pressure ≤100 mm Hg), or tachypnea (respiratory rate ≥22/min). It should be noted that qSOFA is a rapid screen for predictors of mortality rather than a means to identify sepsis specifically and has limited prognostic utility in human studies.^{97–99} Several studies have evaluated modified qSOFA criteria in dogs with variable results. In a critically ill population of dogs, qSOFA was not predictive of mortality in contrast to plasma lactate concentration, but that population had few dogs with sepsis.¹⁰⁰ When used to assess dogs with surgical sepsis where the overall mortality rates were >30%, qSOFA was associated with duration of hospitalization and mortality.^{101,102} In a study of 45 dogs with suspected or confirmed infection admitted to an ICU where the mortality rate was 42%, a composite score of mentation, heart rate, and venous carbon dioxide (PvCO₂) was predictive of outcome.¹⁰³ These results suggest that incorporating organ dysfunction into severity assessments has merit, recognizing that modifications to parameters or criteria to account for biological variation between species may be required.

The 2016 Sepsis-3 process also redefined septic shock in a data-driven manner that incorporated mortality assessments into the clinical criteria. Within the Sepsis-3 consensus panel, there was unanimous agreement that septic shock should reflect a more severe illness with a much higher likelihood of death than sepsis alone. Retrospective databases were then used to determine clinical parameters that identified the subset of patients with high mortality. In a very large registry dataset collected by the Surviving Sepsis Campaign ($n = 28,150$),⁹⁰

inclusion of lactate in the classification criteria enabled clear mortality stratification, as follows:

Fluid resistant hypotension requiring vasopressors + lactate >4 mmol/L = 49.7%;

Fluid resistant hypotension requiring vasopressors + lactate >2 mmol/L = 42.3%;

Fluid resistant hypotension requiring vasopressors (lactate normal) = 30.1%;

Hyperlactatemia >4 mmol/L = 29.9%;

Hyperlactatemia >2 mmol/L = 25.7%.

These cutoffs were evaluated in 2 separate electronic health record data sets and were reproducible. Thus, septic shock is presently defined as a subset of sepsis in which underlying circulatory and cellular/metabolic abnormalities are profound enough to substantially increase mortality. The corresponding clinical criteria are persistent hypotension requiring vasopressors for mean arterial pressure ≥65 mm Hg and a plasma lactate concentration >2 mmol/L after volume resuscitation. With these criteria, the expected mortality for patients in high-income countries with septic shock is >40%, but this figure is unlikely to apply to all middle- and low-income countries.^{104–106}

The Sepsis-3 process has various merits, including an updated pathophysiologic basis, standardized organ dysfunction scoring, clarification of terminology, and the use of large datasets to generate evidence-based clinical correlates. However, it should be acknowledged that the updated definitions have not been universally accepted,^{107,108} with bodies such as the American College of Chest

Physicians publishing statements of opposition.¹⁰⁹ Various criticisms have been leveled at the 2016 definitions including concerns about limited sensitivity in non-ICU settings, a lack of specificity for infection in general, and poor performance of the qSOFA score in diseases that directly cause hypotension, tachypnea, or delirium. For instance, community-acquired pneumonia is a common cause of sepsis in people, but qSOFA is inferior to a validated pneumonia-specific score for mortality prediction.¹¹⁰ Similarly, delayed therapeutic intervention in patients that do not fulfill qSOFA criteria can lead to increased mortality.⁹⁶ As such, while Sepsis-3 represents a paradigm shift in our understanding and identification of sepsis, challenges remain in applying the concepts to human health globally or to veterinary patients.

7 | VETERINARY SEPSIS DEFINITIONS

Before 2016, sepsis in veterinary medicine was typically defined per the premise of the 1991/2001 human consensus definitions, that is, "SIRS plus documented or suspected infection."¹¹¹ Within the literature, and likely within clinical practice, there are variations in what constitutes documentation and suspicion of infection.^{62,112-115} The standard methods described for documenting infection include culture, cytology, and histopathology. For bacterial sepsis, culture and susceptibility testing is ideal to confirm infection, document the organism(s) involved, and identify antimicrobial drug resistance. However, not all organisms can be grown successfully in the laboratory,¹¹⁶ and culture might not adequately resolve polymicrobial infections. Molecular techniques including 16S rRNA metagenomics can increase sensitivity and identification of multiple bacterial species but as yet are not readily available to veterinary clinicians.^{117,118} Nonetheless, molecular techniques designed to detect some pathogens are readily available. Point-of-care tests antigen tests, for canine parvovirus, for example, often have good specificity, but variable sensitivity.¹¹⁹ In some studies and in clinical practice, identification of intracellular organisms within samples collected from anatomic locations typically considered to be sterile is diagnostic for infection. For example, the identification of bacteria within phagolysosomes of neutrophils in aseptically collected abdominal fluid samples is diagnostic of septic peritonitis. Less commonly cytologic diagnosis of infection involves the identification of intracellular pathogens such as hemotropic pathogens, fungal pathogens, or viral or rickettsial inclusions. However, cytology may have limited sensitivity (ie, high false negative rates) and visual identification of organisms does not necessarily indicate an infection is active (ie, identified organisms may be dead). The terms suspected or highly suspected infection are potentially valuable for patient management since they legitimize intensification of diagnostic assessment or therapeutic intervention where strong clinical suspicion for sepsis exists but where samples cannot readily be obtained or initial testing is pending, negative, or inconclusive. However, from a research perspective, the inclusion of cases with suspected but unconfirmed infection in studies of sepsis might bias results by enrolling animals with less severe disease or by including animals that cannot respond to the treat-

ment under study because they do not have sepsis. As such, variation in the stringency of requirements for documentation or confirmation of infection might lead to, or necessitate, the divergence of sepsis diagnostic criteria for clinical trials compared to clinical practice.

Various sets of SIRS criteria have been reported for both dogs and cats with differing sensitivity and specificity (Table 4).¹²⁰⁻¹²³ Many of these sets of criteria have been repeatedly used, but there is no consensus regarding which set to use. Data from human medicine collected prior to 2016 indicate that the 1991 and 2001 sepsis definitions are highly sensitive, but have poor specificity. A study evaluating the diagnostic accuracy of the 1991 and 2001 definitions found the 1991 sepsis definition to be 95% sensitive and 61% specific, while the 2001 definition was 97% sensitive and 58% specific.¹²⁴ Veterinary sepsis definitions derived from these criteria are likely to be similarly affected and legitimate concerns exist regarding construct validity wherein calculations of sensitivity and specificity are predicated on test accuracy that itself may be limited. For instance, in the Hauptman study, the diagnostic criteria for sepsis were evidence of infection and the presence of "systemic illness," a term that was not further described. Questionable construct validity and lack of consistency within the literature create a tension between the demands of patient management and the requirements of clinical trials that cannot adequately be resolved at present. It has been argued that because the primary aim of the SIRS criteria is to identify animals that are systemically unwell and require prompt attention, the exact cutoff points for these parameters are less important than the implication of the derangement, that is, the patient should be thoroughly and urgently assessed. However, a recent study determined that SIRS-positive status was common in small animals presented to the emergency room and primary care, and was only weakly associated with outcome.¹²⁵

Following the publication of Sepsis-3, the human and veterinary definitions are now divergent, further limiting the translation of human sepsis literature to veterinary practice. Moreover, the premise that sepsis is an overexuberant pro-inflammatory response to infection that underpins the use of the SIRS criteria is of questionable validity. Informed by the Sepsis-3 update, the Brazilian Veterinary Emergency and Critical Care Society published consensus definitions for sepsis in 2017,¹²⁸ such that there is now geographic variation in the definition of sepsis in veterinary medicine. For all these reasons, it is apparent that a formal redefinition of sepsis in veterinary medicine is warranted to optimally identify sepsis in the clinic and enable future research endeavors.

To these ends, we have established a steering committee comprising 12 enthusiastic, engaged sepsis experts and identified a larger group of participants to help accomplish this task. In the first phase, we will define, by consensus,⁸³ what we consider sepsis "is," akin to a veterinary dictionary definition.⁴ We will independently generate 12 separate definitions before collating and combining these for iterative refinement via an anonymous Delphi survey.⁹⁰ Following this, we will perform a systematic review of the veterinary literature to answer 2 Population, Exposure, Comparator, Outcome (PECO) format questions. First, we will seek to identify associations between phenotypic factors in animals with infection and negative outcome measures,

TABLE 4 Criteria for the systemic inflammatory response syndrome (SIRS) used in dogs and cats.

Criteria	Dogs (2/4 criteria)			Cats (3/4 criteria)	
	Hauptman 1997 ¹²	de Laforcade 2003 ¹¹	Okano 2002 ¹⁵	Brady 2000 ¹²⁶	DeClue 2011 ¹²⁷
Temperature	>39°C	>39.4°C	>39.7°C	>39.7°C	≥39.7°C
	>102.2°F	>103°F	>103.5°F	>103.5°F	≥103.5°F
	<38°C	<37.8°C	<37.8°C	<37.8°C	≤37.8°C
	<100.4°F	<100°F	<100°F	<100°F	≤100°F
Heart rate	>120/min	>140/min	>160/min	>225/min	≥225/min
	-	-	-	<140/min	≤140/min
Respiratory rate	>20/min	>20/min	>40/min	>40/min	≥40/min
Leukocyte count	>16 × 10 ⁹ /L	>16 × 10 ⁹ /L	>12 × 10 ⁹ /L	>19.5 × 10 ⁹ /L	>19.5 × 10 ⁹ /L
	>16 × 10 ³ /μL	>16 × 10 ³ /μL	>12 × 10 ³ /μL	>19.5 × 10 ³ /μL	>19.5 × 10 ³ /μL
	<6 × 10 ⁹ /L	<6 × 10 ⁹ /L	<4 × 10 ⁹ /L	<5 × 10 ⁹ /L	≤5 × 10 ⁹ /L
	<6 × 10 ³ /μL	<6 × 10 ³ /μL	<4 × 10 ³ /μL	<5 × 10 ³ /μL	≤5 × 10 ³ /μL
Band neutrophils	>3%	>3%	>10%	>5%	≥5%

to identify predictors of sepsis development. Second, in animals with sepsis (however defined), we will attempt to find associations with mortality, to identify predictors of sepsis severity or septic shock. A similar approach was successfully used to identify clinical criteria for pediatric sepsis.^{84,85} We will consider how sepsis should be defined from a scientific perspective to enhance the quality, homogeneity, reproducibility, and generalizability of future research and make recommendations about what data should be collected by future veterinary sepsis studies. The third phase of our work will be to evaluate the diagnostic utility and prognostic value of the clinical criteria established during Phase 2 through retrospective medical record review. Subsequently, we intend to launch a sepsis case registry to prospectively gather multicenter data on small animals with sepsis. We envisage this process will be similar to the Veterinary Committee on Trauma (VetCOT) and Reassessment Campaign on Veterinary CPR (RECOVER) registries for trauma and cardiopulmonary resuscitation.^{129,130} Ultimately, we aspire to reach a point where the data collected prospectively can be used to refine or replace the consensus definitions we initially establish.

8 | SUMMARY

The history of efforts to define, codify, and diagnose sepsis in human and veterinary medicine makes clear that it is not straightforward to describe what sepsis is or to determine how best to identify it. Yet, there is an urgent need to reach a consensus on veterinary definitions of sepsis, to harmonize human and veterinary recommendations, and to provide rational, evidence-based guidance for clinicians. We hope that taking a systematic approach will yield useable definitions and practical clinical criteria that enhance clinical recognition and support research initiatives. Taking a global approach to sepsis in veterinary medicine is important from a One Health perspective and will hopefully lead to widely applicable guidelines and improved patient outcomes.

CONFLICT OF INTEREST STATEMENT

Virginia Sinnott-Stutzman is an assistant editor for the Journal but only participated in the review process as an author. The authors declare no other conflicts of interest.

OFFPRINTS

Offprints will not be available from the authors.

ORCID

Robert Goggs BVSc, PhD, DACVECC, DECVECC <https://orcid.org/0000-0001-7446-6987>

Rodrigo C. Rabelo DVM, DSc <https://orcid.org/0000-0002-8035-687X>

Elizabeth A. Rozanski DVM, DACVECC <https://orcid.org/0000-0003-3233-8930>

Claire R. Sharp DVM, DACVECC <https://orcid.org/0000-0002-1797-9783>

Virginia Sinnott-Stutzman DVM, DACVECC <https://orcid.org/0000-0003-0993-6510>

REFERENCES

- Geroulanos S, Douka ET. Historical perspective of the word "sepsis". *Intensive Care Med.* 2006; 32(12):2077-2077.
- Vincent JL, Abraham E. The last 100 years of sepsis. *Am J Respir Crit Care Med.* 2006;173(3):256-263.
- Berg D, Gerlach H. Recent advances in understanding and managing sepsis. *F1000Res.* 2018;7:F1000.
- Singer M, Deutschman CS, Seymour CW, et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA.* 2016;315(8):801-810.
- Vincent JL, Rello J, Marshall J, et al. International study of the prevalence and outcomes of infection in intensive care units. *JAMA.* 2009;302(21):2323-2329.
- Rudd KE, Johnson SC, Agesa KM, et al. Global, regional, and national sepsis incidence and mortality, 1990–2017: analysis for the Global Burden of Disease Study. *Lancet.* 2020;395(10219):200-211.

7. CDC. *What is Sepsis?* Department of Health and Human Services; 2022.
8. Finfer S, Machado FR. The global epidemiology of sepsis. Does it matter that we know so little? *Am J Respir Crit Care Med.* 2016;193(3):228-230.
9. Machado FR, Azevedo LCP. Sepsis: a threat that needs a global solution. *Crit Care Med.* 2018;46(3):454-459.
10. Parsons KJ, Owen LJ, Lee K, et al. A retrospective study of surgically treated cases of septic peritonitis in the cat (2000-2007). *J Small Anim Pract.* 2009;50(10):518-524.
11. de Laforcade AM, Freeman LM, Shaw SP, et al. Hemostatic changes in dogs with naturally occurring sepsis. *J Vet Intern Med.* 2003;17(5):674-679.
12. Hauptman JG, Walshaw R, Olivier NB. Evaluation of the sensitivity and specificity of diagnostic criteria for sepsis in dogs. *Vet Surg.* 1997;26(5):393-397.
13. King LG. Postoperative complications and prognostic indicators in dogs and cats with septic peritonitis: 23 cases (1989-1992). *J Am Vet Med Assoc.* 1994;204(3):407-414.
14. Greenfield CL, Walshaw R. Open peritoneal drainage for treatment of contaminated peritoneal cavity and septic peritonitis in dogs and cats: 24 cases (1980-1986). *J Am Vet Med Assoc.* 1987;191(1):100-105.
15. Okano S, Yoshida M, Fukushima U, et al. Usefulness of systemic inflammatory response syndrome criteria as an index for prognosis judgement. *Vet Rec.* 2002;150(8):245-246.
16. Ripanti D, Dino G, Piovano G, Farca A. Application of the Sequential Organ Failure Assessment Score to predict outcome in critically ill dogs: preliminary results. *Schweiz Arch Tierheilkd.* 2012;154(8):325-330.
17. Kenney EM, Rozanski EA, Rush JE, et al. Association between outcome and organ system dysfunction in dogs with sepsis: 114 cases (2003-2007). *J Am Vet Med Assoc.* 2010;236(1):83-87.
18. Li Y, Li H, Zhang D. Timing of norepinephrine initiation in patients with septic shock: a systematic review and meta-analysis. *Crit Care.* 2020;24(1):488.
19. Gaieski DF, Mikkelsen ME, Band RA, et al. Impact of time to antibiotics on survival in patients with severe sepsis or septic shock in whom early goal-directed therapy was initiated in the emergency department. *Crit Care Med.* 2010;38(4):1045-1053.
20. Sankar J, Garg M, Ghimire JJ, et al. Delayed administration of antibiotics beyond the first hour of recognition is associated with increased mortality rates in children with sepsis/severe sepsis and septic shock. *J Pediatr.* 2021;233:183-190.
21. Chalfin DB, Trzeciak S, Likourezos A, et al. Impact of delayed transfer of critically ill patients from the emergency department to the intensive care unit. *Crit Care Med.* 2007;35(6):1477-1483.
22. Lin S, Ge S, He W, Zeng M. Association of delayed time in the emergency department with the clinical outcomes for critically ill patients. *QJM.* 2020;114(5):311-317.
23. Laszlo I, Trasy D, Molnar Z, Fazakas J. Sepsis: from pathophysiology to individualized patient care. *J Immunol Res.* 2015;2015:510436.
24. Lyle NH, Pena OM, Boyd JH, Hancock RE. Barriers to the effective treatment of sepsis: antimicrobial agents, sepsis definitions, and host-directed therapies. *Ann N Y Acad Sci.* 2014;1323:101-114.
25. Vandijck D, Decruyenaere JM, Blot SI. The value of sepsis definitions in daily ICU-practice. *Acta Clin Belg.* 2006;61(5):220-226.
26. Rudd KE, Kissoon N, Limmathurotsakul D, et al. The global burden of sepsis: barriers and potential solutions. *Crit Care.* 2018;22(1):232.
27. Rudd KE, Seymour CW, Aluisio AR, et al. Association of the quick sequential (sepsis-related) organ failure assessment (qSOFA) score with excess hospital mortality in adults with suspected infection in low- and middle-income countries. *JAMA.* 2018;319(21):2202-2211.
28. Levy MM, Fink MP, Marshall JC, et al. 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Intensive Care Med.* 2003;29(4):530-538.
29. Otto CM. Sepsis in veterinary patients: what do we know and where can we go? *J Vet Emerg Crit Care.* 2007;17(4):329-332.
30. Alves F, Prata S, Nunes T, et al. Canine parvovirus: a predicting canine model for sepsis. *BMC Vet Research.* 2020;16(1):199.
31. Pierrakos C, Velissaris D, Bisdorff M, et al. Biomarkers of sepsis: time for a reappraisal. *Crit Care.* 2020;24(1):287.
32. Bone RC, Balk RA, Cerra FB, et al. Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine. *Chest.* 1992;101(6):1644-1655.
33. Chakraborty RK, Burns B. Systemic inflammatory response syndrome. In: *StatPearls.* StatPearls Publishing; 2023.
34. Kumar V. Toll-like receptors in sepsis-associated cytokine storm and their endogenous negative regulators as future immunomodulatory targets. *Int Immunopharmacol.* 2020;89(Pt B):107087.
35. Lewis DH, Chan DL, Pinheiro D, et al. The immunopathology of sepsis: pathogen recognition, systemic inflammation, the compensatory anti-inflammatory response, and regulatory T cells. *J Vet Intern Med.* 2012;26(3):457-482.
36. Sands KE, Bates DW, Lancken PN, et al. Epidemiology of sepsis syndrome in 8 academic medical centers. *JAMA.* 1997;278(3):234-240.
37. Rivers E, Nguyen B, Havstad S, et al. Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med.* 2001;345(19):1368-1377.
38. Poeze M, Ramsay G, Gerlach H, et al. An international sepsis survey: a study of doctors' knowledge and perception about sepsis. *Crit Care.* 2004;8(6):R409-R413.
39. Vincent JL. Dear SIRS, I'm sorry to say that I don't like you. *Crit Care Med.* 1997;25:372-374.
40. Marshall JC, Cook DJ, Christou NV, et al. Multiple organ dysfunction score: a reliable descriptor of a complex clinical outcome. *Crit Care Med.* 1995;23(10):1638-1652.
41. Vincent JL, Moreno R, Takala J, et al. The SOFA (sepsis-related organ failure assessment) score to describe organ dysfunction/failure. *Intensive Care Med.* 1996;22(7):707-710.
42. Wilkinson JD, Pollack MM, Ruttimann UE, et al. Outcome of pediatric patients with multiple organ system failure. *Crit Care Med.* 1986;14(4):271-274.
43. Proulx F, Fayon M, Farrell CA, et al. Epidemiology of sepsis and multiple organ dysfunction syndrome in children. *Chest.* 1996;109(4):1033-1037.
44. Doughty L, Carcillo JA, Kaplan S, Janosky J. Plasma nitrite and nitrate concentrations and multiple organ failure in pediatric sepsis. *Crit Care Med.* 1998;26(1):157-162.
45. Leteurtre S, Martinot A, Duhamel A, et al. Development of a pediatric multiple organ dysfunction score: use of two strategies. *Med Decis Making.* 1999;19(4):399-410.
46. Opal SM. Concept of PIRO as a new conceptual framework to understand sepsis. *Pediatr Crit Care Med.* 2005;6(Suppl 3):S55-S60.
47. Rubulotta F, Marshall JC, Ramsay G, et al. Predisposition, insult/infection, response, and organ dysfunction: a new model for staging severe sepsis. *Crit Care Med.* 2009;37(4):1329-1335.
48. Howell MD, Talmor D, Schuetz P, et al. Proof of principle: the predisposition, infection, response, organ failure sepsis staging system. *Crit Care Med.* 2011;39(2):322-327.
49. Caramello V, Macciotta A, Beux V, et al. Validation of the Predisposition Infection Response Organ (PIRO) dysfunction score for the prognostic stratification of patients with sepsis in the Emergency Department. *Med Intensiva.* 2021;45(8):459-469.
50. Russell JA. Genomics and pharmacogenomics of sepsis: so close and yet so far. *Crit Care.* 2016;20(1):185.

51. Cooke GS, Hill AVS. Genetics of susceptibility to human infectious disease. *Nat Rev Genet.* 2001;2(12):967-977.
52. Srinivasan L, Kirpalani H, Cotten CM. Elucidating the role of genomics in neonatal sepsis. *Semin Perinatol.* 2015;39(8):611-616.
53. Houston DM, Ribble CS, Head LL. Risk factors associated with parvovirus enteritis in dogs: 283 cases (1982-1991). *J Am Vet Med Assoc.* 1996;208(4):542-546.
54. Jitpean S, Hagman R, Strom Holst B, et al. Breed variations in the incidence of pyometra and mammary tumours in Swedish dogs. *Reprod Domest Anim.* 2012;47(Suppl 6):347-350.
55. Kuleš J, Horvatić A, Guillemin N, et al. The plasma proteome and the acute phase protein response in canine pyometra. *J Proteomics.* 2020;223:103817.
56. Franco-Martínez L, Horvatić A, Gelemanović A, et al. Changes in the salivary proteome associated with canine pyometra. *Front Vet Sci.* 2020;7:277.
57. Crouser ED, Julian MW, Huff JE, et al. A proteomic analysis of liver mitochondria during acute endotoxemia. *Intensive Care Med.* 2006;32(8):1252-1262.
58. Bauer TR Jr, Pratt SM, Palena CM, et al. Feline leukocyte adhesion (CD18) deficiency caused by a deletion in the integrin $\beta(2)$ (ITGB2) gene. *Vet Clin Pathol.* 2017;46(3):391-400.
59. Felsburg PJ, De Ravin SS, Malech HL, et al. Gene therapy studies in a canine model of X-linked severe combined immunodeficiency. *Hum Gene Ther Clin Dev.* 2015;26(1):50-56.
60. Gu YC, Bauer TR Jr, Ackermann MR, et al. The genetic immunodeficiency disease, leukocyte adhesion deficiency, in humans, dogs, cattle, and mice. *Comp Med.* 2004;54(4):363-372.
61. Perryman LE. Molecular pathology of severe combined immunodeficiency in mice, horses, and dogs. *Vet Pathol.* 2004;41(2):95-100.
62. Goggs R, Letendre JA. Evaluation of the host cytokine response in dogs with sepsis and noninfectious systemic inflammatory response syndrome. *J Vet Emerg Crit Care.* 2019;29(6):593-603.
63. Hagman R. Clinical and molecular characteristics of pyometra in female dogs. *Reprod Domest Anim.* 2012;47(Suppl 6):323-325.
64. Greiner M, Wolf G, Hartmann K. A retrospective study of the clinical presentation of 140 dogs and 39 cats with bacteraemia. *J Small Anim Pract.* 2008;49(8):378-383.
65. Otto CM. Clinical trials in spontaneous disease in dogs: a new paradigm for investigations of sepsis. *J Vet Emerg Crit Care.* 2007;17(4):359-367.
66. Ong RK, Rasis AL, Swindells KL. *Candida albicans* peritonitis in a dog. *J Vet Emerg Crit Care.* 2010;20(1):143-147.
67. Rogers CL, Gibson C, Mitchell SL, et al. Disseminated candidiasis secondary to fungal and bacterial peritonitis in a young dog. *J Vet Emerg Crit Care.* 2009;19(2):193-198.
68. Bernard GR, Vincent JL, Laterre PF, et al. Efficacy and safety of recombinant human activated protein C for severe sepsis. *N Engl J Med.* 2001;344(10):699-709.
69. Roman-Marchant O, Orellana-Jimenez CE, De Backer D, et al. Septic shock of early or late onset: does it matter? *Chest.* 2004;126(1):173-178.
70. Schlechte J, Zucoloto AZ, Yu IL, et al. Dysbiosis of a microbiota-immune metasytem in critical illness is associated with nosocomial infections. *Nat Med.* 2023;29(4):1017-1027.
71. Langley RJ, Wong HR. Early diagnosis of sepsis: is an integrated omics approach the way forward? *Mol Diagn Ther.* 2017;21(5):525-537.
72. Li J, D'Annibale-Tolhurst MA, Adler KB, et al. A myristoylated alanine-rich C kinase substrate-related peptide suppresses cytokine mRNA and protein expression in LPS-activated canine neutrophils. *Am J Respir Cell Mol Biol.* 2013;48(3):314-321.
73. Giunti M, Peli A, Battilani M, et al. Evaluation of CALC-I gene (CALCA) expression in tissues of dogs with signs of the systemic inflammatory response syndrome. *J Vet Emerg Crit Care.* 2010;20(5):523-527.
74. Frangogiannis NG, Mendoza LH, Smith CW, et al. Induction of the synthesis of the C-X-C chemokine interferon-gamma-inducible protein-10 in experimental canine endotoxemia. *Cell Tissue Res.* 2000;302(3):365-376.
75. Vincent JL, Beumier M. Diagnostic and prognostic markers in sepsis. *Expert Rev Anti Infect Ther.* 2013;11(3):265-275.
76. Pierrakos C, Vincent JL. Sepsis biomarkers: a review. *Crit Care.* 2010;14(1):R15.
77. Gunsolus IL, Sweeney TE, Liesenfeld O, Ledebor NA. Diagnosing and managing sepsis by probing the host response to infection: advances, opportunities, and challenges. *J Clin Microbiol.* 2019;57(7):e00425.
78. Schlapbach LJ. Paediatric sepsis. *Curr Opin Infect Dis.* 2019;32(5):497-504.
79. Itenov TS, Murray DD, Jensen JUS. Sepsis: personalized medicine utilizing 'omic' technologies—a paradigm shift? *Healthcare.* 2018;6(3):111.
80. Průcha M, Zazula R, Russwurm S. Sepsis diagnostics in the era of "omics" technologies. *Prague Med Rep.* 2018;119(1):9-29.
81. Eckerle M, Ambroggio L, Puskarich MA, et al. Metabolomics as a driver in advancing precision medicine in sepsis. *Pharmacotherapy.* 2017;37(9):1023-1032.
82. Le Gall JR, Klar J, Lemeshow S, et al. The Logistic Organ Dysfunction system. A new way to assess organ dysfunction in the intensive care unit. *JAMA.* 1996;276(10):802-810.
83. Goldstein B, Giroir B, Randolph A. International pediatric sepsis consensus conference: definitions for sepsis and organ dysfunction in pediatrics. *Pediatr Crit Care Med.* 2005;6(1):2-8.
84. Menon K, Schlapbach LJ, Akech S, et al. Pediatric sepsis definition—a systematic review protocol for the Pediatric Sepsis Definition Taskforce. *Crit Care Explor.* 2020;2(6):e0123.
85. Menon K, Schlapbach LJ, Akech S, et al. Criteria for pediatric sepsis—a systematic review and meta-analysis by the Pediatric Sepsis Definition Taskforce. *Crit Care Med.* 2022;50(1):21-36.
86. Pollack MM, Ruttimann UE, Getson PR. Pediatric risk of mortality (PRISM) score. *Crit Care Med.* 1988;16(11):1110-1116.
87. Kaukonen KM, Bailey M, Pilcher D, et al. Systemic inflammatory response syndrome criteria in defining severe sepsis. *N Engl J Med.* 2015;372(17):1629-1638.
88. Churpek MM, Zdravec FJ, Winslow C, et al. Incidence and prognostic value of the systemic inflammatory response syndrome and organ dysfunctions in ward patients. *Am J Respir Crit Care Med.* 2015;192(8):958-964.
89. Seymour CW, Liu VX, Iwashyna TJ, et al. Assessment of clinical criteria for sepsis: for the Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA.* 2016;315(8):762-774.
90. Shankar-Hari M, Phillips GS, Levy ML, et al. Developing a new definition and assessing new clinical criteria for septic shock: for the Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA.* 2016;315(8):775-787.
91. Hotchkiss RS, Monneret G, Payen D. Sepsis-induced immunosuppression: from cellular dysfunctions to immunotherapy. *Nat Rev Immunol.* 2013;13(12):862-874.
92. Angus DC, van der Poll T. Severe sepsis and septic shock. *N Engl J Med.* 2013;369(9):840-851.
93. Deutschman CS, Tracey KJ. Sepsis: current dogma and new perspectives. *Immunity.* 2014;40(4):463-475.
94. Vincent JL, de Mendonça A, Cantraine F, et al. Use of the SOFA score to assess the incidence of organ dysfunction/failure in intensive care units: results of a multicenter, prospective study. Working group on "sepsis-related problems" of the European Society of Intensive Care Medicine. *Crit Care Med.* 1998;26(11):1793-1800.
95. Goggs RA, Lewis DH. Multiple organ dysfunction syndrome. In: Silverstein DC, Hoppe K, eds. *Small Animal Critical Care Medicine*. 2nd ed. Saunders; 2015:35-46.

96. Machado FR, Cavalcanti AB, Monteiro MB, et al. Predictive accuracy of the quick sepsis-related organ failure assessment score in Brazil. A prospective multicenter study. *Am J Respir Crit Care Med.* 2020;201(7):789-798.
97. Churpek MM, Snyder A, Han X, et al. Quick sepsis-related organ failure assessment, systemic inflammatory response syndrome, and early warning scores for detecting clinical deterioration in infected patients outside the intensive care unit. *Am J Respir Crit Care Med.* 2017;195(7):906-911.
98. Melero-Guijarro L, Sanz-García A, Martín-Rodríguez F, et al. Pre-hospital qSOFA, mSOFA, and NEWS2 performance for sepsis prediction: a prospective, multi-center, cohort study. *Front Med.* 2023;10:1149736.
99. Guarino M, Perna B, De Giorgi A, et al. A 2-year retrospective analysis of the prognostic value of MqSOFA compared to lactate, NEWS and qSOFA in patients with sepsis. *Infection.* 2022;50(4):941-948.
100. Ortolani JM, Bellis TJ. Evaluation of the quick sequential organ failure assessment score plus lactate in critically ill dogs. *J Small Anim Pract.* 2021;62(10):874-880.
101. Stastny T, Koenigshof AM, Brado GE, et al. Retrospective evaluation of the prognostic utility of quick sequential organ failure assessment scores in dogs with surgically treated sepsis (2011-2018): 204 cases. *J Vet Emerg Crit Care.* 2022;32(1):68-74.
102. Donati P, Londoño LA, Tunes M, et al. Retrospective evaluation of the use of quick Sepsis-related Organ Failure Assessment (qSOFA) as predictor of mortality and length of hospitalization in dogs with pyometra (2013-2019): 52 cases. *J Vet Emerg Crit Care.* 2022;32(2):223-228.
103. Lyons B, Stefanovski D, Silverstein D. Development of a novel scoring system to predict mortality in canine patients with infection. *J Vet Emerg Crit Care.* Forthcoming 2024.
104. Machado FR, Assunção MS, Cavalcanti AB, et al. Getting a consensus: advantages and disadvantages of Sepsis 3 in the context of middle-income settings. *Rev Bras Ter Intensiva.* 2016;28(4):361-365.
105. Stephen AH, Montoya RL, Aluisio AR. Sepsis and septic shock in low- and middle-income countries. *Surg Infect.* 2020;21(7):571-578.
106. Adegbite BR, Edoa JR, Ndeze Ndoumba WF, et al. A comparison of different scores for diagnosis and mortality prediction of adults with sepsis in low-and-middle-income countries: a systematic review and meta-analysis. *EClinicalMedicine.* 2021;42:101184.
107. Carneiro AH, Pova P, Gomes JA. Dear Sepsis-3, we are sorry to say that we don't like you. *Rev Bras Ter Intensiva.* 2017;29(1):4-8.
108. Sartelli M, Kluger Y, Ansaloni L, et al. Raising concerns about the Sepsis-3 definitions. *World J Emerg Surg.* 2018;13:6.
109. Simpson SQ. New sepsis criteria: a change we should not make. *Chest.* 2016;149(5):1117-1118.
110. Guo Q, Li H-Y, Song W-D, et al. qSOFA predicted pneumonia mortality better than minor criteria and worse than CURB-65 with robust elements and higher convergence. *Am J Emerg Med.* 2022;52:1-7.
111. Sharp CR. Systemic inflammatory response syndrome, sepsis, and multiple organ dysfunction syndrome. In: Drobatz KJ, Hopper K, Rozanski E, Silverstein DC, eds. *Textbook of Small Animal Emergency Medicine.* 1st ed. John Wiley & Sons; 2019:1030-1037.
112. Burkitt JM, Haskins SC, Nelson RW, Kass PH. Relative adrenal insufficiency in dogs with sepsis. *J Vet Intern Med.* 2007;21(2):226-231.
113. Bentley AM, Otto CM, Shofer FS. Comparison of dogs with septic peritonitis: 1988-1993 versus 1999-2003. *J Vet Emerg Crit Care.* 2007;17(4):391-398.
114. DeClue AE, Sharp CR, Harmon M. Plasma inflammatory mediator concentrations at ICU admission in dogs with naturally developing sepsis. *J Vet Intern Med.* 2012;26(3):624-630.
115. Summers AM, Vezzi N, Gravelyn T, et al. Clinical features and outcome of septic shock in dogs: 37 cases (2008-2015). *J Vet Emerg Crit Care.* 2021;31(3):360-370.
116. Stewart EJ. Growing unculturable bacteria. *J Bacteriol.* 2012;194(16):4151-4160.
117. Rutanga JP, Van Puyvelde S, Heroes AS, et al. 16S metagenomics for diagnosis of bloodstream infections: opportunities and pitfalls. *Expert Rev Mol Diagn.* 2018;18(8):749-759.
118. Yang J, Li Z, Liu Y, Guo S. Tracking bacterial DNA patterns in septic progression using 16s rRNA gene amplicon sequencing analysis. *Int J Clin Exp Pathol.* 2021;14(6):753-767.
119. Walter-Weingärtner J, Bergmann M, Weber K, et al. Comparison of eight commercially available faecal point-of-care tests for detection of canine parvovirus antigen. *Viruses.* 2021;13(10):2080.
120. Staatz AJ, Monnet E, Seim HB 3rd. Open peritoneal drainage versus primary closure for the treatment of septic peritonitis in dogs and cats: 42 cases (1993-1999). *Vet Surg.* 2002;31(2):174-180.
121. Bentley AM, Mayhew PD, Culp WT, Otto CM. Alterations in the hemostatic profiles of dogs with naturally occurring septic peritonitis. *J Vet Emerg Crit Care.* 2013;23(1):14-22.
122. Cortellini S, Pelligand L, Syme H, et al. Neutrophil gelatinase-associated lipocalin in dogs with sepsis undergoing emergency laparotomy: a prospective case-control study. *J Vet Intern Med.* 2015;29(6):1595-1602.
123. Hodgson N, Llewellyn EA, Schaeffer DJ. Utility and prognostic significance of neutrophil-to-lymphocyte ratio in dogs with septic peritonitis. *J Am Anim Hosp Assoc.* 2018;54(6):351-359.
124. Zhao H, Heard SO, Mullen MT, et al. An evaluation of the diagnostic accuracy of the 1991 American College of Chest Physicians/Society of Critical Care Medicine and the 2001 Society of Critical Care Medicine/European Society of Intensive Care Medicine/American College of Chest Physicians/American Thoracic Society/Surgical Infection Society sepsis definition. *Crit Care Med.* 2012;40(6):1700-1706.
125. Spillane AM, Haraschak JL, Gephard SE, et al. Evaluating the clinical utility of the systemic inflammatory response syndrome criteria in dogs and cats presenting to an emergency department. *J Vet Emerg Crit Care.* 2023;33(3):315-326.
126. Brady CA, Otto CM, Van Winkle TJ, King LG. Severe sepsis in cats: 29 cases (1986-1998). *J Am Vet Med Assoc.* 2000;217(4):531-535.
127. DeClue AE, Delgado C, Chang CH, Sharp CR. Clinical and immunologic assessment of sepsis and the systemic inflammatory response syndrome in cats. *J Am Vet Med Assoc.* 2011;238(7):890-897.
128. Castro BGA, Rabelo RC. Sepsis-3: uma análise aplicada à Medicina Veterinária. *J LAVECC.* 2017;9(3):1-12.
129. Hall KE, Rutten JI, Baird TN, et al. ACVECC-Veterinary Committee on Trauma registry report 2017-2019. *J Vet Emerg Crit Care.* 2023;33(3):289-297.
130. Hoehne SN, Balakrishnan A, Silverstein DC, et al. Reassessment Campaign on Veterinary Resuscitation (RECOVER) Initiative small animal CPR registry report 2016-2021. *J Vet Emerg Crit Care.* 2023;33(2):143-155.
131. Delano MJ, Ward PA. The immune system's role in sepsis progression, resolution, and long-term outcome. *Immunol Rev.* 2016;274(1):330-353.
132. Sanz Codina M, Zeitlinger M. Biomarkers predicting tissue pharmacokinetics of antimicrobials in sepsis: a review. *Clin Pharmacokinet.* 2022;61(5):593-617.

How to cite this article: Cortellini S, DeClue AE, Giunti M, et al. Defining sepsis in small animals. *J Vet Emerg Crit Care.* 2024;34:97-109. <https://doi.org/10.1111/vec.13359>