

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

The Interplay Between Respiratory Supercomplexes and ROS in Aging

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Genova, M.L., Lenaz, G. (2015). The Interplay Between Respiratory Supercomplexes and ROS in Aging. *ANTIOXIDANTS & REDOX SIGNALING*, 23(3), 208-238 [10.1089/ars.2014.6214].

Availability:

This version is available at: <https://hdl.handle.net/11585/518458> since: 2015-11-10

Published:

DOI: <http://doi.org/10.1089/ars.2014.6214>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Genova ML, Lenaz G.

The Interplay Between Respiratory Supercomplexes and ROS in Aging.

Antioxid Redox Signal. 2015 Jul 20;23(3):208-38

The final published version is available online at <https://doi.10.1089/ars.2014.6214>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

THE INTERPLAY BETWEEN RESPIRATORY SUPERCOMPLEXES AND ROS IN AGING

Maria Luisa Genova[§] and Giorgio Lenaz

Dipartimento di Scienze Biomediche e Neuromotorie, Alma Mater Studiorum – Università di
Bologna, Via Irnerio 48, 40126, Bologna (Italy)

Running title: Supercomplexes and ROS in Aging

Word count: 15198

Reference numbers: 424

Number of greyscale illustrations: 5

Number of colour illustrations: 0

§ Corresponding address:

Dipartimento di Scienze Biomediche e Neuromotorie, Alma Mater Studiorum – Università di
Bologna, Via Irnerio 48, 40126, Bologna (Italy).

Phone: +39 051 2091214; Fax: +39 051 2091224; email: marialuisa.genova@unibo.it

ABSTRACT

SIGNIFICANCE: The molecular mechanism of aging is still vigorously debated, although a general consensus exists that mitochondria are significantly involved in this process. However, the previously postulated role of mitochondria-derived reactive oxygen species (ROS) as the damaging agents inducing functional loss in aging has fallen out of favour in the recent past. In this review, we critically examine the role of ROS in aging in the light of recent advances on the relationship between mitochondrial structure and function.

RECENT ADVANCES: The functional mitochondrial respiratory chain is now recognized as a reflection of the dynamic association of respiratory complexes in the form of supercomplexes. Besides providing kinetic advantage (channelling), supercomplexes control ROS generation by the respiratory chain, thus providing a means to regulate ROS levels in the cell. Depending upon their concentration, these ROS are either physiological signals essential for the life of the cell or toxic species that damage cell structure and functions.

CRITICAL ISSUES: We propose that under physiological conditions the dynamic nature of supercomplexes reversibly controls the generation of ROS as signals involved in mitochondrial-nuclear communication. During aging there is a progressive loss of control of ROS generation so that their production is irreversibly enhanced, inducing a vicious circle in which signalling is altered and structural damage takes place.

FUTURE DIRECTIONS: A better understanding on the forces affecting supercomplex association would allow the manipulation of ROS generation, directing these species to their physiological signalling role.

1. INTRODUCTION

The molecular mechanism by which aging induces a functional loss of the cells is a pleiotropic process involving several steps; mitochondria, however, undoubtedly play a central role in the mechanism of aging. Somatic mutations in mitochondrial DNA (mtDNA), originating from progressive ROS-induced damage, have been considered the major determinants of energy loss accompanying aging, as a consequence of alterations in the subunits of OXPHOS complexes encoded by that mtDNA. Since the generation of ROS in the cell is largely a task of the mitochondrial respiratory chain, the “mitochondrial theory of aging” (268) is a direct derivation of the “free radical theory of aging” (156).

The perspective of the role of ROS in the aging process has undergone an important change after the realization that ROS are physiological messengers acting through redox modifications in signalling proteins. For this reason it is believed that aging may be, at least in part, the result of alterations of signalling pathways such as those involved in mitochondrial biogenesis and apoptosis, induced by an increasing ROS production. However, the domination of the free radical theory has faltered in recent years, being challenged by a number of studies that seem to contradict it. For example, it was shown by a meta-analysis that antioxidant vitamin supplements such as vitamin A and E, singly or combined, significantly increase mortality in the general population (32, 33). In addition, other findings seem to demonstrate that ROS increase the life span of several organisms, rather than decreasing it (119, 332, 211). The significance of these findings has been, however, challenged (14, 239) and will be discussed in this review.

Indeed mitochondria might also affect longevity by ROS-independent mechanisms. MtDNA mutations and other damages may derive from errors or infidelity in metabolic mechanisms (transcription, translation etc.) (131). Of particular impact has been the observation of genetic manipulations that impair mitochondrial function and accelerate aging

but do not increase ROS (i.e. 384). Nevertheless, though important, this observation does not necessarily contradict the mitochondrial theory (14).

In this respect, understanding the mechanisms by which damaged molecules and structures are repaired or removed as a means to keep mitochondria functional over time may be of fundamental importance for the aging process.

The relative importance and weight that each of these factors exerts in the aging process is at the moment not fully clarified. For this reason it is important to understand the mechanism underlying ROS generation from mitochondria and other sources and their interplay. In particular, it is necessary to define (i) which are the precise sources of ROS involved, (ii) which factors control ROS generation and (iii) what are the immediate and long-term responses within the mitochondria in terms of redox chemistry.

In this review, we discuss evidence that the supramolecular structure of the mitochondrial respiratory chain is a major factor controlling the generation of ROS and is therefore largely involved in the aging process.

2. MITOCHONDRIAL ROS IN PHYSIOLOGY AND PATHOLOGY

Reactive oxygen species (ROS) are recognised today both as important factors physiologically involved in cell signalling, by affecting the oxidoreductive (redox) state of signalling proteins, and as among the major determinants of toxicity in cells and organisms.

ROS is a collective term including oxygen derivatives, either radical or non-radical, that are oxidizing agents and/or are easily converted into radicals. Diatomic oxygen O_2 is a radical because it has two unpaired electrons each located in a different π^* antibonding orbital, but both with the same spin quantum number: this parallel spin is the reason for its low reactivity with non-radical molecules. However, inverting the spin of one of the unpaired

electrons by an energy input converts O_2 into the much more reactive singlet oxygen 1O_2 ; in the excited state, the two electrons may either remain in two different orbitals or form a pair in the same π^* orbital (115). Spin restriction can also be overcome by adding electrons to oxygen one at a time (115).

Transition metals such as iron and copper, when in a free state, have a strong capacity to reduce O_2 , thus generating radical species. If a single electron is supplied to O_2 , it enters one of the π^* orbitals to form an electron pair there, thus leaving only one unpaired electron in the superoxide radical anion $O_2^{\cdot-}$. Addition of another electron to $O_2^{\cdot-}$ gives the peroxide ion, which is a weaker acid and is protonated to hydrogen peroxide H_2O_2 . Addition of two more electrons splits the molecule producing water H_2O . If one single electron is added to H_2O_2 by a reduced metal ion (e.g. Fe^{2+}), the hydroxyl radical $OH\cdot$ is produced by the *Fenton reaction*. The hydroxyl radical is extremely reactive with a half-life of less than 1 ns, thus it reacts close to its site of formation.

Consequently it must be borne in mind that ROS are widely different molecules, each having specific chemical properties that must necessarily reflect in their different biological effects. Thus, efforts should be made to clarify the individual effects of each ROS species in physiological and pathological processes.

The only technique that can detect free radicals directly is ESR (151). However, most ROS persist for very short times *in vivo* and cannot be measured directly. In this case, one solution is to add probes that act as “traps” and intercept ROS to form stable radicals detectable by ESR. A wide range of traps, mostly nitroxides, has been used in animals and cells in culture (151). Another approach to detect transient ROS is based on the use of *probes*, mainly fluorescence probes (151). Dichlorofluorescein diacetate (DCFDA) is widely used for this. It can enter cells where it is converted to the highly fluorescent 2',7'-dichlorofluorescein (DCF) upon cleavage of the acetate groups by intracellular esterases and oxidation by

endogenous ROS. Although DCF is usually considered a probe for peroxides, it reacts slowly with H_2O_2 , and also with superoxide, while reacting mostly with OH^\cdot and peroxynitrite. There are several limitations and artefacts associated with the DCF assay for intracellular H_2O_2 measurement and it is essential to keep these limitations in mind for proper interpretation of data (190).

Measuring the amount of oxidative damage caused by ROS or the antioxidant capacity of body fluids and cells is often taken as an indirect index of oxidative stress. It has to be borne in mind, however, that different biomarkers correlate differently with oxidative stress and that the studies employing fluorescence probes are amenable to artefacts and should be interpreted with caution (151).

2.1. Mitochondrial sources of ROS

ROS arise in cells from exogenous and endogenous sources. Exogenous sources of ROS include UV and visible light, ionizing radiation, drugs and environmental toxins. Among endogenous sources there are: xanthine oxidase, cytochrome P-450 enzymes in the endoplasmic reticulum, peroxisomal flavin oxidases and plasma membrane NADPH oxidases (150). Nevertheless, the mitochondrial respiratory chain in the inner mitochondrial membrane (IMM) is usually considered one of the major sources of ROS, although other enzyme systems in mitochondria can be important contributors to ROS generation (229). Among these, we mention here dihydrolipoamide dehydrogenase (a subunit of the α -ketoglutarate and pyruvate dehydrogenase complexes) (369, 382, 375, 8, 311, 192, 193), monoamine oxidase (46, 256, 85), and mitochondrial nitric oxide synthase (418). The last becomes a direct source of superoxide by an uncoupling mechanism in which it reacts with oxygen and generates O_2^\cdot instead of NO^\cdot (320), particularly in the absence of the cofactor tetrahydrobiopterin BH_4 .

It is worth noting that mitochondria from different tissues may vary conspicuously in their capacity to produce ROS using different substrates (207), and this capacity is also related to animal species and age.

Murphy (282) has carefully analysed the thermodynamics of mitochondrial superoxide production *in vivo*, concluding that the one-electron reduction of O_2 to $O_2^{\cdot -}$ ($E_m = -160$ mV) is thermodynamically favoured by the existing steady-state concentrations of O_2 and $O_2^{\cdot -}$ so that the actual reduction potential E_h may approach 200 mV and over. This high potential would allow O_2 to be theoretically reducible even by redox couples having high E_m . A wide range of electron donors within mitochondria could potentially carry out this reaction. However, only a small proportion of mitochondrial electron carriers is effective. The reason is that electron transfer to oxygen requires the protein-linked potential donors to be at a distance compatible with electron tunnelling, according to the Marcus theory (275).

2.1.1 The respiratory chain

According to Quinlan et al. (323), at least ten different sites of superoxide/ H_2O_2 production in the electron transport chain and associated enzymes (Krebs cycle, β -oxidation etc.) have been identified in mammalian mitochondria. The FMN and CoQ-binding sites of Complex I and the Q_o site (at the outer or positive side) of Complex III are often invoked as the most important mitochondrial superoxide producers, but other sites have also been defined, including glycerol phosphate dehydrogenase, ETF-CoQ reductase, and pyruvate and α -ketoglutarate dehydrogenases. The absolute and relative contribution of each site differs greatly with different substrates (Fig. 1).

INSERT FIGURE 1

The topology of the sites is important because it establishes whether the ROS are produced on the matrix side of the IMM, where they may damage mtDNA, or directly released out of the mitochondria. In the latter case, the ROS can damage other systems within the cell or they may constitute a signalling pathway from mitochondria to the cytoplasm and nucleus.

Most of superoxide is generated at the matrix side of the IMM, as appears from the observation that superoxide is detected in submitochondrial particles (SMP) which have an opposite orientation with respect to mitochondria. A study with suitable spin traps, however, demonstrated the formation of superoxide radical in mitoplasts, indicating that a significant aliquot of this species is released at the outer face of the IMM (370). It is likely that Complex I releases ROS in the matrix while Complex III mostly in the intermembrane space (IMS). The superoxide anion released at the IMS may be directly exported to the cytoplasm through an anion channel related to the Voltage-Dependent Anion Channel, VDAC (153).

Complex I is a major source of superoxide production in several types of mitochondria.

The identification of the oxygen reducing site has been the subject of extensive investigation, and several prosthetic groups in Complex I have been suggested to be the direct reductants of oxygen. These include FMN (118, 206, 100), ubisemiquinone (208, 209), and iron sulphur cluster N2 (125, 228, 104).

Grivennikova and Vinogradov (142) investigated ROS generation by Complex I either isolated or membrane-bound at different NAD^+/NADH levels. At the optimal NADH concentration of 50 μM , Complex I produced both superoxide and hydrogen peroxide at a 0.7 ratio $\text{O}_2^-/\text{H}_2\text{O}_2$. The production of superoxide was attributed presumably to FeS cluster N2, whereas hydrogen peroxide was interpreted as deriving from 2-electron oxidation of fully

reduced FMN. Rotenone, a specific inhibitor of Complex I, enhances ROS formation during forward electron transfer in the respiratory chain (164, 125).

The formation of superoxide in Complex III depends on the peculiar mechanism of electron transfer, the so-called Q-cycle (74, 75). This mechanism involves the biphasic oxidation of ubiquinol: one electron is given to the Rieske iron sulphur cluster and then to cytochromes c_1 and c , while the other electron reduces low potential cytochrome b (b_L). Since the subsequent electron transfer from cytochrome b_L to the high potential b_H occurs against the electrical gradient (from the positive to the negative side), this reaction is strongly retarded when the electrochemical potential is high, as in the controlled state (*State 4*). This retardation, prolonging the lifetime of the semiquinone ($Q\cdot$) at the outer Q_o site, has been interpreted as allowing reaction of $Q\cdot$ with O_2 , thus forming superoxide (276, 184).

Antimycin A (AA), an inhibitor acting at the inner or negative side of the membrane (Q_i site), is known not to completely inhibit electron flow from ubiquinol to cytochrome c . The AA-insensitive reduction of cytochrome c is mediated by superoxide radicals. According to the Q-cycle, AA blocks ubiquinone reduction by cytochrome b_H at Q_i . The antimycin-stimulated production of ROS is inhibited by the inhibitors acting at Q_o site, where ubiquinol reduces both the Rieske iron-sulphur cluster and cytochrome b_L . Thus, we may locate the site of one-electron reduction of oxygen in presence of AA at a component located at Q_o , presumably ubisemiquinone (56).

It has been assumed that superoxide is formed during the physiological operation of the Q-cycle (in which the semiquinone is formed when ubiquinol is oxidized by the Rieske FeS cluster). More recent evidence, however, suggests that the source of the electron to reduce oxygen is the semiquinone formed in the so-called semi-reverse reaction in which cytochrome b_L reduces the fully oxidized quinone. In fact, superoxide formation is stimulated by the presence of oxidized quinone (95) and by mutations preventing the proximity of the

FeS cluster of the Rieske protein to the ubiquinone site (344). This latter condition does not allow interaction of ubisemiquinone with the Rieske cluster and thus favours its reaction with oxygen.

Several observations suggest that also Complex II may be a significant source of ROS (176, 177, 178, 420, 217, 323). Indeed, the highest rate of ROS production in isolated mitochondria occurs with succinate as substrate (327), but this phenomenon is sensitive to rotenone (208, 288, 396) therefore ROS generation is commonly attributed to the contribution of Complex I by the energy-dependent reverse electron transfer from succinate to NAD^+ via the CoQ pool (202, 184, 205, 368). Nevertheless, this view has been challenged by a series of experiments that seems to exclude a significant ROS production catalysed by reverse electron flow through Complex I, thus allowing ROS production by Complex II as the only plausible explanation (273).

It is also worth noting that mitochondria in intact cells oxidize predominantly NAD-linked substrates so that reverse electron transfer cannot be a very important contributor to superoxide formation (273).

The simple passive diffusion of hydrophilic ROS across lipid membranes is limited due to permeability restrictions, but evidence exists that the superoxide produced in the mitochondrial matrix may also be exported in the cytoplasm. A new function of uncoupling proteins UCP2 and UCP3 was suggested (409) by observing that guanine nucleotides decrease the release of superoxide anion, but not of hydrogen peroxide, from respiring heart and muscle mitochondria. Since guanine nucleotides are known inhibitors of UCPs, the authors inferred that these proteins act as anion channels to export dangerous superoxide from the mitochondrial matrix. Accordingly, guanine nucleotides enhanced the damaging effect of ROS on matrix aconitase.

In addition, it has been proved that selected aquaporin homologues (e.g. AQP8) have the capacity to channel H_2O_2 across the IMM (30), thus somewhat challenging the importance of ROS compartmentation with respect to the IMM. These observations, however, clearly point out that ROS export is largely a protective device, potentially allowing ROS detoxification elsewhere in the cell without consuming mitochondrial reduction equivalents (29).

An extensive review of other sites in the respiratory chain which are responsible for ROS generation (e.g. glycerol phosphate dehydrogenase, dihydroorotate dehydrogenase, electron transfer flavoprotein (ETF) and ETF dehydrogenase) can be found in (222).

2.1.2 Redox cycling

Mitochondrial generation of ROS can be implemented by exogenous compounds that interact with the respiratory chain by virtue of their redox properties. Among these are organic compounds and metal ions, in particular heavy metal ions. The toxicity of many of these compounds is best explained by their interaction with mitochondrial respiration and induction of oxidative stress (221).

Being so significantly present in living organisms, Fe is the metal most commonly involved in oxidative stress (103). Iron can exist in different oxidation states, varying from -2 to $+6$. However, within biological systems, Fe is bound to specific metalloproteins and is found in the $+2$ or $+3$ oxidation states. Such change in its redox state is crucial to a large number of reactions in oxidative metabolism and determines Fe participation in potentially cytotoxic reactions. In fact, Fe^{2+} can catalyse the Fenton reaction by which H_2O_2 is reduced to the aggressive OH^\cdot radical. Fe^{3+} can also be reduced to Fe^{2+} after reacting with superoxide anion by the Haber-Weiss reaction; this reaction, however, may not have physiological

significance, due to the presence in the cells of reducers of Fe^{3+} which are more powerful than superoxide (240).

Many physiologically active substances and xenobiotics have electron transfer functionalities, either *per se*, or more usually in their metabolites. These main groups include quinones (or phenolic precursors), metal complexes (or complexors), aromatic nitro compounds (or reduced derivatives), and conjugated imines or iminium species (203). Many of these compounds generate ROS exploiting a mechanism of *redox cycling*.

In vivo redox cycling with oxygen can occur in a catalytic fashion giving rise to a futile cycle that regenerates the parent compound and releases ROS. Redox cycling has been suggested for several drugs, like cocaine, other abused drugs, catecholamines, and several other compounds, besides a large number of organic components of fine-particulate air pollution that initiate a series of cellular reactions and ultimately lead to cell injury (129).

Electron transfer with redox cycling occurs through interference with physiological electron transfer reactions such as microsomal cytochrome P450, xanthine oxidase, and the mitochondrial respiratory chain. For example, adrenaline may undergo oxidation and cyclisation to adrenochrome in a multi-step process in which the main oxidant under physiological conditions is the superoxide anion. On the other hand adrenochrome can be reduced to the corresponding semiquinone by mitochondrial Complex I, thus establishing a redox cycle in which the semiquinone reacts with O_2 producing superoxide and regenerating adrenochrome (123).

2.1.3 The adaptor protein $p66^{\text{Shc}}$.

Special mention must be made of the *adaptor protein* $p66^{\text{Shc}}$, a 66-kDa Src collagen homologue (Shc) protein that is one of three main isoforms encoded by the SHC1 gene ($p46^{\text{Shc}}$, $p52^{\text{Shc}}$, $p66^{\text{Shc}}$) (266). While $p46^{\text{Shc}}$ and $p52^{\text{Shc}}$ isoforms link activated receptor

tyrosine kinases to the Ras pathway, by recruitment of the GRB2/SOS complex, $p66^{\text{Shc}}$ mediates an inhibitory signalling effect on the extracellular signal-regulated kinase (ERK) pathway that is required for actin cytoskeleton polymerization and normal glucose transport control. $p66^{\text{Shc}}$ has been also identified as a sensor of oxidative stress-induced apoptosis. This action requires Ser36 phosphorylation of the protein. Pathways that regulate the cellular response to oxidative stress and life-span by leading to increase of $p66^{\text{Shc}}$ expression involve p53 (385, 299) and serine-threonine kinases, including the PKC- β isoform (319).

Expression of $p66^{\text{Shc}}$ is required for mitochondrial depolarization and release of cytochrome *c* after a variety of pro-apoptotic signals (385). $P66^{\text{Shc}/-}$ cells are resistant to apoptosis (265) and $p66^{\text{Shc}}$ deletion in mice decreases the incidence of aging-associated diseases (285, 261, 336) and prolongs lifespan of animals (265), due to their resistance to oxidative stress.

A fraction of $p66^{\text{Shc}}$ has a mitochondrial localization in the IMS, where it is bound as an inactive form in a high molecular weight complex, including the TIM/TOM protein import system (130). Under stress conditions, however, antioxidant systems are not able to retain $p66^{\text{Shc}}$ in their reduced dimeric state. Pro-apoptotic signals dissociate $p66^{\text{Shc}}$ from the complex and activate it to a tetrameric form (126) that triggers the permeability transition by opening a high conductance channel in the IMM, *the permeability transition pore* (26), which is involved in the events leading to apoptosis. This effect is due to the intrinsic property of $p66^{\text{Shc}}$ to act as a redox protein accepting electrons from cytochrome *c* and directly producing hydrogen peroxide (130). A cytochrome *c*-binding domain near the N-terminus is responsible for this activity (127).

Thus the $p66^{\text{Shc}}$ protein may be not only a redox enzyme but also a ROS sensor and possibly a ROS amplifier in the mitochondrial IMS through an inherent ROS-producing activity.

Since the reaction equilibrium of cytochrome *c* oxidation by $p66^{Shc}$ is low ($K_{eq} = 0.1$), the reaction is thermodynamically favoured when the level of cytochrome *c* reduction is high (130). This means that H_2O_2 production by this mechanism should be enhanced when cytochrome *c* oxidase (the enzyme catalysing cytochrome *c* reoxidation by oxygen) is inhibited. The K_m of cytochrome oxidase for oxygen is very low ($< 1\mu M$), thus allowing its activity even at low oxygen tensions. However, low oxygen tensions promote the activation of hypoxia-inducible factor (HIF-1 α) by a still controversial mechanism (378); HIF-1 α triggers a series of metabolic changes among which is alteration of cytochrome oxidase subunits and activity. Thus the activation of the $p66^{Shc}$ pathway may in part explain the paradoxical enhancement of ROS production during hypoxia (145, 28, 55).

2.2. Upstream control of ROS production

The steady-state level of ROS is determined by the rate and site of their generation, by their nature and lifetime, by their diffusion constants, by the interconversions occurring between different ROS, and finally by the rate of removal by different antioxidant systems.

The generation of ROS by isolated mitochondria accounts for 0.1-0.2% of oxygen consumed and may reach up to 2-3% under particular conditions that may not be found physiologically. The rate and extent of ROS generation greatly varies in different tissues and specifically depends upon the substrate employed: for example, succinate is important for ROS production in brain, heart, kidney, and skeletal muscle, while fatty acids are major generators in kidney and liver (301, 375).

There is increasing evidence that mitochondrial ROS level is physiologically regulated; this is particularly important for their action as cellular signalling molecules (cf. section 2.3.1).

2.2.1 Role of Mitochondrial Membrane Potential

Mitochondrial ROS production is enhanced in State 4 and when the rate of electron transfer is lowered (364). The rationale is in a more reduced state of the respiratory carriers capable of donating electrons to oxygen. Hence, uncoupling and release of excessive membrane proton potential may protect mitochondria from damage induced by excessive free radical production. In rat hepatocytes the futile cycle of proton pumping and proton leak may be responsible for 20-25% of respiration (37); in perfused rat muscle the value is even greater. Uncoupling may be achieved by activating proton leak through *uncoupling proteins* (56) (see above, however, for a different role postulated for uncoupling proteins, cf. 409). In such a way, whilst a tissue may dissipate a conspicuous part of the energy conserved by its mitochondria, it also keeps the mitochondrial respiratory chain under more oxidized conditions preventing the formation of free radicals.

The notion that mild uncoupling may be a protective mechanism that lowers mitochondrial $\Delta\Psi$ and thus alleviates oxidative stress has been challenged on the theoretical basis that the mitochondrial membrane potential within living cells is much lower (100-130 mV) than that measured in isolated mitochondria. Accordingly, no significant change in matrix superoxide occurred after treatment of cerebellar neurons with the uncoupler FCCP (185).

It is worth mentioning that matrix pH *per se* is also an essential factor defining ROS production by the respiratory chain, even in the absence of pH gradient, and that pH increase in the matrix induces the increase in ROS generation (356). This observation is in agreement with a previous demonstration that the rate of superoxide generation increased in conjunction with an increase of medium pH from 7 to 9.2, in the presence of NADH or succinate as substrates and of rotenone or antimycin A as specific enzyme inhibitors (388). Moreover, the decrease of matrix pH induced by the addition of P_i and nigericin is accompanied by an

increase of $\Delta\Psi$, which is expected to stimulate ROS production (356). However, the ROS generation rate decreased, thus indicating that in these circumstances the effect of pH dominates over the opposite effect of $\Delta\Psi$.

2.2.2. Role of Post-Translational Modifications on ROS Production

Events leading to decrease of the rate of electron flow in Complex I lead to overproduction of ROS. Physiological states, such as subunit phosphorylation that inhibits Complex I activity, may modify its ROS generating capacity (326, 248, 346). It is therefore tempting to speculate that endocrine alterations may affect the capacity of ROS formation by hyper- or hypo-phosphorylation of the Complex.

Kadenbach et al. (189) have proposed a different mechanism, centred on cytochrome *c* oxidase. The cAMP-dependent phosphorylation of subunit I of the oxidase greatly increases its sensitivity to allosteric inhibition by ATP. The result of this inhibition is decrease of the H^+/e^- stoichiometry of the enzyme from 1 to 0.5, with resulting decrease of $\Delta\Psi_m$ and, consequently, of ROS generation by the respiratory chain. It was also proposed that stress conditions would induce dephosphorylation of Complex IV with transient increase of membrane potential and a burst of ROS generation by mitochondria. The discovery of a mitochondrial CO_2 -adenylyl cyclase-cAMP-PKA signalosome, conserved from the yeast *S. cerevisiae* to human cells (165) and modulating the allosteric inhibition of Complex IV by ATP, underlies a universal mechanism for metabolic regulation in eukaryotes exposed to environmental fluctuations in temperature, oxygen, and nutrient availability. However, the hormonal signals inducing phosphorylation and dephosphorylation remain to be fully established.

Other protein modifications in the respiratory complexes can modulate ROS production (154), such as thiol oxidation or S-nitrosation or S-glutathiolation of Complex I, or Complex I and II acetylation that is regulated by sirtuins (NAD⁺-dependent deacetylases).

In addition, ROS production by mitochondria is under signalling control through the pathways leading to mitochondrial uptake and activation of *p66^{Shc}* under conditions of oxidative stress (cf. section 2.1.3).

2.2.3 Hypoxia and ROS production

The presence of a pseudo-hypoxic state in aged tissues (cf. Section 3.2) requires a brief discussion of ROS in hypoxia.

A major point of controversy regarding the role of ROS as hypoxic signalling molecules is that it is seemingly paradoxical that a decrease in a required substrate, O₂, would result in an increase in ROS production. However, using new ratiometric fluorescent probes, it proved possible to demonstrate an increased ROS production during hypoxia (23). Furthermore, both DNA and lipid oxidation products accumulate during hypoxia (141).

The first breakthrough suggesting a mitochondrial oxygen sensor of the HIF pathway came with the discovery that ρ⁰ Hep3B cells, which contain no mtDNA and thus no electron transport, are incapable of HIF-1 DNA-binding activity following hypoxia (58). This finding implies that mitochondria are responsible for propagation of the hypoxic signal. Correspondingly, treatment of cells with exogenous H₂O₂ or induction of cellular H₂O₂ production are sufficient to stabilize HIF-1α during normoxia (59). Nevertheless, the activating effect of ROS on HIF-1α (22, 152, 308) is still a controversial issue (cf. section 2.3).

The content of oxidized cytochrome *c* in the IMS seems to be an essential factor in controlling mitochondria-derived ROS (47, 307); in fact, cytochrome *c* in the IMS oxidizes

superoxide produced by Complex III to O_2 , thus preventing generation of H_2O_2 . The loss of cytochrome *c* due to outer membrane permeabilization during hypoxia and ischemia (62) was considered a major factor of ROS generation under pathological conditions (307).

Inhibition of cytochrome *c* oxidase by hypoxia (57) enhances ROS formation by the respiratory chain; this result may be due to the increased membrane potential (423) or by inducing a more reduced state of cytochrome *c*. The reduction of cytochrome *c* facilitates its interaction with $p66^{Shc}$ and further amplification of ROS production (cf. section 2.1.3) while preventing its superoxide scavenging activity. Evidence also exists that ROS generation at mitochondrial Complex III is critical for hypoxia signalling (59, 144, 201, 250) (Fig. 2). More recently, the role of Complex III was confirmed in a transgenic mouse model where deletion of the Rieske protein of Complex III (402) abolishes the hypoxia-induced increase in ROS signalling.

INSERT FIGURE 2

In addition, it has to be pointed out that the enhanced ROS production during hypoxia has not been ascribed to mitochondria only, but also to the plasma membrane NADPH oxidase (cf. 271).

2.3. ROS as signals

It is clear nowadays that, at moderate concentrations, ROS act as second messengers (182, 11, 331, 138, 154, 400, 283, 69) by interfering with the expression of a number of signal transduction pathways and genes. The effects related to ROS signals are double-edged. For example, ROS may be oncogenic and promote proliferation, invasiveness, angiogenesis and metastasis (411), but they may also be anti-oncogenic and promote cell

cycle stasis, senescence and apoptosis (149). Many of the ROS-mediated responses actually protect the cell against oxidative stress and re-establish redox homeostasis. However, there is a threshold above which ROS become harmful and induce oxidative stress (331). The concentration range of ROS can vary conspicuously; moreover it must be kept in mind that the composite term ROS includes several species that have widely different properties. Thus, not all ROS are equally suitable for signal transduction; for instance, the OH· radical is too unspecific to participate in catalysed reactions, while O₂⁻, H₂O₂ and lipid peroxidation derivatives like 4-hydroxy-2-nonenal (HNE) are likely to be employed as signalling molecules.

Intracellular signalling achieved by ROS molecular recognition occurs at the atomic level.

Because some ROS such as H₂O₂ are oxidants, they influence the redox state of signalling proteins through reactions with specific sulfhydryl groups (81) but enzymatic catalysis is usually required to allow the modification to occur under physiological conditions.

A number of transcription factors contain redox-sensitive cysteine residues at their DNA-binding sites (147, 381). Thiols exposed on the surface of mitochondrial proteins (283) are very reactive since the relatively high matrix pH favours the dissociation to thiolate anions; moreover, the high proportion of vicinal thiols facilitates the formation of disulphide bonds acting as redox switches to transduce the response of the protein to redox signalling (282, 283). Such clustering of thiols groups is not casual and is favoured by evolution (255).

The ROS-mediated oxidative modification of proteins is usually followed by other post-translational changes (e.g. phosphorylation, acetylation, ubiquitination, and SUMOylation among others (400)) in the same protein and in other proteins of the signalling cascade.

One notable example of mitochondrial signalling leading to changes in nuclear gene expression (406) is Nrf2 that, in the presence of ROS, is translocated from the cytoplasm to the nucleus. There, Nrf2 binds the antioxidant response element of cytoprotective genes involved in the antioxidant response (e.g. heme oxygenase, NRF-1 and other inducers of mitochondrial biogenesis) (180). Nrf2 dysfunction in ageing likely exacerbates age-related cellular oxidative stress and increases sensitivity to oxidative stress-induced cellular damage.

Signalling proteins modified by ROS include phosphoprotein phosphatases (PTPs), Ras, large G-proteins, serine/threonine kinases of the MAPK families, transcription factors as AP-1, NF κ B, p53 and others. The effect is different with different proteins: PTPs are inhibited, nuclear transcription factors are activated (390). Thus, for example, activation of ERK1/2, Akt and NF κ B promotes cell survival; whereas activation of c-jun N-terminal kinases, p38 kinase and p53 would lead to arrest of the cell cycle and apoptosis.

In addition, ROS appear to activate the hypoxia-inducible factor HIF-1 α (59, 196) by inhibiting the prolyl-4-hydroxylase that addresses the factor to proteolytic digestion (Fig. 2). HIF-1 α becomes stabilized and binds to hypoxia responsive elements in the DNA, stimulating a large array of genes (357, 191) and decreasing the input of reducing equivalents to the respiratory chain (199). These events mimic what happens in anoxia through lack of O₂ required for HIF hydroxylation.

Mitochondrial ROS production is the determinant of preconditioning, by which small doses of a noxious stimulus are required to induce a protective response against future injury. Stimulation of the hypoxia-signalling pathway by hypoxia or by cyanide treatment (71) is a major factor responsible for preconditioning in brain (361, 70), heart (284, 358) and kidney (61).

Mitochondrial ROS can also act as important signals to regulate the inflammatory response by activating the inflammasome (107, 310) and to regulate autophagic processes including mitophagy (350).

2.3.1 Regulation of Mitochondrial ROS in Cell Signalling

Mitochondrial ROS levels reflect the balance between their rate of generation and of removal.

The ROS generation by rat liver mitochondria was investigated under different substrate and inhibitor conditions and different oxygen tensions, with the conclusion that only Complex I may be a significant source of ROS at physiological O₂ concentration (167). In this scenario, the factors directly associated with respiratory activity (e.g. the redox potential of the NAD⁺/NADH couple and the proton-motive force) are powerful regulators of electron leaking (331). These factors in turn are regulated by the redox supply to the chain, by the degree of coupling and by physio-pathological constraints to electron transfer (enzyme phosphorylation, cytochrome *c* removal, Complex IV inhibition, oxygen concentration etc.).

In addition, mitochondria enhance ROS generation in response to external stimuli, such as TNFα (198), hypoxia (59), serum deprivation (215) and oxidative stress itself (ROS-induced ROS release, 422). Several proteins, such as p53, p66^{Shc}, the Bcl-2 family and Romo-1 (11) can control ROS generation and release from mitochondria.

In the last years, interplay between mitochondria and other ROS generating systems has been discovered. This interplay is particularly puzzling between mitochondria and the NADPH oxidases (NOX) family (89). There is accumulating evidence showing that stress of the endoplasmic reticulum mediates the cross-talk between mitochondrial- and NOX-derived ROS by activating the Unfolded Protein Response (UPR) signalling pathway and regulating prosurvival and proapoptotic components (339, 340).

ROS removal by enzymatic systems may be directly involved in regulation of ROS signalling (111). Various isoforms of superoxide dismutase, including mitochondrial Mn-SOD, catalyse the dismutation of superoxide to H_2O_2 and oxygen; whereas catalase, glutathione peroxidase and peroxiredoxins are capable of removing H_2O_2 . In addition ferric cytochrome *c*, by virtue of its capacity to be reduced by superoxide, participates in antioxidant defence in the respiratory chain (307), while an integrated set of thiol systems within the mitochondrial matrix prevents much of the oxidative damage (283) and may transduce redox signals within and through mitochondria. Peroxiredoxins (Prx) act on small dithiol proteins, of which thioredoxin-2 is localized in the mitochondrial matrix (329, 235). Prx3 is the target of most H_2O_2 produced in the mitochondrial matrix (72) whereas Prx5 is less effective in reducing H_2O_2 but has a higher reactivity towards peroxynitrite (387).

Mitochondrial glutathione (mGSH) is a dominant factor in the maintenance of the appropriate mitochondrial redox environment, mainly because of its abundance and versatility to counteract hydrogen peroxide, lipid hydroperoxides, or xenobiotics, by acting as a cofactor of enzymes such as glutathione peroxidase or glutathione-S-transferase (254). The mitochondrial GSH:GSSG ratio is generally greater than 100:1 and is widely used as an indicator of the redox status (186).

Protein glutathionylation may occur through thiol disulphide exchange by protein disulphide isomerase, as a regulatory device for proteins involved in energy metabolism, redox signalling, and apoptosis (415, 77). Glutaredoxin is the primary enzyme responsible for catalysis of de-glutathionylation of protein-mixed disulphides with glutathione. This reversible post-translational modification alters the activity and function of many proteins important in regulation of critical cellular processes (5).

2.4. ROS and mitochondrial quality control

Damaged molecules are recognized by degrading systems; in particular, oxidatively modified proteins are extensively ubiquitinated and directed to the proteasome, a member of the ATP-dependent AAA⁺ proteases, where their complete digestion takes place (345, 380). Inability of the proteasome to digest some proteins and the accumulation of insoluble protein aggregates may lead to profound alterations in cells.

Since mitochondria are a major source of reactive oxygen species, mitochondrial proteins are especially exposed to oxidative modification, so that their elimination is crucial for maintaining the integrity of this organelle (246, 379). Internal mitochondrial proteins may be retrotranslocated to the outer mitochondrial membrane where multiple E3 ubiquitin ligases are present. Cdc48/p97 is recruited to stressed mitochondria, extracts ubiquitinated proteins from the outer membrane and presents them to the proteasome for degradation.

Within the mitochondrial matrix, the ATP-stimulated Lon protease (Pim-1 in yeast) is devoted to the regulatory mechanism of selective degradation of oxidized proteins. For example, during hypoxia, Lon participates in remodelling cytochrome oxidase by selectively degrading the Cox4-1 subunit (116, 392). Failure of protein degradation has been implicated in the age-related accumulation of oxidized proteins (42, 389).

Cellular structures and organelles undergo turnover and after a suitable life time are directed to autophagy by lysosomal digestion (232, 25, 63).

Non-selective autophagy and mitophagy are triggered by ROS in response to nerve growth factor deprivation, rapamycin, or starvation (214, 350). In mouse and human cells, mitophagy is triggered in a mitochondrial fission dependent manner by mild oxidative stress comparable to that occurring during physical activity (113). These conditions increase ROS levels only slightly and are insufficient to trigger non-selective autophagy, thus suggesting the presence of a very specific and selective signalling cascade initiated by ROS. On the other

hand, when non-selective autophagy is induced by starvation or other means, mitochondria hyperfuse and are protected from mitophagy (134, 328).

3. A ROLE OF MITOCHONDRIA AND ROS IN AGING

Most research on the role of mitochondria in aging has been formerly performed using different tissues from aged animals (cf. 220) including domestic mammals and primates (108, 162, 296). The necessity of employing simpler systems that allow the investigation of a number of genetic mutations and/or to exert metabolic manipulations during a short life span (120, 200) has focussed attention more and more on invertebrates. *Drosophila* (306, 236, 238, 289), the housefly *Musca domestica* (67), the nematode *Caenorhabditis elegans* (407, 45, 78), the yeast *S. cerevisiae* (183, 39, 297, 257) and the filamentous fungus *Podospora anserina* (245, 290) have all been usefully employed in research on ageing.

Planarian flatworms (*Tricladia*, *Platyhelminthes*) share a very rare characteristic with human mitochondria, a strong control of oxidative phosphorylation by the phosphorylation system, and for this reason they have been proposed as useful models of mitochondrial disturbances (218). Other species, such as the bivalve *Arctica islandica*, have been proposed as aging models in consideration of their exceptional longevity (281).

Also, naturally occurring mouse mutants, such as the senescence-accelerated mouse (170, 179) and transgenic mice (270), have provided important clues. The “mutator mouse”, having a defect of the proof-reading function of mitochondrial DNA polymerase γ (Poly), is a remarkable model of premature aging due to mtDNA mutations (383, 384, 97, 166, 38).

Finally, although the use of aging *in vitro* of cell cultures as a model for senescence is controversial (160, 99, 83), primary cell cultures from senescent animals represent a useful tool for studies of mitochondria in relation to aging (148, 397, 160). An unprecedented avenue is the use of human induced pluripotent stem cell-based models of aging (241). These

cells can be transplanted into model animals and advance our knowledge of molecular mechanisms of aging and help to develop strategies for treating aging-associated human diseases.

It is also expected that the development of novel and powerful technical approaches will enable investigators to get deeper insights into structure and function of mitochondria and their interactions within the cell, as well as their alterations in disease and aging. The proteomic approach (294, 401), including redox proteomics (94), will become an indispensable tool in unravelling the unknown functions of these organelles. MitoCarta (293), an extensive study by mass spectrometry, GFP tagging, and machine learning has been used to create a mitochondrial compendium of 1098 genes and their protein expression across 14 mouse tissues.

3.1. A CRITICAL ANALYSIS OF THE MITOCHONDRIAL THEORY OF AGING

The mitochondrial theory of aging is based on the following basic assumptions that are linked by a causal relationship (220):

1. Mitochondrial ROS produced during the life time are the major cause of aging
2. Mitochondrial ROS induce structural damage to mitochondrial biomolecules, in particular mtDNA somatic mutations, mostly in post-mitotic tissues.
3. The mtDNA mutations induce damage to the mtDNA-encoded proteins, and consequent decrease of oxidative phosphorylation, culminating in energy failure, metabolic derangement and cell death.

A corollary to the mitochondrial theory is that the decreased electron transfer may cause further ROS generation, establishing a vicious circle of mtDNA damage and ROS production (292).

Although experimental evidence exists, pertaining to each of these points, there are still several controversial aspects and the causal relationships among them are often lacking. The major points to be clarified are the following:

- a) What is the precise role of ROS in aging? In other words, can we determine whether, and how much, are ROS involved in the aging process?
- b) What can we learn from mtDNA genetics in relation to aging and longevity?
- c) What is the relevance for aging of mtDNA deletions and mutations?
- d) Can mtDNA damage be causally correlated with mitochondrial functional defects?
- e) Are mitochondria self-sufficient for the increase and propagation of the damage or is the expression of nuclear genes also necessary for the aging phenotype? In other words, do mitochondria elaborate signals (ROS or others) able to induce the aging process? And how are these signals regulated during the lifespan of an individual?
- f) Finally, can we predict whether, and how much, we can control the aging process by interfering with one or more of the above items?

3.1.1. ROS generation and ROS-induced damage

The continuous generation of ROS and other toxic species induces damage to all the biomolecules; thus it is not necessary that an increase of ROS generation occurs in aging, since it is the damage that would accumulate even at a steady ROS generation with time (14). Nevertheless, a vicious circle (292, 91) can be established if the accumulated damage to the respiratory chain would enhance ROS generation. This event is theoretically expected when electron transfer within a respiratory complex competent in ROS generation is inhibited (164, 125, 35). Exposure of mitochondria *in vitro* to different concentrations of hydrogen peroxide or cumene-hydroperoxide induces a transient increase in ROS production, followed by a steady state without further increase (342). In this case, ROS originate mainly from Complex

It is a consequence of the inhibition of the enzyme, supporting the vicious circle hypothesis. It is clear, however, that these *in vitro* conditions are not fully comparable with the situation *in vivo*.

Many reports demonstrated that the rate of production of ROS from mitochondria increases with age in mammalian tissues (366, 330, 54, 188, 17, 21) as well as in cultured cells from old individuals (174, 216, 373) and during replicative cell senescence (160, 110). An additional factor possibly eliciting an increased ROS production in aging is the postulated spreading of damage from one cell to the surroundings by means of the plasma membrane oxidoreductase (274, 82), thus triggering a chain reaction of oxidative damage.

A few reports have indicated a lack of increase of the rate of ROS generation with aging (155, 244, 343). It has to be noted that such results may be a consequence of the decrease of content and activity of respiratory complexes, so that there is a decreased total number of respiratory units producing ROS, even if each unit has an enhanced ROS generation. For this reason the rate of ROS generation should be related to the content of the enzymes that are responsible for it.

A different aspect that unambiguously relates aging to ROS generation is the strong negative correlation of animal longevity with the rate of mitochondrial ROS generation and with the degree of fatty acid unsaturation of cellular membranes (reviewed by 14). According to Barja (14) only the mitochondrial free-radical theory of aging can explain these correlations and, in particular, the latter correlation is explained by the higher oxidative damage (peroxidizability) that may be induced in unsaturated membrane fatty acids. In tissues of aged animals where there is an increase of oxidized lipids (416, 410, 377, 7), mitochondria exhibit peroxidation of their signature lipid, cardiolipin (CL), with consequent CL depletion and loss of respiratory activity (305, 304, 303, 359, 315).

Complex I is considered to be a major source of ROS involved in aging (cf. section 2.1), since ROS generated by Complex I are released in the matrix and may damage mitochondrial proteins and mtDNA. However, ROS produced by Complex III must also be taken in serious consideration, since they are released in the intramembrane space and may be exported in the cytosol, where they can activate, or interfere with, proteins involved in signal transduction (cf. section 2.3.1).

In high eukaryotes, damaged proteins, protein aggregates and damaged organelles tend to accumulate in aging, in the form of lipofuscin deposits in post-mitotic tissues such as muscle, heart, liver and brain (374, 353, 60, 338); these aggregates are removed by autophagy (272) (cf. section 2.4).

The mitochondrial theory of aging considers mtDNA as the major target of mitochondrial ROS because it is close to the site of mitochondrial ROS production and is less protected than nuclear DNA. Indeed, the damage by an oxidative stress is higher and persists longer in mtDNA than in nuclear DNA (412, 249) and the level of 8-oxo-7,8-dihydro-2'-deoxyguanosine (8-oxodG) in mtDNA (but not in nuclear DNA) negatively correlates with the longevity of several animal species, including humans (16, 159, 163). A recent review (194) summarizes the evidence on mtDNA copy number, deletions and point mutations during aging in humans and mice.

Moreover, the oxidation of mitochondrial proteins and lipids also increases with age (366, 4). Genotoxic intermediates of lipid peroxidation may also have a role in causing age-associated DNA mutations (172). This idea is supported by *in vitro* experiments which show that mtDNA is damaged when mitochondria undergo lipid peroxidation.

In mutant mice deficient in SOD2, mitochondrial oxidative stress specifically promotes glycation of mtDNA and does not affect nuclear DNA or cytosolic proteins (40).

Because DNA glycation can change DNA integrity and gene functions, glycation of mtDNA may play an important role in the decline of mitochondrial functions.

A generally negative correlation has been consistently found between endogenous tissue antioxidants and longevity in animals (313, 295). The general result of these comparative studies agrees with the observation that antioxidants do not increase maximum longevity and lack the capacity to slow down aging, independently of the way in which the antioxidants are manipulated: dietary, or genetically. This disappointing lack of effect has contributed to the criticism of the basic postulates of the mitochondrial theory of aging (cf. 239, 312, 41, 341).

Contrary to this view, Barja (14) correctly reasons that these results are expected, since it is the rate of production of ROS and not their elimination that determines the life expectancy. Moreover, ROS concentration in particular compartments such as mitochondria is much more dependent on the rate of ROS generation than on antioxidants, as the free radical generation source, especially Complex I, is approached at the micro level (15). This is especially important because the main target for aging, mtDNA, is located very close to, or perhaps even in contact with the free radical generation source. This can help explain why lowering the rate of ROS generation instead of increasing levels of antioxidants occurred during the evolution of long-lived species.

An additional factor explaining why chemical antioxidants are inactive against aging is that upon scavenging ROS they may be converted into prooxidant byproducts as semiquinones. Accordingly, we suggest that the fact that the novel potent artificial hydroxylamine scavenger (IAC) can prolong the lifespan of the freshwater annelid *Aeolosoma viride* by 170% arises because, upon scavenging ROS, the molecule is converted into an even more potent nitroxide antioxidant (53).

According to Barja (14), the negative correlation between levels of antioxidants and life span of different species only shows that the level of antioxidant enzymes and of repair enzymes is adjusted to compensate for the proportion of ROS generation; thus, if a species produces lower ROS, it is expected that it has less need for antioxidants and DNA repair enzymes.

In a basic study, Trifunovic et al. (384) showed that expression of a proof-reading deficient Polγ in a homozygous knock-in mouse strain leads to increased levels of somatic mtDNA mutations causing progressive respiratory chain deficiency; the mice develop symptoms strikingly reminiscent of aging. This is the most striking demonstration that mtDNA mutations can cause aging. The rate at which mtDNA mutations reach phenotypic expression differs markedly among tissues, which may be an important factor in determining the tolerance of a tissue to random mitochondrial mutagenesis (394, 395).

In apparent contrast with the mitochondrial theory, the mutator mice were not shown to have an enhanced ROS production (383). It is likely, however, that the lack of enhancement of ROS production results from the exceedingly severe extent of mutation in this mouse strain. Nevertheless we cannot exclude the possibility that a ROS attack is the natural way to induce mutations in other organisms. Indeed, a recent study (242) showed that the level of hydrogen peroxide was the same in the young mutator and control mice, but that hydrogen peroxide increased as the mutator mouse aged. Thus mtDNA mutations increase ROS generation contributing, also in the mutator mice, to the accelerated aging phenotype.

In agreement with the ROS origin of mtDNA defects, hydrogen peroxide induces large scale deletions of mtDNA through formation of double-strand DNA breaks (367). Moreover, some lines of evidence suggest that the frequency of mtDNA deletions is significantly decreased in mice that express the human catalase gene targeted at mitochondria (354). These results demonstrate the importance of mitochondria as a source, and possible

target, of ROS and are consistent with a role of H_2O_2 in the limitation of mouse lifespan. Nevertheless, the effect of increased mitochondrial peroxide detoxification capacity in model animals, where the endogenous H_2O_2 removing systems are likely to be already abundant, is not clearly explained at a molecular level and cannot be taken as a proof that preventing peroxide-mediated mitochondrial damage alone is sufficient to delay the aging phenotype.

Contrary to most evidence, a recent study (181) proposes that oxidative stress is not a major cause of somatic mtDNA mutations. The authors found that many of the features associated with mtDNA mutations in vertebrates are conserved in *Drosophila*, including a comparable somatic mtDNA mutation frequency, an increased frequency of mtDNA mutations with age, and a prevalence of transition mutations. Only a small fraction of the mtDNA mutations detected in young or old animals were G-C to T-A transversions, a signature of oxidative damage (cf. section 3.3). Although these observations are puzzling, they cannot be taken as disproving the overwhelming evidence of a contribution of ROS to the aging process.

The evidence existing on the accumulation of oxidative damage in cells and tissues of aging individuals supports the mitochondrial theory of aging in a straightforward way. Nevertheless, the effects of ROS are not limited to structural damage to cell components. In fact, some ROS are now recognized as inducing redox changes in molecules involved in signalling pathways, and ROS themselves are seen today as possible physiological signals (cf. section 2.3). We may therefore ask whether the structural damage induced by ROS, particularly to mtDNA, is sufficient to explain the aging phenotype, or whether more complex factors, possibly involving the derangement of signalling pathways, are the major driving force for the metabolic failure characterizing aging (cf. section 3.2).

3.1.2. Functional impact of respiratory chain damage

There is overwhelming evidence that the bioenergetic function of mitochondria declines with aging, especially in post-mitotic tissues (386, 66, 372, 227, 404, 93, 139, 363, 362, 298) and this decline may be induced by ROS, as demonstrated by the decrease of respiratory activities in MnSOD-deficient mice (260). It is also well documented that exposure of mitochondria to ROS can affect the respiratory activity via oxidative damage of CL, which is required for the optimal functioning of the enzyme complexes (302). Relevant to this observation is the decline of the content of mitochondrial CL and the increase of oxidized CL with aging (316, 317).

A mosaic pattern of cytochrome oxidase alterations in muscular tissues from elderly individuals has been found by histochemical investigations (277-280, 399). The absent or rare decrease of activity of individual respiratory complexes observed in many studies (155, 6, 273) may reflect the existence of threshold effects (335); moreover, electron transfer activity alone does not reflect the whole bioenergetic capacity of the mitochondria, since also membrane potential and ATP synthesis should be taken in account. In this regard a study (139) exploiting fibroblasts from donors of different ages, carefully exploring the P:O ratios is very informative in detecting a significant fall of phosphorylation efficiency in the elder. However, tissue differences also exist in the susceptibility of mitochondria to ROS-induced damage. For example the ATP content and production declined to 50% with age in the skeletal muscle (gastrocnemius) but underwent no change in the heart of Fischer-344 rats (93).

The altered mitochondrial function is accompanied by decreased mtDNA transcription (52, 117, 106, 18, 19); however compensatory mechanisms may exist, since the levels of Complex I subunits ND1 and ND3 increase in platelets from aged individuals, as observed in our laboratory (262). Other reports also showed decreased mitochondrial protein synthesis and decreased levels of mtDNA-encoded proteins (12, 253, 333).

Mitochondrial oxidative stress in mouse skeletal muscle is increased with age, leading to oxidative DNA damage and alterations in mitochondrial bioenergetic function (251).

Mitochondrial protein synthesis is unimpaired in mtDNA mutator mice (cf. previous section), consistent with the observed minor alterations of steady-state levels of mitochondrial transcripts. Despite that, the stability of several respiratory chain complexes is severely impaired, thus suggesting that the aging phenotype is caused by amino acid substitutions in the mtDNA-encoded protein subunits of the respiratory enzymes. In turn, the alteration of protein sequence leads to respiratory chain deficiency. It is worth mentioning that the mitochondrial respiratory proteins are specifically decreased in abundance in the brains of the mutator mice harbouring extensive mtDNA mutations (158). Those changes were not attributed to decreased transcription but to post-translational effects associated with the Poly mutation.

The notion that the respiratory chain is mainly controlled at the level of Complex I (393) suggests that the main alterations due to aging must be found at the level of this enzyme (414, 20, 227). In agreement with this observation, NADH-dependent State 3 respiration is decreased in aged rats (226, 227, 393).

The protein components of Complex I were found to be reduced or absent in aged motor neurons from elderly subjects (337), leading to neuron loss and sarcopenia. The activity of Complex I was investigated in human platelets from young and aged individuals (263). In this case, the most striking result was the decrease of rotenone sensitivity of the enzyme. This study demonstrates that bioenergetic alterations may exist not only in cells of post-mitotic tissues but also in cells directly deriving from mitotic divisions where it is usually assumed that selection washes away the damaged cells (44). Nevertheless, if cells can meet their energy needs by glycolysis, they will survive and maintain the mitochondrial alterations.

A recent study (360) demonstrated that fibroblasts from centenarians display significantly lower Complex I-driven ATP synthesis and higher production of H₂O₂ in comparison with 75 years old subjects. Despite these changes, bioenergetics of the cells from centenarians appeared to operate normally, due to the significant increase of mitochondrial mass, allowing total ATP synthesis per cell to be unchanged. This lack of functional consequences appears to be due to a decreased mitophagy, induced by the presence of hyperfused, elongated mitochondria. The overall data indicate that longevity is characterized by a preserved bioenergetic function likely attained by a successful mitochondria remodelling that can compensate for functional defects through an increase in mass, i.e. a sort of mitochondrial "hypertrophy".

3.2. ROS signalling in aging

Recent investigations attempt to relate the mitochondrial changes with the cellular environment, and the cross-talk between mitochondrial and nuclear genome is receiving increased attention. The availability of techniques such as serial analysis of gene expression (391, 10, 168) has been applied to aging studies in *C. elegans* (187) and in rodent heart (9).

Also the yeast model has provided some important clues. Yeast is characterized by the retrograde response triggered by mitochondrial dysfunction (43) and activating specific signalling proteins. These proteins then migrate to the nucleus and induce numerous nuclear genes coding for metabolic enzymes and stress proteins (183). The result is an extension of yeast longevity.

Changes in the redox status of cellular components by oxidative stress during aging are considered the cause of the observed increased contents or DNA-binding activities of such transcription factors as NF- κ B, AP-1 and HIF-1 (213, 50, 197), of heat-shock proteins (51) and of heme oxygenase (49). Their increased activity is considered a compensatory

mechanism for cellular protection and may depend either upon direct alteration of the factor or indirectly through activation of related transduction pathways (197).

A decrease of NAD^+ and of the NAD^+/NADH ratio is a hallmark of aging (36); such decrease may be due to decreased mitochondrial respiration but also to the enhanced demand of NAD^+ -requiring enzymes such as PARP and sirtuins. The decline in nuclear NAD^+ during aging in mice leads to the accumulation of HIF-1 α under normoxic conditions, simulating the Warburg effect in cancer cells (133) (Figure 3).

Deleting SIRT1 accelerates this process, whereas raising NAD^+ levels in old mice restores mitochondrial function to that of a young mouse in a SIRT1-dependent manner. Thus, a pseudohypoxic state that disrupts PGC-1 α/β -independent nuclear-mitochondrial communication contributes to the decline in mitochondrial function with age, a process that is apparently reversible. The activation of HIF1 α leads to selective loss of mtDNA-encoded, but not nuclear-encoded, OXPHOS subunits. It is important to note that this unbalance disrupts OXPHOS before the accumulation of evident DNA damage (133).

It must be considered in this connection, however, that also ROS appear to activate HIF1 α (see Section 2.2). In addition, HIF-1 α elevation can be elicited not only via NAD^+ /sirtuins-mediated stabilization, but also via translation induced by TOR (Target Of Rapamycin), therefore it is suggested that the pseudohypoxic state is not necessarily caused by mitochondrial dysfunction, but rather driven by TOR hyperfunction (230) (see below).

INSERT FIGURE 3

In accordance with the importance of oxidative stress in activation of redox-sensitive transcription factors, caloric restriction, the main known factor recognized to delay aging (13, 109), was found to prevent their activation (197). Similarly, hepatocytes from old

mice (233) and rats (175) showed reduced activation of ERK by H₂O₂, and the effect was suppressed by caloric restriction (175). Available data on redox-responsive transcription factors suggest that their uncontrolled activation in aging could lead to serious chronic pathogenic conditions characterized by what has been called “molecular inflammation” (65) or inflammaging (112).

We now know that the longevity response to dietary restriction is actively regulated by nutrient-sensing pathways involving the TOR, AMP kinase, sirtuins and insulin/insulin-like growth factor (IGF-1) signalling in a variety of model organisms (195). In nutrient replete conditions, organisms develop, grow, and age quickly. When nutrients become scarce as with dietary restriction, growth and development decline, stress response pathways become induced and organisms live longer.

According to Blagosklonny (34), aging is not a program, but it is a quasi-program, a useless and unintentional continuation of developmental programs. Similarly, cellular senescence is a continuation of cellular growth. The same TOR pathway, which drives developmental growth, later drives aging and its associated diseases. Actually, TOR has been proposed to be the central regulator of aging (34, 230, 403). Under nutrient-rich conditions TOR promotes growth by phosphorylating several factors, including the ribosomal protein S6K1, related with protein synthesis, whereas under starvation conditions its inhibition slows down cell proliferation and activates cell maintenance pathways involved in autophagy, apoptosis, and mitochondrial stress responses. Indeed TOR inhibition by starvation or by rapamycin has been shown to prolong the life span of several organisms (258). Reducing TOR activity increases autophagic flux, enhances mitochondrial membrane potential, reduces reactive oxygen species within the cell, and increases replicative life span. These effects appear to be mediated in part by an interaction between p62/SQSTM1 and Keap1 which allows nuclear accumulation of NRF2, increased expression of NRF1, and increased

expression of nuclear-encoded mitochondrial genes, such as the mitochondrial transcription factor A, and mitochondrial-encoded genes involved in oxidative phosphorylation (231, 31).

The TOR-centric theory of aging is not necessarily in contrast with the traditional view of the mitochondrial theory: antiaging drugs such as rapamycin, metformin, berberine, resveratrol, vitamin D3, 2-deoxyglucose, and acetylsalicylic acid, attenuate the level of constitutive TOR signalling (79). In parallel, they suppress the level of constitutive DNA damage induced by endogenous ROS. While the primary target of each of these agents may be different, the data obtained on several human cancer cell lines, WI-38 fibroblasts and normal lymphocytes suggest common downstream mechanism in which the decline in mTOR/S6K1 signalling and translation rate is coupled with a reduction of oxidative phosphorylation and ROS that leads to decreased oxidative DNA damage.

In this context, the adaptor protein $p66^{Shc}$ (Section 2.1.4) could be a link between ROS and mTOR pathway (300). With its double identity as generator of mitochondrial oxidant species and as signalling adaptor in the insulin receptor cascade, $p66^{Shc}$ has drawn major attention as a negative determinant of life span and healthy longevity in mammals.

In one scenario, $p66^{Shc}$ action on S6K may lead to increased mitochondrial metabolism and, as a consequence, to a rise of mitochondrial ROS (385), as observed in cells where $p66^{Shc}$ is overexpressed. In other words, TOR/S6K may mediate, at least in part, the pro-oxidant action of $p66^{Shc}$. Ablation of $p66^{Shc}$, by leading to reduced responsiveness of S6K to nutrients, creates a Rapamycin-like (although presumably milder) signalling block that conceivably promotes animal longevity, like caloric restriction.

More intriguingly, ROS may act upstream of the $p66$ /S6K module, since $p66^{Shc}$ not only generates ROS, but is also stimulated by oxidants (265). For instance, in fibroblasts exposed to oxidative stress, PI3K/AkT activation by ROS is mediated, at least to some extent, by $p66^{Shc}$ (286). AkT can, in turn, activate mTOR. ROS are also generated in mitochondria in

response to energy substrates; these species may increase the phosphorylation/expression level of $p66^{Shc}$, thereby promoting its (redox-independent) stimulatory action on S6K. This would represent an intriguing alternative route for nutrients to signal, *via* mitochondria, ROS and $p66^{Shc}$, to the mTOR/S6K cascade.

Interestingly, oxidative stress and DNA damage can also induce the expression of miR-210, a micro-RNA that negatively regulates mitochondrial respiration under hypoxic conditions by limiting the functionality of the respiratory chain in many different cell types. In fact, at least two members of Complex I and II (i.e. NDUFA4 and SDHD, respectively) are targets of miR-210, with consequential transmembrane potential reduction, ROS production increase and induction of an apoptotic-like mitochondrial phenotype (321). Another senescence-induced miRNA that preferentially localizes to mitochondria, miR-494, was found to be even more effective than miR-210 in inducing a senescent phenotype in fibroblasts upon pre-miRNA transfection, thus indicating a role for these molecules as transducers of ROS or DNA damage associated signalling (102). Since these miRNAs can induce ROS production, a self-sustaining ROS vicious cycle that involves these two miRNAs could be hypothesized.

The analysis of young and old mouse brains further confirmed the role of miRNAs in reducing the mitochondrial activity during the senescence process and aging. The profiling of both miRNAs and proteins showed 70 differentially regulated miRNAs of which 27 were predicted to target subunits of Complex III, Complex IV and ATPase, as confirmed by protein profiling (234), supporting the fact that miRNA-driven mitochondrial dysfunction and ROS increase are common traits shared by aging tissues as well as senescent cells (212)

3.2.1. Beneficial effects of ROS in aging: a sort of preconditioning

Some studies suggest that ROS are beneficial and not detrimental for life (119, 332, 211) when they serve as molecular signals to ultimately induce endogenous defence mechanisms culminating in increased stress resistance and longevity (76, 424, 332, 247), an adaptive response more specifically named mitochondrial hormesis or mitohormesis (376, 48, 417).

Knockdown of Complex I, III, IV, or V has been shown to increase lifespan in both *C. elegans* (90, 413) and *Drosophila* (68), whereas knockdown of Complex II does not extend lifespan (204). Accordingly, mutation of Complexes I or III subunits also increases lifespan, despite concomitantly increasing superoxide levels (105, 413), whereas the mutation of Complex II subunits that also elevates oxidative stress accelerates aging (176). It is possible that a specific ROS-mediated signal activates a prosurvival program that overcomes the deleterious effects of increased oxidative damage (73) caused by mutations in Complexes I or III, but that this prosurvival program cannot compensate for alterations in the Krebs cycle caused by mutations in Complex II.

Hydrogen peroxide enhances chronological lifespan in caloric-restricted budding yeast cells in parallel with a decrease in superoxide anions. This decrease is caused by induction by hydrogen peroxide of the activity of the cytosolic SOD1 and the mitochondrial SOD2, respectively (264). The induction of SOD activity by hydrogen peroxide is consistent with the earlier demonstration that the lifespan of budding yeast is extended by SOD overexpression (101, 157). These findings are also consistent with earlier reports that sublethal concentrations of hydrogen peroxide induce transcription of both the *SOD1* and *SOD2* genes as well as an increase in levels of the corresponding proteins (132).

A transient increase in intracellular hydrogen peroxide has also been implicated in lifespan extension associated with impaired insulin/IGF-1 signalling in *C. elegans*. Such

impairment disrupts glucose uptake and upregulates superoxide dismutases and other oxidative stress defences, leading to a subsequent decline in overall levels of ROS (419).

According to these results, the caloric restriction induces an initial stress response with an increase of ROS generation. The consequent induction of protective mechanisms, mainly antioxidant enzymes, results in an increased antioxidant defence and a lower level of ROS and ROS-induced damage (332). This interpretation agrees with the finding that the time necessary for caloric restriction to lower ROS production is of several months (140). Irrespective of the interpretation, it is clear that aging is linked to ROS-induced damage, and we can combat aging by eventually decreasing ROS levels in mitochondria.

It has been recently shown that increased lifespan in *C. elegans* is promoted by either dietary restriction or Complex I inhibition, but these two ways are not additive (351), indicating that impaired Complex I activity mimics dietary restriction. Consistent with the concept of mitohormesis, Complex I inhibition transiently increases mitochondrial formation of ROS that activate PMK-1/p38 MAP kinase and SKN-1/NRF-2. Sirtuins are also involved in ROS signalling (352): a sirtuin-dependent nicotinamide-N-methyltransferase generates 1-methylnicotinamide (MNA) from nicotinamide. MNA serves as a substrate for a newly identified aldehyde oxidase to generate hydrogen peroxide, which acts as a mitohormetic reactive oxygen species signal to promote *C. elegans* longevity.

Another aspect of defence against mitochondrial stress is the mitochondrial UPR (161, 309) that may be triggered by inhibition of the TOR pathway by rapamycin or by caloric restriction (403). The mtUPR is analogous to the better known endoplasmic reticulum UPR (see Section 2.3.1). The primary purpose of this initial stress response at low ROS concentration is to improve the folding environment of newly synthesized mitochondrial proteins and enhance mitochondrial functions.

3.3. Modifications not due to ROS: Are ROS really involved in aging?

Evidence has been presented that only a small fraction of the mtDNA mutations accumulating in *Drosophila* upon aging are G-C to T-A transversions, a signature of ROS oxidative damage to the bases (181). Moreover, loss-of-function mutations in the mitochondrial superoxide dismutase, Sod2 have no detectable influence on the somatic mtDNA mutation frequency, suggesting that mutations arise primarily from errors that occur during mtDNA replication.

These results are in line with the “Damage Theory” of Gladyshev (132) who proposes that infidelity, heterogeneity, and imperfectness of each and every biological process may be responsible for the inevitable accumulation of by-products and other damage forms. Thus, according to this view, aging is indeed due to damage to cellular structures, but the damage is not, or only in part, due to oxidative modifications by ROS.

The results, however, cannot be reconciled with the evidence shown in Section 3.1.1 that mtDNA mutations are proportional to the levels of 8-oxodG and to the extent of oxidative stress. It is however possible that both mechanisms of mtDNA mutagenesis exist. It has also to be borne in mind that mtDNA damage may not be the initial step leading to the aging phenotype, whilst lipid and protein damage by ROS may initially trigger the events leading to aging. We will more deeply discuss this aspect in Section 4.

ROS-dependent and ROS-independent mechanisms regulating lifespan through the mitochondrial respiratory chain have been reviewed (355), with particular emphasis on the metabolic consequences of changing the NAD^+/NADH ratio by changing the rate of respiration. Aging is characterized by a low NAD^+/NADH ratio, due to decrease of mitochondrial respiration (365, 36) (cf. Section 3.2).

If NADH is not reoxidized to NAD^+ by Complex I, there is a shortage of NAD^+ available for the conversion of glyceraldehyde-3-phosphate into 1,3-biphosphoglycerate

causing accumulation of dihydroxyacetone, which decomposes into methylglyoxal, the main initiator of nonenzymatic glycation (24, 323).

Furthermore, the ratio of NAD^+/NADH , together with AMP/ATP levels, is a major sensor of the energetic and redox state of the cell (171). Based on the information provided by these sensors, cells can choose between running under a proaging or prosurvival program. Dietary restriction that counteracts aging is characterized by a change shifting the NAD^+/NADH ratio to the oxidized form (237). Different experimental reports support the notion that it is possible to regulate longevity by manipulating the levels of NADH and NAD^+ .

In addition, both sirtuins and poly(ADP-ribose) polymerase use NAD^+ or some of its metabolites as cofactors. This implies that a change in the ratio of NAD^+/NADH elicited by a change in the electron transfer chain activity would alter the activity of these enzymes, thus activating or repressing a prosurvival genetic program (143).

Again, we wish to point out that even these mechanisms, that are apparently not dependent on ROS, are however the consequence of a deficiency of mitochondrial respiration that is largely a consequence of ROS attack (Section 3.1.2).

4. A UNIFYING HYPOTHESIS INVOLVING SUPERCOMPLEX DESTABILIZATION IN AGING

Contrary to the view of a random organization of the respiratory chain complexes prevailing in the last decades of the past century (146), evidence has now accumulated that a large proportion of the mitochondrial respiratory chain complexes in a variety of organisms is arranged in supramolecular assemblies called supercomplexes or respirasomes (348, 3, 224).

The natural assembly of the respiratory complexes I, III, and IV into supramolecular stoichiometric entities, such as $\text{I}_1\text{III}_2\text{IV}_{0-4}$, is not just a mere structural feature

but has deep functional implications on the properties of the respiratory chain (reviewed in 224). The most striking and obvious consequence of supercomplex dissociation is loss of enzymatic channelling (121, 122). Another functional consequence of disruption of the supercomplex (SC) core structure I_1III_2 is loss of stability of Complex I. Flux control analysis in aerobic respiration in coupled liver mitochondria (393) showed that Complex I has little control in young rats but very high control in the old animals, meaning that defective operation induced by aging at the level of this enzyme is reflected on the entire OXPHOS process.

There is growing awareness that the free complexes can co-exist with supercomplexes. In this context, an integrated model, *the plasticity model*, has been proposed for the organization of the mitochondrial electron transport chain (3). The previous opposed models, solid vs. fluid, would be two possible extreme situations of a dynamic range of different molecular associations between respiratory complexes. The plasticity model and the dynamics of mitochondrial supercomplexes are widely discussed (1) in a recent review.

A fundamental prediction of the plasticity model is that, *in vivo*, the mitochondrial respiratory chain should be able to work both when supercomplexes are present and when the formation of supercomplexes is prevented. Indeed, several studies *in vitro* support the view that electron transfer in the respiratory chain can occur in absence of supercomplexes though with lower efficiency (122). Among the factors able to modulate supercomplex association are the mitochondrial membrane potential and posttranslational changes of the individual complexes, although these factors and the mechanisms through which they act are still unknown. In the reconstructed models of the $I_1III_2IV_1$ supercomplex (96), only limited apparent interaction sites exist between neighbouring complexes. Some of the gap volumes may be lipid filled. It is worth noting that CL, the signature phospholipid of mitochondria, can also integrate into the structure of the respiratory complexes and act as a critical determinant

of the respiratory function. Consistent with this observation, impaired CL synthesis gives rise to more labile respiratory supercomplexes (cf. Section 4.1.2) in yeast and in human patients (267, 259). Another recent advance is the discovery of non-OXPHOS proteins that appear to adhere to and seal the individual respiratory complexes to form stable assemblages and to prevent electron leakage (408).

We first suggested an implication of supercomplex disorganization as the missing link between oxidative stress and energy failure (223). A dissociation of supercomplex association was proposed to occur under conditions of oxidative stress. The consequent loss of facilitated electron channelling leads to resumption of a less efficient random diffusional behaviour, with electron transfer depending upon the collisional encounters of the free ubiquinone molecules with the partner complexes. Dissociation of supercomplexes might have further deleterious consequences, such as disassembly of complex I and III subunits and loss of electron transfer and/or proton translocation (223); the consequent alteration of electron transfer may elicit further induction of ROS generation. Following this line of thought, the different susceptibility of different types of cells and tissues to ROS damage may be a consequence of the extent and tightness of supercomplex organisation in their respiratory chains. Supercomplex stability depends on phospholipids content and composition of their mitochondrial membranes. It would be interesting to show if a correlation exists between longevity of a species and tightness of supercomplex association, although no data presently exist on this issue. Supercomplex dissociation may have deep metabolic consequences, as depicted in the scheme in Figure 4. An initial enhanced ROS generation due to different possible reasons and originating in different regions of the cell besides mitochondria (229) would induce supercomplex disorganization eventually leading to possible decrease of Complex I assembly. Both the lack of efficient electron channelling and the loss of Complex I

would decrease NAD-linked respiration and ATP synthesis. In the following sections we briefly summarize the experimental evidence pertaining to this hypothesis.

INSERT FIGURE 4

4.1 Supercomplex dissociation and aging

4.1.1. *Evidences for supercomplex dissociation in aging*

Analysis of the occurrence of respiratory supercomplexes comprised of various stoichiometries of complexes I, III and IV reveals age-related variations, suggesting that destabilization of their supramolecular organization may be crucial for the development of the aging-phenotype (84, 135). In cardiac mitochondria from old rats (136), supercomplexes of the highest molecular weight decline to the greatest extent with age. It is noteworthy that this supercomplex disintegration is not caused by age-associated decrements in a particular enzyme component. In another study in mitochondria from rat skeletal muscle (243), proteomic analysis revealed an age-associated increase of the heavier supercomplexes, which was hypothesized to be a compensation for the significant loss of the smaller supercomplex I_1III_2 .

Also in mitochondria of brain cortex from aged rats (30-months), profound adverse changes in the supramolecular organization of the respiratory chain complexes as well as of the FoF1 ATP synthase were evidenced (114); notably, the overall decline with age (40%) in the supercomplexes containing Complex I is caused to large extent by the pronounced decline (58%) of abundance of the supercomplex I_1III_2 .

Studies of the effect of heart failure on cardiac mitochondria (334) showed that electron transfer in absence of supercomplex organization is decreased even if activity of the individual complexes is normal. In proteoliposomes enriched in supercomplex I_1III_2 (at

protein to phospholipid ratio 1:1), electron transfer between Complex I and Complex III (NADH-cytochrome *c* oxidoreductase) is more efficient than predicted by random behaviour on the basis of the activities of the individual complexes (124). However, when the supercomplex is dissociated into its individual enzyme components (e.g. by lipid dilution of the proteoliposomes to a ratio of 1:30) the rate of NADH-cytochrome *c* oxidoreductase is much lower. Similar results were obtained by dissociation of the supercomplexes by detergents (252). This is a demonstration that supercomplex formation indeed enhances the rate of electron transfer above that occurring via a ubiquinone pool in the membrane (27, 124, 252), whereas loss of supercomplex organization induces less efficient pool activity of the quinone.

The progeroid profile of the mutator mice harbouring a mutation in the Poly proof-reading domain and hence exhibiting multiple mtDNA mutations (Section 3) is accompanied by decreased respiration and impeded assembly of respiratory complexes (98). Despite that only Complex IV subunits were significantly reduced, the steady state levels and activities of the other complexes, particularly Complex I, were secondarily affected and strongly decreased. This is in line with the idea that a decreased assembly of Complex IV may affect the stability of Complex I through disruption of the supercomplex organization (80). Nevertheless, although mtDNA deletions and concomitant loss of respiratory activity have been shown to occur in an age-dependent fashion in the rat kidney, no significant differences were found when comparing the protein abundance of individual respiratory complexes and the supercomplex profiles between young and old mitochondria (287). It must be noted that in this latter study the respiratory activities were only marginally decreased but it is possible that kidney mitochondria are not yet compromised at the age studied (24 months).

4.1.2 Peroxidised phospholipids prevent supercomplex formation

Lipids impose on membrane proteins the correct conformation for optimal activity (219, 173) and a correct template for membrane protein topogenesis (92).

The phospholipids in closest vicinity to the protein surface, as well as those in the free bilayer, are actually highly mobile and free to exchange, but crystallographic studies have shown the presence of a few tightly bound CL molecules in each of the crystal structures of Complex III (210) and Complex IV (291). These results suggest that CL is an integral component of these proteins, and that its presence is critical to protein folding and function. There are now extensive indications that CL stabilises respiratory supercomplexes as well as the individual complexes (405, 302).

The availability of a CL-lacking yeast mutant provided the opportunity to demonstrate that mitochondrial membranes still contained the III₂-IV₂ supercomplex, but that it was significantly less stable than supercomplexes in the parental strain (421, 318).

Mutations of tafazzin, an acyltransferase involved in the synthesis of mature tetralinoleyl CL (169), result in Barth syndrome, a cardio-skeletal myopathy with neutropenia, characterised by respiratory chain dysfunction. The CL defect in Barth syndrome results in destabilisation of the supercomplexes by weakening the interactions between respiratory complexes (259, 137).

Flux control analysis (124) showed that the maintenance of a supercomplex I-III in proteoliposomes is abolished if lipid peroxidation is induced by 2,2'-azobis-(2-amidinopropane) dihydrochloride before reconstitution. Evidently, the distortion of the lipid bilayer induced by peroxidation and the alteration of the tightly bound phospholipids determine dissociation of the supercomplex originally present in the preparation.

Moreover, mitochondrial-mediated ROS production affects complex I and IV activity through CL peroxidation in beef heart submitochondrial particles (303, 304). CL liposomes added exogenously to mitochondria from aged rats almost completely restored the

activity of these enzyme complexes to the values of young control animals (316, 317). Neither other major phospholipid components of mitochondria, such as phosphatidylcholine and phosphatidylethanolamine, nor peroxidised CL could replace this effect of CL. Altogether, these results indicate that the defect in complex I and IV activities observed in mitochondria from aged rats could be ascribed, at least in part, to oxidative damage to mitochondrial CL.

4.1.3. Loss of supercomplexes decreases the stability of Complex I and enhances generation of ROS

Analysis of the state of supercomplexes in patients with an isolated deficiency of single complexes (349) and in cultured cell models harbouring cytochrome b mutations (2, 80, 87, 259) provided evidence that the formation of respirasomes affects the stability of Complex I. From these findings, supercomplex assembly emerged as a necessary step for respiration, its defect setting the threshold for respiratory impairment in mtDNA mutant cells.

Genetic alterations leading to a loss of Complex III prevented respirasome formation and led to secondary loss of Complex I, and therefore primary Complex III assembly deficiencies presented as Complex III/I defects. Conversely, Complex III stability was not influenced by the absence of Complex I.

The disassembly of Complex I does not appear to result from altered biogenesis assembly (1). The available evidence would favour the following course of events: misassembled Complex III prevents formation of supercomplex; the lack of the supercomplex association induces an enhanced ROS generation from Complex I (see below), with consequent damage to neighbour molecules and to Complex I itself (86, 88) which is vulnerable to oxidative stress both directly and through the lipid peroxidation, particularly of CL (303).

The possibility that loss of supercomplex organisation may enhance ROS generation by the respiratory chain has been advanced on theoretical grounds (223, 224, 84): a tighter organisation of the respiratory enzymes may hide auto-oxidisable prosthetic groups hindering their reaction with oxygen. Alternatively, slowing electron flow in the chain may keep the prosthetic groups in a more reduced form allowing them to interact with oxygen.

The molecular structure of the individual complexes would not seem to allow a close apposition of the matrix arm of Complex I, where the prosthetic groups are localised, with either Complex III or IV (347, 314). However, the actual shape of the I₁III₂IV₁ supercomplex from bovine heart (347) suggests a slightly different conformation of Complex I in the supercomplex, with the matrix arm showing a higher bending towards the membrane (and presumably Complex III). This observation agrees with the notion that Complex I may undergo important conformational changes (325). Moreover, the observed destabilisation of Complex I in absence of supercomplex may render the 51 kDa subunit containing the FMN more “loose” allowing it to interact with oxygen.

It was suggested (301) that supercomplex organization of Complex I within the chain prevents excessive superoxide production on oxidation of NAD-linked substrates because the efficient channelling helps maintaining the chain in the oxidized state. On the other hand, on succinate oxidation, the backward electron flow keeps the centres in Complex I more reduced favouring production of superoxide. In relation to this it is interesting to note that Complex II does not form a respirasome.

Additional circumstantial evidence on the role of supercomplex organisation comes from the observation that high mitochondrial membrane potential elicits ROS generation, while uncoupling strongly reduces ROS production (222, 184). Although other explanations may be given to these observations (cf. section 2.2.1), they are compatible with the suggestion

(322) that high membrane potential may dissociate the supercomplexes into the individual units.

A recent study (252) gave the first direct demonstration that loss of supercomplex organization causes an enhancement of ROS generation by Complex I (Fig. 5). This is clearly evident both in reconstituted proteoliposomes undergoing mild detergent treatment in the presence of dodecyl maltoside (DDM), where the dissociation of Complex I from the respirasome is accompanied by a three/four-fold increase of ROS generation, and in mitochondrial membranes. In the latter case, the existence of endogenous systems operating to reduce ROS levels in the mitochondrial sample (i.e. mitochondrial glutathione peroxidase, Mn-superoxide dismutase and non-enzymatic endogenous antioxidants) might have counteracted the dramatic effects of the complete dissociation of Complex I, thus leading to a two-fold only increase of the measured ROS production.

A further demonstration of the enhanced ROS generation due to supercomplex disruption was similarly obtained in a model system of reconstituted binary Complex I/Complex III at high lipid to protein ratio (30:1) (252) where formation of the supercomplex I_1III_2 is prevented. Likewise, the generation of superoxide is several folds higher than in the same system reconstituted at a 1:1 ratio, which is rich in supercomplexes. It is worth noting that in the experiments reported in the above mentioned study, ROS production is investigated in the presence of inhibitors (mucidin and rotenone) that prevent electron transfer to any possible acceptor. For this reason the redox centres in Complex I are maximally reduced both in the situations where supercomplexes are maintained and in the situations where Complex I is free. We therefore exclude the possible reasoning that facilitation of electron flow by substrate channelling within the respirasome could maintain the redox components of the complexes in the oxidised state and limit ROS formation.

INSERT FIGURE 5

The *in vitro* study quoted above (252) is supported by several observations in cellular and animal models linking together supercomplex dissociation and enhanced ROS production. In particular, very recent observations (86) show that diminished stability of SC and Complex I is associated with increased levels of ROS in mouse lung fibroblasts lacking the Rieske iron-sulphur protein of Complex III and hence devoid of the supercomplexes containing Complex I.

Also in mouse fibroblasts expressing the activated form of the k-ras oncogene, a strong decrease of high molecular weight supercomplexes correlated with higher ROS generation in comparison with wild type fibroblasts (225). Moreover, enhanced ROS generation and oxidative stress were found in yeast mutants lacking the supercomplex assembly factor Rcf1 and thus devoid of supercomplexes III-IV (64, 371, 398). Since the yeast *S. cerevisiae* lacks Complex I, in this case we may consider the origin of the extra ROS being presumably Complex III.

Recently, the biochemical properties of cybrids carrying a human cytochrome b missense mutation m.15579A>G were analysed (128). This mutation causes a dramatic reduction in the Complex III activity and in Complex III-driven mitochondrial ATP synthesis, but does not induce disassembly of the supercomplex containing Complex I. However, the mutation enhances superoxide production, as indicated by direct measurements in mitochondria and by the imbalance of glutathione homeostasis in intact cybrids. The amounts of CIII dimer and III₂IV₁ were reduced, whereas those of I₁III₂IV_n slightly increased, suggesting that the deleterious effects of p.278Y>C mutation on cytochrome b are palliated when Complex III is assembled into the supercomplexes I₁III₂IV_n, in contrast to when it is

found alone. In this case it is likely that the enhanced ROS formation is the result of derangement of the Q-cycle in Complex III.

It is tempting to suggest that physiological changes in the dynamic equilibrium of respiratory supercomplexes with isolated complexes may be aimed to regulation of ROS levels in the cell, in view of the well documented role of ROS in cellular redox signalling.

4.2. Concluding remarks: is there an evidence for the vicious circle?

The model we have just proposed poses supercomplex dissociation as having a double influence in relation to ROS. On the one hand, ROS contribute to dissociate supercomplexes, but on the other hand, supercomplex dissociation enhances ROS generation. This means that if these events are not tightly controlled they may initiate a vicious circle of ROS generation. Is this the series of events that conduce to aging? We have discussed in the previous sections the controversial existence of a vicious circle of ROS generation and mitochondrial failure during aging. Although the mitochondrial theory of aging does not require that a vicious circle be operative (14), there is wide evidence that ROS generation increases in aging (Section 3). However, exceptions exist to this rule, the most striking being the lack of increase of ROS in the mutator mouse, in which extensive mitochondrial DNA mutations lead to mitochondrial defect and premature aging (383). Nevertheless the reason for such a lack of ROS generation may be in the excessive disruption of the respiratory chain. It is obvious that completely non-functional complexes cannot catalyse reactions, nor generate ROS.

Taken together, the observations collected in this review locate supercomplex dissociation in a physiological signalling network that can be easily altered and lead to a catastrophic event when the generation of ROS loses control.

We may envisage supercomplex association/dissociation to occur under physiological conditions, according to the plasticity model (3), in response to such stimuli as mitochondrial membrane potential and protein phosphorylation/dephosphorylation of the respiratory complexes. The ensuing changes in ROS generation modulate the ROS-dependent signalling pathways. These changes are reversible and are kept under strict control by changes in the initial stimuli.

We propose that the primary event responsible for aging is the structural damage induced by ROS in mitochondria, as predicted by the original mitochondrial theory of aging. This structural damage is likely to initiate preferentially within the membrane proteins and lipids, and may be influenced by the signalling pathways that sense the nutrition state of the organism, via a modulation of mitochondrial activity and ROS generation. Retrograde signals starting from mitochondria may also induce compensatory mechanisms attempting to counteract the ROS generation and consequent damage. MtDNA mutations, although present, may not necessarily be an early phenomenon in the aging process.

A possible series of events might be the following (Fig. 4). Progressive damage is induced by ROS to the mitochondrial membrane lipids and proteins. The level of ROS is affected by such factors as the nutrition state and the activity of the mTOR and insulin/IGF pathways. Direct protein damage and increased CL peroxidation hamper supercomplex association (124); this leads to further increase of ROS generation. ROS at low concentration may induce a retrograde response inhibiting TOR pathway and thus leading to a protective life-extending response, but at higher concentration they may induce further damage with loss of coordination of the signalling pathways. Mutations in mtDNA at a later stage would make the overall process irreversible and lead to the final aging phenotype.

LIST OF ABBREVIATIONS

AA, Antimycin A

AOX, alternative oxidase

BHM, bovine heart mitochondria

CL, cardiolipin

CoQ, Coenzyme Q (ubiquinone)

COX, cytochrome oxidase

DDM, dodecyl maltoside

DHO, dihydroorotate

ERK, extracellular signal-regulated kinase

ETF, electron transfer flavoprotein

GSH, glutathione

α GP, glycerol-3-phosphate

HIF-1 α , hypoxia-inducible factor

I, NADH-ubiquinone oxidoreductase

II, succinate-ubiquinone oxidoreductase

IGF-1, insulin-like growth factor

III, ubiquinol-cytochrome *c* oxidoreductase

IMM, inner mitochondrial membrane

IMS, intermembrane space

IV, cytochrome *c* oxidase

MNA, 1-methylnicotinamide

mtDNA, mitochondrial DNA

ND, alternative NAD(P)H dehydrogenases

NOX, NADPH NOX, NADPH oxidases

P66, cytochrome p66

Poly, mitochondrial DNA polymerase γ

Prx, Peroxiredoxins

PTPs, phosphoprotein phosphatases

Q, ubiquinone (Coenzyme Q)

ROS, reactive oxygen species

SC, supercomplex

SMP, submitochondrial particles

SOD, Superoxide dismutase

TOR, Target Of Rapamycin

UPR, Unfolded Protein Response

ACKNOWLEDGMENTS

The recent work from our laboratory that is quoted in this review has been supported by MIUR (grant number PRIN 20107Z8XBW_003).

We would like to thank Professor Harold Baum (Professor Emeritus at King's College London, UK) for critically reading and correcting the manuscript.

We apologize with the authors of the many papers that have not been cited because of the lack of space allowed.

AUTHOR DISCLOSURE STATEMENT

No competing financial interests exist.

REFERENCES

1. Acín-Pérez R, and Enriquez JA. The function of the respiratory supercomplexes: the plasticity model. *Biochim Biophys Acta* 1837: 444-450, 2014.
2. Acín-Pérez R, Bayona-Bafaluy MP, Fernández-Silva P, Moreno-Loshuertos R, Pérez-Martos A, Bruno C, Moraes CT, and Enriquez JA. Respiratory complex III is required to maintain complex I in mammalian mitochondria. *Mol Cell* 13: 805-815, 2004.
3. Acín-Pérez R, Fernández-Silva P, Peleato ML, Pérez-Martos A, and Enriquez JA. Respiratory active mitochondrial supercomplexes. *Mol Cell* 32: 529-539, 2008.
4. Agarwal S, and Sohal RS. Differential oxidative damage to mitochondrial proteins during aging. *Mech Ageing Dev* 85: 55-63, 1995.
5. Allen EM, and Mieyal JJ. Protein-thiol oxidation and cell death: regulatory role of glutaredoxins. *Antioxid Redox Signal* 17: 1748-1763, 2012.
6. Allen RG, Keogh BP, Tresini M, Gerhrad GS, Volker C, Pignolo RJ, Horton J, and Cristofalo VJ. Development and age-associated differences in electron transport potential and consequences for oxidant generation. *J Biol Chem* 272: 24805-24812, 1997.
7. Almadia-Pagán PF, de Costa J, Mendiola P, and Tocher DR. Comp Biochem Physiol Changes in tissue and mitochondrial membrane composition during rapid growth, maturation and aging in rainbow trout, *Oncorhynchus mykiss*. *B Biochem Mol Biol* 161: 404-412, 2012.
8. Ambrus A, Tretter L, and Adam-Vizi V. Inhibition of the alpha-ketoglutarate dehydrogenase-mediated reactive oxygen species generation by lipoic acid. *J Neurochem* 109 Suppl 1: 222-229, 2009.

9. Anisimov SV, and Boheler KR. Aging-associated changes in cardiac gene expression: large scale transcriptome analysis. *Adv Gerontol* 11: 67-75, 2003.
10. Anisimov SV. Serial Analysis of Gene Expression (SAGE): 13 years of application in research. *Curr Pharm Biotechnol* 9: 338-350, 2008.
11. Bae YS, Oh H, Rhee SG, and Yoo YD. Regulation of reactive oxygen species generation in cell signaling. *Mol Cells* 32: 491-509, 2011.
12. Bailey PJ, and Webster GC. Lowered rates of protein synthesis by mitochondria isolated from organisms of increasing age. *Mech Ageing Dev* 24: 233-241, 1984.
13. Barja G. Aging in vertebrates, and the effect of caloric restriction: a mitochondrial free radical production-DNA damage mechanism? *Biol Rev* 79: 235-251, 2004.
14. Barja G. Updating the mitochondrial free radical theory of aging: an integrated view, key aspects, and confounding concepts. *Antioxid Redox Signal* 19: 1420-1445, 2013.
15. Barja G, Cadenas S, Rojas C, López-Torres M, and Pérez-Campo R. A decrease of free radical production near critical sites as the main cause of maximum longevity in animals. *Comp Biochem Physiol* 108B: 501-512, 1994.
16. Barja G, and Herrero A. Oxidative damage to mitochondrial DNA is inversely related to maximum life span in the heart and brain of mammals. *FASEB J* 14: 312-318, 2000.
17. Barogi S, Baracca A, Cavazzoni M, Parenti Castelli G, and Lenaz G. Effect of the oxidative stress induced by adriamycin on rat hepatocyte bioenergetics during ageing. *Mech Ageing Dev* 113: 1-21, 2000.
18. Barrientos A, Casademont J, Cardellach F, Ardite E, Estivill X, Urbano-Márquez A, Fernández-Checa JC, and Nunes V. Qualitative and quantitative changes in skeletal

- muscle mtDNA and expression of mitochondrial-encoded genes in the human aging process. *Biochem Mol Med* 62: 165-171, 1997.
19. Barrientos A, Casademont J, Cardellach F, Estivill X, Urbano-Marquez A, and Nunes V. Reduced steady-state levels of mitochondrial RNA and increased mitochondrial DNA amount in human brain with aging. *Brain Res Mol Brain Res* 52: 284-289, 1997.
 20. Barrientos A, and Moraes CT. Titrating the effects of mitochondrial complex I impairment in the cell physiology. *J Biol Chem* 274: 16188-16197, 1999.
 21. Bejma J, and Ji LL. Aging and acute exercise enhance free radical generation in rat skeletal muscle. *J Appl Physiol* 87: 465-470, 1999.
 22. Bell EL, Klimova T, and Chandel NS. Targeting the mitochondria for cancer therapy: regulation of hypoxia-inducible factor by mitochondria. *Antioxid Redox Signal* 10: 635-640, 2008.
 23. Bell EL, Klimova TA, Eisenbart J, Schumacker PT, and Chandel NS Mitochondrial reactive oxygen species trigger hypoxia-inducible factor-dependent extension of the replicative life span during hypoxia. *Mol Cell Biol* 27: 5737–5745, 2007.
 24. Bento CF, Marques F, Fernandes R, and Pereira P. Methylglyoxal alters the function and stability of critical components of the protein quality control. *PLoS One* 5: e13007, 2010.
 25. Bergamini E. Autophagy: a cell repair mechanism that retards ageing and age-associated diseases and can be intensified pharmacologically. *Mol Aspects Med* 27: 403-410, 2006.
 26. Bernardi P. The mitochondrial permeability transition pore: a mystery solved? *Front Physiol* 4: article 95, 2013.

27. Bianchi C, Fato R, Genova ML, Parenti Castelli G, and Lenaz G. Structural and functional organization of Complex I in the mitochondrial respiratory chain. *Biofactors* 18: 3-9, 2003.
28. Bianchi G, Di Giulio C, Rapino C, Antonucci A, and Castaldi A. p53 and p66 proteins compete for hypoxia-inducible factor 1 alpha stabilization in young and old rat hearts exposed to intermittent hypoxia. *Gerontology* 52: 17-23, 2006.
29. Bienert GP, Schjoerring JK, and Jahn TP. Membrane transport of hydrogen peroxide. *Biochim Biophys Acta* 1758: 994-1003, 2006.
30. Bienert GP, Møller AL, Kristiansen KA, Schulz A, Møller IM, Schjoerring JK, and Jahn TP. Specific aquaporins facilitate the diffusion of hydrogen peroxide across membranes. *J Biol Chem* 282: 1183-1192, 2007.
31. Bitto A, Lerner CA, Nacarelli T, Crowe E, Torres C, and Sell C. p62/SQSTM1 at the interface of aging, autophagy, and disease. *Age (Dordr)* 36: 9626, 2014.
32. Bjelakovic G, Nikolova D, Gluud C. Meta-regression analyses, meta-analyses, and trial sequential analyses of the effects of supplementation with beta-carotene, vitamin A, and vitamin E singly or in different combinations on all-cause mortality: do we have evidence for lack of harm? *PLoS One* 8: e74558, 2013.
33. Bjelakovic G, Nikolova D, Gluud LL, Simonetti RG, and Gluud C. Mortality in randomized trials of antioxidant supplements for primary and secondary prevention: systematic review and meta-analysis. *JAMA* 297: 842-857, 2007.
34. Blagosklonny MV. Answering the ultimate question "what is the proximal cause of aging?". *Aging (Albany NY)* 4: 861-877, 2012.
35. Boveris A, and Cadenas E. Mitochondrial production of superoxide anions and its relationship to the antimycin insensitive respiration. *FEBS Lett* 54: 311-314, 1975.

36. Braidy N, Guillemin GJ, Mansour H, Chan-Ling T, Poljak A, and Grant R. Age related changes in NAD⁺ metabolism oxidative stress and Sirt1 activity in wistar rats. *PLoS One* 6: e19194, 2011.
37. Brand MD. Uncoupling to survive? The role of mitochondrial inefficiency in ageing. *Exp Gerontol* 35: 811-820, 2000.
38. Bratic A, and Larsson NG. The role of mitochondria in aging. *J Clin Invest* 123: 951-957, 2013.
39. Braun RJ, and Westermann B. Mitochondrial dynamics in yeast cell death and aging. *Biochem Soc Trans* 39: 1520-1526, 2011.
40. Breyer V, Weigel I, Huang TT, and Pischetsrieder M. Endogenous mitochondrial oxidative stress in MnSOD-deficient mouse embryonic fibroblasts promotes mitochondrial DNA glycation. *Free Radic Biol Med* 52: 1744-1749, 2012.
41. Buffenstein R, Edrey YH, Yang T, and Mele J. The oxidative stress theory of aging: embattled or invincible? Insights from non-traditional model organisms. *Age* 30: 99-109, 2008.
42. Bulteau AL, Szweda LI, and Friguet B. Mitochondrial protein oxidation and degradation in response to oxidative stress. *Exp Gerontol* 41: 653-657, 2006.
43. Butow RA, and Avadhani NG. Mitochondrial signaling: the retrograde response. *Mol Cell* 14: 1-15, 2004.
44. Byrne E, Dennet X, and Trounce I Oxidative energy failure in post-mitotic cells: a major factor in senescence. *Rev Neurol* 147: 532-535, 1991.
45. Cabreiro F, and Gems D. Worms need microbes too: microbiota, health and aging in *Caenorhabditis elegans*. *EMBO Mol Med* 5: 1300-1310, 2013.
46. Cadenas E, and Davies KJA. Mitochondrial free radical generation, oxidative stress, and aging. *Free Radic Biol Med* 29: 222-230, 2000.

47. Cai J, and Jones DP. Superoxide in apoptosis. Mitochondrial generation triggered by cytochrome c loss. *J Biol Chem* 273: 11401-11404, 1998.
48. Calabrese EJ, Iavicoli I, and Calabrese V. Hormesis: its impact on medicine and health. *Hum Exp Toxicol* 32: 120-152, 2013.
49. Calabrese V, Scapagnini G, Colombrita C, Ravagna A, Pennini G, Giuffrida Stella AM, Galli F, and Butterfield DA. Redox regulation of heat shock protein expression in aging and neurodegenerative disorders associated with oxidative stress: a nutritional approach. *Amino Acids* 25: 437-444, 2003.
50. Calabrese V, Scapagnini G, Giuffrida Stella AM, Bates TE, and Clark JB. Mitochondrial involvement in brain function and dysfunction: relevance to aging, neurodegenerative disorders and longevity. *Neurochem Res* 26: 739-764, 2001.
51. Calabrese V, Scapagnini G, Ravagna A, Colombrita C, Spadaro F, Butterfield DA, and Giuffrida Stella AM. Increased expression of heat shock proteins in rat brain during aging: relationship with mitochondrial function and glutathione redox state. *Mech Ageing Dev* 125: 325-335, 2004.
52. Calleja M, Peña P, Ugalde C, Ferreiro C, Marco R, and Garesse R. Mitochondrial DNA remains intact during *Drosophila* aging, but the levels of mitochondrial transcripts are significantly reduced. *J Biol Chem* 268: 18891-18897, 1993.
53. Canistro D, Boccia C, Falconi R, Bonamassa B, Valgimigli L, Vivarelli F, Soleti A, Genova ML, Lenaz G, Sapone A, Zaccanti F, Abdel-Rahman SZ, and Paolini M. Redox-based flagging of the global network of oxidative stress greatly promotes longevity. *J Gerontol A Biol Sci Med Sci*, 2014. [Epub ahead of print]
54. Capel F, Buffiere C, Patureau Mirand P, and Mosoni L. Differential variation of mitochondrial H₂O₂ release during aging in oxidative and glycolytic muscles in rats. *Mech Ageing Dev* 125: 367-373, 2004.

55. Carraro, F, Pucci, A, Pellegrini, M, Pelicci, PG, Baldari, CT, and Naldini, A. p66Shc is involved in promoting HIF-1-alpha accumulation and cell death in hypoxic T cells. *J Cell Physiol* 211: 439-447, 2007.
56. Casteilla L, Rigoulet, M, and P énicaud, L. Mitochondrial ROS metabolism: modulation by uncoupling proteins. *IUBMB Life* 52: 181-188, 2001.
57. Chandel N, Budinger GR, Kemp RA, and Schumacker PT. Inhibition of cytochrome-c oxidase activity during prolonged hypoxia. *Am J Physiol* 268: L918-925, 1995.
58. Chandel NS, Maltepe E, Goldwasser E, Mathieu CE, Simon MC, and Schumacker PT. Mitochondrial reactive oxygen species trigger hypoxia-induced transcription. *Proc Natl Acad Sci USA* 95: 11715–11720, 1998.
59. Chandel NS, McClintock DS, Feliciano CE, Wood TM, Melendez JA, Rodriguez AM, and Schumacker PT. Reactive oxygen species generated at mitochondrial complex III stabilize hypoxia-inducible factor-1alpha during hypoxia: a mechanism of O₂ sensing. *J Biol Chem* 275: 25130–25138, 2000.
60. Chaudhary KR, El-Sikhry H, and Seubert JM. Mitochondria and the aging heart. *J Geriatr Cardiol* 8: 159-167, 2011.
61. Chen CF, Tsai SY, Ma MC, and Wu MS. Hypoxic preconditioning enhances renal superoxide dismutase levels in rats. *Physiol* 552(Pt 2): 561-569, 2003.
62. Chen Q, Moghaddas S, Hoppel CL, and Lesnefsky EJ. Ischemic defects in the electron transport chain increase the production of reactive oxygen species from isolated rat heart mitochondria. *Am J Physiol Cell Physiol* 294: C460-466, 2008.
63. Chen Y, and Gibson SB. Is generation of reactive oxygen species a trigger for autophagy? *Autophagy* 4: 246-248, 2008.

64. Chen Y-C, Taylor EB, Dephoure N, Heo J-M, Tonhato A, Papandreou I, Nath N, Denko NC, Gygi SP and Rutter J. Identification of a Protein Mediating Respiratory Supercomplex Stability. *Cell Metab* 15: 348-360, 2012.
65. Chung HY, Kim HJ, Kim KW, Choi JS, and Yu BP. Molecular inflammation hypothesis of aging based on the anti-aging mechanism of calorie restriction. *Micros. Res Techniq* 59: 264–272, 2002.
66. Cooper JM, Mann VM, and Schapira AH. Analyses of mitochondrial respiratory chain function and mitochondrial DNA deletion in human skeletal muscle: effect of ageing. *J Neurol Sci* 113(1): 91-98, 1992.
67. Cooper TM, Mockett RJ, Sohal BH, Sohal RS, and Orr WC. Effect of caloric restriction on life span of the housefly, *Musca domestica*. *FASEB J* 18: 1591-1593, 2004.
68. Copeland JM, Cho J, Lo T, Jr, Hur JH, Bahadorani S, Arabyan T, Rabie J, Soh J, and Walker DW. Extension of *Drosophila* life span by RNAi of the mitochondrial respiratory chain. *Curr Biol* 19: 1591–1598, 2009.
69. Corcoran A, and Cotter TG. Redox regulation of protein kinases. *FEBS J* 280: 1944-1965, 2013.
70. Correia SC, Carvalho C, Cardoso S, Santos RX, Santos MS, Oliveira CR, Perry G, Zhu X, Smith MA, and Moreira PI. Mitochondrial preconditioning: a potential neuroprotective strategy. *Front Aging Neurosci* 2: 138, 2010.
71. Correia SC, Santos RX, Cardoso SM, Santos MS, Oliveira CR, and Moreira PI. Cyanide preconditioning protects brain endothelial and NT2 neuron-like cells against glucotoxicity: role of mitochondrial reactive oxygen species and HIF-1 α . *Neurobiol Dis* 45: 206-218, 2012.

72. Cox AG, Winterbourn CC, and Hampton MB. Mitochondrial peroxiredoxin involvement in antioxidant defence and redox signalling. *Biochem J* 425: 313-325, 2009.
73. Cristina D, Cary M, Lunceford A, Clarke C, and Kenyon C. A regulated response to impaired respiration slows behavioral rates and increases lifespan in *Caenorhabditis elegans*. *PLoS Genet* 5: e1000450, 2009.
74. Crofts AR, Lhee S, Crofts SB, Cheng J, and Rose S. Proton pumping in the bc1 complex: a new gating mechanism that prevents short circuits. *Biochim Biophys Acta* 1757: 1019-1034, 2006.
75. Crofts AR. The cytochrome bc1 complex: function in the context of structure. *Annu Rev Physiol* 66: 689-733, 2004.
76. Cypser JR, Tedesco P, and Johnson TE. Hormesis and aging in *Caenorhabditis elegans*. *Exp Gerontol* 41: 935-939, 2006.
77. Dalle-Donne I, Colombo G, Gagliano N, Colombo R, Giustarini D, Rossi R, and Milzani A. S-glutathiolation in life and death decisions of the cell. *Free Radic Res* 45: 3-15, 2011.
78. Dancy BM, Sedensky MM, and Morgan PG. Analysis of aging in *Caenorhabditis elegans*. *Exp Gerontol* 56: 245-255, 2014.
79. Darzynkiewicz Z, Zhao H, Halicka HD, Li J, Lee YS, Hsieh TC, and Wu JM. In search of antiaging modalities: Evaluation of mTOR- and ROS/DNA damage-signaling by cytometry. *Cytometry A* 85: 386-399, 2014.
80. D'Aurelio M, Gajewski CD, Lenaz G, and Manfredi G. Respiratory chain supercomplexes set the threshold for respiration defects in human mtDNA mutant cybrids. *Hum Mol Genet* 15: 2157-2269, 2006.

81. D'Autréaux B, and Toledano MB. ROS as signalling molecules: mechanisms that generate specificity in ROS homeostasis. *Nat Rev Mol Cell Biol* 8: 813-824, 2007.
82. De Grey AD. The reductive hotspot hypothesis of mammalian aging: membrane metabolism magnifies mutant mitochondrial mischief. *Eur. J. Biochem* 269: 2003-2009, 2002.
83. De Magalhaes JP. From cells to ageing: a review of models and mechanisms of cellular senescence and their impact on human ageing. *Exp Cell Res* 300: 1-10, 2004.
84. Dencher NA, Frenzel M, Reifschneider NH, Sugawa M, and Krause F. Proteome alterations in rat mitochondria caused by aging. *Ann NY Acad Sci* 1100: 291-298, 2007.
85. Di Lisa F, Canton M, Menabò R, Kaludercic N, and Bernardi P. Mitochondria and cardioprotection. *Heart Fail Rev* 12: 249-260, 2007.
86. Diaz F, Enriquez JA, and Moraes CT. Cells lacking Rieske iron-sulfur protein have a reactive oxygen species-associated decrease in respiratory complexes I and IV. *Mol Cell Biol* 32: 415-429, 2012.
87. Diaz F, Fukui H, Garcia S, and Moraes CT. Cytochrome c oxidase is required for the assembly/stability of respiratory complex I in mouse fibroblasts. *Mol Cell Biol* 26: 4872-4881, 2006.
88. Diaz F, Garcia S, Padgett KR, and Moraes CT. A defect in the mitochondrial complex III, but not complex IV, triggers early ROS-dependent damage in defined brain regions. *Hum Mol Genet* 21: 5066-5077, 2012.
89. Dikalov S. Cross talk between mitochondria and NADPH oxidases. *Free Radic Biol Med* 51: 1289-1301, 2011.

90. Dillin A, Hsu AL, Arantes-Oliveira N, Lehrer-Graiwer J, Hsin H, Fraser AG, Kamath RS, Ahringer J, and Kenyon C. Rates of behavior and aging specified by mitochondrial function during development. *Science* 298: 2398–2401, 2002.
91. Dlasková A, Hlavatá L, and Jezek P. Oxidative stress caused by blocking of mitochondrial complex I H(+) pumping as a link in aging/disease vicious cycle. *Int J Biochem Cell Biol* 40: 1792-1805, 2008.
92. Dowhan W, and Bogdanov M. Lipid-dependent membrane protein topogenesis. *Annu Rev Biochem* 78: 515-540, 2009.
93. Drew B, Phaneuf S, Dirks A, Selman C, Gredilla R, Lezza A, Barja G, and Leeuwenburgh C. Effects of aging and caloric restriction on mitochondrial energy production in gastrocnemius muscle and heart. *Am J Physiol Regul Integr Comp Physiol* 284: R474-480, 2003.
94. Dröse S, Brandt U, and Wittig I. Mitochondrial respiratory chain complexes as sources and targets of thiol-based redox-regulation. *Biochim Biophys Acta* 1844: 1344-1354, 2014.
95. Dröse S, and Brandt U. The mechanism of mitochondrial superoxide production by the cytochrome bc1 complex. *J Biol Chem* 283: 21649-21654, 2008.
96. Dudkina NV, Kudryashev M, Stahlberg H, and Boekema EJ. Interaction of complexes I, III, and IV within the bovine respirasome by single particle cryoelectron tomography. *Proc Natl Acad Sci USA* 108: 15196–15200. 2011.
97. Edgar D, Larsson NG, and Trifunovic A. Point mutations are causing progeroid phenotypes in the mtDNA mutator mouse. *Cell Metab* 11(1): 1, 2010.
98. Edgar D, Shabalina I, Camara Y, Wredenberg A, Calvaruso MA, Nijtmans L, Nedergaard J, Cannon B, Larsson NG, and Trifunovic A. Random point mutations

- with major effects on protein-coding genes are the driving force behind premature aging in mtDNA mutator mice. *Cell Metab* 10: 131-138, 2009.
99. Effros RB. From Hayflick to Walford: the role of T cell replicative senescence in human aging. *Exp Gerontol* 39: 885-890, 2004.
 100. Esterházy D, King MS, Yakovlev G, and Hirst J. Production of reactive oxygen species by complex I (NADH:ubiquinone oxidoreductase) from *Escherichia coli* and comparison to the enzyme from mitochondria. *Biochemistry* 47: 3964-3971, 2008.
 101. Fabrizio P, Battistella L, and Vardavas R, Gattazzo C, Liou LL, Diaspro A, Dossen JW, Gralla EB, and Longo VD. Superoxide is a mediator of an altruistic aging program in *Saccharomyces cerevisiae*. *J Cell Biol* 166: 1055-1067, 2004.
 102. Faraonio R, Salerno P, Passaro F, Sedia C, Iaccio A, Bellelli R, Nappi TC, Comegna M, Romano S, Salvatore G, Santoro M, and Cimino F. A set of miRNAs participates in the cellular senescence program in human diploid fibroblasts. *Cell Death Differ* 19: 713–721, 2012.
 103. Farina M, Avila DS, da Rocha JB, and Aschner M. Metals, oxidative stress and neurodegeneration: A focus on iron, manganese and mercury. *Neurochem Int* 62: 575-594, 2013.
 104. Fato R, Bergamini C, Bortolus M, Maniero AL, Leoni S, Ohnishi T, and Lenaz G. Differential effects of Complex I inhibitors on production of reactive oxygen species. *Biochim Biophys Acta* 1787: 384-392, 2009.
 105. Feng J, Bussiere F, and Hekimi S. Mitochondrial electron transport is a key determinant of life span in *Caenorhabditis elegans*. *Dev Cell* 1: 633–644, 2001.
 106. Fernandez-Silva P, Petruzzella V, Fracasso F, Gadaleta MN, and Cantatore P. Reduced synthesis of mtRNA in isolated mitochondria of senescent rat brain. *Biochem Biophys Res Commun* 176: 645-653, 1991.

107. Finkel T. Signal transduction by mitochondrial oxidants. *J Biol Chem* 287: 4434-4440, 2012.
108. Firląg M, Kamaszewski M, Gaca K, and Bałasińska B. Age-related changes in the central nervous system in selected domestic mammals and primates. *Postepy Hig Med Dosw (Online)* 67: 269-275, 2013.
109. Fontana L, Partridge L, and Longo VD. Extending healthy life span-from yeast to humans. *Science* 328: 321-326, 2010.
110. Ford JH. Saturated fatty acid metabolism is key link between cell division, cancer, and senescence in cellular and whole organism aging. *Age (Dordr)* 32: 231-237, 2010.
111. Fourquet S, Huang ME, D'Autreaux B, and Toledano MB. The dual functions of thiol-based peroxidases in H₂O₂ scavenging and signaling. *Antioxid Redox Signal* 10: 1565-1576, 2008.
112. Franceschi C, Capri M, Monti D, Giunta S, Olivieri F, Sevini F, Panourgia MP, Invidia L, Celani L, Scurti M, Cevenini E, Castellani GC, and Salvioli S. Inflammaging and anti-inflammaging: a systemic perspective on aging and longevity emerged from studies in humans. *Mech Ageing Dev* 128: 92-105, 2007.
113. Frank M, Duvezin-Caubet S, Koob S, Occhipinti A, Jagasia R, Petcherski A, Ruonala MO, Priault M, Salin B, and Reichert AS. Mitophagy is triggered by mild oxidative stress in a mitochondrial fission dependent manner. *Biochim Biophys Acta* 1823: 2297-2310, 2012.
114. Frenzel M, Rommelspacherr H, Sugawa MD, and Dencher NA Ageing alters the supramolecular architecture of OxPhos complexes in rat brain cortex. *Exp Gerontol* 45: 563-572, 2010.
115. Fridovich I. Oxygen: how do we stand it? *Med Princ Pract* 22: 131-137, 2013.

116. Fukuda R, Zhang H, Kim JW, Shimoda L, Dang CV, and Semenza GL. HIF-1 regulates cytochrome oxidase subunits to optimize efficiency of respiration in hypoxic cells *Cell* 129: 111–122, 2007.
117. Gadaleta MN, Petruzzella V, Renis M, Fracasso F, and Cantatore P. Reduced transcription of mitochondrial DNA in the senescent rat. Tissue dependence and effect of L-carnitine. *Eur J Biochem* 187: 501-506, 1990.
118. Galkin A, and Brandt U. Superoxide radical formation by pure complex I (NADH:ubiquinone oxidoreductase) from *Yarrowia lipolytica*. *J Biol Chem* 280: 30129-30135, 2005.
119. Gems D, and Doonan R. Antioxidant defense and aging in *C. elegans*: is the oxidative damage theory of aging wrong? *Cell Cycle* 8: 1681-1687, 2009.
120. Gems D, and Partridge L. Genetics of longevity in model organisms: debates and paradigm shifts. *Annu Rev Physiol* 75: 621-644, 2013.
121. Genova ML, and Lenaz G. A critical appraisal of the role of respiratory supercomplexes in mitochondria. *Biol Chem* 394: 631-639, 2013.
122. Genova ML, and Lenaz G. Functional role of mitochondrial respiratory supercomplexes. *Biochim Biophys Acta* 1837: 427-443, 2014.
123. Genova ML, Abd-Elsalam NM, Mahdy el SM, Bernacchia A, Lucarini M, Pedulli GF, and Lenaz G. Redox cycling of adrenaline and adrenochrome catalysed by mitochondrial Complex I. *Arch Biochem Biophys* 447: 167-173, 2006.
124. Genova ML, Baracca A, Biondi A, Casalena G, Faccioli M, Falasca AI, Formiggini G, Sgarbi G, Solaini G, and Lenaz G. Is supercomplex organization of the respiratory chain required for optimal electron transfer activity? *Biochim Biophys Acta* 1777: 740-746, 2008.

125. Genova ML, Ventura B, Giuliano G, Bovina C, Formiggini G, Parenti Castelli G, and Lenaz G. The site of production of superoxide radical in mitochondrial Complex I is not a bound ubiquinone but presumably iron-sulfur cluster N2. *FEBS Lett* 505: 364-368, 2001.
126. Gertz M, Fischer F, Wolters D, and Steegborn C. Activation of the lifespan regulator p66Shc through reversible disulfide bond formation. *Proc Natl Acad Sci USA* 105: 5705-5709, 2008.
127. Gertz M, and Steegborn C. The Lifespan-regulator p66Shc in mitochondria: redox enzyme or redox sensor? *Antioxid Redox Signal* 13: 1417-1428, 2010.
128. Ghelli A, Tropeano CV, Calvaruso MA, Marchesini A, Iommarini L, Porcelli AM, Zanna C, De Nardo V, Martinuzzi A, Wibrand F, Vissing J, Kurelac I, Gasparre G, Selamoglu N, Daldal F, and Rugolo M. The cytochrome b p.278Y>C mutation causative of a multisystem disorder enhances superoxide production and alters supramolecular interactions of respiratory chain complexes. *Hum Mol Genet* 22: 2141-2151, 2013.
129. Ghio AJ, Carraway MS, and Madden MC. Composition of air pollution particles and oxidative stress in cells, tissues, and living systems. *J Toxicol Environ Health B Crit Rev* 15: 1-21, 2012.
130. Giorgio M, Migliaccio E, Orsini F, Paolucci D, Moroni M, Contursi C, Pelliccia G, Luzi L, Minacci S, Marcaccio M, Pinton P, Rizzuto R, Bernardi P, Paolucci F, and Pelicci PG. Electron transfer between cytochrome c and p66Shc generates reactive oxygen species that trigger mitochondrial apoptosis. *Cell* 122: 221-233, 2005.
131. Gladyshev VN. The free radical theory of aging is dead. Long live the damage theory! *Antioxid Redox Signal* 20: 727-731, 2014.

132. Godon C, Lagniel G, and Lee J, Buhler JM, Kieffer S, Perrot M, Boucherie H, Toledano MB, and Labarre J. The H₂O₂ stimulon in *Saccharomyces cerevisiae*. *J Biol Chem* 273: 22480–22489, 1998.
133. Gomes AP, Price NL, Ling AJ, Moslehi JJ, Montgomery MK, Rajman L, White JP, Teodoro JS, Wrann CD, Hubbard BP, Mercken EM, Palmeira CM, de Cabo R, Rolo AP, Turner N, Bell EL, and Sinclair DA. Declining NAD(+) induces a pseudohypoxic state disrupting nuclear-mitochondrial communication during aging. *Cell* 155: 1624-1638, 2013.
134. Gomes LC, Di Benedetto G, and Scorrano L. During autophagy mitochondria elongate, are spared from degradation and sustain cell viability. *Nat Cell Biol* 13: 589-598, 2011.
135. Gómez LA, and Hagen TM. Age-related decline in mitochondrial bioenergetics: does supercomplex destabilization determine lower oxidative capacity and higher superoxide production? *Semin Cell Dev Biol* 23: 758-767, 2012.
136. Gómez LA, Monette JS, Chavez JD, Maier CS, and Hagen TM. Supercomplexes of the mitochondrial electron transport chain decline in the aging rat heart. *Arch Biochem Biophys* 490: 30-35, 2009.
137. Gonzalez F, D'Aurelio M, Boutant M, Moustapha A, Puech JP, Landes T, Arnauné-Pelloquin L, Vial G, Taleux N, Slomianny C, Wanders RJ, Houtkooper RH, Bellenguer P, Møller IM, Gottlieb E, Vaz FM, Manfredi G, and Petit PX. Barth syndrome: cellular compensation of mitochondrial dysfunction and apoptosis inhibition due to changes in cardiolipin remodeling linked to tafazzin (TAZ) gene mutation. *Biochim Biophys Acta* 1832: 1194-1206, 2013.
138. Gough DR, and Cotter TG. Hydrogen peroxide: a Jekyll and Hyde signalling molecule. *Cell Death Dis* 2: e213, 2011.

139. Greco M, Villani G, Mazzucchelli F, Bresolin N, Papa S, and Attardi G. Marked aging-related decline in efficiency of oxidative phosphorylation in human skin fibroblasts. *FASEB J* 17: 1706-1708, 2004.
140. Gredilla R, López-Torres M, and Barja G. Effect of time of restriction on the decrease in mitochondrial H₂O₂ production and oxidative DNA damage in the heart of food-restricted rats. *Microsc Res Tech* 59: 273-277, 2002.
141. Grishko V, Solomon M, Breit JF, Killilea DW, Ledoux SP, Wilson GL, and Gillespie MN. Hypoxia promotes oxidative base modifications in the pulmonary artery endothelial cell VEGF gene. *FASEB J* 15: 1267–1269, 2001.
142. Grivennikova VG, and Vinogradov AD. Partitioning of superoxide and hydrogen peroxide production by mitochondrial respiratory complex I. *Biochim Biophys Acta* 1827: 446-454, 2013.
143. Guarente L. Sirtuins and calorie restriction. *Nat Rev Mol Cell Biol* 13: 207, 2012.
144. Guzy RD, Hoyos B, Robin E, Chen H, Liu L, Mansfield KD, Simon MC, Hammerling U, and Schumacker PT. Mitochondrial complex III is required for hypoxia-induced ROS production and cellular oxygen sensing. *Cell Metab* 1: 401–408, 2005.
145. Guzy RD, and Schumacker PT. Oxygen sensing by mitochondria at Complex III: the paradox of increased reactive oxygen species during hypoxia. *Exp Physiol* 91: 807-819, 2006.
146. Hackenbrock CR, Chazotte B, and Gupte SS. The random collision model and a critical assessment of diffusion and collision in mitochondrial electron transport. *J Bioenerg Biomembr* 18: 331-368, 1986.
147. Haddad JJ. Antioxidant and prooxidant mechanisms in the regulation of redox(y)-sensitive transcription factors. *Cell Signal* 14: 879-897, 2002.

148. Hall DM, Sattler GL, Zhang HJ, Oberley LW, Pitot HC, Kregel KC. Aging lowers steady-state antioxidant enzyme and stress protein expression in primary hepatocytes. *J Gerontol A Biol Sci Med Sci* 56: B259-267, 2001.
149. Halliwell B. Oxidative stress and cancer: have we moved forward? *Biochem J* 401: 1-11, 2007.
150. Halliwell B, and Gutteridge J. Free Radicals in Biology and Medicine, 4th Ed, Oxford UK, Oxford University Press, 2007
151. Halliwell B, and Whiteman M. Measuring reactive species and oxidative damage in vivo and in cell culture: how should you do it and what do the results mean? *British Journal of Pharmacology* 142: 231-255, 2004.
152. Hamanaka RB, and Chandel NS. Mitochondrial reactive oxygen species regulate cellular signaling and dictate biological outcomes. *Trends Biochem Sci* 35: 505-513, 2010.
153. Han D, Antunes F, Canali R, Rettori D, and Cadenas E. Voltage-dependent anion channels control the release of the superoxide anion from mitochondria to cytosol. *J Biol Chem* 278: 5557-5563, 2003.
154. Handy DE, and Loscalzo J. Redox regulation of mitochondrial function. *Antioxid Redox Signal* 16: 1323-1367, 2012.
155. Hansford RG, Hogue BA, and Mildaziene V. Dependence of H₂O₂ formation by rat heart mitochondria on substrate availability and donor age. *J Bioenerg Biomembr* 29: 89-95, 1997.
156. Harman DJ. Aging: a theory based on free radical and radiation chemistry. *Gerontol* 11: 298-300, 1956.
157. Harris N, Bachler M, Costa V, Mollapour M, Moradas-Ferreira P, and Piper PW. Overexpressed Sod1p acts either to reduce or to increase the lifespans and stress

- resistance of yeast, depending on whether it is Cu(2+)-deficient or an active Cu, Zn-superoxide dismutase. *Aging Cell* 4: 41–52, 2005.
158. Hauser DN, Dillman AA, Ding J, Li Y, and Cookson MR. Post-translational decrease in respiratory chain proteins in the polg mutator mouse brain. *PLoS One* 9(4): e94646, 2014.
 159. Hayakawa M, Hattori K, Sugiyama S, and Ozawa T. Age-associated oxygen damage and mutations in mitochondrial DNA in human hearts. *Biochem Biophys Res Commun* 189: 979-985, 1992.
 160. Hayflick L. Living forever and dying in the attempt. *Exp Gerontol* 38: 1231-1241, 2003
 161. Haynes CM, and Ron D. The mitochondrial UPR - protecting organelle protein homeostasis. *J Cell Sci* 123(Pt 22): 3849-3855, 2010.
 162. Head E. A canine model of human aging and Alzheimer's disease. *Biochim Biophys Acta* 1832: 1384-1389, 2013.
 163. Helbock HJ, Beckman KB, and Ames BN. 8-Hydroxydeoxyguanosine and 8-hydroxyguanine as biomarkers of oxidative DNA damage. *Methods Enzymol* 300: 156-166, 1999.
 164. Herrero A, and Barja G. Localization of the site of oxygen radical generation inside the Complex I of heart and nonsynaptic brain mammalian mitochondria. *J Bioenerg Biomembr* 32: 609-616, 2000.
 165. Hess KC, Liu J, Manfredi G, Mühlischlegel FA, Buck J, Levin LR, and Barrientos A. A mitochondrial CO₂-adenylyl cyclase-cAMP signalosome controls yeast normoxic cytochrome c oxidase activity. *FASEB J* 28: 4369-4380, 2014.
 166. Hiona A, Sanz A, Kujoth GC, Pamplona R, Seo AY, Hofer T, Someya S, Miyakawa T, Nakayama C, Samhan-Arias AK, Servais S, Barger JL, Portero-Otín M, Tanokura M, Prolla TA, and Leeuwenburgh C. Mitochondrial DNA mutations induce

- mitochondrial dysfunction, apoptosis and sarcopenia in skeletal muscle of mitochondrial DNA mutator mice. *PLoS One* 5(7): e11468, 2010.
167. Hoffman DL, and Brookes PS. Oxygen sensitivity of mitochondrial reactive oxygen species generation depends on metabolic conditions. *J Biol Chem* 284: 16236-16245, 2009.
 168. Horan MP. Application of serial analysis of gene expression to the study of human genetic disease. *Hum Genet* 126: 605-614, 2009.
 169. Horvath SE, and Daum G. Lipids of mitochondria. *Prog Lipid Res* 52: 590-614, 2013.
 170. Hosokawa M. A higher oxidative status accelerates senescence and aggravates age-dependent disorders in SAMP strains of mice. *Mech Ageing Dev* 123: 1553-1561, 2002.
 171. Houtkooper RH, Canto C, Wanders RJ, and Auwerx J. The secret life of NAD⁺: an old metabolite controlling new metabolic signaling pathways. *Endocr Rev* 31: 194-223, 2010.
 172. Hruszkewycz AM. Lipid peroxidation and mtDNA degeneration. A hypothesis. *Mutat Res* 275: 243-248, 1992.
 173. Hunte C, and Richers S. Lipids and membrane protein structures. *Curr Opin Struct Biol* 18: 406-411, 2008.
 174. Hutter E, Unterluggauer H, Uberall F, Schramek H, and Jansen-Durr P. Replicative senescence of human fibroblasts: the role of Ras-dependent signaling and oxidative stress. *Exp Gerontol* 37: 1165-1174, 2002.
 175. Ikeyama N, Kokkonen G, Martindale JL, Wang XT, Gorospe M, and Holbrook NJ. Effects of aging and calorie restriction of Fischer 344 rats on hepatocellular response to proliferative signals. *Exp Gerontol* 38: 431-439, 2003.

176. Ishii N, Fujii, M, Hartman, PS, Tsuda, M, Yasuda, K, Senoo-Matsuda, N, Yanase, S, Ayusawa, D, and Suzuki, K. A mutation in succinate dehydrogenase cytochrome b causes oxidative stress and ageing in nematodes. *Nature* 394: 694-697, 1998.
177. Ishii T, Miyazawa M, Hartman PS, and Ishii N. Mitochondrial superoxide anion (O(2)(-)) inducible "mev-1" animal models for aging research. *BMB Rep* 44: 298-305, 2011.
178. Ishii T, Miyazawa M, Onouchi H, Yasuda K, Hartman PS, and Ishii N. Model animals for the study of oxidative stress from complex II. *Biochim Biophys Acta* 1827: 588-597, 2013.
179. Ito K. Frontiers of model animals for neuroscience: two prosperous aging model animals for promoting neuroscience research. *Exp Anim* 62: 275-280, 2013.
180. Itoh K, Wakabayashi N, Katoh Y, Ishii T, Igarashi K, Engel JD, and Yamamoto M. Keap1 represses nuclear activation of antioxidant responsive elements by Nrf2 through binding to the amino-terminal Neh2 domain. *Genes Dev* 13: 76-86, 1999.
181. Itsara LS, Kennedy SR, Fox EJ, Yu S, Hewitt JJ, Sanchez-Contreras M, Cardozo-Pelaez F, and Pallanck LJ. Oxidative stress is not a major contributor to somatic mitochondrial DNA mutations. *PLoS Genet* 10(2): e1003974, 2014.
182. Janssen-Heininger YM, Mossman BT, Heintz NH, Forman HJ, Kalyanaraman B, Finkel T, Stamler JS, Rhee SG, and van der Vliet A. Redox-based regulation of signal transduction: principles, pitfalls, and promises. *Free Radic Biol Med* 45: 1-17, 2008.
183. Jazwinski SM. The retrograde response: when mitochondrial quality control is not enough. *Biochim Biophys Acta* 1833: 400-409, 2013.
184. Jezek P, and Hlavata L. Mitochondria in homeostasis of reactive oxygen species in cell, tissues, and organism. *Int. J. Biochem. Cell Biol* 37: 2478-2503, 2005.

185. Johnson-Cadwell LI, Jekabsons MB, Wang A, Polster BM, and Nicholls DG. 'Mild Uncoupling' does not decrease mitochondrial superoxide levels in cultured cerebellar granule neurons but decreases spare respiratory capacity and increases toxicity to glutamate and oxidative stress. *J Neurochem* 101: 1619-1631, 2007.
186. Jones DP. Redox potential of GSH/GSSG couple: assay and biological significance. *Methods Enzymol* 348: 93-112, 2002.
187. Jones SJM, Riddle DL, Pouzyrev AT, Velculescu VE, Hillier LD, Eddy SR, Stricklin SL, Baillie DL, Waterston R, and Marra MA. Changes in gene expression associated with developmental arrest and longevity in *Caenorhabditis elegans*. *Genome Res* 11: 1346-1352, 2001.
188. Judge S, Jang YM, Smith A, Hagen T, Leeuwenburgh C. Age-associated increases in oxidative stress and antioxidant enzyme activities in cardiac interfibrillar mitochondria: implications for the mitochondrial theory of aging. *FASEB J* 19: 419-421, 2005.
189. Kadenbach B, Ramzan R, and Vogt S. High efficiency versus maximal performance-the cause of oxidative stress in eukaryotes: a hypothesis. *Mitochondrion* 13: 1-6, 2013.
190. Kalyanaraman B, Darley-Usmar V, Davies KJA, Dennery PA, Forman HJ, Grisham MB, Mann GE, Moore K, Roberts LJ, and Ischiropoulos H. Measuring reactive oxygen and nitrogen species with fluorescent probes: challenges and limitations. *Free Radic Biol Med* 52: 1-6, 2012.
191. Kaluz S, Kaluzova M, and Stanbridge EJ. Rational design of minimal hypoxia-inducible enhancers. *Biochem Biophys Res Commun* 370: 613-618, 2008.
192. Kareyeva AV, Grivennikova VG, Cecchini G, and Vinogradov AD. Molecular identification of the enzyme responsible for the mitochondrial NADH-supported

- ammonium-dependent hydrogen peroxide production. *FEBS Lett* 585: 385-389, 2011.
193. Kareyeva AV, Grivennikova VG, and Vinogradov AD. Mitochondrial hydrogen peroxide production as determined by the pyridine nucleotide pool and its redox state. *Biochim Biophys Acta* 1817: 1879-1885, 2012.
 194. Kazachkova N, Ramos A, Santos C, and Lima M. Mitochondrial DNA damage patterns and aging: revising the evidences for humans and mice. *Aging Dis* 4: 337-350, 2013.
 195. Kenyon CJ. The genetics of ageing. *Nature* 464(7288): 504-512, 2010.
 196. Kietzmann T, and Gorlach, A. Reactive oxygen species in the control of hypoxia-inducible factor-mediated gene expression. *Seminars Cell Dev Biol*, 16: 474-486, 2005.
 197. Kim HJ, Jung KJ, Yu BP, Cho CG, Choi JS, and Chung HY. Modulation of redox-sensitive transcription factors by calorie restriction during aging. *Mech Ageing Dev* 123: 1589-1595, 2002.
 198. Kim JJ, Lee SB, Park JK, and Yoo YD. TNF- α -induced ROS production triggering apoptosis is directly linked to Romo1 and Bcl-X(L). *Cell Death Differ* 17: 1420-1434, 2010.
 199. Kim JW, Tchernyshyov I, Semenza GL, and Dang CV. HIF-1-mediated expression of pyruvate dehydrogenase kinase: a metabolic switch required for cellular adaptation to hypoxia. *Cell Metab* 3: 177-185, 2006.
 200. Kishi S. Using zebrafish models to explore genetic and epigenetic impacts on evolutionary developmental origins of aging. *Transl Res* 163: 123-135, 2014.

201. Korde AS, Yadav VR, Zheng YM, and Wang YX. Primary role of mitochondrial Rieske iron-sulfur protein in hypoxic ROS production in pulmonary artery myocytes. *Free Radic Biol Med* 50: 945-952, 2011.
202. Korshunov SS, Skulachev VP, and Starkov AA. High protonic potential actuates a mechanism of production of reactive oxygen species in mitochondria. *FEBS Lett* 416: 15-18, 1997.
203. Kovacic P, and Cooksy AL. Unifying mechanism for the toxicity and addiction by abused drugs: electron transfer and reactive oxygen species. *Med Hypoth* 64: 357-366, 2005.
204. Kuang J, and Ebert PR. The failure to extend lifespan via disruption of complex II is linked to preservation of dynamic control of energy metabolism. *Mitochondrion* 12: 280–287, 2012.
205. Kushnareva Y, Murphy AN, and Andreyev A. Complex I-mediated reactive oxygen species generation: modulation by cytochrome c and NAD(P)+ oxidation-reduction state. *Biochem J* 368: 545-553, 2002.
206. Kussmaul L, and Hirst. The mechanism of superoxide production by NADH:ubiquinone oxidoreductase (complex I) from bovine heart mitochondria. *J Proc Natl Acad Sci USA* 103: 7607-7612, 2006.
207. Kwong LK, and Sohal RS. Substrate and site specificity of hydrogen peroxide generation in mouse mitochondria. *Arch Biochem Biophys* 350: 118-126, 1998.
208. Lambert AJ, and Brand MD. Inhibitors of the quinone-binding site allow rapid superoxide production from mitochondrial NADH:ubiquinone oxidoreductase (complex I). *J Biol Chem* 279: 39414-39420, 2004.
209. Lambert AJ, Buckingham JA, Boysen HM, and Brand MD. Diphenyleneiodonium acutely inhibits reactive oxygen species production by mitochondrial complex I

- during reverse, but not forward electron transport. *Biochim Biophys Acta* 1777: 397-403, 2008.
210. Lange C, Nett JH, Trumpower BL, and Hunte C. Specific roles of protein-phospholipid interactions in the yeast cytochrome bc₁ complex structure. *EMBO J* 20: 6591-6600, 2001
 211. Lapointe J, and Hekimi S. When a theory of aging ages badly. *Cell Mol Life Sci* 67: 1-8, 2010.
 212. Lauri A, Pompilio G, and Capogrossi MC. The mitochondrial genome in aging and senescence. *Ageing Res Rev* 18C:1-15, 2014.
 213. Lavrovsky Y, Chatterjee B, Clark RA, and Roy AK. Role of redox-regulated transcription factors in inflammation, aging and age-related diseases. *Exp Gerontol* 35: 521-532, 2000.
 214. Lee J, Giordano S, and Zhang J. Autophagy, mitochondria and oxidative stress: cross-talk and redox signalling. *Biochem J* 441: 523–540, 2012.
 215. Lee SB, Kim JJ, Kim TW, Kim BS, Lee MS, and Yoo YD. Serum deprivation-induced reactive oxygen species production is mediated by Romo1. *Apoptosis* 15: 204-218, 2010.
 216. Lee YH, Lee JC, Moon HJ, Jung JE, Sharma M, Park BH, Yi HK, Jhee EC. Differential effect of oxidative stress on the apoptosis of early and late passage human diploid fibroblasts: implication of heat shock protein 60. *Cell Biochem Funct* 26: 502-508, 2008.
 217. Lemarie A, Huc L, Pazarentzos E, Mahul-Mellier AL, and Grimm S. Specific disintegration of complex II succinate:ubiquinone oxidoreductase links pH changes to oxidative stress for apoptosis induction. *Cell Death Differ* 18: 338-349, 2011.

218. Lemieux H, and Warren BE. An animal model to study human muscular diseases involving mitochondrial oxidative phosphorylation. *J Bioenerg Biomembr* 44: 503-512, 2012.
219. Lenaz G. Lipid fluidity and membrane protein dynamics. *Biosci Rep* 7: 823-837, 1987.
220. Lenaz G. Role of mitochondria in oxidative stress and ageing. *Biochim Biophys Acta* 1366: 53-67, 1998.
221. Lenaz G. Mitochondrial and reactive oxygen species. Which role in physiology and pathology? *Adv Exp Med Biol* 942: 93–136, 2012.
222. Lenaz G. Role of mitochondria in the generation of reactive oxygen species. In: *Handbook on Reactive Oxygen Species (ROS)*, edited by Suzuki M, and Yamamoto S. New York: Nova Biomedical, 2014, pp 1-108.
223. Lenaz G, and Genova ML. Kinetics of integrated electron transfer in the mitochondrial respiratory chain: random collisions vs. solid state electron channeling. *Am J Physiol Cell Physiol* 292: C1221-1239, 2007.
224. Lenaz G, and Genova ML. Structure and organization of mitochondrial respiratory complexes: a new understanding of an old subject. *Antioxid Redox Signal* 12: 961-1008, 2010.
225. Lenaz G, Baracca A, Barbero G, Bergamini C, DalmonTE ME, del Sole M, Faccioli M, Falasca A, Fato R, Genova ML, Sgarbi G, and Solaini G. Mitochondrial respiratory chain super-complex I-III in physiology and pathology. *Biochim Biophys Acta* 1797: 633-640, 2010.
226. Lenaz G, Bovina C, D'Aurelio M, Fato R, Formiggini G, Genova ML, Giuliano G, Merlo Pich M, Paolucci U, Parenti Castelli G, and Ventura B. Role of mitochondria in oxidative stress and aging. *Ann NY Acad Sci USA* 959, 199-213, 2002.

227. Lenaz G, D'Aurelio M, Merlo Pich M, Genova ML, Ventura B, Bovina C, Formaggini G, and Parenti Castelli G. Mitochondrial bioenergetics in aging. *Biochim Biophys Acta* 1459: 397-404, 2000.
228. Lenaz G, Fato, R, Genova, ML, Bergamini, C, Bianchi, C and Biondi, A. Mitochondrial Complex I: structural and functional aspects. *Biochim Biophys Acta* 1757: 1406-1420, 2006.
229. Lenaz G, and Strocchi P. Reactive oxygen species in the induction of toxicity. In: *General and Applied Toxicology*, edited by Ballantyne B, Marrs T, and Syversen T. Chichester, UK: Wiley, 2009, vol. 1, chapter 15.
230. Leontieva OV, and Blagosklonny MV. M(o)TOR of pseudo-hypoxic state in aging: rapamycin to the rescue. *Cell Cycle* 13: 509-515, 2014.
231. Lerner C, Bitto A, Pulliam D, Nacarelli T, Konigsberg M, Van Remmen H, Torres C, and Sell C. Reduced mammalian target of rapamycin activity facilitates mitochondrial retrograde signaling and increases life span in normal human fibroblasts. *Aging Cell* 12: 966-77, 2013.
232. Levine B, and Klionsky DJ. Development by self-digestion: molecular mechanisms and biological functions of autophagy. *Dev Cell* 6: 463-477, 2004.
233. Li J, and Holbrook NJ. Common mechanisms for declines in oxidative stress tolerance and proliferation with aging. *Free Radic Biol Med* 35: 292-299, 2003.
234. Li, N., Bates, D.J., An, J., Terry, D.A., Wang, E., 2011. Up-regulation of key microRNAs, and inverse down-regulation of their predicted oxidative phosphorylation target genes, during aging in mouse brain. *Neurobiol. Aging* 32: 944-955, 2011.
235. Lillig CH, and Holmgren A. Thioredoxin and related molecules-from biology to health and disease. *Antioxid Redox Signal* 9: 25-47, 2007.

236. Lim HY, Bodmer R, and Perrin L. Dietary restriction in *Drosophila*. *Exp Gerontol* 41: 1213-1216, 2006.
237. Lin SJ, Ford E, Haigis M, Liszt G, and Guarente L. Calorie restriction extends yeast life span by lowering the level of NADH. *Genes Dev* 18: 12–16, 2004.
238. Linford NJ, and Pletcher SD. Mitochondria and ageing in *Drosophila*. *Curr Biol* 19: R895-898, 2009.
239. Liochev SI. Reflections on the theories of aging, of oxidative stress and of science in general. Is it time to abandon the free radical (oxidative stress) theory of aging? *Antioxid Redox Signal* 2014 Aug 12. [Epub ahead of print]
240. Liochev SI and Fridovich I. Superoxide and iron: partners in crime. *IUBMB Life* 48: 157-161, 1999.
241. Liu GH, Ding Z, and Izpisua Belmonte JC. PSC technology to study human aging and aging-related disorders. *Curr Opin Cell Biol* 24: 765-774, 2012.
242. Logan A, Shabalina IG, Prime TA, Rogatti S, Kalinovich AV, Hartley RC, Budd RC, Cannon B, and Murphy MP. In vivo levels of mitochondrial hydrogen peroxide increase with age in mtDNA mutator mice. *Aging Cell* 13: 765-768, 2014.
243. Lombardi A, Silvestri E, Cioffi F, Senese R, Lanni A, Goglia F, de Lange P, and Moreno M. Defining the transcriptomic and proteomic profiles of rat ageing skeletal muscle by the use of a cDNA array, 2D- and Blue native-PAGE approach. *J Proteomics* 72: 708-721, 2009.
244. López-Torres M, Gredilla R, Sanz A, and Barja G. Influence of aging and long-term caloric restriction on oxygen radical generation and oxidative DNA damage in rat liver mitochondria. *Free Radic Biol Med* 32: 882-889, 2002.

245. Lorin S, Dufour E, and Sainsard-Chanet A. Mitochondrial metabolism and aging in the filamentous fungus *Podospora anserina*. *Biochim Biophys Acta* 1757: 604-610, 2006.
246. Luce K, Weil AC, and Osiewacz HD. Mitochondrial protein quality control systems in aging and disease. *Adv Exp Med Biol* 694: 108-125, 2010.
247. Ludovico P, and Burhans WC. Reactive oxygen species, ageing and the hormesis police. *FEMS Yeast Res* 14: 33-39, 2014.
248. Maj MC, Raha S, Myint T, and Robinson BH. Regulation of NADH/CoQ oxidoreductase: do phosphorylation events affect activity? *Protein J* 23: 25-32, 2004.
249. Mandavilli BS, Santos JH, and Van Houten B. Mitochondrial DNA repair and aging. *Mut Res* 509: 127-151, 2002.
250. Mansfield D, Guzy RD, Pan Y, Young RM, Cash TP, Schumacker PT, and Simon MC. Mitochondrial dysfunction resulting from loss of cytochrome c impairs cellular oxygen sensing and hypoxic HIF- α activation. *Cell Metab* 1: 393-399, 2005.
251. Mansouri A, Muller FL, Liu Y, Ng R, Faulkner J, Hamilton M, Richardson A, Huang TT, Epstein CJ, and Van Remmen H. Alterations in mitochondrial function, hydrogen peroxide release and oxidative damage in mouse hind-limb skeletal muscle during aging. *Mech Ageing Dev* 127: 298-306; 2006.
252. Maranzana E, Barbero G, Falasca AI, Lenaz G, and Genova ML. Mitochondrial respiratory supercomplex association limits production of reactive oxygen species from Complex I. *Antioxid Redox Signal* 19: 1469-1480, 2013.
253. Marcus DL, Ibrahim NG, and Freedman ML. Age-related decline in the biosynthesis of mitochondrial inner membrane proteins. *Exp Gerontol* 17: 333-341, 1982.

254. Mar íM, Morales A, Colell A, Garc ía-Ruiz C, and Fern ández-Checa JC. Mitochondrial glutathione, a key survival antioxidant. *Antioxid Redox Signal* 11: 2685-2700, 2009.
255. Marino SM, and Gladyshev VN. Cysteine function governs its conservation and degeneration and restricts its utilization on protein surfaces. *J Mol Biol* 404: 902-916, 2010.
256. Maurel A, Hernandez, C, Kunduzova, O, Bompert, G, Cambon, C, Parini A, and Frances B. Age-dependent increase in hydrogen peroxide production by cardiac monoamine oxidase A in rats. *Am J Physiol Heart and Circulation Physiol* 284: H1460-H1467, 2003.
257. Mazzoni C, Giannattasio S, Winderickx J, and Ludovico P. Yeast stress, aging, and death. *Oxid Med Cell Longev* 2013: 684395, 2013.
258. McCormick MA, Tsai SY, Kennedy BK. TOR and ageing: a complex pathway for a complex process. *Philos Trans R Soc Lond B Biol Sci* 366(1561): 17-27, 2011.
259. McKenzie M, Lazarou M, Thorburn DR, and Ryan MT. Mitochondrial respiratory chain supercomplexes are destabilized in Barth Syndrome patients. *J Mol Biol* 361: 462-469, 2006.
260. Melov S, Coskun, P, Patel, M, Tuinstra, M, Cottrell, B, Jun, AS, Zastawny, TH, Dizdaroglu, M, Goodman, SI, Huang, TT, Miziorko, H, Epstein, CJ, and Wallace, DC. Mitochondrial disease in superoxide dismutase 2 mutant mice. *Proc Natl Acad Sci USA* 96: 846-851, 1999.
261. Menini S, Amadio L, Oddi G, Ricci C, Pesce C, Pugliese F, Giorgio M, Migliaccio E, Pelicci PG, Iacobini C, and Pugliese G. Deletion of p66Shc longevity gene protects against experimental diabetic glomerulopathy by preventing diabetes-induced oxidative stress. *Diabetes* 55: 1642-1650, 2006.

262. Merlo Pich M, Raule N, Catani L, Fagioli ME, Faenza I, Cocco L, and Lenaz G. Increased transcription of mitochondrial genes for Complex I in human platelets during ageing. *FEBS Lett* 558: 19-22, 2004.
263. Merlo Pich M, Bovina C, Formiggini G, Cometti GG, Ghelli A, Parenti Castelli G; Genova ML; Marchetti M; Semeraro S; and Lenaz G. Inhibitor sensitivity of respiratory complex I in human platelets: a possible biomarker of ageing. *FEBS Lett* 380: 176-178, 1996.
264. Mesquita A, Weinberger M, Silva A, Sampaio-Marques B, Almeida B, Leão C, Costa V, Rodrigues F, Burhans WC, and Ludovico P. Caloric restriction or catalase inactivation extends yeast chronological lifespan by inducing H₂O₂ and superoxide dismutase activity. *Proc Natl Acad Sci USA* 107: 15123-15128, 2010.
265. Migliaccio E, Giorgio M, Mele S, Pelicci G, Reboldi P, Pandolfi PP, Lanfranccone L, and Pelicci PG. The p66shc adaptor protein controls oxidative stress response and life span in mammals. *Nature* 402: 309-313, 1999.
266. Migliaccio E, Mele S, Salcini AE, Pelicci G, Lai KM, Superti-Furga G, Pawson T, Di Fiore PP, Lanfranccone L, and Pelicci PG. Opposite effects of the p52shc/p46shc and p66shc splicing isoforms on the EGF receptor-MAP kinase-fos signalling pathway. *EMBO J* 16: 706-716, 1997.
267. Mileykovskaya E, and Dowhan W. Cardiolipin-dependent formation of mitochondrial respiratory supercomplexes. *Chem Phys Lipids* 179: 42-48, 2014.
268. Miquel J, Economos AC, Fleming J, and Johnson JE Jr. Mitochondrial role in cell aging. *Exp Gerontol* 15: 575-591, 1980.
269. Mirò O, Casademont J, Casals E, Perea M, Urbano-Marquez A, Rustin P, and Cardellach F. Aging is associated with increased lipid peroxidation in human

- hearts, but not with mitochondrial respiratory chain enzyme defects. *Cardiovasc Res* 47: 624-663, 2000.
270. Miskin R, Masos T, Yahav S, Shinder D, and Globerson A. AlphaMUPA mice: a transgenic model for increased life span. *Neurobiol. Aging* 20: 555-564, 1999.
 271. Mittal M, Gu XQ, Pak O, Pamenter ME, Haag D, Fuchs DB, Schermuly RT, Ghofrani HA, Brandes RP, Seeger W, Grimminger F, Haddad GG, and Weissmann N. Hypoxia induces Kv channel current inhibition by increased NADPH oxidase-derived reactive oxygen species. *Free Radic Biol Med* 52: 1033-1042, 2012.
 272. Mizushima N, Levine B, Cuervo AM, and Klionsky DJ. Autophagy fights disease through cellular self-digestion. *Nature* 451(7182): 1069-1075, 2008.
 273. Moreno-Sánchez R, Hernández-Esquivel L, Rivero-Segura NA, Marín-Hernández A, Neuzil J, Ralph SJ, and Rodríguez-Enríquez S. Reactive oxygen species are generated by the respiratory complex II-evidence for lack of contribution of the reverse electron flow in complex I. *FEBS J* 280: 927-938, 2013.
 274. MorréDM, Lenaz G, and MorréDJ. Surface oxidase and oxidative stress propagation in aging. *J Exp Biol* 203: 1513-1521, 2000.
 275. Moser CC, Page CC, and Dutton LP. Tunneling in PSII. *Photochem Photobiol Sci* 4: 933-939, 2005
 276. Muller, FL, Roberts, AG, Bowman, NK Kramer, DM. Architecture of the Qo site of the cytochrome bc1 complex probed by superoxide production. *Biochemistry* 42: 6493-6499, 2003.
 277. Müller-Höcker J, Schneiderbanger K, Stefani FH, and Kadenbach B. Progressive loss of cytochrome c oxidase in the human extraocular muscles in ageing-a cytochemical-immunohistochemical study. *Mutat Res* 275: 115-124, 1992.

278. Müller-Höcker J, Seibel P, Schneiderbanger K, and Kadenbach B. Different in situ hybridization patterns of mitochondrial DNA in cytochrome c oxidase-deficient extraocular muscle fibres in the elderly. *Virchows Arch A Pathol Anat Histopathol* 422: 7-15, 1993.
279. Müller-Höcker J. Cytochrome c oxidase deficient fibres in the limb muscle and diaphragm of man without muscular disease: an age-related alteration. *J Neurol Sci* 100: 14-21, 1990.
280. Müller-Höcker J. Cytochrome-c-oxidase deficient cardiomyocytes in the human heart- an age-related phenomenon. A histochemical ultracytochemical study. *Am J Pathol* 134: 1167-1173, 1989.
281. Munro D, and Blier PU. The extreme longevity of *Arctica islandica* is associated with increased peroxidation resistance in mitochondrial membranes. *Aging Cell* 11: 845-855, 2012.
282. Murphy MP. How mitochondria produce reactive oxygen species. *Biochem J* 417: 1-13, 2009.
283. Murphy MP. Mitochondrial thiols in antioxidant protection and redox signaling: distinct roles for glutathionylation and other thiol modifications. *Antioxid Redox Signal* 16: 476-495, 2012.
284. Murry CE, Jennings RB, and Reimer KA. Preconditioning with ischemia: a delay of lethal cell injury in ischemic myocardium. *Circulation* 74: 1124-1136, 1986.
285. Napoli C, Martin-Padura I, De Nigris F, Giorgio M, Mansueto G, Somma P, Condorelli M, Sica G, De Rosa G, and Pelicci PG. Deletion of the p66Shc longevity gene reduces systemic and tissue oxidative stress, vascular cell apoptosis, and early atherogenesis in mice fed a high-fat diet. *Proc Natl Acad Sci USA* 100: 2112-2116, 2003.

286. Nemoto S, and Finkel T. Redox regulation of forkhead proteins through a p66shc-dependent signalling pathway. *Science* 295: 2450-2452, 2002.
287. O'Toole JF, Patel HV, Naples CJ, Fujioka H, and Hopple CL Decreased cytochrome c mediates an age-related decline of oxidative phosphorylation in rat kidney mitochondria. *Biochem J* 427: 105–112, 2010.
288. Ohnishi ST, Ohnishi, T, Muranaka, S, Fujita, H, Kimura, H, Uemura, K, Yoshida, K and Utsumi, K. A possible site of superoxide generation in the complex I segment of rat heart mitochondria. *J Bioenerg Biomembr* 37: 1-15, 2005.
289. Orr WC, Radyuk SN, and Sohal RS. Involvement of redox state in the aging of *Drosophila melanogaster*. *Antioxid Redox Signal* 19: 788-803, 2013.
290. Osiewacz HD. Mitochondrial quality control in aging and lifespan control of the fungal aging model *Podospora anserina*. *Biochem Soc Trans* 39: 1488-1492, 2011.
291. Ozawa T, Tanaka M, and Wakabayashi T. Crystallization of mitochondrial cytochrome oxidase. *Proc Natl Acad Sci USA* 179: 7175-7179, 1982.
292. Ozawa T. Genetic and functional changes in mitochondria associated with aging. *Physiol Rev* 77: 425-464, 1997.
293. Pagliarini DJ, Calvo SE, Chang B, Sheth SA, Vafai SB, Ong SE, Walford GA, Sugiana C, Boneh A, Chen WK, Hill DE, Vidal M, Evans JG, Thorburn DR, Carr SA, and Mootha VK. A mitochondrial protein compendium elucidates complex I disease biology. *Cell* 134: 112-123, 2008.
294. Pagliarini DJ, and Rutter J. Hallmarks of a new era in mitochondrial biochemistry. *Genes Dev* 27: 2615-2627, 2013.
295. Pamplona R, and Constantini D. Molecular and structural antioxidant defenses against oxidative stress in animals. *Am J Physiol* 301: R843–R863, 2011.

296. Pamplona R, and Barja G. An evolutionary comparative scan for longevity-related oxidative stress resistance mechanisms in homeotherms. *Biogerontology* 12: 409-435, 2011.
297. Pan Y. Mitochondria, reactive oxygen species, and chronological aging: a message from yeast. *Exp Gerontol* 46: 847-852, 2011.
298. Pang CY, Ma YS, and Wei YU. MtDNA mutations, functional decline and turnover of mitochondria in aging. *Front Biosci* 13: 3661-3675, 2008.
299. Pani G, and Galeotti T. Role of MnSOD and p66shc in mitochondrial response to p53. *Antioxid Redox Signal* 15: 1715-1727, 2011.
300. Pani G. P66SHC and ageing: ROS and TOR? *Aging (Albany NY)* 2: 514-518, 2010.
301. Panov A, Dikalov S, Shalbuyeva N, Hemendinger R, Greenamyre JT, and Rosenfeld J. Species- and tissue-specific relationships between mitochondrial permeability transition and generation of ROS in brain and liver mitochondria of rats and mice. *Am J Physiol Cell Physiol* 292: C708-718, 2007.
302. Paradies G, Paradies V, De Benedictis V, Ruggiero FM, and Petrosillo G. Functional role of cardiolipin in mitochondrial bioenergetics. *Biochim Biophys Acta* 1837: 408-417, 2014.
303. Paradies G, Petrosillo G, Pistolese M, and Ruggiero FM. Reactive oxygen species affect mitochondrial electron transport complex I activity through oxidative cardiolipin damage. *Gene* 286: 135-141, 2002.
304. Paradies G, Petrosillo G, Pistolese M, and Ruggiero FM. Reactive oxygen species generated by the mitochondrial respiratory chain affect the complex III activity via cardiolipin peroxidation in beef-heart submitochondrial particles. *Mitochondrion* 1: 151-159, 2001.

305. Paradies G, Ruggiero FM, Petrosillo G, and Quagliariello E. Peroxidative damage to cardiac mitochondria: cytochrome oxidase and cardiolipin alterations. *FEBS Lett* 424: 155-158, 1998.
306. Partridge L. Aging: fruit flies break the chain to a longer life. *Aging Cell* 10: 5-9. 2011.
307. Pasdois P, Parker JE, Griffiths EJ, and Halestrap AP. The role of oxidized cytochrome c in regulating mitochondrial reactive oxygen species production and its perturbation in ischaemia. *Biochem J* 436: 493-505, 2011.
308. Patten DA, Lafleur VN, Robitaille GA, Chan DA, Giaccia AJ, and Richard DE. Hypoxia-inducible factor-1 activation in nonhypoxic conditions: the essential role of mitochondrial-derived reactive oxygen species. *Mol Biol Cell* 21: 3247-3257, 2010.
309. Pellegrino MW, Nargund AM, and Haynes CM. Signaling the mitochondrial unfolded protein response. *Biochim Biophys Acta* 1833: 410-416, 2013.
310. Pelletier M, Lepow TS, Billingham LK, Murphy MP, and Siegel RM. New tricks from an old dog: mitochondrial redox signaling in cellular inflammation. *Semin Immunol* 24: 384-392, 2012.
311. Pepelina TY, Chertkova RV, Ostroverkhova TV, Dolgikh DA, Kirpichnikov MP, Grivennikova VG, and Vinogradov AD. Site-directed mutagenesis of cytochrome c: reactions with respiratory chain components and superoxide radical. *Biochemistry (Mosc)* 74: 625-632, 2009.
312. Perez VI, Bokov A, Van Remmen H, Mele J, Ran Q, Ikeno Y, and Richardson A. Is the oxidative stress theory of aging dead? *Biochim Biophys Acta* 1790: 1005–1014, 2009.

313. Pérez-Campo R, López-Torres M, Cadenas S, Rojas C, and Barja G. The rate of free radical production as a determinant of the rate of aging: evidence from the comparative approach. *J Comp Physiol B* 168: 149–158, 1998.
314. Peters K, Dudkina NV, Jänsch L, Braun HP, and Boekema EJ. A structural investigation of complex I and I+III₂ supercomplex from *Zea mays* at 11-13 Å resolution: assignment of the carbonic anhydrase domain and evidence for structural heterogeneity within complex I. *Biochim Biophys Acta* 1777: 84-93, 2008.
315. Petrosillo G, De Benedictis V, Ruggiero FM, Paradies G. Decline in cytochrome c oxidase activity in rat-brain mitochondria with aging. Role of peroxidized cardiolipin and beneficial effect of melatonin. *J Bioenerg Biomembr* 45: 431-440, 2013.
316. Petrosillo G, Matera M, Casanova G, Ruggiero FM, and Paradies G. Mitochondrial dysfunction in rat brain with aging Involvement of complex I, reactive oxygen species and cardiolipin. *Neurochem Int* 53: 126-131, 2008.
317. Petrosillo G, Matera M, Moro N, Ruggiero FM, and Paradies G. Mitochondrial complex I dysfunction in rat heart with aging: critical role of reactive oxygen species and cardiolipin. *Free Radic Biol Med* 46: 88-94, 2009.
318. Pfeiffer K, Gohil V, Stuart RA, Hunte C, Brandt U, Greenberg ML, and Schägger H. Cardiolipin stabilizes respiratory chain supercomplexes. *J Biol Chem* 278: 52873-52880, 2003.
319. Pinton P, Rimessi, A, Marchi, S, Orsini, F, Migliaccio, E, Giorgio, M, Contursi, C, Minucci, S, Mantovani, F, Wieckowski, MR, Del Sal, G, Pelicci, PG, and Rizzuto, R. Protein kinase C β and prolyl isomerase 1 regulate mitochondrial effects of the life-span determinant p66Shc. *Science* 315: 659-663, 2007.

320. Pou S, Keaton, L, Surichamorn, W, Rosen, and GM. Mechanism of superoxide generation by neuronal nitric oxide synthase. *J Biol Chem* 274: 9573-9580, 1999.
321. Puissegur MP, Mazure NM, Bertero T, Pradelli L, Grosso S, Robbe-Sermesant K, Maurin T, Lebrigand K, Cardinaud B, Hofman V, Fourre S, Magnone V, Ricci JE, Pouyssegur J, Gounon P, Hofman P, Barbry P, and Mari B. miR-210 is overexpressed in late stages of lung cancer and mediates mitochondrial alterations associated with modulation of HIF-1 activity. *Cell Death Differ* 18: 465–478, 2011.
322. Quarato G, Piccoli C, Scrima R, and Capitanio N. Variation of flux control coefficient of cytochrome c oxidase and of the other respiratory chain complexes at different values of protonmotive force occurs by a threshold mechanism, *Biochim Biophys Acta* 1807: 1114–1124, 2011.
323. Quinlan CL, Perevoshchikova IV, Hey-Mogensen M, Orr AL, and Brand MD. Sites of reactive oxygen species generation by mitochondria oxidizing different substrates. *Redox Biol* 1: 304-312, 2013.
324. Rabbani N, and Thornalley PJ. The critical role of methylglyoxal and glyoxalase 1 in diabetic nephropathy. *Diabetes* 63: 50-52, 2014.
325. Radermacher M, Ruiz T, Clason T, Benjamin S, Brandt U, and Zickermann V. The three-dimensional structure of complex I from *Yarrowia lipolytica*: a highly dynamic enzyme. *J Struct Biol* 154: 269-279, 2006.
326. Raha S, Myint, AT, Johnstone, L, Robinson, BH. Control of oxygen free radical formation from mitochondrial complex I: roles for protein kinase A and pyruvate dehydrogenase kinase. *Free Radic Biol Med* 32: 421-430, 2002.
327. Ralph SJ, Moreno-Sánchez R, Neuzil J, and Rodríguez-Enríquez S. Inhibitors of succinate: quinone reductase/Complex II regulate production of mitochondrial

- reactive oxygen species and protect normal cells from ischemic damage but induce specific cancer cell death. *Pharm Res* 28: 2695-2730, 2011.
328. Rambold AS, Kostecky B, Elia N, and Lippincott-Schwartz J. Tubular network formation protects mitochondria from autophagosomal degradation during nutrient starvation. *Proc Natl Acad Sci USA* 108: 10190-10195, 2011.
 329. Rhee SG, Chae, HZ, and Kim K. Peroxiredoxins: a historical overview and speculative preview of novel mechanisms and emerging concepts in cell signalling. *Free Radic Biol Med* 38: 1543-1552, 2005.
 330. Richter C. Oxidative damage to mitochondrial DNA and its relationship to ageing. *Int. J. Biochem. Cell Biol* 27: 647-653, 1995.
 331. Rigoulet M, Yoboue ED, and Devin A. Mitochondrial ROS generation and its regulation: mechanisms involved in H₂O₂ signaling. *Antioxid Redox Signal* 14: 459-468, 2011.
 332. Ristow M, and Schmeisser S. Extending life span by increasing oxidative stress. When a theory of aging ages badly. *Free Radic Biol Med* 51: 327-336, 2011.
 333. Rooyackers OE, Adey DB, Ades PA, and Nair KS. Effect of age on in vivo rates of mitochondrial protein synthesis in human skeletal muscle. *Proc Natl Acad Sci USA* 93: 15364-15369, 1996.
 334. Rosca M, Vazquez E, Kerner J, Parland W, Chandler M, Stanley W, Sabbah H, and Hoppel C. Cardiac mitochondria in heart failure: decrease in respirasomes and oxidative phosphorylation. *Cardiovasc Res* 80: 30-39, 2008.
 335. Rossignol R, Faustin B, Rocher C, Malgat M, Mazat JP, and Letellier T. Mitochondrial threshold effects. *Biochem J* 370: 751-762, 2003.
 336. Rota M, LeCapitaine, N, Hosoda, T, Boni, A, De Angelis, A, Padin-Iruegas, ME, Esposito, G, Vitale, S, Urbanek, K, Casarsa, C, Giorgio, M, Luscher, TF, Pelicci,

- PG, Anversa, P, Leri, A, and Kajstura, J. Diabetes promotes cardiac stem cell aging and heart failure, which are prevented by deletion of the p66Shc gene. *Circulation Res* 99: 42-52, 2006.
337. Rygiel KA, Grady JP, and Turnbull DM. Respiratory chain deficiency in aged spinal motor neurons. *Neurobiol Aging* 35: 2230-2238, 2014.
338. Salminen A, Ojala J, Kaarniranta K, Haapasalo A, Hiltunen M, and Soininen H. Astrocytes in the aging brain express characteristics of senescence-associated secretory phenotype. *Eur J Neurosci* 34: 3-11, 2011.
339. Santos CX, Nabeebaccus AA, Shah A, Camargo LL, Filho SV, and Lopes LR. ER stress and Nox-mediated ROS signaling in the peripheral vasculature: potential role in hypertension. *Antioxid Redox Signal* 20: 121-134, 2014.
340. Santos CX, Tanaka LY, Wosniak J, and Laurindo FR. Mechanisms and implications of reactive oxygen species generation during the unfolded protein response: roles of endoplasmic reticulum oxidoreductases, mitochondrial electron transport, and NADPH oxidase. *Antioxid Redox Signal* 11: 2409-2427, 2009.
341. Sanz A, and Stefanatos RKA. The mitochondrial free radical theory of aging: a critical view. *Curr Aging Sci* 1: 10-21, 2008.
342. Sanz A, Caro P, Gómez J, and Barja G. Testing the vicious cycle theory of mitochondrial ROS production: effects of H₂O₂ and cumene hydroperoxide treatment on heart mitochondria. *J Bioenerg Biomembr* 38: 121-127, 2006.
343. Sanz A, Caro P, Ibañez J, Gómez J, Gredilla R, and Barja G. Dietary restriction at old age lowers mitochondrial oxygen radical production and leak at complex I and oxidative DNA damage in rat brain. *J Bioenerg Biomembr* 37: 83-90, 2005.
344. Sarewicz M, Borek A, Cieluch E, Swierczek M, and Osyczka A Discrimination between two possible reaction sequences that create potential risk of generation of

- deleterious radicals by cytochrome bc_1 Implications for the mechanism of superoxide production. *Biochim Biophys Acta* 1797: 1820-1827, 2010.
345. Sauer RT, and Baker TA. AAA+proteases: ATP-fueled machines of protein destruction. *Annu Rev Biochem* 80: 587-612, 2011.
 346. Scacco S, Petruzzella V, Bestini E, Luso A, Papa F, Bellomo F, Signorile A, Torraco A, and Papa S. Mutations in structural genes of complex I associated with neurological diseases. *Ital J Biochem* 55: 254-262, 2006.
 347. Schäfer E, Dencher NA, Vonck J, and Parcej DN. Three-dimensional structure of the respiratory chain supercomplex I₁III₂IV₁ from bovine heart mitochondria. *Biochemistry* 46: 12579-12585, 2007.
 348. Schagger H. Respiratory chain supercomplexes of mitochondria and bacteria. *Biochim Biophys Acta* 1555: 154-159, 2002.
 349. Schagger H, de Coo R, Bauer MF, Hofmann S, Godinot C, and Brandt U. Significance of respirasomes for the assembly/stability of human respiratory chain complex I. *J Biol Chem* 279: 36349-36353, 2004.
 350. Scherz-Shouval R, and Elazar Z. Regulation of autophagy by ROS: physiology and pathology. *Trends Biochem Sci* 36: 30-38, 2011.
 351. Schmeisser S, Priebe S, Groth M, Monajembashi S, Hemmerich P, Guthke R, Platzer M, and Ristow M. Neuronal ROS signaling rather than AMPK/sirtuin-mediated energy sensing links dietary restriction to lifespan extension. *Mol Metab* 2: 92-102, 2013.
 352. Schmeisser K, Mansfeld J, Kuhlmann D, Weimer S, Priebe S, Heiland I, Birringer M, Groth M, Segref A, Kanfi Y, Price NL, Schmeisser S, Schuster S, Pfeiffer AF, Guthke R, Platzer M, Hoppe T, Cohen HY, Zarse K, Sinclair DA, and Ristow M.

- Role of sirtuins in lifespan regulation is linked to methylation of nicotinamide. *Nat Chem Biol* 9: 693-700, 2013.
353. Schmucker DL. Age-related changes in liver structure and function: Implications for disease ? *Exp Gerontol* 40: 650-659, 2005.
 354. Schriener SE, Linford NJ, Martin GM, Treuting P, Ogburn CE, Emond M, Coskun PE, Ladiges W, Wolf N, Van Remmen H, Wallace DC, and Rabinovitch PS. Extension of murine life span by overexpression of catalase targeted to mitochondria. *Science* 308: 1909-1911, 2005.
 355. Scialo F, Mallikarjun V, Stefanatos R, and Sanz A. Regulation of lifespan by the mitochondrial electron transport chain: reactive oxygen species-dependent and reactive oxygen species-independent mechanisms. *Antioxid Redox Signal* 19: 1953-1969, 2013.
 356. Selivanov VA, Zeak JA, Roca J, Cascante M, Trucco M, and Votyakova TV. The role of external and matrix pH in mitochondrial reactive oxygen species generation. *J Biol Chem* 283: 29292-29300, 2008.
 357. Semenza GL. Oxygen-dependent regulation of mitochondrial respiration by hypoxia-inducible factor 1. *Biochem J* 405: 1-9, 2007.
 358. Semenza GL. Oxygen sensing, homeostasis, and disease. *N Engl J Med* 365: 537-547, 2011.
 359. Sen T, Sen N, Jana S, Khan FH, Chatterjee U, and Chakrabarti S. Depolarization and cardiolipin depletion in aged rat brain mitochondria: relationship with oxidative stress and electron transport chain activity. *Neurochem Int* 50: 719-725, 2007.
 360. Sgarbi G, Matarrese P, Pinti M, Lanzarini C, Ascione B, Gibellini L, Dika E, Patrizi A, Tommasino C, Capri M, Cossarizza A, Baracca A, Lenaz G, Solaini G, Franceschi C, Malorni W, and Salvioli S. Mitochondria hyperfusion and elevated autophagic

- activity are key mechanisms for cellular bioenergetic preservation in centenarians. *Aging (Albany NY)* 6: 296-310, 2014.
361. Sharp FR, Ran R, Lu A, Tang Y, Strauss KI, Glass T, Ardizzone T, and Bernaudin M. Hypoxic preconditioning protects against ischemic brain injury. *NeuroRx* 1: 26-35, 2004.
 362. Short KR, Bigelow ML, Kahl J, Singh R, Coenen-Schimke J, Raghavakaimal S, and Nair KS. Decline in skeletal muscle mitochondrial function with aging in humans. *Proc Natl Acad Sci USA* 102: 5618-5623, 2005.
 363. Singh KK. Mitochondrial dysfunction is a common phenotype in aging and cancer. *Ann NY Acad Sci USA* 1019: 260-264, 2004.
 364. Skulachev VP. Role of uncoupled and non-coupled oxidations in maintenance of safely low levels of oxygen and its one-electron reductants. *Q Rev Biophys* 29: 169-202, 1996.
 365. Sohal RS, Arnold L, and Orr WC. Effect of age on superoxide dismutase, catalase, glutathione reductase, inorganic peroxides, TBA-reactive material, GSH/GSSG, NADPH/NADP⁺ and NADH/NAD⁺ in *Drosophila melanogaster*. *Mech Ageing Dev* 56: 223–235, 1990.
 366. Sohal RS, Ku HH, Agarwal S, Forster MJ, and Lal H. Oxidative damage, mitochondrial oxidant generation and antioxidant defenses during aging and in response to food restriction in the mouse. *Mech Ageing Dev* 74: 121-133, 1994.
 367. Srivastava S, and Moraes CT. Double-strand breaks of mouse muscle mtDNA promote large deletions similar to multiple mtDNA deletions in humans. *Hum Mol Genet* 14: 893-902, 2005.
 368. Starkov AA, and Fiskum, G. Regulation of brain mitochondrial H₂O₂ production by membrane potential and NAD(P)H redox state. *J Neurochem* 86: 1101-1107, 2003.

369. Starkov AA, Fiskum, G, Chinopoulos, C, Lorenzo, BJ, Browne, SE, Patel, MS, and Beal MF. Mitochondrial alpha-ketoglutarate dehydrogenase complex generates reactive oxygen species. *J Neuroscience* 24: 7779-7788, 2004.
370. St-Pierre J, Buckingham, JA, Roebuck, SJ and Brand, MD. Topology of superoxide production from different sites in the mitochondrial electron transfer chain. *J Biol Chem* 277: 44784-44790, 2002.
371. Strogolova V, Furness A, Robb-McGrath M, Garlich J, and Stuart RA. Rcf1 and Rcf2, members of the hypoxia induced gene 1 protein family, are critical components of the mitochondrial cytochrome bc1-cytochrome c oxidase supercomplex. *Mol Cell Biol* 32: 1363-1373, 2012.
372. Sugiyama S, Takasawa M, Hayakawa M, and Ozawa T. Changes in skeletal muscle, heart and liver mitochondrial electron transport activities in rats and dogs of various ages. *Biochem Mol Biol Int* 30: 937-944, 1993.
373. Suski JM, Karkucinska-Wieckowska A, Lebiedzinska M, Giorgi C, Szczepanowska J, Szabadkai G, Duszynski J, Pronicki M, Pinton P, and Wieckowski MR. p66Shc Aging Protein in Control of Fibroblasts Cell Fate. *Int J Mol Sci* 12: 5373-5389, 2011.
374. Szweda PA, Camouse M, Lundberg KC, Oberley TD, and Szweda LI. Aging, lipofuscin formation, and free radical-mediated inhibition of cellular proteolytic systems. *Ageing Res Rev* 2: 383-405, 2003.
375. Tahara EB, Navarete FD, and Kowaltowski AJ. Tissue-, substrate-, and site-specific characteristics of mitochondrial reactive oxygen species generation. *Free Radic Biol Med* 46: 1283-1297, 2009.
376. Tapia PC. Sublethal mitochondrial stress with an attendant stoichiometric augmentation of reactive oxygen species may precipitate many of the beneficial alterations in

- cellular physiology produced by caloric restriction, intermittent fasting, exercise and dietary phytonutrients: "Mitohormesis" for health and vitality. *Med Hypotheses* 66: 832-843, 2006.
377. Tatarková Z, Kuka S, Račay P, Lehotský J, Dobrota D, Mištuna D, and Kaplán P. Effects of aging on activities of mitochondrial electron transport chain complexes and oxidative damage in rat heart. *Physiol Res* 60: 281-289, 2011.
378. Taylor CT. Mitochondria and cellular oxygen sensing in the HIF pathway. *Biochem J* 409: 19-26, 2008.
379. Taylor EB, and Rutter J. Mitochondrial quality control by the ubiquitin-proteasome system. *Biochem Soc Trans* 39: 1509-1513, 2011.
380. Tomko Jr RJ, and Hochstrasser M. Molecular Architecture and Assembly of the Eukaryotic Proteasome. *Annu Rev Biochem* 82: 415-445, 2013.
381. Trachootham D, Lu W, Ogasawara MA, Nilsa RD, and Huang P. Redox regulation of cell survival. *Antioxid Redox Signal* 10: 1343-1374, 2008.
382. Tretter L, and Adam-Vizi, V. Generation of reactive oxygen species in the reaction catalyzed by alpha-ketoglutarate dehydrogenase. *J Neurosci* 24: 7771-7778, 2004.
383. Trifunovic A, Hansson A, Wredenberg A, Rovio AT, Dufour E, Khvorostov I, Spelbrink JN, Wibom R, Jacobs HT, and Larsson NG. Somatic mtDNA mutations cause aging phenotypes without affecting reactive oxygen species production. *Proc Natl Acad Sci USA* 102: 17993-17998, 2005.
384. Trifunovic A, Wredenberg A, Falkenberg M, Spelbrink JN, Rovio AT, Bruder CE, Bohlooly-Y M, Gidlöf S, Oldfors A, Wibom R, Törnell J, Jacobs HT, and Larsson NG. Premature ageing in mice expressing defective mitochondrial DNA polymerase. *Nature* 429(6990): 417-423, 2004.

385. Trinei M, Berniakovich I, Beltrami E, Migliaccio E, Fassina A, Pelicci P, and Giorgio M. P66Shc signals to age. *Aging* 1: 503-510, 2009.
386. Trounce I, Byrne E, and Marzuki S. Decline in skeletal muscle mitochondrial respiratory chain function: possible factor in ageing. *Lancet* 1(8639): 637-639, 1989.
387. Trujillo M, Ferrer-Sueta G, and Radi R. Kinetic studies on peroxynitrite reduction by peroxiredoxins. *Methods Enzymol* 441: 173-196, 2008.
388. Turrens JF, and Boveris A. Generation of superoxide anion by the NADH dehydrogenase of bovine heart mitochondria. *Biochem J* 191: 421-427, 1980.
389. Ugarte N, Petropoulos I, and Friguet B. Oxidized mitochondrial protein degradation and repair in aging and oxidative stress *Antioxid Redox Signal* 13: 539-549, 2009.
390. Valko M, Rhodes, CJ, Moncol, J, Izakovic, M, and Mazur M. Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chemico-Biological Interactions* 160: 1-40, 2006.
391. Velculescu VE, Zhang L, Vogelstein B, and Kinzler KW. Serial analysis of gene expression. *Science* 270: 484-487, 1995.
392. Venkatesh S, Lee J, Singh K, Lee I, and Suzuki CK. Multitasking in the mitochondrion by the ATP-dependent Lon protease. *Biochim Biophys Acta* 1823: 56-66, 2012.
393. Ventura B, Genova ML, Bovina C, Formiggini G, and Lenaz G. Control of oxidative phosphorylation by Complex I in rat liver mitochondria: implications for aging. *Biochim Biophys Acta* 1553: 249-260, 2002.
394. Vermulst M, Bielas JH, Kujoth GC, Ladiges WC, Rabinovitch PS, Prolla TA, and Loeb LA. Mitochondrial point mutations do not limit the natural lifespan of mice. *Nat Genet* 39: 540-543, 2007.

395. Vermulst M, Wanagat J, Kujoth GC, Bielas JH, Rabinovitch PS, Prolla TA, and Loeb LA. DNA deletions and clonal mutations drive premature aging in mitochondrial mutator mice. *Nat Genet* 40: 392-394, 2008.
396. Vinogradov AD, Grivennikova VG. Generation of superoxide-radical by the NADH:ubiquinone oxidoreductase of heart mitochondria. *Biochemistry (Mosc)* 70: 120-127, 2005.
397. Von Zglinicki T, Burkle A, and Kirkwood TBL. Stress, DNA damage and ageing – an integrative approach. *Exp Gerontol* 36: 1049-1062, 2001.
398. Vukotic M, Oeljeklaus S, Wiese S, Vögtle F-N, Meisinger C, Meyer HE, Zieseniss A, Katschinski DM, Jans DC, Jakobs S, Warscheid B, Rehling P, and Deckers M. Rcf1 Mediates Cytochrome Oxidase Assembly and Respirasome Formation, Revealing Heterogeneity of the Enzyme Complex. *Cell Metab* 15: 336-347, 2012.
399. Wanagat J, Cao Z, Pathare P, and Aiken JM. Mitochondrial DNA deletion mutations colocalize with segmental electron transport system abnormalities, muscle fiber atrophy, fiber splitting, and oxidative damage in sarcopenia. *FASEB J* 15: 322-332, 2001.
400. Wang Y, Yang J, and Yi J. Redox sensing by proteins: oxidative modifications on cysteines and the consequent events. *Antioxid Redox Signal* 16: 649-657, 2012.
401. Warda M, Kim HK, Kim N, Ko KS, Rhee BD, and Han J. A matter of life, death and diseases: mitochondria from a proteomic perspective. *Expert Rev Proteomics* 10: 97-111, 2013.
402. Waypa GB, Marks JD, Guzy RD, Mungai PT, Schriewer JM, Dokic D, Ball MK, and Schumacker PT. Superoxide generated at mitochondrial complex III triggers acute responses to hypoxia in the pulmonary circulation. *Am J Respir Crit Care Med* 187: 424-432, 2013.

403. Wei Y, Zhang YJ, Cai Y, and Xu MH. The role of mitochondria in mTOR-regulated longevity. *Biol Rev Camb Philos Soc* 2014 Mar 28. [Epub ahead of print]
404. Wei YH, and Lee HC. Oxidative stress, mitochondrial DNA mutation, and impairment of antioxidant enzymes in aging. *Exp Biol Med* 227: 671-682, 2002.
405. Wenz T, Hielscher R, Hellwig P, Schägger H, Richers S, and Hunte C. Role of phospholipids in respiratory cytochrome bc(1) complex catalysis and supercomplex formation. *Biochim Biophys Acta* 1787: 609-616, 2009.
406. Whelan SP, and Zuckerbraun BS. Mitochondrial signaling: forwards, backwards, and in between. *Oxid Med Cell Longev* 2013: 351613, 2013.
407. Wilkinson DS, Taylor RC, and Dillin A. Mitochondrial respiration and reactive oxygen species in *C. elegans*. *Methods Cell Biol* 107: 353-381, 2012.
408. Winge DR. Sealing the mitochondrial respirasome. *Mol Cell Biol* 32: 2647-2652, 2012.
409. Wojtczak L, Lebedzińska M, Suski JM, Więckowski MR, and Schönfeld P. Inhibition by purine nucleotides of the release of reactive oxygen species from muscle mitochondria: indication for a function of uncoupling proteins as superoxide anion transporters. *Biochem Biophys Res Commun* 407: 772-776, 2011.
410. Wong YT, Ruan R, and Tay FE. Relationship between levels of oxidative DNA damage, lipid peroxidation and mitochondrial membrane potential in young and old F344 rats. *Free Radic Res* 40: 393-402, 2006.
411. Wu WS. The signaling mechanism of ROS in tumor progression. *Cancer Metastasis Rev* 25: 695-705, 2006.
412. Yakes FM, and Van Houten B. Mitochondrial DNA damage is more extensive and persists longer than nuclear DNA damage in human cells following oxidative stress. *Proc Natl Acad Sci USA* 94: 514-519, 1997.

413. Yang W, and Hekimi S. A mitochondrial superoxide signal triggers increased longevity in *Caenorhabditis elegans*. *PLoS Biol* 8: e1000556, 2010.
414. Yen TC, Chen YS, King KL, Yeh SH, and Wei YH. Liver mitochondrial respiratory functions decline with age. *Biochem Biophys Res Commun* 165: 944-1003, 1989.
415. Yin F, Sancheti H, and Cadenas E. Mitochondrial thiols in the regulation of cell death pathways. *Antioxid Redox Signal* 17: 1714-1727, 2012.
416. Yu BP. Membrane alteration as a basis of aging and the protective effects of calorie restriction. *Mech Ageing Dev* 126: 1003-1010, 2005.
417. Yun J, and Finkel T. Mitohormesis. *Cell Metab* 19: 757-766, 2014.
418. Zaobornyj T, and Ghafourifar P. Strategic localization of heart mitochondrial NOS: a review of the evidence. *Am J Physiol Heart Circ Physiol* 303: H1283-1293, 2012.
419. Zarse K, Schmeisser S, Groth M, Priebe S, Beuster G, Kuhlow D, Guthke R, Platzer M, Kahn CR, and Ristow M. Impaired insulin/IGF1 signaling extends life span by promoting mitochondrial L-proline catabolism to induce a transient ROS signal. *Cell Metab* 15: 451-465, 2012.
420. Zhang L, Yu, L, and Yu CA. Generation of superoxide anion by succinate cytochrome c reductase from bovine heart mitochondria. *J Biol Chem*, 273: 33972-33976, 1998.
421. Zhang M, Mileykovskaya E, and Dowhan W. Gluing the respiratory chain together. Cardiolipin is required for supercomplex formation in the inner mitochondrial membrane. *J Biol Chem* 277: 43553-43556, 2002.
422. Zorov DB, Juhaszova M, and Sollott SJ. Mitochondrial ROS-induced ROS release: an update and review. *Biochim Biophys Acta* 1757: 509-517, 2006.
423. Zuckerbraun BS, Chin BY, Bilban M, d'Avila JC, Rao J, Billiar TR, and Otterbein LE. Carbon monoxide signals via inhibition of cytochrome c oxidase and generation of mitochondrial reactive oxygen species. *FASEB J* 21: 1099-1106, 2007.

424. Zuin A, Castellano-Esteve D, Ayté J, and Hidalgo E. Living on the edge: stress and activation of stress responses promote lifespan extension. *Aging (Albany NY)* 2: 231-237, 2010.

FIGURE LEGEND




Figure 1 - A schematic drawing of the respiratory chain depicting the protein complexes and their substrates. Complex I is depicted as a component of the I₁III₂IV₁ supercomplex; whereas Complex III and Complex IV are also shown in their free form. The white arrows represent the sources of superoxide at different sites in relation to the inner mitochondrial membrane. I, NADH-ubiquinone oxidoreductase; II, succinate-ubiquinone oxidoreductase; III, ubiquinol-cytochrome *c* oxidoreductase; IV, cytochrome oxidase; NDi and NDe, internal and external alternative NAD(P)H dehydrogenases; AOX, alternative oxidase; αGP, glycerol-3-phosphate; ETF, electron transfer flavoprotein; DHO, dihydroorotate; CoQ, Coenzyme Q (ubiquinone); C, cytochrome *c*; P66, cytochrome p66 (see text for details).




Figure 2 - Mitochondrial ROS regulate the cellular response to hypoxia. Hypoxia induces the production of mitochondrial ROS that inhibit the activity of prolyl hydroxylases (PHD2) and stabilize HIFα. Under normoxic conditions, hydroxylation tags HIFα for recognition by the Von Hippel-Lindau tumor suppressor protein and subsequent proteosomal degradation. Mitochondrial ROS generated during hypoxia also regulate increases in cellular calcium uptake and lead to activation of AMPK, allowing increased cellular energy conservation. AMPK phosphorylates the α-subunit of the Na/K ATPase leading to endocytosis. Figure taken from Hamanaka and Chandel (152), with permission from Elsevier.




Figure 3 - Nuclear-mitochondrial communication and its decline during aging. Nuclear NAD⁺ levels regulate mitochondria via a PGC-1 α -independent pathway that ensures the correct stoichiometry of OXPHOS subunits, but over time, a chronic pseudohypoxic response is activated, inhibiting OXPHOS. Mitochondrial encoded genes were *ND1*, *Cytb*, *COX1*, and *ATP6*. Reprinted from Gomes et al. (133), with permission from Elsevier.




Figure 4 – Scheme showing how the loss of supercomplex organisation may be involved in a vicious circle of oxidative stress and energy failure. ROS production by Complex I is enhanced as a consequence of supercomplex disassembling. Membrane phospholipid peroxidation, mtDNA damage and subsequent misassembly of the respiratory complexes with further loss of supercomplex organisation may occur due to enhanced mitochondrial oxidative

stress, thus perpetuating the vicious circle. Depending on the amount produced, ROS can operate as signalling molecules from mitochondria to the nucleus. See text for explanations.

ROS production




Figure 5 - Production of ROS by mitochondrial Complex I in different situations where supercomplexes are maintained or disassembled. The percent value of ROS production measured in all the samples listed in the table (*lower panel*) is plotted in the graph (*upper panel*) in descending order of the corresponding amount of Complex I-containing supercomplexes. The ratio of bound Complex I vs. total Complex I was determined by densitometric analysis of immunoblots obtained after 2D BN/SDS-PAGE as described in Maranzana et al. (252). The NADH-stimulated production of ROS was measured as the relative fluorescence intensity of dichlorofluorescein in the presence of 1.8 μ M mucidin and 4 μ M rotenone, and expressed as percentage value of the corresponding reference samples: ^aproteoliposomes (cf. text for details), ^bSMP (submitochondrial particles) and ^cBHM (bovine heart mitochondria) respectively assayed in the presence of 1.8 μ M mucidin only, as in Maranzana et al. (252). In the case of BHM, the existence of endogenous systems operating to reduce ROS levels in the mitochondrial sample might have counteracted the dramatic effects of the complete dissociation of Complex I, thus leading to a two-fold only increase of the measured ROS production. DIL., dilution at high lipid to protein ratio (30:1 w:w); DDM, dodecyl maltoside; KSCN, potassium thiocyanate.