This is the final peer-reviewed accepted manuscript of

Sabia, Alessandra; Baldisserotto, Costanza; BIONDI, STEFANIA; Marchesini, Roberta; Tedeschi, Paola; Maietti, Annalisa; Giovanardi, Martina; Ferroni, Lorenzo; Pancaldi, Simonetta: Re-cultivation of *Neochloris oleoabundans* in exhausted autotrophic and mixotrophic media: the potential role of polyamines and free fatty acids. APPLIED MICROBIOLOGY AND BIOTECHNOLOGY, 99. ISSN 0175-7598

DOI: 10.1007/s00253-015-6908-3

The final published version is available online at:

<http://dx.doi.org/10.1007/s00253-015-6908-3>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Re-cultivation of *Neochloris oleoabundans* in exhausted autotrophic and mixotrophic media: the potential role of polyamines and free fatty acids

Alessandra Sabia¹ · Costanza Baldisserotto¹ · Stefania Biondi² · Roberta Marchesini¹ · Paola Tedeschi³ · Annalisa Maietti³ · Martina Giovanardi¹ · Lorenzo Ferroni¹ · Simonetta Pancaldi¹

Abstract *Neochloris oleoabundans* (Chlorophyta) is widely considered one of the most promising microalgae for biotechnological applications. However, the large-scale production of microalgae requires large amounts of water. In this perspective, the possibility of using exhausted growth media for the re-cultivation of *N. oleoabundans* was investigated in order to simultaneously make the cultivation more economically feasible and environmentally sustainable. Experiments were performed by testing the following media: autotrophic exhausted medium (E+) and mixotrophic exhausted medium after cultivation with glucose (EG+) of *N. oleoabundans* cells grown in a 20-L photobioreactor (PBR). Both exhausted media were replenished with the same amounts of nitrate and phosphate as the control brackish medium (C). Growth kinetics, nitrate and phosphate consumption, photosynthetic pigments content, photosynthetic efficiency, cell morphology, and lipid production were evaluated. Moreover, the free fatty acid (FFA) composition of exhausted media and the polyamine (PA) concentrations of both algae and media were analyzed in order to test if some molecules, released into the medium, could influence algal growth and metabolism. Results showed that

N. oleoabundans can efficiently grow in both exhausted media, if appropriately replenished with the main nutrients (E+ and EG+), especially in E+ and to the same extent as in C medium. Growth promotion of *N. oleoabundans* was attributed to PAs and alteration of the photosynthetic apparatus to FFAs. Taken together, results show that recycling growth medium is a suitable solution to obtain good *N. oleoabundans* biomass concentrations, while providing a more sustainable ecological impact on water resources.

Keywords *Neochloris oleoabundans* · Recycling culture media · Photosynthetic apparatus · Biomass production · Polyamines · Free fatty acids

Introduction

Over the last decades, microalgae have received much attention as a promising high-potential feedstock for biodiesel (Chisti 2007; Li et al. 2008; Borowitzka and Moheimani 2013) and as an attractive raw material for the production of a wide range of high-value bioproducts (Molina Grima et al. 2003; Mata et al. 2010). They are photosynthetic microorganisms with numerous key features, which potentially make them a more appreciable renewable source than terrestrial plants (Richmond 2004; Hu et al. 2008; Ndimba et al. 2013). For instance, these microorganisms have faster growth rates, higher photosynthetic efficiencies, higher rates of carbon dioxide fixation, and higher biomass productivities compared with plants (Richmond 2004; Gouveia et al. 2009; Harun et al. 2010; Mata et al. 2010). Moreover, microalgae can thrive in non-arable lands being able to grow in waters having different salinity levels and chemical composition (Smith et al. 2010; Borowitzka and Moheimani 2013).

✉ Simonetta Pancaldi
simonetta.pancaldi@unife.it

¹ Department of Life Sciences and Biotechnology, University of Ferrara, Corso Ercole I d'Este 32, 44121 Ferrara, Italy

² Department of Biological, Geological and Environmental Sciences, University of Bologna, via Imerio 42, 40126 Bologna, Italy

³ Department of Chemical and Pharmaceutical Science, University of Ferrara, via Fossato di Mortara 17, 44121 Ferrara, Italy

The environmental and economical impact of large-scale production of microalgae has been widely discussed, with the aim of making the process increasingly sustainable (Sheehan 2009; Solomon 2010). In this perspective, microalgal cultivation systems are crucial factors to be taken into account (Smith et al. 2010; Stephens et al. 2010; Borowitzka and Moheimani 2013). A wide variety of systems has been described by Mata et al. (2010) and recently by Borowitzka and Moheimani (2013). Microalgae can be cultivated in open pond reactors or closed photobioreactors (PBRs), depending on algal strain, issue of research, type of desired products, and environmental conditions (Chisti 2007; Mata et al. 2010; Borowitzka and Moheimani 2013). Although open ponds are the most commonly used culture systems, due to their low cost of production and operation, recently closed PBRs are receiving much attention for the possibility of producing valuable compounds (Pulz 2001; Grobbelaar 2009; Smith et al. 2010). However, algal cultures require large amounts of water in both culture systems (Lam and Lee 2012; Borowitzka and Moheimani 2013). Considering that water is becoming a scarce natural resource, water demand represents an important factor when performing large-scale cultivation (Batan et al. 2013; Farooq et al. 2014). In order to reduce the high production costs and, at the same time, to make microalgal production more environmentally sustainable, recycling culture medium has been proposed as a possible solution (Yang et al. 2011; Hady-Romdhane et al. 2012; Farooq et al. 2014), especially in large-scale culture systems (Borowitzka 2005; Lam and Lee 2012; Zhu et al. 2013). However, the feasibility of recycling growth medium to re-cultivate microalgae has been tested only in a few cases (Lívanský et al. 1996; Rodolfi et al. 2003; Hady-Romdhane et al. 2013; Zhu et al. 2013). It is reported that the recycling medium could have negative effects on biomass productivity, due to the release of inhibitory secondary metabolites (Richmond 2004). Harmful metabolites are, in fact, released during microalgal growth under physiological stress (Ikawa 2004; Moheimani and Borowitzka 2006). Free fatty acids (FFAs), and substances derived from the photooxidation of unsaturated fatty acids, are the most common metabolites with inhibitory effects on microalgae (Ikawa 2004; Wu et al. 2006; Bosma et al. 2008; Stephens et al. 2010).

In the present work, the suitability of recycling growth media for the re-cultivation of *Neochloris oleoabundans* (*Chlorophyta*) was investigated. This species was chosen because it is considered one of the most promising oil-rich microalgae, due to its capability to accumulate lipids when grown under nitrogen starvation (Tornabene et al. 1983; Chisti 2007; Li et al. 2008; Pruvost et al. 2009) or mixotrophically, in the presence of glucose, or carbon-rich wastes, as organic carbon sources (Giovanardi et al. 2013; Baldissarotto et al. 2014). Present results clearly indicate that *N. oleoabundans* can efficiently grow in its exhausted growth

medium, if suitably replenished with the main nutrients. Based on this encouraging result, and with the aim of gaining further knowledge on the morpho-physiological aspects and biotechnological applications of this algal strain, the following features were analyzed: (1) growth kinetics in parallel to nitrate and phosphate consumption; (2) photosynthetic pigment content; (3) PSII maximum quantum yield; (4) cell morphology, with special attention to intracellular lipid accumulation, (5) FFA accumulation in recycled growth media, and (6) poly-amine (PA) concentration both inside cells and in the corresponding culture media. PAs were determined in order to understand if these plant growth regulators could be responsible for the growth promotion of *N. oleoabundans* in recycled growth medium. It is known, in fact, that PAs, together with other plant growth regulators, have stimulatory effects on algal growth and metabolism, and are involved in mitigating various types of biotic and abiotic stress (Tate et al. 2013).

Materials and methods

Algal strains and culture condition

The strain used in this study was the *Chlorophyta N. oleoabundans* UTEX 1185 (syn. *Ettlia oleoabundans*) (*Sphaeropleales, Neochloridaceae*), obtained from the Culture Collection of Algae of the University of Texas (UTEX, Austin, Texas, USA; www.utex.org).

N. oleoabundans was cultivated in axenic liquid brackish medium (BM) (Baldissarotto et al. 2012) in a coaxial 20-L capacity PBR (M2M Engineering, Grazzanise, Caserta, Italy). Algae were cultivated autotrophically in BM for 25 days or mixotrophically, by addition of 2.5 g L⁻¹ glucose, for 8 days, according to previously described protocols (Baldissarotto et al. 2014; Giovanardi et al. 2014). For autotrophic cultivation, cells were inoculated into the PBR to obtain an initial cell density of about 0.6 × 10⁶ cells mL⁻¹, while for mixotrophic cultivation initial cell density was higher (10 × 10⁶ cells mL⁻¹). Culture conditions in the PBR were 24 ± 1 °C; sterile air injection at the bottom of the PBR, with 0.5/3.5 h bubbling/static cycles; irradiance 65 μmol_{photons} m⁻² s⁻¹ of PAR (16:8 h light/dark photoperiod). Light was supplied with inner cool-white fluorescent Philips tubes.

Algal growth and morphology in the PBR were monitored as described below.

Preparation of growth media

Experiments were performed by testing the following media:

- C Freshly prepared BM medium (control);
- E Autotrophic exhausted medium;

- EG Mixotrophic exhausted medium after cultivation with glucose;
- E+ Autotrophic exhausted medium replenished with nitrate and phosphate concentrations as for BM medium;
- EG+ Mixotrophic exhausted medium after cultivation with glucose and replenished with nutrients as in E+.

In order to obtain the exhausted media for the recycling experiment, *N. oleoabundans* was grown inside the PBR under the culture conditions described in the previous section. For E+ medium preparation, about 500 mL of autotrophic algal culture in the stationary phase of growth (at the 15th day of cultivation in the PBR) were centrifuged at 2000×g for 10 min in order to separate the medium from the algae, thus obtaining an autotrophic exhausted medium (E). The medium appeared pale-yellow and was free from algae, bacteria, and protozoa; its optical density at a wavelength of 750 nm (OD₇₅₀) was 0.02. For EG+ medium, at the 8th day of cultivation in the PBR, the same aliquots of algal culture were harvested by centrifugation (2000×g for 10 min). In this case, the mixotrophic exhausted medium (EG) was straw-yellow in color and presented a weak bacterial contamination due to the glucose addition (OD₇₅₀ = 0.009). Glucose was proved to be totally consumed during the previous cultivation inside the PBR by 3,5-dinitrosalicylic (DNS; Sigma-Aldrich, Gallarate, Milan, Italy) acid assay, according to Giovanardi et al. (2014). In both exhausted media, pH was not adjusted. After determining the nitrate and phosphate concentrations of the two exhausted media (see “Nitrate and phosphate analyses”), KNO₃ and K₂HPO₄ were axenically added to reach final concentrations of 0.2 and 0.02 g L⁻¹, respectively, i.e., the typical concentration of those components in BM, thus obtaining the replenished exhausted media (E+ and EG+) used for experiments.

Experimental design

When autotrophic cultures of *N. oleoabundans*, grown in BM in the PBR, reached a cell density of 10 × 10⁶ cells mL⁻¹ (after about 9 days of cultivation), aliquots of cells were inoculated into 300-mL Erlenmeyer flasks (150 mL total volume) containing C, E+, or EG+ media, to obtain an initial cell density of about 0.5 × 10⁶ cells mL⁻¹. The cultures were placed in a growth chamber (24 ± 1 °C, 80 μmol_{photons} m⁻² s⁻¹ of PAR with a 16:8 h light/darkness photoperiod) and cultivated with continuous shaking at 80 rpm, without external CO₂ supply. Experiments were performed in triplicate. Aliquots of cultures (cells and media) were collected at different times of cultivation up to 25 days, depending on the analysis.

Analyses

Growth evaluation

Aliquots of cell samples cultivated in control and exhausted media were counted at 0, 3, 7, 12, 17, and 25 days of cultivation using a Thoma hemocytometer (HBG, Giessen, Germany) under a light microscope (Zeiss, model Axiophot), and growth curves were obtained.

The growth rate (μ , number of divisions per day) during the exponential phase was calculated with the following equation:

$$\mu(\text{div day}^{-1}) = (\log_2 N_1 - \log_2 N_0) / (t_1 - t_0),$$

where μ is the growth rate, N_1 the cell number at time t_1 , N_0 the cell number at time 0, and $t_1 - t_0$ the time interval (days) (Giovanardi et al. 2013).

In parallel to growth, pH was also periodically monitored on small aliquots of samples.

Nitrate and phosphate analyses

After 12 and 25 days of cultivation, samples of C, E+, and EG+ media were harvested by centrifugation to analyze nitrate and phosphate concentrations. These nutrients were quantified colorimetrically using a flow-injection autoanalyzer (FlowSys, Systea, Roma, Italy).

Photosynthetic pigment extraction and quantification

For photosynthetic pigment analysis, cell samples were collected by centrifugation after 0, 7, 17, 21, and 25 days of cultivation. Extraction of photosynthetic pigments was performed according to Baldissarotto et al. (2014). The extracts were measured with a Pharmacia Ultrospec 2000 UV-Vis spectrophotometer (1-nm bandwidth; Amersham Biosciences, Piscataway, NJ, USA) at 666 (chlorophyll *a* (Chl*a*)), 653 (chlorophyll *b* (Chl*b*)), and 470 nm (carotenoids (Car)). Quantification was performed according to equations reported in Wellburn (1994).

PAM fluorimetry

The PSII maximum quantum yield of algae was determined at the same cultivation times considered for growth kinetics measurements. A pulse amplitude-modulated fluorometer (ADC Bioscientific Ltd., Hoddesdon, Hertfordshire, UK) was used to determine the in vivo chlorophyll fluorescence of PSII. The PSII maximum quantum yield is reported as F_v/F_m ratio, i.e., $(F_m - F_0)/F_m$, where variable fluorescence is $F_v = (F_m - F_0)$, F_m is the maximum fluorescence and F_0 is the initial fluorescence of samples (Lichtenthaler et al. 2005). This measurement is considered a valid method to probe the

maximum quantum yield of photochemistry in PSII (Kalaji et al. 2014). Moreover, it is useful to estimate the physiological state of plants and microorganisms also under nutrient stress (White et al. 2011). Samples were prepared as reported in Ferroni et al. (2011) after 15 min of dark incubation.

Light and fluorescence microscopy

For microscopic observations, cell samples were routinely collected throughout the cultivation period inside the PBR and during the experiment with exhausted media. Aliquots of samples were observed using a microscope (Zeiss, model Axiophot) with conventional and fluorescent attachments. The light source for chlorophyll fluorescence observation was a HBO 100 W pressure mercury vapor lamp (filter set, BP436/10, LP470). Pictures of cells were taken with a Canon IXUS 110 IS digital camera (12.1 megapixels), mounted on the ocular lens through a Leica DC150 system (Leica Camera AG, Solms, Germany).

Lipid staining

During experiments, the intracellular presence of lipids was evaluated by staining cells with the fluorochrome Nile red (9-diethylamino-5H-benzo[α]phenoxazine-5-one, 0.5 mg dissolved in 100 mL acetone; Sigma-Aldrich, Gallarate, Milan, Italy), as described in Giovanardi et al. (2014). After incubation at 37 °C in darkness for 15 min, cells were observed with the microscope described above at an excitation wavelength of 485 nm (filter set, BP485, LP520). Photographs were taken with the camera described above.

Transmission electron microscopy

Transmission electron microscopy (TEM) observations were made on cells harvested by centrifugation (600×g, 10 min) at the 12th day of cultivation. Cells were fixed, post-fixed, and dehydrated as reported in Baldisserotto et al. (2012). Embedding in resin and staining procedures were performed as previously described (Pancaldi et al. 2002). Sections were observed with a Hitachi H800 electron microscope (Hitachi, Tokyo, Japan) at the Electron Microscopy Centre, University of Ferrara.

Extracellular free fatty acids analysis

To analyze the FFA composition of the exhausted E and EG media used to prepare E+ and EG+ replenished media, aliquots of *N. oleoabundans* cultures grown autotrophically and mixotrophically in the PBR were collected at the 15th and at the 8th days of cultivation, respectively. Samples were centrifuged at 2000×g for 10 min in order to separate

the medium from algae. Fifty milliliters of media obtained from the autotrophic (E) and mixotrophic (EG) cultures were freeze-dried for analysis. The extracellular FFA composition was determined in duplicate by gas chromatography-mass spectrometry (GC-MS). For extraction, samples were dissolved in 3 mL of hexane, sonicated for 15 min, and extracted overnight. Fatty acid methyl esters were prepared by transesterification with 1.5 mL of 5 % of sodium hydroxide in methanol solution. Sample volumes of 1 μ L were injected into the GC-MS apparatus, which consisted of a Varian Saturn 2100 MS/MS ion trap mass spectrometer. Separations were performed using a Zebron ZB-WAX Phenomenex capillary column (60 cm in length, 0.25 mm i.d.) supplied with helium carrier gas at 1 mL min⁻¹ constant flow. The injector temperature was 250 °C, and the oven temperature program was the following: start 100 °C for 2 min, ramp to 200 °C at 10 °C/min, and hold for 108 min. The MS acquisitions were performed by full-scan mode.

Determination of polyamines

To analyze the free PA composition of *N. oleoabundans* cells cultivated autotrophically and mixotrophically in PBR and in the corresponding E and EG media, samples were collected as described for FFA analyses. Freeze-dried E or EG media and algae (50 and 20 mg, respectively) were extracted in cold 4 % perchloric acid, kept for 1 h on ice, and then centrifuged at 15,000×g for 15 min. Aliquots (200 μ L) of the supernatants and standard solutions of putrescine (Put), spermidine (Spd), and spermine (Spm) were derivatized with dansyl chloride (Scaramagli et al. 1999). Dansylated derivatives were extracted with toluene, taken to dryness and resuspended in acetonitrile. PAs were separated and quantified by HPLC using a reverse-phase C₁₈ column (Spherisorb ODS2, 5- μ m particle diameter, 4.6 × 250 mm, Waters, Wexford, Ireland) and a programmed acetonitrile/water step gradient (flow rate, 1 mL min⁻¹) on a Jasco system (Jasco Corp., Tokyo, Japan) consisting of a PU-1580 pump, an LG-1580-02 ternary gradient unit, a DG-1580-53 three-line degasser, and a FP-1529 fluorescence detector, linked to an autosampler (AS 2055 Plus).

Data analysis

Data were processed with Graphpad Prism 6 (Graph Pad Software, San Diego CA, USA). In each case, means $\pm$ standard deviations for *n* number of samples are given. Statistical analyses for comparison of means were carried out using ANOVA, followed by Student's *t* test (significance level, 0.05).

Results

Growth and morphology of *N. oleoabundans* cultivated autotrophically and mixotrophically in a 20-L PBR

Growth kinetics and cell morphology of *N. oleoabundans* cultivated autotrophically and mixotrophically in a 20-L PBR are shown in Fig. 1. Autotrophic cells rapidly entered the exponential phase starting on the 2nd–3rd day up to the 7th–9th day of cultivation (Fig. 1a). After this period, cultures had a short late exponential phase and then entered the stationary phase of growth, reaching a final cell density of about 22×10^6 cells mL^{-1} at the end of the cultivation time (25 days). During the entire period, cells showed the typical morphology of *N. oleoabundans* grown in BM medium, i.e., almost spherical, with a cell diameter of 3–5 μm (Fig. 1b, d). One cup-shaped chloroplast, containing a large pyrenoid, was present and emitted an intense red fluorescence due to the presence of chlorophyll (Fig. 1b, c). On the 25th day of cultivation, cells maintained their normal features and dimensions, but accumulated some lipid globules inside the cytoplasm, as revealed by Nile red staining (Fig. 1d, e).

Fig. 1 a e Growth kinetics (a) and morphology (b e) of *Neochloris oleoabundans* cells autotrophically cultivated in a 20 L PBR in BM medium. Light (b, d) and fluorescence microscope observations (c, e) of *N. oleoabundans* cells after 3 days (b, c) and 25 days (d, e) of autotrophic cultivation. **f j** Growth kinetics (f) and cell morphology (g j) of *N. oleoabundans* cultivated in a 20 L PBR in BM medium under mixotrophic conditions (supplemented with 2.5 g L^{-1} of glucose). Light (g, i) and fluorescence microscope observations (h, j) of *N. oleoabundans* cells after 3 days (g, h) and 8 days (i, j) of mixotrophic cultivation. In both graphs, curves are constructed on a $\log_2$ scale and data are means $\pm$ SD ($n = 3$). In all micrographs, bars = 2 μm

During the first 4 days of mixotrophic growth in the PBR in the presence of glucose, cells showed an evident increase in cell density, reaching values of ca. 33×10^6 cells mL^{-1} (Fig. 1f). Subsequently, cells entered the stationary phase of growth, and then cell density leveled off to values of about 36×10^6 cells mL^{-1} up to the end of the cultivation period, i.e., 8 days. At the beginning of cultivation, both cell shape and size, and chloroplast features in mixotrophic cultures (Fig. 1g, h) were similar to those of autotrophic cultures (Fig. 1b, c). Starting on the 3rd day and up to the end of cultivation, however, mixotrophic cells showed peculiar features. In fact, the chloroplast lost its characteristic cup shape and translucent globules, which tended to occupy almost the entire cell volume, accumulated inside the cytoplasm (Fig. 1i). Nile red staining confirmed the lipidic nature of these droplets (Fig. 1j).

Extracellular FFA composition in E and EG media

The extracellular FFA composition in the exhausted autotrophic (E) and mixotrophic (EG) media of *N. oleoabundans* is shown in Table 1. While EG medium comprised both

Table 1 Extracellular free fatty acids composition in the exhausted E and EG media of *Neochloris oleoabundans* grown in 20 L PBR, after 15 and 8 days of cultivation, respectively

Fatty acids	Abbreviation	E medium	EG medium
Myristic acid	C14:0	2.16 ± 0.09	2.09 ± 0.04
Palmitic acid	C16:0	33.54 ± 0.01	34.43 ± 2.80
Stearic acid	C18:0	64.31 ± 0.10	56.56 ± 2.71
Oleic acid	C18:1ω9	nd	5.93 ± 1.47

Values are expressed as percentage of total fatty acids and are means ± SD ($n = 2$)

saturated and mono-unsaturated FFAs, E medium contained only saturated FFAs. In fact, both samples contained similar percentages of saturated myristic acid (C14:0) and palmitic acid (C16:0), while the percentage of stearic acid (C18:0) was different between the two samples. In E medium, stearic acid was present at about 64 %, while in EG medium at about 56 %; however, C18:0 represented the main saturated FFA for both samples. It is noteworthy that only EG medium contained about 6 % of the monoenoic oleic acid (C18:1ω9).

Polyamine content in E and EG media and in the corresponding *N. oleoabundans* cells cultivated in a 20-L PBR

As shown in Fig. 2a, the PA composition of *N. oleoabundans* cells cultivated mixotrophically in the PBR were characterized by higher putrescine, spermine, and, especially, spermidine levels compared with cells cultivated autotrophically. In fact, Spm, Put, and Spd concentrations in mixotrophic cells were ca. 2, 30, and 57 times higher, respectively, than those found in autotrophic cells ($p < 0.001$). Conversely, the PA composition in E and EG medium was not significantly different. In fact, both media contained similar amounts of Put and Spm. Interestingly, spermidine was present in high concentration in both media (119 ± 16.91 and 101 ± 5.00 pmol mL⁻¹ in E and EG, respectively; Fig. 2b).

Fig. 2 Polyamine concentrations **a** in *Neochloris oleoabundans* cells cultivated in a 20 L PBR autotrophically and mixotrophically at 15 and 8 days of cultivation, respectively, and **b** in the corresponding exhausted

Growth kinetics of *N. oleoabundans* in exhausted and control growth media

As shown in Fig. 3a, the growth of *N. oleoabundans* in E+ medium was promoted for the entire experiment relative to controls grown on BM medium; on the contrary, growth in EG+ medium was similar to that of the C. All samples entered the exponential phase very soon (during the first 3 days of cultivation), despite some differences in growth rates (Fig. 3b). The highest μ values were observed for algae cultivated in the two exhausted media (E+ and EG+, 1.29 and 1.18 div day⁻¹, respectively; control, 0.98 div day⁻¹; $p < 0.01$ in both cases relative to C) (Fig. 3b). Subsequently, from the 3rd day onwards, no relevant differences were observed between C and EG+ samples, as both of them showed similar growth kinetics and cell densities (Fig. 3a). Conversely, E+ samples reached and maintained the highest cell density; the major difference as compared with the other samples was observed starting on the 7th day, with a cell density of about 16×10^6 cells mL⁻¹ (45 and 50 % more than EG+ and C samples, respectively; $p < 0.01$ in both cases relative to E+ samples) at the same time point. After the 17th day, all samples entered the stationary phase, reaching, at the end of the experiment, a final cell density of $25\text{--}30 \times 10^6$ cells mL⁻¹. During the experiments, aliquots of samples were periodically harvested to measure the pH of the culture media. As observed in the Electronic supplementary material (Fig. S1), initial pH was very different between control medium (about 7.0) and exhausted media (about 9.5 in both cases). pH gradually increased up to 9.2 in the control medium, while in E+ and EG+ media, the pH varied between 9.5 and 8.0 without any obvious trend.

Consumption of nitrate and phosphate by *N. oleoabundans* cultivated in exhausted and control growth media

The consumption of nitrate and phosphate by *N. oleoabundans* cells in the course of the experiment is depicted in Fig. 4. At

E and EG growth media. Autotrophic cells (white), mixotrophic cells (dark), E+ (light gray), and EG+ medium (dark gray). Values are means ± SD ($n = 3$)

Fig. 3 **a** Growth kinetics of *Neochloris oleoabundans* in BM medium (filled circles), E+ medium (empty squares), and EG+ medium (empty diamonds). **b** Growth rates, calculated during the exponential phase (0–3 days time interval of cultivation), of cells grown in BM (white), E+ (light gray), and EG+ (dark gray) media. The growth curve is constructed on a log₂ scale, and data are means ± SD (*n* = 3)

time 0 (inoculation), the exhausted replenished growth media contained approximately 2 mM NO₃⁻ and 0.115 mM PO₄²⁻, i.e., the typical concentration of those components in BM medium. By the end of the experiment, cells had consumed practically all nitrate and phosphate present in the growth media. However, nutrient consumption by C, E+, and EG+ algal cultures followed different trends, since C samples maintained higher nitrate and phosphate concentrations than samples in exhausted media. In fact, after 12 days of cultivation, cells grown in E+ and EG+ had consumed about 34 and 68 % of nitrate, respectively, while C only 15 %. Nitrate concentration decreased from 1.43 mM at time 0 for all culture media to 0.94 mM for E+, 0.45 mM for EG+, and 1.21 mM for C media (*p* < 0.001 in both treated samples relative to C; Fig. 4a). Differently, during the first 12 days of growth, phosphate exhibited a dramatic decrease, with a reduction by cells cultivated in EG+ and E+ of 94 and 98 %, respectively, while in C medium it was only about 60 %. In fact, phosphate concentration decreased from about 0.086 mM at time 0 for all culture media

to 0.002 for E+, 0.005 for EG+, and 0.035 mM for C media (*p* < 0.001 in both treated samples relative to C; Fig. 4b).

Photosynthetic pigment content in *N. oleoabundans* cultivated in exhausted and control media

During the experiment, a gradual increase in photosynthetic pigment content was observed in all samples (Fig. 5). Chl *a* in algae grown in C medium showed an evident increase up to the end of the experiment, reaching an average concentration of 0.5 nmol 10⁶ cells (Fig. 5a). A similar increasing trend was also observed in E+ samples, though with values 10–15 % lower than those of controls. From the 17th day of cultivation, Chl *a* content of E+ samples was significantly lower than in C (32 %; *p* < 0.01). Conversely, in EG+ samples, Chl *a* content showed a slightly increasing trend as compared with C and E+ samples; this led to a final value of 0.35 nmol 10⁶ cells at the end of the experiment (32 and 21 % lower than C and E+ samples, respectively; Fig. 5a). An increasing trend in Chl *b* content was also

Fig. 4 Time course variations in the concentrations of nitrate (**a**) and of phosphate (**b**) in culture media of *Neochloris oleoabundans* during 25 days of cultivation on BM medium (white), E+ medium (light gray), and EG+ medium (dark gray). Values are means ± SD (*n* = 3)

observed for all samples, without significant differences between C and E+ samples (Fig. 5b). Similar to Chla content, EG+ cells contained lower quantities of Chlb than C and E+ samples (15–25 and 20–35 % lower than C and E+ samples, respectively). More evident differences were observed for Car content (Fig. 5c). In fact, C samples showed an evident increase up to the end of the experiment, always containing higher quantities compared with the other samples (20–45 and 15–30 % more than E+ and EG+ samples, respectively; $p < 0.05$). It is noteworthy that EG+ and E+ cells shared a similar trend of Car concentration during the experiment (Fig. 5c).

Maximum quantum yield of PSII of *N. oleoabundans* cultivated in exhausted and control media

The variations in PSII maximum quantum yield measured during the experiments are shown in Fig. 6. C, E+, and EG+ samples showed a slight increase of the F_V/F_M ratio during the first 3 days of growth, reaching values of ca. 0.70. Subsequently, C cells maintained stable values around 0.70–0.75, while the F_V/F_M ratio of E+ and EG+ samples decreased drastically down to values below 0.50 at the end of experiment. In EG+ cells, the decrease was dramatic already from the 3rd day of cultivation, reaching the lowest value after 17 days (ca. 0.30 for EG+ vs 0.75 for C and 0.50 for E+; $p < 0.01$ in both cases); thereafter, samples maintained stable values of ca. 0.35 until the end of the experiment. On the contrary, in E+ cells, F_V/F_M started to decrease very strongly only from the 7th day of cultivation until the 17th day (from about 0.70 at day 7 down to 0.50 at day 17). During the subsequent experimental times (21th and 25th day), the F_V/F_M ratio of E+ cells remained stable around 0.45.

Morphological observations of *N. oleoabundans* cultivated in exhausted and control media

Light microscopy and Nile red staining

Light microscopy of both control and treated samples showed that *N. oleoabundans* maintained similar cell morphology and dimensions throughout the experiment (Figs. 7 and 8). In fact, cells were almost spherical with a cell diameter of 3–5 μm . One cup-shaped chloroplast, containing a large pyrenoid, was present inside the cells (Fig. 7a, c, and e). Moreover, the chloroplast emitted an intense red fluorescence due to Chl (Fig. 7b, d, and f). Interestingly, after 25 days of cultivation, while cell size and shape remained substantially unchanged, all algal

Fig. 5 Time course variations of Chla (a), Chlb (b), and Car content (c) in *Neochloris oleoabundans* cells grown in BM (filled circles), E+ (empty squares) and EG+ (empty diamonds) media during the 25 days of cultivation. Values are means $\pm$ SD ($n = 3$)

Fig. 6 Time course variations of PSII maximum quantum yield (F_v/F_m ratio) in *Neochloris oleoabundans* cells grown in BM (filled circles), E+ (empty squares), and EG+ (empty diamonds) media during the 25 days of cultivation. Values are means $\pm$ SD ($n = 3$)

samples showed some translucent granulation at the cytoplasmic level (Fig. 8a, c, and e). In order to understand the nature of those granulations and to investigate if the recycled growth media could promote the production of lipid globules, all samples were periodically stained with the lipid-specific fluorochrome Nile red. The reaction was positive only at the end of the experiment, and the translucent globules were then unequivocally identified as lipid droplets (Fig. 8b, d, and f).

TEM observations

To investigate the morphological and cytological changes induced by cultivation in recycled growth media, the ultrastructure of C, E+, and EG+ samples was observed by TEM. At 12 days of cultivation, most of the C cell volume was occupied by a characteristic cup-shaped chloroplast. Inside the plastid, one large pyrenoid, which was crossed by two elongated and appressed thylakoids, was present (Fig. 9a). In particular, the organelle contained starch in the shape of a shell around the pyrenoid and showed the typical thylakoid organization (Fig. 9b). Typically featured chloroplasts, as described for cells grown in C, were observed in cells grown in E+ medium (Fig. 9c); however, thylakoids were more appressed than in C (Fig. 9d). EG+ samples showed a more strongly altered chloroplast structure (Fig. 9e), as compared with C and E+ samples. Photosynthetic membranes showed different degrees of thylakoid appression: some portions of thylakoid membranes were appressed while others were loose and sometimes swollen (Fig. 9f, g). In addition, large portions of the stroma were free of thylakoids and some plastoglobules in proximity of the thylakoid membranes were also visible (Fig. 9g, h). Finally, in EG+ samples, the pyrenoid lost its round shape and appeared malformed (Fig. 9e).

Fig. 7 Light and fluorescence microscopy observations of *Neochloris oleoabundans* cells at 12 days of cultivation. **a** Control cells and **b** the relative fluorescence of the chloroplast; **c** cell grown in E+ medium and **d** the relative fluorescence of the chloroplast; **e** cell grown in EG+ medium and **f** the relative fluorescence of the chloroplast. In all micrographs, bars = 2 μ m

Discussion

The recycling of culture medium has been proposed as a possible solution in order to reduce water consumption for algal cultivation, thereby making the process more economically feasible and environmentally sustainable (Yang et al. 2011; Hadj-Romdhane et al. 2013; Farooq et al. 2014). Yang et al. (2011) estimated that the large-scale cultivation of the microalga *Chlorella vulgaris* in recycled culture medium could reduce water use by about 84 %. Present results clearly suggest that *N. oleoabundans* can also efficiently grow in exhausted growth media, especially in the autotrophic medium replenished with nitrate and phosphate (E+). Observing the consumption trend of these main nutrients (Fig. 4), it is clear that in the exhausted media their concentration decreased faster with respect to controls in parallel to the enhanced cell growth. Moreover, phosphate was consumed faster than nitrate by all samples, probably because of its 100 times lower concentration with respect to the latter nutrient (0.2 g L⁻¹ of nitrate vs 0.002 g L⁻¹ of phosphate; Baldisserotto et al. 2012). Thus, in order to make the use of a recycled culture medium feasible, exhausted media should be replenished with

Fig. 8 Light and epifluorescence pictures of *Neochloris oleoabundans* cells after 25 days of growth. **a** Control cells and **b** relative Nile red staining observation, **c** cells grown in E+ medium and **d** relative Nile red staining observation, and **e** cells grown in E+ medium and **f** relative Nile red staining observation in EG+ medium. Translucent globules are indicated with *arrows*. In all micrographs, bars = 2 μm

the correct ratio of nitrate and phosphate, as they are the main nutrients that guarantee cell growth (Stephens et al. 2010). This aspect was preliminarily verified by employing unmodified exhausted growth media (E and EG) to recultivate *N. oleoabundans*. In fact, cells showed a slight growth, reaching, at the end of the cultivation period, low cell densities (ca. 11 vs 32×10^6 cells mL^{-1} of C cultures) (data not shown). These cells accumulated intracellular lipids throughout the experiment (data not shown) as a consequence of the limitation of nitrate and phosphate (Mata et al. 2010; Popovich et al. 2012). These results are consistent with those of Zhu et al. (2013), who found that *C. zoofingensis* cultivated in recycled medium, without nitrogen and phosphorus, displayed enhanced lipid production compared with cultures with full nutrients.

Fig. 9 Transmission electron micrographs of *Neochloris oleoabundans* cells at 12 days of cultivation. **a** Cell grown in BM medium and **b** detail of its chloroplast showing the typical thylakoid membranes organization (*white arrow*). **c** Cell grown in E+ medium and **d** detail of its chloroplast with quite compact and appressed (*white arrow*) thylakoids. **e** Cell grown in EG+ medium and **f** **h** details of its chloroplast. The presence of intermediate stages of thylakoid membranes are evident in EG+ cells, showing a general disorganization of thylakoid membranes (**f** **h**), which appeared wavy, loose, and sometimes swollen (**f**, **g**). **h** Some plastoglobules in proximity of thylakoid membranes are also observed (*arrowheads*). P pyrenoid, N nucleus, M mitochondrion. Bars, 0.5 μm (**a**, **c**, **e**) and 0.05 μm (**b**, **d**, **f**, **g**, **h**)

In order to understand if the growth promotion of *N. oleoabundans* cultivated in E+ and EG+, relative to C, was influenced by some molecules released from the algae,

a characterization of autotrophic (E) and mixotrophic (EG) media was performed. First of all, pH of the control and exhausted media was monitored. In particular, an increasing trend was observed in control, which reached the pH of the exhausted media at the end of the experiment. This was not surprising, considering that during the microalgal growth, pH usually increases because of the consumption of CO₂ during the photosynthetic process (Zhang et al. 2014). However, considering that in E⁺ and EG⁺ pH fluctuated around their initial values, or at least gradually decreased, it is not possible to attribute to pH any evident role in the cell growth promotion. Moheimani and Borowitzka (2006) and Stephens et al. (2010) reported that molecules released from cells can positively/negatively alter cell metabolism and biomass production.

Here we show that both exhausted media contained the main PAs (putrescine, spermidine, and spermine). It is known that these plant growth regulators are involved in a great variety of developmental processes in plant cells, e.g., cell division and protein synthesis (Kaur-Sawhney et al. 2003; Kuznetsova et al. 2006). PA biosynthetic pathways have also been studied in green algae (Cohen et al. 1984; Theiss et al. 2002; Fuell et al. 2010). Although algae produce “unusual” PAs, such as homospermidine and thermospermine (Hamana et al. 2013), the more common PAs (Put, Spd, and Spm), when added to the media, promoted growth and metabolism in *C. vulgaris* (Czerpak et al. 2003). PAs are also known to alleviate the effect of biotic and abiotic stress in plants as well as algae (Tate et al. 2013).

In the present work, a very strong increase in Put and, especially, Spd concentrations was observed in *N. oleoabundans* mixotrophic cells as compared with the autotrophic ones, both cultivated in a 20-L PBR. The growth-promoting role of PAs could explain the faster (0–4 days) and stronger cell density increase (ca. 33×10^6 cell mL⁻¹) in mixotrophic cultures relative to the autotrophic ones (0–9 days; ca. 22×10^6 cell mL⁻¹). The higher PA concentrations in mixotrophically grown cells also suggest that, under these culture conditions, algal cells may be better protected from stress-inducing factors, e.g., bacterial contamination. To our knowledge, the presence of PAs released into algal growth medium in PBRs has not been documented before. The release of PAs (predominantly Spd) from autotrophic and mixotrophic cells into their respective (E and EG) exhausted media was not significantly different. However, the presence of these plant growth regulators seems to contribute to the promotion of *N. oleoabundans* growth in both media replenished with nitrate and phosphate. In fact, during the entire experiment, a higher cell density was observed in E⁺ relative to C medium. Moreover, the EG⁺ samples showed a promotion of cell density, albeit similar to that of C.

Both autotrophic and mixotrophic exhausted media also contained FFAs, which could have induced a change in the normal metabolism of *N. oleoabundans*. This assumption is

demonstrated by the more rapid consumption of nitrate and phosphate, the decreased photosynthetic pigments content and the strong decline in the F_V/F_M ratio observed in cells grown in E⁺ and EG⁺ media. These results corroborate the assumption that microalgae release metabolites in high cell density cultures (Richmond 2004) or as a consequence of stressful conditions (Ikawa 2004; Wu et al. 2006). Even if the role of FFAs is currently under debate (Ikawa 2004; Stephens et al. 2010), recent studies showed that they can strongly inhibit growth or exert cytotoxic effects on microalgae (Kogteva and Bezuglov 1998; Wu et al. 2006; Bosma et al. 2008; Stephens et al. 2010). In fact, the presence of these metabolites, and their oxidative products, negatively affect biomass productivity, especially when the microalgae are cultivated in recycled medium (Lívanský et al. 1996; Rodolfi et al. 2003). Granum et al. (2002) reported that the accumulation of intracellular lipids in microalgae enhanced the release of FFAs into the culture medium, and Harun et al. (2010) observed that this release was caused by cell lysis.

In this work, FFA composition in autotrophic (E) and mixotrophic (EG) media after *N. oleoabundans* cultivation was similar. The only main difference was represented by the presence of the monoenoic oleic acid (C18:1 ω 9) in EG medium. Oleic acid is one of the major lipid components in lipid-enriched *N. oleoabundans* grown under N-stress conditions (Popovich et al. 2012). It is reported that mixotrophic cultivation alters N/C ratio, inducing similar lipid production as N-depleted autotrophic cultures (Giovanardi et al. 2014). Probably for this reason, the release of oleic acid occurred only in the mixotrophic growth medium. Moreover, Wu et al. (2006) observed an altered plasma membrane permeability due to the toxic effects of FFAs in two *Chlorophyta* (*C. vulgaris* and *Monoraphidium contortum*) and in a cyanobacterium (*Anabaena* P-9). Then, in addition to the lower concentration with respect to nitrate in BM medium, the rapid consumption of phosphate in both exhausted media could be linked also to an alteration in membrane permeability, due to the activity of FFAs (Ikawa 2004; Wu et al. 2006). In addition, FFAs can cause inhibition of the PSII and PSI electron transport chains (Siegenthaler 1973) and disorganization of thylakoids (Wu et al. 2006). Indeed, *N. oleoabundans* cells in E⁺ and EG⁺ were characterized by alterations of the photosynthetic apparatus. These cells, in fact, contained less photosynthetic pigments than C cells and exhibited a drastic decrease of the PSII maximum quantum yield (F_V/F_M). These variations are linked to an alteration of photosynthetic efficiency, especially as regards PSII (White et al. 2011). The decreased photosynthetic pigment contents, observed in cells grown in E⁺ and EG⁺, also reflected the changes in thylakoid membrane arrangement (Baldiasserotto et al. 2012). Therefore, correct thylakoid organization is necessary to

maintain optimal photosynthetic activity, and this is often influenced by culture conditions (Nevo et al. 2012). In addition, the presence of some plastoglobules in chloroplasts of algal cells, grown on exhausted media, could be another indicator of an alteration of photosynthetic membranes (Besagni and Kessler 2013). On the contrary, *N. oleoabundans* cells grown in C medium showed stable and normal F_v/F_m values throughout the experiment (Giovanardi et al. 2014), and the typical assembly of thylakoid membranes (Nevo et al. 2012; Baldisserotto et al. 2012; Giovanardi et al. 2014).

The presence of lipid droplets, observed only at the end of the experiment in C, E+, and EG+ cells, is probably related to aging of the microalgal cultures (Hu et al. 2008; Baldisserotto et al. 2012), rather than to the use of recycled culture media.

In conclusion, results presented here demonstrate that recycling autotrophic and mixotrophic growth media is a suitable solution to obtain high cell density cultures of the microalga *N. oleoabundans*. For this reason, they represent a contribution for improving the scale-up of microalgal cultivation while providing a more sustainable ecological impact on water resources. However, further studies are needed to deepen knowledge on the nature and the specific role of some molecules that are released into the growth media, in order to obtain useful information for the advancement of the biotechnological use of this strain.

Acknowledgments This work was financially supported by grants from the University of Ferrara. We are grateful to Dr. Immacolata Maresca of the University of Ferrara for technical assistance in freeze drying of samples.

Conflict of interest The authors declare they have no conflict of interest.

References

- Baldisserotto C, Ferroni L, Giovanardi M, Pantaleoni L, Boccaletti L, Pancaldi S (2012) Salinity promotes growth of freshwater *Neochloris oleoabundans* UTEX 1185 (*Sphaeropleales*, *Neochloridaceae*): morpho physiological aspects. *Phycologia* 51: 700–710
- Baldisserotto C, Giovanardi M, Ferroni L, Pancaldi S (2014) Growth, morphology and photosynthetic responses of *Neochloris oleoabundans* during cultivation in a mixotrophic brackish medium and subsequent starvation. *Acta Physiol Plant* 36:461–472
- Batan L, Quinn CJ, Bradley TH (2013) Analysis of water footprint of a photobioreactor microalgae biofuel production system from blue, green and lifecycle perspectives. *Algal Research* 2:196–203
- Besagni C, Kessler F (2013) A mechanism implicating plastoglobules in thylakoid disassembly during senescence and nitrogen starvation. *Planta* 237:463–470
- Borowitzka MA (2005) Culturing microalgae in outdoor ponds. In: Anderson RA (ed) *Algal culturing techniques*. Elsevier Academic, London, pp. 205–218
- Borowitzka MA, Moheimani NR (2013) Sustainable biofuels from algae. *Mitig Adapt Strateg Glob Change* 18:13–25
- Bosma R, Miazek K, Willemsen SM, Vermuë MH, Wijffels RH (2008) Growth inhibition of *Monodus subterraneus* by free fatty acids. *Biotechnol Bioeng* 101:1108–1114
- Chisti Y (2007) Biodiesel from microalgae. *Biotech Adv* 25:294–306
- Czerpak R, Bajguz A, Pietrowska A, Dobrogowska R, Matejczyk M, Weislawski W (2003) Biochemical activity of di- and polyamines in the green alga *Chlorella vulgaris* Beijerinck (*Chlorophyceae*). *Acta Soc Bot Pol* 72:19–24
- Cohen E, Arad SM, Heimer YH, Mizrahi Y (1984) Polyamine biosynthetic enzymes in the cell cycle of *Chlorella*. *Plant Physiol* 74:385–388
- Farooq W, Suh WI, Park MS, Yang J (2014) Water use and its recycling in microalgal cultivation for biofuel application. *Bioresour Technol*. doi:10.1016/j.biortech.2014.10.140
- Ferroni L, Baldisserotto C, Giovanardi M, Pantaleoni L, Morosinotto T, Pancaldi S (2011) Revised assignment of room temperature chlorophyll fluorescence emission bands in single living cells of *Chlamydomonas reinhardtii*. *J Bioenerg Biomembr* 43:163–173
- Fuell C, Elliott KA, Hanfrey CC, Franceschetti M, Michael AJ (2010) Polyamine biosynthetic diversity in plants and algae. *Plant Physiol Biochem* 48:513–520
- Giovanardi M, Ferroni L, Baldisserotto C, Tedeschi P, Maietti A, Pantaleoni L, Pancaldi S (2013) Morpho physiological analyses of *Neochloris oleoabundans* (*Chlorophyta*) grown mixotrophically in a carbon rich waste product. *Protoplasma* 250:161–174
- Giovanardi M, Baldisserotto C, Ferroni L, Longoni P, Cella R, Pancaldi S (2014) Growth and lipid synthesis promotion in mixotrophic *Neochloris oleoabundans* (*Chlorophyta*) cultivated with glucose. *Protoplasma* 251:115–125
- Gouveia L, Marques AE, Lopes da Silva T, Reis A (2009) *Neochloris oleoabundans* UTEX #1185: a suitable renewable lipid source for biofuel production. *J Ind Microbiol Biotechnol* 36:821–826
- Granum E, Kirkvold S, Mykkestad SM (2002) Cellular and extracellular production of carbohydrates and amino acids by the marine diatom *Skeletonema costatum*: diel variations and effects of N depletion. *Mar Ecol Prog Ser* 242:83–94
- Grobbelaar JU (2009) Factors governing algal growth in photobioreactors: the “open” versus “closed” debate. *J Appl Phycol* 21:489–492
- Hadj Romdhane F, Jaouen P, Pruvost J, Grizeau D, Van Vooren G, Bourseau P (2012) Development and validation of a minimal growth medium for recycling *Chlorella vulgaris* culture. *Bioresour Technol* 123:366–374
- Hadj Romdhane F, Zheng X, Jaouen P, Pruvost J, Grizeau D, Croué JP, Bourseau P (2013) The culture of *Chlorella vulgaris* in a recycled supernatant: effects on biomass production and medium quality. *Bioresour Technol* 132:285–292
- Hamana K, Niitsu M, Hayashi H (2013) Occurrence of homospermidine and thermospermine as a cellular polyamine in unicellular chlorophyte and multicellular charophyte green algae. *J Gen Appl Microbiol* 59:313–319
- Harun R, Singh M, Forde GM, Danquah MK (2010) Bioprocess engineering of microalgae to produce a variety of consumer products. *Renew Sust Energ Rev* 14:1037–1047
- Hu Q, Sommerfeld M, Jarvis E, Ghirardi M, Posewitz M, Seibert M, Darzins A (2008) Microalgal triacylglycerols as a feedstocks for biofuel production: perspective and advances. *Plant J* 54:621–639
- Ikawa M (2004) Algal polyunsaturated fatty acids and effects on plankton ecology and other organism. *UNH Center for Freshwater Biology Research* 6:17–44
- Kalaji HM, Schansker G, Ladle RJ, Vasilij Goltsev V, Bosa K, Allakhverdiev SI, Brestic M, Bussotti F, Calatayud A, Dąbrowski P, Elsheery NI, Ferroni L, Guidi L, Hogewoning SW, Jajoo A, Misra AN, Nebauer SG, Pancaldi S, Penella C, Poli D, Pollastrini M,

- Romanowska Duda ZB, Rutkowska B, Serôdio J, Suresh K, Szulc W, Tambussi E, Yannicari M, Zivcak M (2014) Frequently asked questions about in vivo chlorophyll fluorescence: practical issues. *Photosynth Res* 122:121–158
- Kaur Sawhney R, Tiburcio AF, Altabella T, Galston AW (2003) Polyamines in plants: an overview. *J Cell Mol Biol* 2:1–12
- Kogteva GS, Bezuglov VV (1998) Unsaturated fatty acids as endogenous bioregulators. *Biochem Mosc* 63:4–12
- Lam MK, Lee KT (2012) Potential of using organic fertilizer to cultivate *Chlorella vulgaris* for biodiesel production. *Appl Energy* 94:303–308
- Li Y, Horsman M, Wang B, Wu N, Lan CQ (2008) Effect on nitrogen sources on cell growth and lipid accumulation of green alga *Neochloris oleoabundans*. *Appl Microbiol Biotechnol* 81:629–636
- Lichtenthaler HK, Buschmann C, Knapp M (2005) How to correctly determine the different chlorophyll fluorescence parameters and the chlorophyll fluorescence decrease ratio RFD of leaves with the PAM fluorometer. *Photosynthetica* 43:379–393
- Livanský K, Dědič K, Bínová J, Tichý V, Novotný P, Doucha J (1996) Influence of the nutrient solution recycling on the productivity of *Scenedesmus obliquus*, utilization of nutrients and water in outdoor cultures. *Algological Studies/Archiv für Hydrobiologie* 81:105–113
- Mata TM, Martins AA, Caetano NS (2010) Microalgae for biodiesel production and other applications: a review. *Renew Sust Energy Rev* 14:217–232
- Moheimani NR, Borowitzka MA (2006) (2006) the long term culture of the coccolithophore *Pleurochrysis carterae* (*Haptophyta*) in outdoor raceway ponds. *J Appl Phycol* 18:703–712
- Molina Grima E, Belarbi H, Acién Fernández FG, Robles Medina A, Chisti Y (2003) Recovery of microalgal biomass and metabolites: process options and economics. *Biotech Adv* 20:491–515
- Ndimba BK, Ndimba RJ, Johnson TS, Waditee Sirisattha R, Baba M, Sirisattha S, Shiraiwa Y, Rakwal R, Agrawal GK, Rakwal R (2013) Biofuels as a sustainable energy source: an update of the applications of proteomics in bioenergy crops and algae. *J Prot* 93:234–244
- Nevo R, Charuni D, Tsabari O, Reich Z (2012) Composition, architecture and dynamics of the photosynthetic apparatus in higher plants. *Plant J* 70:157–176
- Pancaldi S, Baldisserotto C, Ferroni L, Bonora A, Fasulo MP (2002) Room temperature microspectrofluorimetry as a useful tool for studying the assembly of the PSII chlorophyll protein complexes in single living cells of etiolated *Euglena gracilis* Klebs during the greening process. *J Exp Bot* 53:1753–1763
- Popovich CA, Damiani MC, Constenla D, Martínez AM, Giovanardi M, Pancaldi S, Leonardi PI (2012) *Neochloris oleoabundans* grown in natural enriched seawater for biodiesel feedstock: evaluation of its growth and biochemical composition. *Bioresource Technol* 114:287–293
- Pruvost J, Van Vooren G, Cogne G, Legrand J (2009) Investigation of biomass and lipids production with *Neochloris oleoabundans* in photobioreactor. *Bioresource Technol* 100:5988–5995
- Pulz O (2001) Photobioreactors: production systems for phototrophic microorganisms. *Appl Microbiol Biotechnol* 57:287–293
- Richmond A (2004) Handbook of microalgal culture: biotechnology and applied phycology. Blackwell Sciences Ltd., Oxford
- Rodolfi L, Zittelli GC, Barsanti L, Rosati G, Tredici MR (2003) Growth medium recycling in *Nannochloropsis* sp. mass cultivation. *Biomol Eng* 20:243–248
- Scaramagli S, Biondi S, Capitani F, Gerola P, Altamura MM, Torrigiani P (1999) Polyamine conjugate levels and ethylene biosynthesis: in verse relationship with vegetative bud formation in tobacco thin layers. *Physiol Plant* 105:366–375
- Sheehan JJ (2009) Biofuels and the conundrum of sustainability. *Curr Opin Biotechnol* 20:318–324
- Siegenthaler PA (1973) Change in pH dependence and sequential inhibition of photosynthetic activity in chloroplasts by unsaturated fatty acids. *Biochim Biophys Acta* 305:153–162
- Smith VH, Sturm BSM, deNoyelles FJ, Billings SA (2010) The ecology of algal biodiesel production. *Trends Ecol Evo* 25:301–309
- Solomon BD (2010) Biofuels and sustainability. *Ann N Y Acad Sci* 1185:119–134
- Stephens E, Ross IL, Mussgnug JH, Wagner LD, Borowitzka MA PC, Kruse O, Hankamer B (2010) Future prospects of microalgal biofuel production systems. *Trends Plant Sci* 15:554–564
- Tate JJ, Gutierrez Wing MT, Rush KA, Benton MG (2013) The effects of plant growth substances and mixed cultures on growth and metabolite production of green algae *Chlorella* sp.: a review. *J Plant Growth Regul* 32:417–428
- Tomabene TG, Holzer G, Lien S, Burris N (1983) Lipid composition of the nitrogen starved green alga *Neochloris oleoabundans*. *Enzyme Microbiol Tech* 5:435–440
- Theiss C, Bohley P, Voigt J (2002) Regulation by polyamines of ornithine decarboxylase activity and cell division in the unicellular green alga *Chlamydomonas reinhardtii*. *Plant Physiol* 128:1470–1479
- Wellburn AR (1994) The spectral determination of chlorophylls a and b, as well as total carotenoids, using various solvents with spectrophotometer of different resolution. *J Plant Physiol* 144:307–313
- White S, Anandraj A, Bux F (2011) PAM fluorometry as a tool to assess microalgal nutrient stress and monitor cellular neutral lipids. *Bioresource Technol* 102:1675–1682
- Wu JT, Chiang YR, Huang WY, Jane WN (2006) Cytotoxic effects of free fatty acids on phytoplankton algae and cyanobacteria. *Aquat Toxicol* 80:338–345
- Yang J, Xu M, Zhang X, Hu Q, Sommerfeld M, Chen Y (2011) Lifecycle analysis on biodiesel production from microalgae: water footprint and nutrients balance. *Bioresource Technol* 102:159–165
- Zhang Q, Wang T, Hong Y (2014) Investigation of initial pH effects on growth of an oleaginous microalgae *Chlorella* sp. HQ for lipid and nutrient uptake. *Water Sci Technol* 70:712–719
- Zhu LD, Takala J, Hiltunen E, Wang ZM (2013) Recycling harvest water to cultivate *Chlorella zoofingensis* under nutrient limitation for biodiesel production. *Bioresource Technol* 144:14–20