

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Characterization of alkaline phosphatase activity in seminal plasma and in fresh and frozen-thawed stallion spermatozoa

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Bucci, D., Giaretta, E., Spinaci, M., Rizzato, G., Isani, G., Mislei, B., et al. (2016). Characterization of alkaline phosphatase activity in seminal plasma and in fresh and frozen-thawed stallion spermatozoa. THERIOGENOLOGY, 85(2), 288-295 [10.1016/j.theriogenology.2015.09.007].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/541780> since: 2019-07-17

*Published:*

DOI: <http://doi.org/10.1016/j.theriogenology.2015.09.007>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Bucci D, Giaretta E, Spinaci M, Rizzato G, Isani G, Mislei B, Mari G, Tamanini C, Galeati G. Characterization of alkaline phosphatase activity in seminal plasma and in fresh and frozen-thawed stallion spermatozoa. *Theriogenology*. 2016 Jan 15;85(2):288-295.e2.

The final published version is available online at:

<https://doi.org/10.1016/j.theriogenology.2015.09.007>

© [2016]. This manuscript version is made available under the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) 4.0 International License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

**Characterization of alkaline phosphatase activity in seminal plasma and in fresh and frozen-thawed stallion spermatozoa.**

Diego Bucci<sup>1\*</sup>; Elisa Giaretta<sup>1</sup>, Marcella Spinaci<sup>1</sup>, Giovanni Rizzato<sup>2</sup>, Gloria Isani<sup>1</sup>, Beatrice Mislei<sup>2</sup>, Gaetano Mari<sup>1,2</sup>, Carlo Tamanini<sup>1</sup>, Giovanna Galeati<sup>1</sup>.

<sup>1</sup> DIMEVET Department of Veterinary Medical Sciences, Via Tolara di sopra, 50, 40064 Ozzano dell'Emilia, BO, Italy; <sup>2</sup>INFA National institute of assisted reproduction, Via Gandolfi 16, 40057 Cadriano, BO Italy.

\*Corresponding Author: Dr Diego Bucci, Via Tolara di sopra, 50, 40064 Ozzano dell'Emilia, BO; mail [diego.bucci3@unibo.it](mailto:diego.bucci3@unibo.it). Tel 00390512097904.

## **Abstract**

Alkaline phosphatase (AP) has been studied in several situations to elucidate its role in reproductive biology of the male from different mammalian species; at present its role in horse sperm physiology is not clear. The aim of the present work was to measure AP activity in seminal plasma and sperm extracts from freshly ejaculated as well as in frozen-thawed stallion spermatozoa and to verify whether relationship exists between AP activity and sperm quality parameters.

Our data on 40 freshly ejaculated samples from 10 different stallions demonstrate that the main source of AP activity is seminal plasma, while sperm extracts contribution is very low. In addition we demonstrated that AP activity at physiological pH (7.0) is significantly lower than that observed at pH 8.0, including the optimal AP pH (pH 10.0). AP did not exert any effect on sperm oocyte interaction assessed by heterologous oocyte binding assay. Additionally, we observed a thermal stability of seminal plasma AP, concluding it is similar to that of bone isoforms. Positive correlations were found between seminal plasma AP activity and sperm concentration, while a negative correlation was present between both spermatozoa extracts and seminal plasma AP activity and seminal plasma protein content.

A significant decrease in sperm extract AP activity was found in frozen thawed samples compared with freshly ejaculated ones (n=21), concomitantly with the decrease in sperm quality parameters. The positive correlation between seminal plasma AP activity measured at pH 10 and viability of frozen-thawed spermatozoa suggests that seminal plasma AP activity could be utilized as an additional predictive parameter for stallion sperm freezability.

In conclusion, we provide some insights in AP activity in both seminal plasma and sperm extracts and describe a decrease in AP after freezing and thawing.

## **Keywords**

Horse sperm  
Alkaline phosphatase  
Seminal plasma  
Sperm extracts  
Freezing and thawing

## **1. Introduction**

47

48 Alkaline Phosphatase (AP) is an enzyme that catalyzes the detachment of phosphate groups from  
49 several substrates [1]; it is present in male genital tract fluids and its activity has been revealed in  
50 semen from various mammalian species [2]. The presence of AP in male genital secretions suggests  
51 it could have a role in mammalian reproduction and in particular in sperm metabolism [3], even if a  
52 definite, clear conclusion has not been reached so far..

53 In bull seminal plasma the activity of the enzyme has been correlated to fertility and sperm  
54 concentration [4], while in stallion it resulted a promising tool to determine whether ejaculation  
55 failure is due to either azoospermia or a blockage along the genital tract [5] and it was linked to sperm  
56 quality [6].

57 Other recent studies on AP in stallion seminal plasma furnish data on the different levels of activity  
58 in separated semen fractions [7], as well as on its possible role as an indicator of sperm longevity in  
59 fractionated semen [8].

60 The above-mentioned studies evaluated the activity in seminal plasma after removing spermatozoa,  
61 or in the whole ejaculate, thus indirectly giving information on AP activity of the sperm cells [5]. In  
62 a recent study on boar semen [9] we measured the activity of AP from spermatozoa as well as the  
63 functional change it undergoes after sperm capacitation; the overall conclusion was that it could play  
64 a role in the control of sperm function. There are no data on AP activity in sperm extracts after  
65 freezing-thawing process, excepting for those on frozen ram spermatozoa by Salamon and Maxwell  
66 [10] who reported a loss in phosphatase activity after freezing and thawing, thus indicating that the  
67 enzyme activity could be used as an additional parameter for detecting cryoinjuries.

68 Based on the questions raised by the aforementioned studies, the present work was aimed at:  
69 describing AP kinematic properties (thermal inactivation test) in stallion seminal plasma.

70

71 determining the possible correlations between AP and sperm quality parameters and sperm function  
72 as well as its possible predictive role in freezability of stallion semen.

73

74

## 75 **2. Materials and methods**

76

77 The experiment was approved by the Ethic-scientific Committee of Alma Mater Studiorum,  
78 University of Bologna

79 All the reagents were obtained from Sigma Chemical Co. (St. Louis, MO, USA), unless otherwise  
80 specified.

81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114

*2.1. Semen collection and preparation*

Forty ejaculates from 10 fertile stallions (9 trotter breed and one Connemara), aging from 5 to 25 years, were sampled. Stallions were housed individually and semen was sampled between February and April 2014 by an artificial vagina (Missouri model) equipped with a disposable liner with an in line filter; stallions were collected on a phantom and in presence of an oestrous mare.

Semen concentration was evaluated using a hemocytometer chamber (Thoma).

Two aliquots of semen were kept undiluted: the first one was used to obtain seminal plasma, the second one to obtain spermatozoa for alkaline phosphatase activity assay. A third aliquot of semen was diluted in Kenney extender at a concentration of  $40 \times 10^6$  spermatozoa /mL for semen quality analysis (sperm motility, viability, mitochondrial membrane integrity).

Twenty-one ejaculates, collected in the same period of time from four of the aforementioned stallions, were used for analysis of AP activity in frozen-thawed spermatozoa in comparison with fresh ones. Part of the ejaculates was immediately diluted in Kenney extender for subsequent freezing procedure, while part was kept undiluted to obtain seminal plasma and spermatozoa for AP activity determination.

*2.2. AP activity in seminal plasma, freshly ejaculated and frozen-thawed spermatozoa*

Protein concentration in seminal plasma and sperm extracts was measured by Bradford's method [11] using bovine serum albumin as the protein standard.

Seminal plasma was obtained by two subsequent centrifugations of the undiluted sperm aliquot at  $12,000 \times g$  for 15 min at 4°C. The resultant supernatant was observed at the microscope to check the absence of either spermatozoa or their fragments.

Sperm cells ( $2 \times 10^9$  spz/mL) (both freshly ejaculated and frozen-thawed) were washed twice in PBS ( $900 \times g$  for 3 min at RT) Subsequently the sperm pellet was sonicated in PBS and subsequently centrifuged at  $12,000 \times g$  for 15 min at 4°C; finally the pellet was discarded and the supernatant analyzed for AP activity.

The enzyme activity was assayed as described in [9]. Briefly, AP activity was measured by a spectrophotometric assay that monitors the absorbance change at 405 nm as para-nitrophenylphosphate (pNPP, colourless) is converted to para-nitrophenol (yellow). AP was assayed in 50 mM TrisHCl buffer, at 25°C at different pH: 7.0, 8.0 and 10.0. Fifty  $\mu$ L of seminal plasma or sperm extracts were used and the absorbance at 405 nm was measured during 1 min and 30 min

115 respectively before the addition of pNPP to evaluate the non enzymatic variations of absorbance; no  
116 interferences were recorded in any case. In the case of seminal plasma analyzed at pH 10.0, it was  
117 diluted 1:10 in Tris buffer. The reaction was started by the addition of 10 mM pNPP and each  
118 measurement was run in duplicate and averaged. Enzyme activity was expressed as specific activity,  
119 nmol/min/mg of protein.

120 In order to better characterize the seminal plasma isoform, AP thermo-inhibition was carried out at  
121 60°C, as described by [15] with some modifications. The samples were kept at 60°C for 0'', 15'',  
122 30'', 1', 1'15'', 1'30'', 1'45'', 2', 2'30'', 3' then placed in ice and the activity was measured within  
123 30'' at pH 8.0. The same assay was also performed on the thermo-stable AP from bovine intestinal  
124 mucosa as a control.

125

### 126 *2.3. Evaluation of sperm viability, motility and mitochondrial activity*

127

128 Twenty-five µL of semen were incubated with 2 µL of a 300 µM solution of propidium iodide (PI),  
129 and 2 µL of a 10 µM solution of SYBR green-14, obtained from the live/dead sperm viability kit  
130 (Molecular Probes, Inc., Eugene, OR, USA) for 5 min at 37 °C in the dark. Ten µL of the stained  
131 suspensions were placed on clean microscope slides, carefully overlaid with coverslips, and at least  
132 200 spermatozoa per sample were observed under a Nikon Eclipse E 600 epifluorescence microscope  
133 (Nikon Europe BV, Badhoevedop, The Netherlands). Spermatozoa stained with SYBR-14 but not  
134 with PI were considered as viable (SYBR-14- PI-). Spermatozoa both SYBR-14+ and PI+ and those  
135 SYBR-14-/PI+ were considered with damaged membranes or dead.

136 For each sample, an aliquot (25 µL) of semen was incubated with 2 µL of a 300 µM propidium iodide  
137 (PI) stock solution, 2 µL of a 10 µM SYBR green-14 stock solution, both obtained from the live/dead  
138 sperm viability kit (Molecular Probes, Inc.) and 2 µL of a 150 µM JC-1 solution, for 20 min at 37°C  
139 in the dark. Ten µL of the sperm suspension were then placed on a slide, and at least 200 spermatozoa  
140 per sample were scored using the above described microscope. Spermatozoa stained with SYBR  
141 green-14 and not stained with PI were considered as viable. Spermatozoa SYBR positive and PI  
142 positive and those SYBR negative / PI positive were considered as cells with non-intact membrane  
143 or dead. JC-1 monomers emit a green fluorescence in mitochondria with low membrane potential,  
144 while emitting a bright red-orange fluorescence in case of multimer formation (J-aggregates) in  
145 mitochondria with high membrane potential. When an orange fluorescence was present in the mid  
146 piece, live spermatozoa were considered to have functional active mitochondria (SYBR+/PI-/JC-1+).  
147 Motility was measured using a computer-assisted sperm analysis system (CASA, Hamilton Thorne,  
148 IVOS Ver. 12, standard equine set up). Thousand-1000 cells were evaluated on each sample diluted

149 to  $30 \times 10^6$  sperm/mL using a fixed height Leja Chamber SC 20-01-04-B,(Leja, The Netherlands).  
150 Sperm motility endpoints assessed were: percent of total motile spermatozoa (TM), percent of  
151 progressive spermatozoa (PM), curvilinear velocity (VCL) and mean velocity (VAP). The setting  
152 parameters of the program were the followings: frames per second 60, number of frames 45, threshold  
153 path velocity 30 microns/sec, straightness threshold 50.

154

#### 155 *2.4. Effect of AP on heterologous oocyte binding*

156

157 Porcine oocytes maturation was performed as reported by [12]. Briefly, ovaries were obtained from  
158 pre-pubertal gilts at a local abbatoir and transported to the laboratory within 1 h. Cumulus–oocyte  
159 complexes (COCs) were aspirated from 4 to 6 mm follicles using a 18 gauge needle attached to a 10  
160 mL disposable syringe . Under a stereomicroscope, intact COCs were selected and transferred into a  
161 petri dish (35 mm, Nunclon, Denmark) prefilled with 2 mL of modified PBS supplemented with 0.4%  
162 BSA. After three washes in NCSU 37 supplemented with 5.0 mg/mL insulin, 0.57 mM cysteine, 10  
163 ng/mL epidermal growth factor (EGF), 50 mM  $\beta$ -mercaptoethanol and 10% porcine follicular fluid  
164 (IVM medium), groups of 50 COCs were transferred to a Nunc 4-well multidish containing 500  $\mu$ L  
165 of the same medium per well and cultured at 39 °C in a humidified atmosphere of 5% CO<sub>2</sub>/7% O<sub>2</sub> in  
166 air. For the first 22 h of in vitro maturation the medium was supplemented with 1.0 mM db-cAMP,  
167 10 IU/mL, eCG (Folligon, Intervet, Boxmeer, The Netherlands) and 10 IU/mL hCG (Corulon,  
168 Intervet). For the last 22 h COCs were transferred to fresh maturation medium.

169 Heterologous oocyte binding assay was conducted as described by [13]. Briefly, semen collected  
170 from three of the above described stallions was washed twice in modified Tyrode's medium pH 7.4  
171 for 2.5 min at 800 x g; the pellet was resuspended in the same medium at  $1 \times 10^6$  spermatozoa /mL and  
172 aliquots of 500  $\mu$ l were capacitated in presence or absence of AP (1.25 or 2.5 IU/mL) for 1 h at 38.5°  
173 C in 95% humidity and 5% CO<sub>2</sub> atmosphere. Subsequently, 50 matured denuded oocytes were added  
174 to each well. After 1 h of co-incubation at 38.5° C in 95% humidity and 5% CO<sub>2</sub> in air, the oocytes  
175 were washed three times in PBS 0.4% BSA with a wide bore glass pipette, fixed in 4%  
176 paraformaldehyde for 15 min at room temperature in the dark and stained with 8,9  $\mu$ M Hoechst  
177 33342. Cells were washed twice in PBS, and individually placed in droplets of Vectashield (Vector  
178 Laboratories, Burlingame, CA, USA) on a slide, and covered with a coverslip. The number of  
179 spermatozoa attached to each oocyte was assessed by using the above described microscope and was  
180 expressed as mean number of spermatozoa per oocyte.

181

#### 182 *2.5. Sperm cryopreservation*



183

184 After dilution in Kenney extender, semen was centrifuged in glass conical tube at 600 g for 20', the  
185 supernatant partially removed and the sperm pellet resuspended in Heitland's extender supplemented  
186 with 3% egg yolk and 3% glycerol [14] to a final concentration of  $150 \times 10^6$  sperm/mL. Resuspended  
187 semen was loaded in 0.5 mL straws, kept at +4°C for two h and then frozen on liquid nitrogen vapors  
188 for 20'6 cm above the liquid nitrogen level.

189

## 190 2.6. Statistical analysis

191

192 Data were analyzed using R version 3.0.3. (Copyright © 2014, The R Foundation for Statistical  
193 Computing) and significance was set at  $p < 0.05$  unless otherwise specified.

194 Data were assayed for normal distribution using Shapiro-Wilk test; in case of non-normal distribution,  
195 data were handled by a log transformation to match normality. Subsequently an ANOVA test was  
196 used to assess differences between the AP activity at pH 7.0, 8.0 and 10.0. The paired samples t test  
197 was used for comparison of the sperm quality parameters and AP activity in pre and post thawing  
198 samples.

199 To test correlation between sperm quality parameters and AP activity, Spearman's rank test was used.  
200 As for the oocyte binding assay, data were tested by a general mixed effects model, setting the  
201 treatment as a fixed effect and the horse as a random effect .Linear regression was used to examine  
202 relationship between AP activity in seminal plasma and post-thawing sperm viability [22].  
203 Results are expressed as mean  $\pm$  standard deviation.

204

205

## 206 3. Results

207

208 Sperm concentration, motility (total motility TM, progressive motility PM, path velocity VAP,  
209 curvilinear velocity VCL), viability and mitochondrial membrane integrity as well as seminal plasma  
210 and sperm extracts protein concentration are reported in Table 1.

211 Sperm quality parameters from each stallion are presented in supplementary file 1 (S1).

212 AP activity in both seminal plasma and sperm extracts is significantly different depending on the  
213 three analyzed pH. A significant increase is evident from pH 7.0 to 8.0 ( $p < 0.001$ ) as well as from pH  
214 8.0 to pH 10.0 ( $p < 0.001$ ), as reported in Table 2.

215 Different AP activities in both seminal plasma and sperm extract were found among stallions, and  
216 data from single subjects are resumed in supplementary file 1 (S1).

217 The thermal inhibition test, carried out on seminal plasma AP activity, showed a decrease due to heat  
 218 exposure in a time dependent manner, while the activity of the intestinal AP isoform was not affected  
 219 by incubation at 60 °C. In fact, seminal plasma AP lost 45% activity after 1 min and 15 sec at 60°C,  
 220 reaching the 80% after 1 min and 45 sec and was completely inactivated after 3 min. (Figure 1).  
 221 AP activity in frozen-thawed sperm extracts significantly decreased ( $p<0.01$  for pH 7.0 and 8.0 and  
 222  $p<0.05$  for pH 10 measurements) in comparison with the relative freshly ejaculated sample. Results  
 223 on AP activity are resumed in Table 2.  
 224 AP activity in sperm extracts from freshly ejaculates and frozen-thawed spermatozoa from each  
 225 stallion are presented in supplementary file 2 (S2).  
 226 Sperm quality parameters significantly changed after freezing (Table 3). Motility parameters (TM,  
 227 PM, VAP and VCL) significantly ( $p<0.01$ ) decreased after freezing and the same trend was evident  
 228 in viability and mitochondrial activity ( $p<0.01$ ); on the contrary, sperm protein content was not  
 229 affected by the freezing process.  
 230 Results on sperm quality parameters from each stallion are presented in Supplementary file 2 (S2).  
 231 The oocyte binding assay did not show any difference between control group and those treated with  
 232 different AP concentrations (1.25 and 2.5 IU/mL).  
 233 The control group scored  $13.85\pm14.4$  sperm bound per oocyte, while AP 1.25 scored  $15.17\pm13.04$   
 234 and AP 2.5  $14.44\pm14.26$ .  
 235 Spearman's correlations between AP activity in seminal plasma and sperm extracts and sperm quality  
 236 parameters in freshly ejaculated spermatozoa are resumed in Table 4.  
 237 Seminal plasma AP activities measured at pH 7.0, 8.0 and 10.0 were positively correlated. In addition,  
 238 sperm extracts AP activity measured at pH 7.0, 8.0 and 10.0 showed positive correlation. AP activity  
 239 in seminal plasma was positively correlated with sperm concentration and negatively with seminal  
 240 plasma protein concentration.  
 241 Table 5 resumes the correlation between AP activity in sperm extracts and sperm quality parameters  
 242 in frozen-thawed stallion spermatozoa. While in fresh sperm extracts no correlation was found  
 243 between motility and AP activity from sperm extracts, in frozen-thawed samples positive correlations  
 244 between AP activity in sperm extracts at pH 7.0 and 10.0 and total motility and progressive motility  
 245 were recorded.  
 246 Correlations between freshly ejaculated stallion sperm AP activity (in seminal plasma and sperm  
 247 extracts) and sperm quality parameters and frozen-thawed ones are presented in Supplementary file  
 248 3 (S3).  
 249 AP activity in seminal plasma measured at pH 10.0 and 7.0, as well as total motility and progressive  
 250 motility in freshly ejaculated spermatozoa were correlated with sperm viability after thawing.

251 Multiple regression generated a statistical model with only one explanatory variable (AP activity in  
252 seminal plasma measured at pH 10.0) explaining 26.5% (Adjusted  $R^2$ ) of the variation in sperm  
253 viability after thawing.

254

#### 255 4. Discussion

256

257 This study was aimed at determining alkaline phosphatase activity in stallion seminal plasma and  
258 sperm extracts, and at defining AP relationship with some sperm quality parameters.

259 Our results are reported as specific activity, expressed as nmoles of substrate transformed in 1 min  
260 for milligram of protein in the sample. This is, in our opinion, the best way to express the activity, as  
261 it permits to delineate the substrate consumption in relation to the effective protein content of the  
262 sample, that can considerably differ among both stallions (see supplementary file S1) (inter subject  
263 variability) and samples from the same subject (intra subject variability). Therefore, this measurement  
264 is standardized and is not affected by the different concentrations of protein (and enzyme) in the  
265 ejaculate.

266 To better define the properties of seminal plasma AP, we performed a thermal inhibition test, as the  
267 enzyme's isoforms show significantly different sensitiveness to heat. AP in horse is easily and rapidly  
268 inhibited by heat, as reported in Figure 1. Recently, we observed that AP from boar seminal plasma  
269 is quite more resistant, losing 100% activity only after 10 min at 60°C [9]; on the basis of these data  
270 and those reported for boar and humans [15,16], we could infer that seminal plasma AP in stallion is  
271 very similar to bone isoforms.

272 This is the first report that approached the study of AP activity respecting the physiological  
273 environment of stallion semen. The enzyme activity at pH 7.0 is very low, in both seminal plasma  
274 and sperm extracts. This fact seems to be conflicting with other reports [5, 6], stating that AP activity  
275 in stallion semen is very high. Anyway, our data on AP activity reveal that it parallels the increase  
276 of pH, thus confirming that, at least at pH 8.0 and 10.0, AP activity in stallion semen is very high.  
277 The intriguing question risen from these data regards the role of AP in seminal plasma and sperm  
278 extracts of physiological stallion semen pH (near 7.0). Turned and McDonnell [5] studied the activity  
279 of the enzyme in extracts from testis, epididymis and ampullae, as well as from ampullary fluid and  
280 epididymal fluid. The highest activity was found in fluids from both epididymis and ampullae, thus  
281 letting to hypothesize that the enzyme is secreted in these compartments; in particular, epididymal  
282 fluid activity is 50 fold the mean activity registered in seminal plasma.

283 In a previous work we hypothesized that AP could play a role in maintaining porcine spermatozoa  
284 quiescent, preventing a premature capacitation [9]. In horse sperm, on the basis of ours and other's

[5] data, AP likely plays a role on spermatozoa metabolism and activity during their transit from testicles to the ampullae, while sperm dilution in seminal plasma dramatically reduces the enzyme activity and therefore its possible involvement in sperm metabolism. It should also be stressed that spermatozoa, during natural mating, are diluted in seminal plasma for a brief period of time, after which they enter the female genital tract. In this region, the environment is completely different and spermatozoa gradually lose seminal plasma to begin the capacitation process that lead to sperm-oocyte fusion and fertilization [17]. This process, in the mare, takes from 0.5 to 4 h.

To reinforce this hypothesis, the heterologous oocyte binding assay showed no difference between untreated spermatozoa under capacitating condition and spermatozoa treated with AP at two different doses. We may infer that the addition of AP to a capacitating medium with a neutral pH (7.4) is not effective in modifying heterologous oocyte binding, considered a good parameter to assess sperm capacitation in stallion [13].

AP activity of seminal plasma is significantly higher than that of sperm extracts ( at all pH tested, 7.0, 8.0 and 10.0), with a minimum mean ratio respectively of 5:1 at pH 10.0 and a maximum of 9:1 at pH 8.0. This is not surprising: other Authors [5] showed that the contribution of spermatozoa to the whole ejaculate's activity is very low if compared to that of seminal plasma. Those researchers, anyway, obtained an indirect indication of AP activity from spermatozoa, as they subtracted the activity of unprocessed semen samples to that of seminal plasma, while in our study we pointed out the effective AP activity from sperm cell extracts

The role of AP attached to sperm surface is more intriguing: in boar spermatozoa we demonstrated a significant decrease in AP activity after in vitro capacitation [9]. It should be highlighted that AP activity at physiological pH in boar (8.0) is similar to that observed in this study in stallion at the same pH that is significantly higher than that registered at pH 7.0. In that study we demonstrated that AP exerts a down regulatory activity in tyrosine protein phosphorylation during capacitation [9]. This action is intriguing, as under capacitating conditions, boar semen protein phosphorylation increases significantly, and it lets us to hypothesize that AP should be hashed down to permit capacitation in pig [9]. We do not have any experimental evidence that AP plays a role in stallion sperm capacitation, as it is deducible from heterologous oocyte binding, and its role (if any) during sperm transit in the male genital tract does not seem to be important. Recently, other Authors [18] showed that stallion spermatozoa acquire capacitation by interacting with tubal epithelial cells: in particular, those Authors demonstrated that tyrosine phosphorylation of spermatozoa is positively correlated to the increase in intracellular pH, strictly associated with the microenvironment induced by epithelial cells. Nevertheless, further studies are necessary to determine the possible role of AP during the capacitation process.

319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352

.  
There is a great range of AP activity in both seminal plasma and sperm extracts; we found significant differences among horses in AP seminal plasma activity at pH 7.0 and 8.0, while no difference was recorded either at pH 10.0 or in AP activity from sperm extracts (see supplementary file S1). The same trend between AP activity in seminal plasma and sperm extracts has been observed in boar [9]. The individual variation is well known in horse and pig, thus it is not surprising that mean values of our samples are quite dispersed.

The relationship between sperm quality parameters and AP activity was investigated by correlation test. As already described by Pesch and colleagues [6], we found a positive correlation between seminal plasma AP activity and sperm concentration. A high sperm concentration results from a lower secretion of seminal plasma by the accessory glands and, therefore, it is concomitant with a lower dilution of epididymal and testicle secretions which are rich of AP [5]. On the other hand, in our study, sperm extracts AP activity does not correlate with sperm concentration. We registered a negative correlation between seminal plasma protein concentration and AP activity in both seminal plasma and sperm extracts, excepting for sperm extracts activity at pH 7.0. These findings are difficult to interpret and further studies are necessary to understand if they could be related to AP origin (secretion from ampullae and epididymis). No correlations between AP activity in seminal plasma and/or sperm extracts and viability were found, while a negative correlation between AP activity in seminal plasma and sperm extracts, measured at pH 10.0, and the two sperm velocity parameters, VCL and VAP, was recorded. This finding maybe deserves further insights, as it seems to indicate that high AP activity is related with lower velocity of the spermatozoa, but evidence of a similar effect at physiological pH is still lacking, thus letting the question open.

The second part of our work aimed at determining the possible involvement of AP activity in cryopreservation of stallion sperm, with particular attention to sperm cryo-damage. Our results on sperm quality parameters (motility, viability and mitochondrial activity) are consistent with those reported by others [19; 20], thus delineating a significant loss in sperm quality after freezing and thawing procedure.

We demonstrated that AP activity in sperm extracts significantly decreases after freezing and thawing; similar results were reported for ram spermatozoa by Salamond and Maxwell [10], who observed that the phosphatase activity of ram semen after freezing and thawing decreases significantly. However, those Authors did not specify whether it was alkaline or acid one. Alkaline phosphatase activity is related to protein dephosphorylation and it possibly plays an important role in

353 regulating this process, at least in pig [9]. Evidence exists that tyrosine phosphorylation is activated  
354 after freezing and thawing of stallion spermatozoa and that the response of stallion sperm to  
355 capacitation stimuli is different after freezing and thawing [21]. In addition, tyrosine phosphorylation  
356 is one of the aspects of the so called cryo-capacitation [19], and its increase after freezing and thawing  
357 may be due to AP inhibition. The intimate mechanism of AP inactivation is still unclear: can be the  
358 inactivation a consequence of the membrane damage that lead to a loss of enzyme from the sperm  
359 surface? Cryoelution of surface proteins?

360 As for the correlation between frozen-thawed sperm quality parameters and AP activity of relative  
361 sperm extracts, only measurements at pH 10.0 and 7.0 (but not at pH 8.0) showed some correlations.  
362 A strong positive correlation between AP activity (at pH 7.0 and 10.0) and TM and PM is evident;  
363 this correlation is not present in the freshly ejaculated spermatozoa. We may suppose that this change  
364 is due to a perturbation of both motility parameters and AP activity because of the freezing-thawing  
365 process.

366 We also correlated AP activity in freshly ejaculated cells with sperm quality parameters of the relative  
367 frozen sample (presented in supplementary file S3). This correlation could possibly be useful for  
368 predicting sperm freezability based on AP activity in seminal plasma. The multiple linear regression  
369 showed that a part of the variability in sperm viability after thawing could be related to AP activity  
370 measured at pH 10, and that this parameter could be used for predicting, at least in part, the  
371 freezability of stallion semen. The linear relationship, in fact, links the amount of AP activity  
372 (measured at pH 10) in seminal plasma of the fresh ejaculate with the viability of post-thawing  
373 spermatozoa indicating a direct proportionality between the parameters.

374 In conclusion, our data demonstrate for the first time the presence of AP on stallion sperm surface  
375 and describe the enzyme activity in both seminal plasma and sperm extracts at different pH.

376 Measurements at pH 7.0 indicates that AP in the ejaculate is not very active, letting thus unknown its  
377 role in the ejaculate. In contrast, the activity on sperm surface, when reaching the utero tubal junction,  
378 could be enhanced by the higher pH in this part of the female genital tract. In addition, we found some  
379 interesting correlations between AP and sperm quality parameters. Finally, we observed a decrease  
380 of AP activity after freezing-thawing process, which parallels that of both viability and sperm  
381 motility.

382

### 383 **Acknowledgements**

384

385 The Authors wish to thank Dr Marco Tamburrini and Allevamento Folli srl (Mordano, BO, Italy) for  
386 their precious collaboration and Mrs Cinzia Cappannari for her valuable technical support.

387 Work supported by an University of Bologna RFO Grant. EG was financially supported by  
388 “Fondazione Sfameni”.  
389

390 **References**

391

392 [1] Price PA, Toroian D, Chan WS. (2009) Tissue-nonspecific alkaline phosphatase is required for  
393 the calcification of collagen in serum: a possible mechanism for biomineralization. *J Biol Chem*,  
394 2009; 284: 4594–4604.

395

396 [2] Bell DJ, Lake PE. A comparison of phosphomonoesterase activities in the seminal plasmas of the  
397 domestic cock, turkey tom, boar, bull, buck rabbit and of man. *J Reprod Fertil* 1962; 3: 363-368.

398

399 [3] Mann T. The biochemistry of semen and of the male reproductive tract. [1954] London: Methuen  
400 and Co. Ltd.

401

402 [4] Stallcup OT. Acid and alkaline phosphatase activity in bovine semen as related to fertility. *J Dairy*  
403 *Sci* 1965; 48: 752-754.

404

405 [5] Turner RMO, McDonnell SM. Alkaline phosphatase in stallion semen: characterization and  
406 clinical applications. *Theriogenology* 2003; 60: 1-10.

407

408 [6] Pesch S, Bergmann M, Bostedt H. Determination of some enzymes and macro- and microelements  
409 in stallion seminal plasma and their correlations to semen quality. *Theriogenology* 2006; 66: 307-  
410 313.

411

412 [7] Kareskoski M, Reilas T, Sankari S, Andersson M, Güvenc K, Katila T. Alkaline and acid  
413 phosphatase,  $\beta$ -glucuronidase and electrolyte levels in fractionated stallion ejaculates. *Reprod*  
414 *Domest Anim* 2010; 45(6): e369–374.

415

416 [8] Kareskoski M, Sankari S, Johannisson A, Kindahl H, Andersson M, Katila, T. The association of  
417 the presence of seminal plasma and its components with sperm longevity in fractionated stallion  
418 ejaculates. *Reprod Domest Anim* 2011; 46(6): 1073–1081.

419

420 [9] Bucci D, Isani G, Giaretta E, Spinaci M, Tamanini C, Ferlizza E, Galeati G. Alkaline phosphatase  
421 in boar sperm function. *Andrology* 2014; 2(1): 100–106.

422



423 [10] Salamon S, Maxwell WMC. Frozen storage of ram semen II. Causes of low fertility after cervical  
 424 insemination and methods of improvement. *Anim Reprod Scie* 1995; 38: 1-36.  
 425

426 [11] Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of  
 427 protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976; 72: 248-254.  
 428

429 [12] Galeati G, Vallorani C, Bucci D, Bernardini C, Tamanini C, Parmeggiani A, Spinaci M. Daidzein  
 430 does affect progesterone secretion by pig cumulus cells but it does not impair oocytes IVM.  
 431 *Theriogenology* 2011; 74(3): 451-457.  
 432

433 [13] Balao da Silva CM, Spinaci M, Bucci D, Giaretta E, Peña FJ, Mari G, Galeati G. Effect of sex  
 434 sorting on stallion spermatozoa: Heterologous oocyte binding, tyrosine phosphorylation and  
 435 acrosome reaction assay. *Anim Reprod Sci* 2013; 141(1-2): 68-74  
 436

437 [14] Heitland AV, Jasko DJ, Squires EL, Graham JK, Pickett BW, Hamilton C. Factors affecting  
 438 motion characteristics of frozenthawed stallion spermatozoa. *Eq Vet J* 1996; 28: 47-53.  
 439

440 [15] Iyer SK, Daron HH, Aull JL. Purification and properties of alkaline phosphatase from boar  
 441 seminal plasma. *J Reprod Fertil* 1988; 82: 657-664.  
 442

443 [16] Lewin LM, Golan R, Soffer Y, Kaufman S, Yulzary Y, Zaidman J. Alkaline phosphatase in  
 444 human semen: an investigation using enzyme inhibitors and gel electrophoresis. *Eur J Clin Chem*  
 445 *Clin Biochem* 1993; 31: 811-814.  
 446


447 [17] Katila T. Post-mating inflammatory response of the uterus. *Reprod Domest Anim* 2012; 47(5):  
 448 31-41.  
 449

450 [18] Leemans B, Gadella BM, Sostaric E, Nelis H, Stout TAE, Hoogewijs M, Van Soom A. Oviduct  
 451 Binding and Elevated Environmental pH Induce Protein Tyrosine Phosphorylation in Stallion  
 452 Spermatozoa. *Biol Reprod* 2014, 91(1) 13: 1-12.  
 453

454 [19] Sieme H, Harrison RAP, Petrunkina AM. Cryobiological determinants of frozen semen quality,  
 455 with special reference to stallion. *Anim Reprod Sci* 2008; 107: 276-292.  
 456

- 457 [20] Peña FJ, Macías García B, Samper JC, Aparicio IM, Tapia JA, Ortega Ferrusola C. Dissecting  
458 the molecular damage to stallion spermatozoa: the way to improve current cryopreservation  
459 protocols? *Theriogenology* 2011; 76: 1177-1186.  
460
- 461 [21] Pommer AC, Rutland J, Meyers SA. Phosphorylation of protein tyrosine residues in fresh and  
462 cryopreserved stallion spermatozoa under capacitating condition. *Biol reprod* 2003; 68: 1208-1214.  
463
- 464 [22] Holt C., Holt W.V., Moore H.D.M, Reed H.C.B., Curnock R.M. Objectively measured boar  
465 sperm motility parameters correlate with the outcomes of on farm insemination: results of two fertility  
466 trials. *J Androl* 1997; 18: 312-323.  
467  
468

469 Figure 1. Thermal inactivation of seminal plasma AP activity.  
470



471  
472  
473 AP activity was inhibited by heat (60°C) for different time periods. Error bars represent 1 SD.  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492

493 Table 1. Sperm quality parameters in freshly ejaculated spermatozoa. Results are expressed as mean  
 494  $\pm$  SD (n=40).

Parameter	mean $\pm$ SD
Concentration ( $10^6$ /mL)	197 $\pm$ 125
Total Motility %	69.3 $\pm$ 20.0
Progressive motility %	45.0 $\pm$ 12.7
Average velocity (VAP) ( $\mu$ m/sec)	102.0 $\pm$ 22.5
Curvilinear velocity (VCL) ( $\mu$ m/sec)	190.1 $\pm$ 41.8
% Viability	71.4 $\pm$ 13.8
Concentration ( $10^6$ /mL)	197 $\pm$ 125
Seminal plasma protein (mg/mL)	12.4 $\pm$ 10.4
Sperm extracts protein (mg/mL)	2.3 $\pm$ 1.0

497

498 Table 2. AP activity in seminal plasma and spermatozoa extract at different pH.

499

	Specific activity (nmol/min/ $\mu$ g protein)		
	pH 7.0	pH 8.0	pH 10.0
AP activity Seminal plasma (n=40)	11.1 $\pm$ 10.5 <sup>a</sup>	68.2 $\pm$ 48.5 <sup>b</sup>	503.2 $\pm$ 419.4 <sup>c</sup>
AP activity Sperm extracts (n=40)	2.7 $\pm$ 2.3 <sup>a</sup>	7.5 $\pm$ 6.9 <sup>b</sup>	102.9 $\pm$ 78.7 <sup>c</sup>
AP activity pre-freezing (n=21)	2.6 $\pm$ 2.0 <sup>*</sup>	6.8 $\pm$ 4.7 <sup>*</sup>	58.3 $\pm$ 46.2 <sup>#</sup>
AP activity after freezing/thawing (n=21)	0.9 $\pm$ 0.6 <sup>*</sup>	3.0 $\pm$ 1.6 <sup>*</sup>	19.9 $\pm$ 11.6 <sup>#</sup>

500

501 Data are expressed as mean  $\pm$  SD. Lowercase letters superscript indicate significant differences for  
 502  $p < 0.001$ . \* indicates significant differences within a column for  $p < 0.01$ ; # indicates significant  
 503 differences within a column for  $p < 0.05$ .

504

505

506

507

508

509

Table 4. Sperm quality parameters in freshly ejaculated and frozen/thawed spermatozoa.

Parameter	Freshly ejaculated sperm	Frozen/thawed sperm
TM	73.8±18.2	28.8±15.5*
PM	47.4±11.2	22.5±13.4*
VAP	107.1±16.2	88.2±14.6*
VCL	195.9±30.0	157.9±25.5*
% JC1 positive live	63.1±13.9	31.8±12.5*
% Viability	74.8±12.6	41.1±13.1*
Protein sperm extracts (mg/mL)	2.6±1.1	2.2±1.1

Data are expressed as mean ± sd; n=21. \* indicate significant difference in row for p<0.001.

515 Table 5. Spearman's correlations between sperm quality parameters and AP activity in seminal plasma and sperm extracts.  
516

	AP_P7	AP_P8	AP_P10	AP_S7	AP_S8	AP_S10	Conc	PIProt	SPProt	TM	PM	VAP	VCL	Live	Livepos
AP_P7	1														
AP_P8	0.95*	1													
AP_P10	0.90*	0.88*	1												
AP_S7	0.14	0.17	0.27	1											
AP_S8	0.49*	0.51*	0.47*	0.74*	1										
AP_S10	0.37	0.49 <sup>#</sup>	0.37	0.60*	0.91*	1									
Conc	0.59*	0.48*	0.53*	-0.05	0.01	-0.02	1								
PIProt	-0.56*	-0.60*	-0.50*	-0.13	-0.53*	-0.65*	0.18	1							
SPProt	0.49 <sup>#</sup>	-0.25	-0.33	-0.09	-0.14	-0.61*	0.05	0.33 <sup>#</sup>	1						
TM	-0.29	0.08	0.19	0.07	0.04	-0.27	0.31	0.33	0.10	1					
PM	0.19	0.35	0.50 <sup>#</sup>	0.14	0.19	-0.21	0.33	0.03	-0.02	0.83*	1				
VAP	-0.33	-0.33	-0.42 <sup>#</sup>	-0.27	-0.50	-0.59*	0.13	0.52*	0.17	0.41 <sup>#</sup>	0.30	1			
VCL	-0.46 <sup>#</sup>	-0.36	-0.52*	-0.25	-0.53 <sup>#</sup>	-0.54 <sup>#</sup>	0.14	0.49*	0.19	0.16	-0.04	0.89	1		
Live	0.20	0.18	0.18	-0.24	0.03	-0.11	0.10	0.12	-0.04	0.63*	0.61*	0.29	0.02	1	
Livepos	-0.09	-0.08	-0.12	-0.25	-0.26	-0.46 <sup>#</sup>	0.01	0.40 <sup>#</sup>	0.04	0.48*	0.38 <sup>#</sup>	0.39 <sup>#</sup>	0.26	0.60*	1

517  
518 Data represent Spearman's rho coefficient; \* indicates  $p < 0.01$ ; <sup>#</sup> indicates  $p < 0.05$ .

519 Parameters are reported as follows: AP\_P7 AP activity in seminal plasma pH7; AP\_P8 AP activity in seminal plasma pH8; AP\_P10 AP activity in  
520 seminal plasma pH10; AP\_S7 AP activity in sperm extracts pH7; AP\_S8 AP activity in sperm extracts pH8; AP\_S10 AP activity in sperm extracts  
521 pH10; Conc: sperm concentration; PIProt: Seminal plasma proteins; SPProt: sperm extracts protein; TM: total motility; PM: progressive motility;  
522 VAP: average path velocity; VCL: curvilinear velocity; Live: % live spermatozoa; Livepos: % live spermatozoa with high mitochondrial membrane  
523 activity.

524

525

526 Table 6. Spearman's correlation between AP activity in sperm extracts and sperm quality parameters in frozen/thawed spermatozoa.  
527

	AP_S7	AP_S8	AP_S10	Conc	SPProt	TM	PM	VAP	VCL	Live	Livepos
AP_S7	1										
AP_S8	-0,0992	1									
AP_S10	0,5767 <sup>#</sup>	0,3299	1								
Conc	-0,1808	-0,037	-0,0515	1							
SPProt	0,2149	-	-0,0283	0,0455	1						
		0,6305 <sup>*</sup>									
TM	0,5717 <sup>#</sup>	-0,1129	0,5988 <sup>#</sup>	0,2861	0,1482	1					
PM	0,5994 <sup>*</sup>	-0,1334	0,6102 <sup>*</sup>	0,2861	0,17	0,9928 <sup>*</sup>	1				
VAP	0,1948	-0,1817	0,1214	0,1215	0,4448 <sup>#</sup>	0,5629 <sup>*</sup>	0,5672 <sup>*</sup>	1			
VCL	0,0847	-0,0234	-0,0368	-0,0468	0,3012	0,3225	0,3108	0,8912 <sup>*</sup>	1		
Live	0,0785	-0,0208	-0,0221	0,3494	-0,1535	0,6665 <sup>*</sup>	0,658 <sup>*</sup>	0,3852	0,2442	1	
Livepos	0,2841	-0,1326	0,2966	0,0091	-0,0338	0,7666 <sup>*</sup>	0,7581 <sup>*</sup>	0,5138 <sup>#</sup>	0,3455	0,6753 <sup>*</sup>	1

528  
529 Data represent Spearman's rho coefficient; \* indicate p<0.01; # indicate p<0.05.

530 Parameters are reported as follow: AP\_S7 AP activity in sperm extracts pH7; AP\_S8 AP activity in sperm extracts pH8; AP\_S10 AP activity in sperm  
531 extracts pH10; Conc: sperm concentration; SPProt: sperm extracts protein; TM: total motility; PM: progressive motility; VAP: average path velocity;  
532 VCL: curvilinear velocity; Live: % live spermatozoa; Livepos: % live spermatozoa with high mitochondrial membrane activity.

533  
534  
535  
536  
537

538 S1. AP activity in seminal plasma and spermatozoa extract and sperm quality parameters in freshly ejaculated stallion spermatozoa.

Horse	n	AP_P7	AP_P8	AP_P10	AP_S7	AP_S8	AP_S10	PIProt	SPProt	TM	PM	VAP	VCL	Live	Livepos
1	1	15.7±NA	49.7±NA	291.2±NA	NaN	5.2±NA	NaN	11.6±NA	3.0±NA	NaN	NaN	NaN	NaN	41.2±NA	41.2±NA
2	2	53.3±NA	218.6±NA	1181.4±NA	2.2±2.8	7.1±3.8	102.7±69.2	4.7±NA	2.4±0.7	9.0± NA	6.0±NA	70.0± NA	170.3± NA	60.8±28.1	14.5±14.0
3	8	6.2±3.4	33.5±15.23	204.8±128.3	1.7±0.8	3.81.2±	49.6±43.5	19.4±13.1	2.5±1.3	77.0±7.2	47.0±4.6	121.4±14.7	223.6±23.3	67.0±10.3	62.2±11.3
4	2	16.9±0.7	81.0±4.16	334.2±135.9	1.4±1.6	10.9±2.7	144.6±53.0	5.5±1.0	1.5±0.5	71.0± NA	55.0±NA	98.9± NA	162.0± NA	82.8±2.0	73.7±14.8
5	9	9.7±8.9	47.3±32.84	481.5±455.2	2.6±1.3	5.6±5.1	76.5±53.6	19.5±13.0	2.6±1.1	75.5±10.3	46.0±8.2	103.3±19.0	195.8±42.2	79.2±9.0	71.0±7.8
6	6	23.0±13.3	107.8±53.28	746.0±405.4	3.6±3.3	10.3±9.5	87.1±88.8	8.5±2.8	2.1±1.2	88.3±3.9	58.2±1.2	112.9±10.6	197.2±23.6	80.7±8.7	70.5±8.6
7	2	17.5±1.4	85.6±19.59	578.0±328.7	8.6±NA	19.7±22.0	183.0±198.3	6.2±1.3	2.0±1.1	66.0± NA	52.0±NA	70.0± NA	139.2± NA	80.5±3.3	72.6±3.7
8	2	29.3±3.1	127.0±6.92	1077.3±604.3	2.3±3.1	8.3±6.3	95.9±77.1	9.1±0.2	2.3±0.5	65.0± NA	43.0±NA	78.0± NA	134.4± NA	76.0±12.5	71.0±19.5
9	6	9.7±7.2	56.9±42.35	410.7±467.8	1.6±1.2	5.7±2.6	170.8± NA	5.6±1.2	2.0±0.8	45.6±13.2	31.4±7.6	99.4±8.8	194.2±30.1	56.2±10.1	46.9±10.8
10	2	8.0±1.0	62.3±27.73	171.3±72.9	3.3±NA	12.4±3.5	180.3±25.0	5.48±0.4	2.6±0.9	53.0± NA	39.0±NA	33.8± NA	68.3± NA	72.1±0.1	47.3±10.3

539

540 Parameters are reported as follow: n: number of ejaculates; AP\_S7 AP activity in sperm extracts pH7; AP\_S8 AP activity in sperm extracts pH8;  
541 AP\_S10 AP activity in sperm extracts pH10; Conc: sperm concentration; PIProt: Seminal plasma proteins; SPProt: sperm extracts protein; TM: total  
542 motility; PM: progressive motility; VAP: average path velocity; VCL: curvilinear velocity; Live: % live spermatozoa; Livepos: % live spermatozoa  
543 with high mitochondrial membrane activity.

544

545



546 S2. AP activity and sperm quality parameters in freshly ejaculated and frozen/thawed spermatozoa.

547

Horse	n	AP_S7	AP_S8	AP_S10	SPProt	TM	PM	VAP	VCL	Live	Livepos
3	4	2.65±1.85	6.24±5.96	33.98±17.82	3.09±0.97	79.25±5.91	49.00±1.41	120.55±16.77	219.95±25.30	75.38±3.30	61.06±12.58
3FT	4	1.01±0.31	2.17±0.85	15.52±4.35	3.25±1.22	30.25±7.93	23.25±5.56	103.10±9.64	190.80±10.58	39.73±5.36	35.14±3.11
5	6	2.32±0.85	6.38±3.00	52.10±42.02	2.98±1.15	79.00±7.01	48.83±7.47	98.78±19.45	180.18±34.21	83.28±2.28	71.45±6.11
5FT	6	0.81±0.54	2.59±0.70	17.46±3.86	2.65±0.70	32.17±16.19	24.00±13.43	95.42±12.45	166.33±23.12	44.43±15.43	30.77±13.66
6	6	3.22±3.06	8.26±6.98	62.58±41.45	2.18±1.20	88.33±3.93	58.17±1.17	112.92±10.61	197.15±23.62	80.68±8.68	70.53±8.63
6FT	6	1.14±0.78	3.05±1.64	26.43±17.83	2.05±1.09	39.33±13.32	33.00±11.78	85.03±6.14	144.67±12.53	48.79±13.71	38.21±13.45
9	5	1.62±1.24	5.94±2.87	170.83±17.82	2.07±0.91	45.60±13.20	31.40±7.57	99.40±8.85	194.20±30.06	57.00±11.09	45.82±11.69
9FT	5	0.44±0.52	4.07±2.46	12.22±4.35	1.19±0.64	11.00±4.58	7.40±4.51	71.32±9.23	137.24±17.55	29.15±4.53	22.74±11.56

548

549 Parameters are reported as follow: n: number of ejaculates; AP\_S7 AP activity in sperm extracts pH7; AP\_S8 AP activity in sperm extracts pH8;  
550 AP\_S10 AP activity in sperm extracts pH10; Conc: sperm concentration; PlProt: Seminal plasma proteins; SPProt: sperm extracts protein; TM: total  
551 motility; PM: progressive motility; VAP: average path velocity; VCL: curvilinear velocity; Live: % live spermatozoa; Livepos: % live spermatozoa  
552 with high mitochondrial membrane activity.

553

554

555  
556  
557

S3. Spearman's correlation between AP activity and sperm quality parameters in freshly ejaculated and frozen/thawed spermatozoa.

	AP_P 7	AP_P 8	AP_P 10	AP_S 7	AP_S 7F	AP_S 8	AP_S 8F	AP_S 10	AP_S1 0F	Conc	Conc F	PIPr ot	SPPr ot	SPPr otF	TM	TMF	PM	PMF	VAP	VAP F	VCL	VCLF	Live	LiveF	Lipos	Lipos F
AP_P7	1																									
AP_P8	0.99	1																								
AP_P10	0.94	0.94	1																							
AP_S7	0.49	0.45	0.38	1																						
AP_S7F	-0.09	-0.10	-0.11	-0.19	1																					
AP_S8	0.63	0.59	0.55	0.77	-0.03	1																				
AP_S8F	0.40	0.45	0.34	0.27	-0.10	0.44	1																			
AP_S10	0.62	0.60	0.6	0.3	-0.30	0.60	0.50	1																		
AP_S10F	0.07	0.04	0.10	-0.11	0.58	0.20	0.33	0.16	1																	
Conc	0.63	0.59	0.56	0.39	0.02	0.33	0.12	0.43	0.12	1																
ConcF	0.60	0.55	0.55	0.56	-0.18	0.51	-0.04	0.09	-0.05	0.64	1															
PIProt	-0.60	-0.70	-0.63	-0.07	0.15	-0.34	-0.37	-0.44	0.07	0.001	-0.02	1														
SPProt	-0.47	-0.50	-0.56	-0.01	0.11	-0.21	-0.15	-0.64	-0.05	-0.12	0.12	0.58	1													
SPProtF	-0.59	-0.63	-0.59	-0.15	0.21	-0.40	-0.63	-0.65	-0.03	-0.07	0.05	0.80	0.51	1												
TM	0.33	0.19	0.28	0.47	0.27	0.25	-0.31	-0.07	0.43	0.44	0.60	0.25	-0.06	0.36	1											
TMF	0.32	0.21	0.33	0.18	0.57	0.12	-0.11	0.004	0.60	0.47	0.29	0.28	-0.17	0.15	0.70	1										
PM	0.53	0.36	0.55	0.30	0.03	0.12	-0.02	0.22	0.30	0.54	0.48	0.13	-0.19	0.13	0.80	0.69	1									
PMF	0.31	0.22	0.34	0.18	0.60	0.12	-0.13	-0.01	0.61	0.49	0.29	0.26	-0.21	0.17	0.71	0.99	0.68	1								
VAP	-0.22	-0.27	-0.34	-0.04	0.40	-0.27	-0.21	-0.31	0.10	0.15	-0.15	0.38	0.11	0.38	0.36	0.49	0.36	0.46	1							
VAPF	-0.14	-0.23	-0.09	0.29	0.19	0.14	-0.18	-0.33	0.12	0.12	0.12	0.60	0.10	0.44	0.40	0.56	0.31	0.57	0.33	1						
VCL	-0.35	-0.30	-0.46	-0.01	0.41	-0.19	-0.07	-0.35	0.11	0.079	-0.23	0.29	0.23	0.26	0.10	0.28	-0.01	0.25	0.87	0.16	1					
VCLF	-0.24	-0.31	-0.16	0.22	0.08	0.16	-0.02	-0.25	-0.04	-0.01	-0.05	0.53	0.12	0.30	0.12	0.32	0.05	0.31	0.36	0.89	0.32	1				
Live	0.16	0.03	0.16	-0.06	0.45	0.002	-0.24	0.37	0.46	0.21	0.21	0.35	-0.10	0.28	0.53	0.68	0.52	0.67	0.22	0.32	-0.04	0.08	1			
LiveF	0.51	0.37	0.55	0.44	0.08	0.28	-0.02	0.05	-0.02	0.26	0.35	-0.01	-0.35	-0.2	0.58	0.67	0.63	0.66	0.23	0.39	0.01	0.24	0.49	1		

Livepos	-0.22	-0.30	-0.26	-0.29	0.41	-0.27	-0.17	-0.44	0.53	0.01	0.12	0.55	0.37	0.50	0.45	0.47	0.40	0.46	0.33	0.34	0.13	0.17	0.61	0.19	1	
LiveposF	0.15	0.08	0.16	0.14	0.28	-0.01	-0.13	-0.24	0.30	0.18	0.01	0.09	-0.26	-0.0	0.45	0.77	0.44	0.76	0.48	0.51	0.32	0.35	0.32	0.68	0.34	1

Data represent Spearman's rho coefficient; yellow underline indicate  $p < 0.01$ ; <sup>green</sup> underline indicate  $p < 0.05$ .

Parameters are reported as follow: AP\_P7 AP activity in seminal plasma pH7; AP\_P8 AP activity in seminal plasma pH8; AP\_P10 AP activity in seminal plasma pH10; AP\_S7 AP activity in sperm extracts pH7; AP\_S8 AP activity in sperm extracts pH8; AP\_S10 AP activity in sperm extracts pH10; SPProt: sperm extracts protein; TM: total motility; PM: progressive motility; VAP: average path velocity; VCL: curvilinear velocity; Live: % live spermatozoa; Lipos: % live spermatozoa with high mitochondrial membrane activity. "F" letter following the parameters name refers to frozen/thawed spermatozoa.