PUFA and Oxidative Stress. Differential Modulation of the Cell Response by DHA

Mattia Di Nunzio ^{1,*}, Veronica Valli ², Alessandra Bordoni ^{1,2}

¹ Interdepartmental Centre for Industrial Agri-Food Research, University of Bologna, piazza Goidanich, 60 Cesena (FC) - 47521, Italy; E-Mail: mattia.dinunzio@unibo.it (M.D.N.); alessandra.bordoni@unibo.it (A.B.)

² Department of Agri-Food Science and Technology, University of Bologna, piazza Goidanich, 60 - 47521 Cesena (FC) - 47521, Italy; E-Mails: veronica.valli9@unibo.it (V.V); alessandra.bordoni@unibo.it (A.B.)

* Author to whom correspondence should be addressed; E-Mail: mattia.dinunzio@unibo.it; Tel.: +39-0547-338957; Fax: +39-0547-382348.

Abstract:

Although an increased dietary intake of long-chain n-3 PUFA is considered an effective preventive strategy, a theoretical concern related to the possible increase of lipid peroxidation induced by a PUFA-rich diet still remains. In this study, the effects of different PUFA (linoleic, α -linolenic, arachidonic, eicosapentaenoic, and docosahexaenoic acid) on cytotoxicity, lipid oxidation, and modulation of antioxidant defenses were evaluated in HepG2 cells submitted to an oxidative stress (H_2O_2). Results clearly evidenced that all supplemented PUFA but DHA enhanced cell susceptibility to H_2O_2 . Overall, our results underline that PUFA cannot be considered as a single category but as individual compounds, and research on mechanisms of action and preventive effects should deal with the individual fatty acids, particularly in the case of DHA.

Keywords: HepG2 cells; hydrogen peroxide; cytotoxicity; antioxidant enzymes

1. Introduction

There are converging opinions among experts, organizations and health professionals that a recommendation for a daily individual consumption of 500 mg of eicosapentaenoic (EPA)/docosahexaenoic (DHA) acids would provide health benefits (Salem & Eggersdorfer, 2015). An increased dietary intake of n-3 long-chain polyunsaturated fatty acids (n-3 LC-PUFA), EPA and DHA, is considered an effective strategy in the prevention and modulation of cardiovascular (Ohnishi & Saito, 2013; Cao et al., 2015) and neurological diseases (Patrick & Ames, 2015), for the optimization of cognitive performance (Muldoon et al., 2014) and brain development (Brenna & Carlson, 2014).

Since DHA and other PUFA are highly oxidizable molecules (Miyashita, 2014), a theoretical concern remains on their use at high concentration in the prevention of chronic diseases having the oxidative stress as one of the underlying mechanisms.

Besides being prone to oxidation, PUFA have been reported to activate the stress-signaling pathway controlling the genetic program used by cells to induce the synthesis of proteins responsible for detoxification and counteraction of the oxidative stress (Ma, 2013). In this way, PUFA can modulate the activity of antioxidant enzymes (Wang et al., 2004) to ensure survival and reduced status.

Supplementing cultured cells with individual PUFA we previously observed diverse effects on oxidation and modulation of antioxidant defenses. The different effects were not simply related to the length of the carbon chain, or to the number and position of double bonds in the fatty acid (Di Nunzio et al., 2011). In particular, DHA supplementation did not cause any adverse effect, suggesting that its use as preventive dietary strategy does not induce a higher sensitivity to oxidation. It is conceivable that the overall effects of the different PUFA are related to their individual ability to increase the intracellular concentration of pro- or antioxidants.

In this light, the concomitant oxidative status of the cell could have a role in its response to PUFA supplementation, and the differential effect of DHA than other PUFA needs to be confirmed in cells submitted to an oxidative insult. To evaluate it, in the present work, we supplemented HepG2 cells with **linoleic acid (18:2n-6, LA)**, **α -linolenic acid (18:3n-3, ALA)**, **arachidonic acid (20:4n-6, ARA)**, **eicosapentaenoic acid (20:5n-3, EPA)**, and **docosahexaenoic acid (22:6n-3, DHA)**, then a mild oxidative stress **reflecting physiological condition** was induced (Halliwell et al., 2000; González-Flecha et al., 1993; Khassaf et al., 2001). HepG2 cells were used as model system since liver is a major organ attacked by reactive oxygen species (ROS) (Sánchez-Valle et al., 2012), and parenchymal cells are primary cells subjected to oxidative stress induced injury in the liver. Cells were supplemented with physiological concentration (60 μ M) of n-6 and n-3 PUFA, and cell survival, lipid oxidation, and antioxidant defenses were evaluated after exposure to hydrogen peroxide.

2. Experimental Section

2.1. Chemical

Dulbecco's modified Eagle's medium (DMEM), and Dulbecco's phosphate-buffered saline (DPBS) were purchased from Lonza (Milan, Italy). All chemicals and solvents were of the highest analytical grade and were purchased from Sigma-Aldrich (Milan, Italy).

2.2. HepG2 cells tissue culture

HepG2 human hepatoma cells were maintained at 37°C, 95% air, 5% CO₂ in DMEM supplemented with 10% (v/v) fetal calf serum, 100 U/ml penicillin and 100 μ g/ml streptomycin (Di Nunzio et al., 2010). Once a week cells were split 1:20 into a new 75cm² flask, and medium was refreshed after 72 h.

2.3. Fatty acid supplementation and oxidative stress

Cells were seeded in 6-well plates, and after 24 hours, at 75-80% confluence, they were incubated for 21 h with the different fatty acids at 60 μ M concentration. Free fatty acids were dissolved in 100%

isopropanol, and complexed to bovine serum albumin (BSA). Fatty acid-BSA complexes were prepared fresh each time, and BSA final concentration was 0.5% in serum free DMEM. Unsupplemented (US) cells received corresponding amounts of BSA and isopropanol, which final concentration in the media was kept below 1% (v/v). LA, ALA, ARA, EPA, and DHA were used for cell supplementation. After 21 h incubation cells were washed twice with warm DPBS, and then exposed for 20 or 60 min to 0.2 mM H₂O₂ in Earle's Balanced Salt Solution (EBSS) (116 mM NaCl, 5.4 mM KCl, 0.8 mM NaH₂PO₄, 26 mM NaHCO₃, 2.38 mM CaCl₂, 0.39 mM MgSO₄). Some unsupplemented cells received EBSS only (control cells). Then EBSS medium was collected, centrifuged at 400 g for 3 min, and used to evaluate LDH release and TBARS concentration. Cells were washed twice with warm DPBS, and processed for further analysis.

2.4. Lactate dehydrogenase (LDH) release

LDH activity in the EBSS was assessed following spectrophotometrically for 1 min the rate of NADH oxidation at 340 nm (Di Nunzio et al., 2013), and enzyme activity was expressed as mUnits/ml.

2.5. Thiobarbituric acid reactive substances (TBARS) concentration

TBARS, the end products of lipid peroxidation, were assayed in EBSS as previously reported by Valli *et al.* (Valli et al., 2012), and their concentration was expressed as relative fluorescence units (RFU) normalized for mg protein in the corresponding well.

2.6. Cell viability

Cell viability was evaluated by the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay, based on the cleavage of the yellow tetrazolium salt MTT to purple formazan crystal by metabolic active cells (Danesi et al., 2011). Results were expressed as percent of control cell viability, assigned as 100%.

2.7. Conjugated diene (CD) level

Cellular level of CD containing lipids was assayed as reported by Di Nunzio *et al.* (Di Nunzio et al., 2011), and expressed as percent of the value detected in control cells, assigned to 100%.

2.8. Cytoplasmic total antioxidant activity (TAA)

Cytoplasmic TAA was measured as reported by Di Nunzio *et al.* (Di Nunzio et al., 2013), based on the ability of the antioxidant molecules in the sample to reduce the radical cation of ABTS. Values were normalized for protein content in the sample, and expressed as micromoles of trolox equivalent (TE)/mg protein.

2.9. Reduced glutathione (GSH) content

Cytoplasmic GSH content was determined as previously described (Valli et al., 2012), normalized for protein content in the sample, and expressed as nmoles of GSH/mg protein.

2.10. Antioxidant enzymes activity

Cytoplasmic superoxide dismutase (SOD) activity was assayed using a commercial kit (Cayman Chemical, Michigan, USA) following the manufacture instructions. SOD activity (Units) was then normalized for mg of protein in the sample. Cytoplasmic catalase (CAT) activity was assayed spectrophotometrically at 240 nm following the extinction of H₂O₂ (Danesi et al., 2006), normalized for protein content in the sample, and expressed as Units/mg protein. Cytoplasmic glutathione-S-transferase (GST) activity was assayed as previously reported by Di Nunzio *et al.* (Di Nunzio et al., 2011). GST activity (nmol/min) was then normalized for protein content in the sample. Cytoplasmic glutathione peroxidase (GPx) activity was assayed using a commercial kit (Cayman Chemical, Michigan, USA) following the manufacture instructions. GPx activity (nmol/min) was then normalized for protein content in the sample.

2.11. Protein content

Protein content was determined according to Bradford (Bradford, 1976), using BSA as standard.

2.12. Statistical analysis

Data are reported as mean $\pm$ SD of at least six samples derived from three independent cell cultures. Statistical analysis was by the one way ANOVA with Dunnett as post-test.

3. Results

To evidence the detrimental effect of H₂O₂ addition, control cells were compared to stressed cells, both US and fatty acid supplemented. Then a statistical comparison was performed among stressed cells, to verify the differential effect of the single fatty acids, and to evidence whether the diverse supplementation could protect or exacerbate the outcomes of the oxidative stress.

3.1. H₂O₂-induced cell oxidative damage

The H₂O₂-induced cell damage was evaluated by measuring cell viability, LDH release, TBARS level and CD content (figure 1). As shown in figure 1A, in all cells H₂O₂ exposure caused a significant reduction of viability with respect to controls. Among stressed cells, viability decreased in all supplemented cells, but DHA.

The oxidative stress caused a significant increase in LDH activity and TBARS concentration in the medium of all cells except the US and DHA supplemented ones (figure 1B and 1C, respectively). Notwithstanding the increase of TBARS in almost all stressed cells, the level of CD, the first products of lipid peroxidation, increased in LA and ARA supplemented cells only (figure 1D).

3.2. Modulation of antioxidant defenses

137 To evaluate the effect of PUFA supplementation on the cell response to an oxidative stress,
138 cytosolic TAA and GSH content (figure 2), and the activity of SOD, CAT, GST and GPx enzymes
139 (figure 3) were determined.

140 Compared to controls, H₂O₂ exposure caused a significant decrease of cytosolic TAA in US and LA
141 supplemented cells only (figure 2A), and no modification in GSH content apart from a significant
142 increase in DHA supplemented cells (figure 2B). Comparing stressed cells, TAA appeared higher in
143 ARA and DHA supplemented hepatocytes, the latter also showing a higher GSH concentration.

144 Compared to controls, upon H₂O₂ exposure a significant increase of SOD activity was detected in
145 LA and n-3 PUFA supplemented cells (figure 3A), and CAT activity increased in all cells (figure 3B).
146 GST activity was not modified in US cells, while it significantly increased in all supplemented cells
147 but DHA ones (figure 3C). Furthermore, a significant increase in GPx activity was detected in LA
148 supplemented cells only (figure 3D). Comparing stressed cells, a differential influence of the
149 supplemented fatty acids on enzyme activities was evidenced. SOD activity being higher in EPA and
150 DHA supplemented, CAT in ARA and EPA supplemented, GST in ARA supplemented, and GPx in
151 LA supplemented cells than in US ones.

152 Since cellular oxidative status may be influenced by nonlethal oxidative stress exposure in a time
153 depending manner (Brunk et al., 1995), SOD, CAT, GST and GPx activities were also evaluated after
154 20 min H₂O₂ exposure (figure 4A, 4B, 4C and 4D respectively).

155 Compared to controls, upon the shorter H₂O₂ exposure time SOD activity increased in ARA and
156 EPA supplemented cells only, and no significant modification in CAT and GPx activity was detected.
157 GST activity was reduced in all supplemented cells except DHA and US ones. Comparing stressed
158 cells a differential effect of the supplementation on SOD, CAT and GST activities was still detected,
159 while no differences were evidenced in GPx activity.

160 **4. Discussion**

161 n-6 and n-3 PUFA are well known for their physiological functions and preventive actions (Liu et
162 al., 2015; Pelliccia et al., 2013) that are ascribable to different mechanisms including regulation of
163 membrane functionality, cell signaling, and gene expression (Robichaud & Surette, 2015; Righi et al.,
164 2011; Bordoni et al., 2007). In contrast to their health effects, PUFAs are known to be prone to rapid
165 peroxidation, which has been shown to impair membrane functions and inactivate proteins and
166 enzymes, eventually leading to various disorders and diseases (Csala et al., 2015). To explain this
167 apparent oxymoron, it has been speculated that the lipid peroxidation occurring in PUFA-rich cells
168 causes a low level, sub-toxic stress that could induce a protective adaptations over time (i.e. hormesis),
169 contributing to PUFA health effects. Actually, n-3 PUFA have been shown to up-regulate the
170 antioxidant systems in various cells (Giordano & Visioli, 2014), so counteracting the oxidative stress
171 (Zanatta et al., 2014; Tourtas et al., 2012). Although different PUFA share common chemical
172 characteristics, in a previous work we evidenced that they differently modulate the activity of
173 antioxidant enzymes, and have a different impact on cell oxidative status (Di Nunzio et al., 2011). A
174 concomitant oxidative stress could have an impact on the effects of PUFA, this representing an
175 important issue since a higher intake of PUFA, particularly n-3 PUFA, is suggested for the
176 prevention/modulation of a wide range of diseases associated with an increased production of ROS

177 (Fischer & Maier, 2015; Reuter et al., 2010; Pashkow, 2011; Salomone et al., 2016). To further
178 elucidate it, in the present work we have supplemented different PUFA to HepG2 cells, which were
179 then submitted to a mild oxidative stress.

180 Results herein reported clearly evidence that the cell response to a subsequent oxidative stress can
181 be modulated by a preceding PUFA supplementation.

182 All supplemented fatty acids except DHA enhance cell susceptibility to H₂O₂. This could be due to
183 an increased lipid peroxidation inducing disturbance of fine structures, alteration of integrity, fluidity,
184 and permeability, and functional loss of biomembranes, and generates potentially toxic products
185 (Ayala et al., 2014), as indicated by the increase in TBARS concentration in the medium of all
186 supplemented cells but DHA supplemented ones. Since CD containing lipids, the first products of lipid
187 peroxidation, are usually quickly cleaved to their corresponding breakdown products (Yin et al., 2011),
188 the higher CD concentration in n-6 PUFA than n-3 PUFA supplemented hepatocytes could indicate a
189 longer time lasting lipoperoxidation.

190 Although we already observed a PUFA sensitizing effect in basal condition (Di Nunzio et al.,
191 2011), results herein reported cannot be simply considered as its consequence but as an additive effect
192 due to the concomitant presence of PUFA and H₂O₂. It can be clearly evidenced comparing present
193 and previous data; as example, in basal condition LDH release was significantly higher in ARA
194 supplemented cells only (Di Nunzio et al., 2011), while it is significantly increased in all supplemented
195 cells except DHA ones after H₂O₂ addition, the increase being higher in ARA supplemented cells. It is
196 worth noting that DHA supplementation has no adverse effect either in basal condition (Di Nunzio et
197 al., 2011) or after the oxidative insult.

198 This differential modulation of the cell response to the oxidative stress is in part ascribable to the
199 modulation of the antioxidant enzymes activities. The effect of PUFA on the antioxidant enzymes
200 status have been extensively studied leading to controversial results. To the Authors knowledge,
201 previous studies have been performed using fish oil (Avramovic et al., 2012; Romieu et al., 2008) or
202 other oils (Umesha & Naidu, 2015; Taranu et al., 2014; Haggag et al., 2014) containing different
203 combinations of PUFA, this not allowing to extrapolate the effect of the single fatty acid. Results
204 herein reported clearly evidence that each fatty acid has specific modulatory effects on antioxidant
205 enzymes, and promotes a specific profile of antioxidant enzyme activities that cannot be predicted
206 using oils containing different proportion of PUFA. The importance of the proportion of PUFA in a
207 mixture has been also shown by Molinar-Toribio *et al.* (Molinar-Toribio et al., 2015) who evidenced in
208 rat supplemented with EPA plus DHA mixtures that the activity of antioxidant enzymes increases with
209 magnitudes depending on the EPA:DHA ratio. Notably, the highest DHA concentration reduced the
210 oxidative stress.

211 Our study confirms that exposure time to H₂O₂ modulates anti-oxidant enzymes activity, as
212 previously reported by Shull *et al.* (Shull et al., 1991) and Franco *et al.* (Franco et al., 1999). In
213 addition, our results evidence that PUFA supplementation exacerbates this time-depending
214 modulation. Assuming that the observed increased activity of antioxidant enzymes is based on an
215 increased transcription of the corresponding encoding genes and an increased synthesis of the
216 enzymatic proteins, a lag time between the insult and the cell response is conceivable. This can explain
217 the less evident changes in antioxidant enzymes activities after 20 than 60 min H₂O₂ exposure. In
218 addition, the weaker cell response after shorter time of exposure could be due to the lower intracellular

219 concentration of H₂O₂. In fact, H₂O₂ diffusion across the plasma membrane is low during the first 10
220 minutes of exposure (Alexandre et al., 2006).

221 Oxidative stress underlies a complex array of metabolic alterations. Under this condition,
222 derangement of cell biochemistry and function is known to involve misregulation of key components
223 responsible for cell signaling, such as membrane receptors, kinases, phosphatases, and transcription
224 factors (Leonarduzzi et al., 2011; Brigelius-Flohé et al., 2011). In this view, the overall differential
225 effect of supplemented DHA could rely on different mechanisms. Among them, a different extent of
226 incorporation/metabolism of the supplemented fatty acid inside the cell. Although all supplemented
227 PUFA are readily incorporated by HepG2 cells at the expense of monounsaturated fatty acids, in a
228 previous work we observed different conversion rate (Di Nunzio et al., 2010). LA and ALA are further
229 metabolized upon cellular uptake, and EPA is partially desaturated/elongated to DHA. On the contrary,
230 no DHA retro-conversion is detected. Therefore the effects of all PUFA except DHA could be
231 ascribable to both the supplemented PUFA and its derivatives. The differential metabolism of
232 supplemented PUFA could also impact on the modification of membrane composition, and
233 consequently functionality, since the majority of PUFA is incorporated in the cell membrane
234 phospholipids (Abbott et al., 2012). The oxidative stress may selectively impair cell PUFA
235 concentration, which in turn may modulate the effect of the oxidative stress (Zaloga et al., 2006).
236 Oxidative condition modifies fatty acid composition in neonatal rat cardiomyocytes (Bordoni et al.,
237 2005), and a slight increase in the saturated and a decrease in monounsaturated and PUFA content has
238 been detected in the liver of pro-oxidant pesticide-treated rats (Nakbi et al., 2010). At membrane level,
239 exposure to an oxidative stress leads not only to fatty acid oxidation but also to phospholipid
240 breakdown due to activation of phospholipase A₂ (Nalbone et al., 1990). DHA released from
241 phospholipids is the substrate for the formation of resolvins and protectins, that act through distinct
242 receptors to unfold biological effects (Calder, 2015) including oxidation preventing effects (Liu et al.,
243 2014; Tian et al., 2015).

244 A higher DHA cellular content may play an important role in the protection from oxidative damage
245 because 4-hydroxyhexenal, an aldehyde derived from DHA peroxidation, up-regulates Nrf2-HO-1
246 antioxidant signaling pathway, which induces antioxidant effects (Gao et al., 2007).

247 Many studies indicate that PUFA, particularly EPA and DHA, control gene expression also by
248 directly governing the transcription/activity of peroxisome proliferator-activated receptor- α
249 (PPAR α) (Di Nunzio et al., 2009; Georgiadi et al., 2012), whose activation may result in enhanced
250 expression/activity of antioxidant enzymes such as CAT and SOD (Toyama et al., 2004; Inoue et al.,
251 2001). PPARs exert their effect on gene transcription by dimerization with the 9-cis retinoic acid
252 receptors (RXRs) (Feige et al., 2005). RXR is also able to bind PUFA, mainly DHA (Crawford et al.,
253 2003). Other PUFAs such as EPA or ARA can activate RXR but with lower efficiency (Egea et al.,
254 2002). Thus, DHA can be considered a more efficient activator of PPAR α (Bordoni et al., 2006).

255 In conclusion, based on reported results each PUFA seems to possess specific pro- and antioxidative
256 characteristics, not predictable on chemical basis such as the length of the carbon chain or the
257 unsaturation degree. According to many *in vivo* (Mas et al., 2010; Patten et al., 2013) and *in vitro*
258 (González-Pérez et al., 2006; Mukherjee et al., 2004) studies, DHA appears the only fatty acid not
259 increasing cell susceptibility to H₂O₂, although its effect is not ascribable to a stronger induction of
260 antioxidant enzymes activity.

Overall, results herein reported clearly evidence that fatty acid supplementation can modify the cell response to an oxidative stress, and the response is dependent on the supplemented fatty acid. We are beginning to understand the physiology and molecular basis of the effects of n-3 and n-6 fatty acids, and thereby gradually encompass the full biological potential of these compounds. Our results underline that PUFA cannot be considered as a single category, but as individual compounds. It appears particularly evident for DHA, and although observed results must be confirmed *in vivo*, this apparent “DHA paradox” represents an important step ahead in the use of DHA supplementation as preventive strategy in the clinical practice.

Disclosure statement

The authors declare no conflict of interest.

Funding statement

This work was partially supported by a grant of Italian MIUR (RFO A.B.), of Italian Ministry of Economic Development (Industria 2015 – MIAOVER50 project). The research leading to these results has received funding from the European Union Seventh Framework Programme (FP7/2007-2013) under grant agreement n° 311876: Pathway-27.

References

1. Abbott SK, Else PL, Atkins TA, Hulbert AJ. 2012. Fatty acid composition of membrane bilayers: importance of diet polyunsaturated fat balance. *Biochim Biophys Acta*. 1818:1309-1317.
2. Alexandre J, Batteux F, Nicco C, Chéreau C, Laurent A, Guillevin L, Weill B, Goldwasser F. 2006. Accumulation of hydrogen peroxide is an early and crucial step for paclitaxel-induced cancer cell death both in vitro and in vivo. *Int J Cancer*. 119:41-48.
3. Avramovic N, Dragutinovic V, Krstic D, Colovic M, Trbovic A, de Luka S, Milovanovic I, Popovic T. 2012. The effects of omega 3 fatty acid supplementation on brain tissue oxidative status in aged wistar rats. *Hippokratia*. 16:241-245.
4. Ayala A, Muñoz MF, Argüelles S. 2014. Lipid peroxidation: production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxid Med Cell Longev*. 2014:360438.
5. Bordini A, Angeloni C, Leoncini E, Danesi F, Maranesi M, Biagi PL, Hrelia S. 2005. Hypoxia/reoxygenation alters essential fatty acids metabolism in cultured rat cardiomyocytes: protection by antioxidants. *Nutr Metab Cardiovasc Dis*. 15:166-73.
6. Bordini A, Astolfi A, Morandi L, Pession A, Danesi F, Di Nunzio M, Franzoni M, Biagi P, Pession A. 2007. N-3 PUFAs modulate global gene expression profile in cultured rat cardiomyocytes. Implications in cardiac hypertrophy and heart failure. *FEBS Lett*. 581:923-929.
7. Bordini A, Di Nunzio M, Danesi F, Biagi PL. 2006. Polyunsaturated fatty acids: From diet to binding to ppars and other nuclear receptors. *Genes Nutr*. 1:95-106.

- 297 8. Bradford MM. 1976. A rapid and sensitive method for the quantitation of microgram quantities
298 of protein utilizing the principle of protein-dye binding. *Anal Biochem.* 72:248-254.
- 299 9. Brenna JT, Carlson SE. 2014. Docosahexaenoic acid and human brain development: evidence
300 that a dietary supply is needed for optimal development. *J Hum Evol.* 77:99-106.
- 301 10. Brigelius-Flohé R, Flohé L. 2011. Basic principles and emerging concepts in the redox control of
302 transcription factors. *Antioxid Redox Signal.* 15:2335-2381.
- 303 11. Brunk UT, Zhang H, Dalen H, Ollinger K. 1995. Exposure of cells to nonlethal concentrations of
304 hydrogen peroxide induces degeneration-repair mechanisms involving lysosomal destabilization.
305 *Free Radic Biol Med.* 19:813-822.
- 306 12. Calder PC. 2015. Marine omega-3 fatty acids and inflammatory processes: Effects, mechanisms
307 and clinical relevance. *Biochim Biophys Acta.* 1851:469-484.
- 308 13. Cao Y, Lu L, Liang J, Liu M, Li X, Sun R, Zheng Y, Zhang P. 2015. Omega-3 Fatty Acids and
309 Primary and Secondary Prevention of Cardiovascular Disease. *Cell Biochem Biophys.* 72:77-81.
- 310 14. Crawford MA, Golfetto I, Ghebremeskel K, Min Y, Moodley T, Poston L, Phylactos A, Cunnane
311 S, Schmidt W. 2003. The potential role for arachidonic and docosahexaenoic acids in protection
312 against some central nervous system injuries in preterm infants. *Lipids.* 38:303-315.
- 313 15. Csala M, Kardon T, Legeza B, Lizák B, Mandl J, Margittai É, Puskás F, Száraz P, Szelényi P,
314 Bánhegyi G. 2015. On the role of 4-hydroxynonenal in health and disease. *Biochim Biophys*
315 *Acta.* 1852:826-838.
- 316 16. Danesi F, Ferioli F, Caboni MF, Boschetti E, Di Nunzio M, Verardo V, Valli V, Astolfi A,
317 Pession A, Bordoni A. 2011. Phytosterol supplementation reduces metabolic activity and slows
318 cell growth in cultured rat cardiomyocytes. *Br J Nutr.* 106:540-548.
- 319 17. Danesi F, Malaguti M, Di Nunzio M, Maranesi M, Biagi PL, Bordoni A. 2006. Counteraction of
320 adriamycin-induced oxidative damage in rat heart by selenium dietary supplementation. *J Agric*
321 *Food Chem.* 54:1203-1208.
- 322 18. Di Nunzio M, Danesi F, Bordoni A. 2009. n-3 PUFA as regulators of cardiac gene transcription:
323 a new link between PPAR activation and fatty acid composition. *Lipids.* 44:1073-1079.
- 324 19. Di Nunzio M, Toselli M, Verardo V, Caboni MF, Bordoni A. 2013. Counteraction of oxidative
325 damage by pomegranate juice: influence of the cultivar. *J Sci Food Agric.* 93:3565-3573.
- 326 20. Di Nunzio M, Valli V, Bordoni A. 2011. Pro- and anti-oxidant effects of polyunsaturated fatty
327 acid supplementation in HepG2 cells. *Prostaglandins Leukot Essent Fatty Acids.* 85:121-127.
- 328 21. Di Nunzio M, van Deursen D, Verhoeven AJ, Bordoni A. 2010. n-3 and n-6 Polyunsaturated
329 fatty acids suppress sterol regulatory element binding protein activity and increase flow of non-
330 esterified cholesterol in HepG2 cells. *Br J Nutr.* 103:161-167.
- 331 22. Egea PF, Mitschler A, Moras D. 2002. Molecular recognition of agonist ligands by RXRs. *Mol*
332 *Endocrinol.* 16:987-997.
- 333 23. Feige JN, Gelman L, Tudor C, Engelborghs Y, Wahli W, Desvergne B. 2005. Fluorescence
334 imaging reveals the nuclear behavior of peroxisome proliferator-activated receptor/retinoid X
335 receptor heterodimers in the absence and presence of ligand. *J Biol Chem.* 280:17880-17890.
- 336 24. Fischer R, Maier O. 2015. Interrelation of oxidative stress and inflammation in
337 neurodegenerative disease: role of TNF. *Oxid Med Cell Longev.* 2015:610813.

25. Franco AA, Odom RS, Rando TA. 1999. Regulation of antioxidant enzyme gene expression in response to oxidative stress and during differentiation of mouse skeletal muscle. *Free Radic Biol Med.* 27:1122-1132.
26. Gao L, Wang J, Sekhar KR, Yin H, Yared NF, Schneider SN, Sasi S, Dalton TP, Anderson ME, Chan JY, Morrow JD, Freeman ML. 2007. Novel n-3 fatty acid oxidation products activate Nrf2 by destabilizing the association between Keap1 and Cullin3. *J Biol Chem.* 282:2529-2537.
27. Georgiadi A, Boekschoten MV, Müller M, Kersten S. 2012. Detailed transcriptomics analysis of the effect of dietary fatty acids on gene expression in the heart. *Physiol Genomics.* 44:352-361.
28. Giordano E, Visioli F. 2014. Long-chain omega 3 fatty acids: molecular bases of potential antioxidant actions. *Prostaglandins Leukot Essent Fatty Acids.* 90:1-4.
29. González-Flecha B, Cutrin JC, Boveris A. 1993. Time course and mechanism of oxidative stress and tissue damage in rat liver subjected to in vivo ischemia-reperfusion. *J Clin Invest.* 91:456-464.
30. González-Pérez A, Planagumà A, Gronert K, Miquel R, López-Parra M, Titos E, Horrillo R, Ferré N, Deulofeu R, Arroyo V, Rodés J, Clària J. 2006. Docosahexaenoic acid DHA blunts liver injury by conversion to protective lipid mediators: protectin D1 and 17S-hydroxy-DHA. *FASEB J.* 20:2537-2539.
31. Haggag Mel-S, Elsanhoty RM, Ramadan MF. 2014. Impact of dietary oils and fats on lipid peroxidation in liver and blood of albino rats. *Asian Pac J Trop Biomed.* 4:52-58.
32. Halliwell B, Clement MV, Long LH. 2000. Hydrogen peroxide in the human body. *FEBS Letters.* 486:10-13.
33. Inoue I, Goto S, Matsunaga T, Nakajima T, Awata T, Hokari S, Komoda T, Katayama S. 2001. The ligands/activators for peroxisome proliferator-activated receptor alpha (PPARalpha) and PPARgamma increase Cu²⁺, Zn²⁺-superoxide dismutase and decrease p22phox message expressions in primary endothelial cells. *Metabolism.* 50:3-11.
34. Khassaf M, Child RB, McArdle A, Brodie DA, Esanu C, Jackson MJ. 2001. Time course of responses of human skeletal muscle to oxidative stress induced by nondamaging exercise. *J Appl Physiol.* 90:1031-1035.
35. Leonarduzzi G, Sottero B, Testa G, Biasi F, Poli G. 2011. New insights into redox-modulated cell signaling. *Curr Pharm Des.* 17:3994-4006.
36. Liu JJ, Green P, John Mann J, Rapoport SI, Sublette ME. 2015. Pathways of polyunsaturated fatty acid utilization: implications for brain function in neuropsychiatric health and disease. *Brain Res.* 1597:220-246.
37. Liu M, Boussetta T, Makni-Maalej K, Fay M, Driss F, El-Benna J, Lagarde M, Guichardant M. 2014. Protectin DX, a double lipoxygenase product of DHA, inhibits both ROS production in human neutrophils and cyclooxygenase activities. *Lipids.* 49:49-57.
38. Ma Q. 2013. Role of nrf2 in oxidative stress and toxicity. *Annu Rev Pharmacol Toxicol.* 53: 401-426.
39. Mas E, Woodman RJ, Burke V, Puddey IB, Beilin LJ, Durand T, Mori TA. 2010. The omega-3 fatty acids EPA and DHA decrease plasma F2-isoprostanes: Results from two placebo-controlled interventions. *Free Radic Res.* 44, 983-990.

- 379 40. Miyashita K. Paradox of omega-3 PUFA oxidation. 2014. *Eur J Lipid Sci Technol*. 116:1268-
380 1279.
- 381 41. Molinar-Toribio E, Pérez-Jiménez J, Ramos-Romero S, Romeu M, Giralt M, Taltavull N,
382 Muñoz-Cortés M, Jáuregui O, Méndez L, Medina I, Torres JL. 2015. Effect of n-3 PUFA
383 supplementation at different EPA:DHA ratios on the spontaneously hypertensive obese rat model
384 of the metabolic syndrome. *Br J Nutr*. 113:878-887.
- 385 42. Mukherjee PK, Marcheselli VL, Serhan CN, Bazan NG. 2004. Neuroprotectin D1: a
386 docosahexaenoic acid-derived docosatriene protects human retinal pigment epithelial cells from
387 oxidative stress. *Proc Natl Acad Sci USA*. 101:8491-8496.
- 388 43. Muldoon MF, Ryan CM, Yao JK, Conklin SM, Manuck SB. 2014. Long-chain omega-3 fatty
389 acids and optimization of cognitive performance. *Mil Med*. 179:95-105.
- 390 44. Nakbi A, Tayeb W, Grissa A, Issaoui M, Dabbou S, Chargui I, Ellouz M, Miled A, Hammami
391 M. 2010. Effects of olive oil and its fractions on oxidative stress and the liver's fatty acid
392 composition in 2,4-Dichlorophenoxyacetic acid-treated rats. *Nutr Metab (Lond)*. 7:80.
- 393 45. Nalbone G, Grynberg A, Chevalier A, Leonardi J, Termine E, Lafont H. 1990. Phospholipase A
394 activity of cultured rat ventricular myocyte is affected by the nature of cellular polyunsaturated
395 fatty acids. *Lipids*. 25:301-306.
- 396 46. Ohnishi H, Saito Y. 2013. Eicosapentaenoic acid EPA reduces cardiovascular events:
397 relationship with the EPA/arachidonic acid ratio. *J Atheroscler Thromb*. 20:861-877.
- 398 47. Pashkow FJ. 2011. Oxidative Stress and Inflammation in Heart Disease: Do Antioxidants Have a
399 Role in Treatment and/or Prevention? *Int J Inflam*. 2011:514623.
- 400 48. Patrick RP, Ames BN. 2015. Vitamin D and the omega-3 fatty acids control serotonin synthesis
401 and action, part 2: relevance for ADHD, bipolar disorder, schizophrenia, and impulsive behavior.
402 *FASEB J*. 29:2207-2222.
- 403 49. Patten AR, Brocardo PS, Christie BR. 2013. Omega-3 supplementation can restore glutathione
404 levels and prevent oxidative damage caused by prenatal ethanol exposure. *J Nutr Biochem*.
405 24:760-769.
- 406 50. Pelliccia F, Marazzi G, Greco C, Franzoni F, Speziale G, Gaudio C. 2013. Current evidence and
407 future perspectives on n-3 PUFAs. *Int J Cardiol*. 170:S3-7.
- 408 51. Reuter S, Gupta SC, Chaturvedi MM, Aggarwal BB. 2010. Oxidative stress, inflammation, and
409 cancer: how are they linked? *Free Radic Biol Med*. 49:1603-1616.
- 410 52. Righi V, Di Nunzio M, Danesi F, Schenetti L, Mucci A, Boschetti E, Biagi P, Bonora S, Tugnoli
411 V, Bordoni A. 2011. EPA or DHA supplementation increases triacylglycerol, but not
412 phospholipid, levels in isolated rat cardiomyocytes. *Lipids*. 46:627-636.
- 413 53. Robichaud PP, Surette ME. 2015. Polyunsaturated fatty acid-phospholipid remodeling and
414 inflammation. *Curr Opin Endocrinol Diabetes Obes*. 22:112-118.
- 415 54. Romieu I, Garcia-Esteban R, Sunyer J, Rios C, Alcaraz-Zubeldia M, Velasco SR, Holguin F.
416 2008. The effect of supplementation with omega-3 polyunsaturated fatty acids on markers of
417 oxidative stress in elderly exposed to PM2.5. *Environ Health Perspect*. 116:1237-1242.
- 418 55. Salem N Jr, Eggersdorfer M. 2015. Is the world supply of omega-3 fatty acids adequate for
419 optimal human nutrition? *Curr Opin Clin Nutr Metab Care*. 18:147-154.

- 420 56. Salomone F, Godos J, Zelber-Sagi S. 2016. Natural antioxidants for non-alcoholic fatty liver
421 disease: molecular targets and clinical perspectives. *Liver Int.* 36:5-20.
- 422 57. Sánchez-Valle V, Chávez-Tapia NC, Uribe M, Méndez-Sánchez N. 2012. Role of oxidative
423 stress and molecular changes in liver fibrosis: a review. *Curr Med Chem.* 19:4850-4860.
- 424 58. Shull S, Heintz NH, Periasamy M, Manohar M, Janssen YM, Marsh JP, Mossman BT. 1991.
425 Differential regulation of antioxidant enzymes in response to oxidants. *J Biol Chem.* 266:24398-
426 24403.
- 427 59. Taranu I, Gras M, Pistol GC, Motiu M, Marin DE, Lefter N, Ropota M, Habeanu M. 2014. ω -3
428 PUFA rich camelina oil by-products improve the systemic metabolism and spleen cell functions
429 in fattening pigs. *PLoS One.* 9:e110186.
- 430 60. Tian Y, Zhang Y, Zhang R, Qiao S, Fan J. 2015. Resolvin D2 recovers neural injury by
431 suppressing inflammatory mediators expression in lipopolysaccharide-induced Parkinson's
432 disease rat model. *Biochem Biophys Res Commun.* 460:799-805.
- 433 61. Tourtas T, Birke MT, Kruse FE, Welge-Lüssen UC, Birke K. 2012. Preventive effects of omega-
434 3 and omega-6 Fatty acids on peroxide mediated oxidative stress responses in primary human
435 trabecular meshwork cells. *PLoS One.* 7:e31340.
- 436 62. Toyama T, Nakamura H, Harano Y, Yamauchi N, Morita A, Kirishima T, Minami M, Itoh Y,
437 Okanoue T. 2004. PPAR α ligands activate antioxidant enzymes and suppress hepatic fibrosis
438 in rats. *Biochem Biophys Res Commun.* 324:697-704.
- 439 63. Umesha SS, Naidu KA. 2015. Antioxidants and antioxidant enzymes status of rats fed on n-3
440 PUFA rich Garden cress *Lepidium Sativum* L seed oil and its blended oils. *J Food Sci Technol.*
441 52:1993-2002.
- 442 64. Valli V, Gómez-Caravaca AM, Di Nunzio M, Danesi F, Caboni MF, Bordonni A. 2012. Sugar
443 cane and sugar beet molasses, antioxidant-rich alternatives to refined sugar. *J Agric Food Chem.*
444 60:12508-12515.
- 445 65. Wang HH, Hung TM, Wei J, Chiang AN. 2004. Fish oil increases antioxidant enzyme activities
446 in macrophages and reduces atherosclerotic lesions in apoE-knockout mice. *Cardiovasc Res.*
447 61:169-176.
- 448 66. Yin H, Xu L, Porter NA. 2011. Free radical lipid peroxidation: mechanisms and analysis. *Chem*
449 *Rev.* 111:5944-5972.
- 450 67. Zaloga GP, Ruzmetov N, Harvey KA, Terry C, Patel N, Stillwell W, Siddiqui R. 2006. (N-3)
451 long-chain polyunsaturated fatty acids prolong survival following myocardial infarction in rats. *J*
452 *Nutr.* 136:1874-1878.
- 453 68. Zanatta AL, Miranda DT, Dias BC, Campos RM, Massaro MC, Michelotto PV Jr, West AL,
454 Miles EA, Calder PC, Nishiyama A. 2014. Fish oil supplementation decreases oxidative stress
455 but does not affect platelet-activating factor bioactivity in lungs of asthmatic rats. *Lipids.* 49:665-
456 675.

457 **Figure 1. Cell viability A), LDH activity B), TBARS concentration C) and CD level D) in basal**
458 **condition and after 1h H₂O₂ exposure.**

459 *Data are means ± SD of at least six samples in each group, coming from three independent cell*
460 *cultures. Statistical analysis was by the one-way ANOVA ($p < 0.001$) using Dunnett's post-test to*
461 *compare: 1) stressed cells both US and supplemented) to controls (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$);*
462 *and 2) supplemented stressed cells to US stressed cells ($^{\circ}p < 0.05$; $^{\circ\circ}p < 0.001$).*

463

464 **Figure 2. TAA A) and GSH level B) in basal condition and after 1h H₂O₂ exposure.**

465 *Data are means ± SD of at least six samples in each group, coming from three independent cell*
466 *cultures. Statistical analysis was performed by the one-way ANOVA ($p < 0.01$) using Dunnett's as post-*
467 *test to compare: 1) stressed cells both US and supplemented) to controls (* $p < 0.05$); and 2)*
468 *supplemented stressed cells to US stressed cells ($^{\circ}p < 0.05$; $^{\circ\circ}p < 0.01$).*

469

470 **Figure 3. SOD A), CAT B), GST C) and GPx D) activity in basal condition and after 1h H₂O₂**
471 **exposure.**

472 *Data are means ± SD of at least six samples in each group, coming from three independent cell*
473 *cultures. Statistical analysis was performed by the one-way ANOVA A, B and C ($p < 0.001$; D $p < 0.01$)*
474 *using Dunnett's as post-test to compare: 1) stressed cells both US and supplemented) to controls*
475 ** $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$); and 2) supplemented stressed cells to US stressed cells ($^{\circ}p < 0.05$;*
476 *$^{\circ\circ}p < 0.01$; $^{\circ\circ\circ}p < 0.001$).*

477

478 **Figure 4 SOD A), CAT B), GST C) and GPx D) activity in basal condition and after 20 min H₂O₂**
479 **exposure.**

480 *Data are means ± SD of at least six samples in each group, coming from three independent cell*
481 *cultures. Statistical analysis was performed by the one-way ANOVA A and B ($p < 0.01$; C $p < 0.001$)*
482 *using Dunnett's as post-test to compare: 1) stressed cells both US and supplemented) to controls*

483 $*p<0.05$; $**p<0.01$; $***p<0.001$); and 2) supplemented stressed cells to US stressed cells $^{\circ}p<0.05$;
484 $^{\circ\circ}p<0.01$; $^{\circ\circ\circ}p<0.001$).