

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Porcine circovirus type 2 detection in in vitro produced porcine blastocysts after virus sperm exposure

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Galeati, G., Zannoni, A., Spinaci, M., Bucci, D., Ostanello, F., Panarese, S., et al. (2016). Porcine circovirus type 2 detection in in vitro produced porcine blastocysts after virus sperm exposure. *ANIMAL SCIENCE JOURNAL*, 87(4), 511-516 [10.1111/asj.12465].

Availability:

This version is available at: <https://hdl.handle.net/11585/550122> since: 2019-07-17

Published:

DOI: <http://doi.org/10.1111/asj.12465>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the peer reviewed version of the following article:

Galeati G, Zannoni A, Spinaci M, Bucci D, Ostanello F, Panarese S, Tamanini C, Sarli G. Porcine circovirus type 2 detection in in vitro produced porcine blastocysts after virus sperm exposure. Anim Sci J. 2016;87(4):511-6.

which has been published in final form at
<https://doi.org/10.1111/asj.12465>

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

Porcine circovirus type 2 detection in *in vitro* produced porcine blastocysts after virus sperm exposure

Giovanna GALEATI*, Augusta ZANNONI, Marcella SPINACI, Diego BUCCI, Fabio

OSTANELLO, Serena PANARESE, Carlo TAMANINI and Giuseppe SARLI

Department of Veterinary Medical Sciences, University of Bologna, Bologna, Italy

Running title: PCV2 and porcine blastocysts

*Corresponding author. Tel.: +39 051 2097910; fax: +39 051 2097899.

Dipartimento di Scienze Mediche Veterinarie (DIMEVET)

Via Tolara di Sopra, 50

40064 Ozzano dell'Emilia, Bologna, Italy

E-mail address: giovanna.galeati@unibo.it

ABSTRACT

This study was aimed at assessing the capability of semen experimentally infected with porcine circovirus type 2 (PCV2) to produce porcine blastocysts PCR positive for PCV2. Embryos were obtained from *in vitro* maturation (IVM) and *in vitro* fertilization (IVF) of porcine oocytes or by parthenogenesis. Sperm suspension was exposed to PCV2b and utilized for IVF. PCV2 spiked semen did not reveal any reduction in sperm viability and motility but its ability to produce infected blastocysts was irrelevant as only one out of 15 blastocysts obtained by IVF were PCV2b; however 2 blastocysts were PCV2a positive. Furthermore, the presence of PCV2 was demonstrated also in embryos obtained by parthenogenesis (1 out of 17 was PCV2b and one PCV2a positive). Even if PCV2 firmly attaches to the surface of spermatozoa, experimentally spiked sperm were not effective in infecting oocytes during IVF and in producing PCR positive embryos. The infected blastocysts we obtained derived most probably from infected oocytes recovered at abattoir.

Keywords

Blastocysts, IVM-IVF, PCV2, pig, spermatozoa.

INTRODUCTION

Porcine circovirus type 2 (PCV2) related conditions cause economic losses to the pig industry worldwide. Apart from post-weaning multisystemic wasting syndrome (PMWS), PCV2 is associated with a number of conditions collectively known as porcine circovirus diseases (PCVD) (Segales et al. 2004, 2005; Chae 2005). A recent line of studies has focused on PCV2-associated reproductive failure, including irregular return to oestrus, failure of pregnancy progressing to abortion, or reduced litter size with PCV2 being detected in aborted and stillborn foetuses (O'Connor et al. 2001; Ladekjaer-Mikkelsen et al. 2001; Sanchez et al. 2001). Despite the fact that PCV2 is now being regarded as a potentially important emerging pathogen in swine reproduction, its possible interaction with oocytes, spermatozoa and embryos has not been deeply investigated.

In general, oocytes may become infected before ovulation by contact with infectious agents present in either granulosa cells or follicular fluid and after fertilization during the passage through the oviduct and the uterus (Bielanski et al. 2004). These Authors showed that only a small proportion of cumulus–oocyte complexes (COCs) recovered from PCV2 seropositive gilts is positive for viral DNA. They suggested that COCs may have the potential for the production of PCV2-contaminated embryos when follicular fluid containing COCs from infected animals is pooled for *in vitro* fertilization (IVF) as already observed for bovine herpesvirus-1 (BHV-1) and bovine viral diarrhea virus (BVDV) during *in vitro* production of bovine embryos (Bielanski et al. 1993).

In contrast, all *in vivo* fertilized, uterine stage, embryos (developmental range: from four cells to hatched blastocysts) from infected sows tested were negative for PCV2 DNA (Bielanski et al. 2004). These authors suggested that this could be due to the fact that,

despite the excretion of some pathogenic agents into the reproductive tract, uterine stage embryos may not be associated with infectious agents and/or they may be rendered free from infectious agents by a simple sequential washing procedure (Stringfellow 1998). Recently PCV2 nucleic acid has been detected in semen samples from healthy and experimentally infected boars but information regarding the fraction in which PCV2 occurs in semen are conflicting. Pal et al. (2008) showed PCV2 DNA to be present mainly in the cell fraction as opposed to the seminal plasma fraction, whereas Kim et al. (2001) found that PCV2 DNA is mainly present in seminal fluid and nonsperm cell fractions and in only few percent of sperm head.

Due to a recent increase in the use of pig as a model in modern biotechnology applications such as *in vitro* fertilization, *in vitro* embryo production and sperm sexing technology, the potential for infectious disease transmission by pig gametes is a topic of considerable interest.

To the best of our knowledge, no studies have been performed to date on the possibility of inducing PCV2 infection through infected sperm cell.

Basing on above mentioned observations, this study was set up to evaluate the ability of PCV2 experimentally infected semen to produce blastocysts PCR positive for PCV2 from IVM (*in vitro maturation*) - IVF (*in vitro fertilization*) oocytes.

MATERIALS AND METHODS

Unless otherwise specified, all the reagents were purchased from Sigma–Aldrich (Milan, Italy).

Virus

The virus used was a PCV2b strain isolated from an outbreak of PMWS in Italy and genotyped as reported by Hesse et al. (2008). It was propagated in circovirus-free PK15 cells and its identity was confirmed by PCR analysis and reactivity with specific PCV2 monoclonal antibodies. The viral suspension titre was determined by growing serial dilutions in circovirus-free PK15 cell monolayers, followed by immunofluorescent labelling for viral antigen.

Semen processing and exposure of boar spermatozoa to PCV2b

The boar used to collect semen had a proven fertility, was housed since five years in our facility (without contact with other pigs) for experimental purposes and was serologically negative to PCV2. Sperm-rich fraction of ejaculate was collected by gloved hand technique and extended in equal volumes of AndrohepTM (Minitub, Tiefenbach, Germany). The semen resulted negative by PCR to PCV2 DNA (Sarli et al. 2012).

Sperm suspension containing 3×10^9 spermatozoa/100 mL of Androhep was incubated for 4 h at 16°C with 10 mL of a PCV2b viral suspension containing $10^{3.9}$ TCID₅₀/mL.

Semen evaluation

To evaluate a potential negative effect of viral suspension on sperm cells, viability of spermatozoa was analyzed before and after the above described incubation with viral suspension. Briefly, 25 µL of semen were incubated with 2 µL of a 300 µM Propidium Iodide (PI) stock solution and 2 µL of a 10 µM SYBR-14 stock solution (both obtained from the Live/dead sperm viability kit; Invitrogen, Eugene, OR, USA), for 5 min at 37°C in the dark. After incubation, 10 µL of sperm suspensions were analyzed with a

Nikon Eclipse epifluorescence microscope using a double-band-pass filter for green and red fluorescence. The spermatozoa with green or red fluorescence on the head were considered live or dead, respectively. At least 200 spermatozoa were scored for each sample.

***In vitro* maturation (IVM) of cumulus-oocyte-complexes**

Ovaries were collected at a local abattoir and transported to the lab within 2 h in a thermos filled with physiological saline at 30-35°C. Cumulus oocyte complexes (COCs) from follicles 3-6 mm in diameter were aspirated using a 18 gauge needle attached to a 10 mL disposable syringe. Under a stereomicroscope, intact COCs were selected and transferred into a petri dish (35 mm, Nunclon, Roskilde, Denmark) prefilled with 2 mL of modified PBS supplemented with 0.4% BSA. Only COCs with complete and dense cumulus oophorus were used. After three washes in NCSU 37 (Petters & Wells 1993) supplemented with 5.0 mg/mL insulin, 0.57 mM cysteine, 10 ng/mL epidermal growth factor, 50 µM β-mercaptoethanol and 10% PCV2-PCR-negative porcine follicular fluid (IVM medium), groups of 50 COCs were transferred to a Nunc 4-well multidish containing 500 µL of the same medium per well and cultured at 39°C in humidified atmosphere of 5% CO₂/7% O₂ in air. For the first 22 h of *in vitro* maturation, the IVM medium was supplemented with 1.0 mM dibutyryl cyclic adenosine monophosphate (db-cAMP), 10 IU/mL eCG (Folligon, Intervet, Boxmeer, The Netherlands) and 10 IU/mL hCG (Corulon, Intervet). For the last 20 h COCs were transferred to fresh maturation medium (Funahashi et al. 1997). Part of the oocytes was washed in PBS, treated with pronase, in order to remove zona pellucida, rinsed twice in nuclease free water, frozen in 12 pools of 50 and stored at -80°C until PCR analysis.

Collection of immature oocytes, cumulus cells and follicular fluid

COCs and follicular fluids from 80-90 follicles, 3-6 mm in diameter, were aspirated and placed in a petri dish. COCs were then transferred to another petri dish containing 500µL of modified PBS supplemented with 0.4% BSA; the remainig follicular fluids were centrifuged and the supernatants were stored at -80°C until PCR analysis.

Oocytes were mechanically denuded from cumulus cells, then treated with pronase and frozen at -80°C. Cumulus cells were counted using a Thoma's hemocytometer then centrifuged and the pellets stored at -80°C until PCR analysis.

***In vitro* fertilization (IVF) of cumulus-oocyte-complexes**

Matured oocytes were fertilized with infected (infected group; INF) and non-infected (control group; CTR) spermatozoa as follows. Sperm suspensions were washed twice in Brackett&Olifant medium (Bracket & Olifant 1975) supplemented with 12% fetal calf serum (FCS; Gibco, Invitrogen, Monza, Italy) and 0.7 mg/mL caffeine (IVF medium). Sperm concentrations were evaluated and 50 matured oocytes, freed from cumulus cells by gentle repeated pipetting, were transferred to 500 µl IVF medium containing 1×10^6 sperm/mL. After 90 min of co-culture, oocytes were transferred to fresh IVF medium and cultured for an additional 18 h period. Oocytes were then washed twice in NCSU-23 and cultured in 500 µL of the same medium. On day 5 post fertilization, 250 µL of medium were replaced with fresh pre-equilibrated NCSU-23 containing 20% (v/v) FCS to reach a final FCS concentration of 10% (v/v). At day 7 post-fertilization, embryos were examined by stereomicroscope and blastocysts were treated with pronase, in order to remove zona pellucida, rinsed twice in nuclease free water, individually frozen in 5

μL water and stored at –80°C until PCR analysis. Aliquots of infected spermatozoa used for the IVF trials were washed twice with PBS and stored at –80°C until PCR analysis.

Parthenogenetic activation

After IVM, oocytes were denuded as described above, washed three times in IVF medium and then parthenogenetically activated according to the method described by Boquest et al. (2002) slightly modified. Briefly, the oocytes were transferred to IVF medium containing 5 mM ionomycin for 5 min, then washed twice and incubated in NCSU-23 containing 2 mM 6-dimethylaminopurine (6-DMAP) for 3 h at 39°C. Presumptive parthenotes were washed twice in NCSU-23 and cultured in groups of 50 in 500 μL of the same medium. On day 5 postactivation, 250 μL of medium were replaced with fresh pre-equilibrated NCSU-23 containing 20% (v/v) FCS to reach a final FCS concentration of 10% (v/v). At day 7 post-activation, percent of blastocysts was determined and treated as described above.

Effect of sperm washing on virus presence in experimentally infected spermatozoa

Sperm suspension containing 3×10^9 spermatozoa/100 mL of Androhep, was incubated for 4 h at 16° C with 10 mL of a PCV2b viral suspension containing $10^{3.9}$ TCID₅₀/mL. Then 3 aliquots of 1 mL of sperm suspension (30×10^6 spermatozoa) were washed with 10 mL PBS 5 times. After that spermatozoa were stored at –80°C until PCR analysis.

PCV2 detection in blastocysts, oocytes, cumulus cells and follicular fluid

Blastocysts and pool of oocytes were lysed as previously described (Spinaci et al. 2007) and 2 μL were used to perform the first round of PCR for PCV2. Cumulus cells (110000

to 880000 cells/pool) were submitted to DNA extraction with ZR Genomic DNA Tissue Miniprep (Zymo Research, Corp. Irvine, CA, USA) according to manufacturer's instruction. Follicular fluid pools were assayed as follows: 200 μ L samples underwent DNA isolation by using a commercially available kit "NucleoSpin Tissue" (Macherey-Nagel, Düren, Germany) in accordance to manufacturer's instruction. Twenty ng of DNA from cumulus cells and follicular fluid were used for first PCR round for PCV2.

The detection of PCV2a or PCV2b was conducted by nested-polymerase chain reaction (n-PCR). In the first round of PCR, the outer primers used were specific for PCV2. A forward primer 5'-CAGTTCGTCACC-3' (designed using the Beacon Designer 2.07 Software, premier Biosoft International, Palo Alto, CA, USA) and reverse primers 5'-CCGCACCTTCGGATATACTGTC -3' primer (Ouradani et al. 2000) were used. Amplification of DNA was achieved by 35 cycles of 95°C for 1 min, 55°C for 1 min and 72°C for 1 min.

In the second PCR run (n-PCR), a common forward primer, 5'-CACGGATATTGTAGTCCTGGT-3' (Ouradani et al. 2000) and different reverse primers sequences, were used to amplify either PCV2a 5'-GGGGGACCAACAAAATCTC-3' or PCV2b (5'-GGGGCTCAAACCCCGCTC-3') (Hesse et al. 2008). Two μ L of first PCR were used to performed the n-PCR. Amplification of DNA was achieved by 35 cycles of 95°C for 1 min, 62°C for 1 min and 72°C for 1 min. The reaction mixtures contained 3 mM $MgCl_2$; 200mM of each dNTP, 0.2 μ M of each primer; 0.5U of Go Taq Polymerase (Promega Corp, Madison, WI, USA). All reactions were performed in duplicate. PCR products were analyzed on 2% agarose gel stained with ethidium bromide under UV light. The expected PCR products of the first and n-PCR were 627 bp or 391 bp respectively. The specificity of

the primers used was demonstrated by PCR amplification of PCV2a and PCV2b control viruses ($10^{3.3}$ TCID₅₀/mL and $10^{3.9}$ TCID₅₀/mL for PCV2a and PCV2b respectively).

The sensitivity of the PCR was estimated through serial dilutions from 1×10^9 to 10^5 DNA copies of reference positive control (PCV2a and PCV2b). The PCR (first round) had sensitivity of 1×10^5 whereas the n-PCR was 10 times more sensitive (1×10^4).

Amplification of β -Actin gene was performed for all sample as internal control as previously described (Forni et al. 2003).

PCV2 detection in sperm samples

Aliquots of sperms (6×10^6 spermatozoa) were resuspended with denaturing buffer (20 mM TrisHCl; EDTA 20mM; DTT 80mM; SDS 4%; NaCl 200mM; proteinase K 20 mg/mL) and incubated for 3h at 55°C then submitted to genomic DNA extraction by using ZR Genomic DNA (Zymo Research, Corp. Irvine, CA, USA) according to manufacturer's instruction. The first round of PCR was performed with 50 ng of extracted DNA and PCR product was analyzed on 2% agarose gel stained with ethidium bromide under UV light.

Experimental design

Four experiments were performed:

Experiment 1: evaluation of PCV2 presence in blastocysts obtained by IVF with *in vitro* infected or control semen;

Experiment 2: evaluation of PCV2 presence in blastocysts obtained by parthenogenesis;

Experiment 3: evaluation of PCV2 presence in *in vitro* matured oocytes and immature oocytes, cumulus cells and follicular fluid;

Experiment 4: evaluation of PCV2 presence in *in vitro* infected spermatozoa after increasing washing steps.

Statistical analysis

Data on the number of blastocyst obtained were pooled from two replicates and analyzed using SPSS 11 (SPSS Inc., Chicago, IL, USA) with chi-squared test; the level of significance was set at $P<0.05$.

RESULTS

Sperm viability

Sperm viability was not affected by the treatment with viral suspension: it was $82.97 \pm 3.24\%$ before incubation and $81.44 \pm 2.8\%$ (mean \pm SE) after incubation.

Experiment 1: evaluation of PCV2 presence in blastocysts obtained by *in vitro* infected semen

Sperm cells exposed for 4 h to PCV2b were PCR positive. Exposure of spermatozoa to PCV2b did not exert any effect on the percentage of embryos that developed to blastocyst stage (Table 1). In CTR group, 4 out of 15 blastocysts tested were PCV2a positive. In INF group, 1 out of 15 blastocysts screened were PCV2b positive and 2 PCV2a positive (Table 2, Fig. 1).

Experiment 2: evaluation of PCV2 presence in blastocysts obtained by parthenogenesis

At day 7 the percent of blastocysts parthenogenetically activated was 23.92%.

One out of 17 blastocysts parthenogenetically activated was PCV2b positive and one PCV2a positive.

Experiment 3: evaluation of PCV2 presence in *in vitro* matured oocytes and immature oocytes, cumulus cells and follicular fluid

None of the twelve pools of fifty *in vitro* matured oocytes was positive to PCV2b or PCV2a. On the basis of these results PCV2b or PCV2a positivity was assayed in six more consistent pools of immature oocytes, cumulus cells and follicular fluid (from 80-90 follicles). The results obtained are presented in Table 3 and Figure 2.

Experiment 4: evaluation of PCV2 presence in *in vitro* infected spermatozoa after increasing washing steps

As shown in Figure 3 a sequential washing procedure was not effective in removing virus from spermatozoa.

DISCUSSION

The purpose of the present work was to evaluate the ability of PCV2 experimentally spiked semen to produce blastocysts PCR positive for PCV2. Embryos were obtained from *in vitro* maturation and fertilization of porcine oocytes from ovaries collected at a local abattoir.

To our knowledge, the present study is the first one in which spermatozoa have been contaminated with PCV2 and utilized for IVF under experimental conditions.

Our results clearly demonstrate that PCV2 firmly attaches to the surface of spermatozoa (as demonstrated in experiment 4) but they don't seem to be effective in infecting the oocytes during IVF and in producing infected embryos.

The infected blastocysts we obtained derived most probably from infected oocytes obtained at abattoir; in fact, 6 out of 30 blastocysts tested were PCV2a positive. Moreover parthenogenetic blastocysts PCV2b or PCV2a positive were obtained. Similar contamination has been observed also by Bureau et al. (2005).

In order to confirm this hypothesis, we evaluated the presence of PCV2a or PCV2b in different pools of fifty *in vitro* matured oocytes but none of the pools was positive. When more consistent oocyte pools were assayed, few of them resulted positive for PCV2a or PCV2b. Moreover, PCV2 positivity was also recorded in follicular fluids and cumulus cells from pool of follicles.

These results confirm that oocyte from ovaries collected at the slaughterhouse can be infected and therefore could generate infected embryos. Why experimentally spiked semen was not effective in producing infected embryos in our study is not clear.

A hypothesis could be that the virus bound to sperm cells could be entrapped, during the zona pellucida crossing, in the ZP network of filaments with pores/holes and meshes or to the presence of carbohydrates and glycoproteins that can function as receptors for viruses. The PCV2 capsid is constituted by capsomeres, proteins with a high affinity for glycosylated proteins (Misinzo et al. 2006). Since porcine ZP is composed of glycoproteins (pZPA, pZPB, pZPC) (Harris et al. 1994), an interaction between the viral capsid and these glycoproteins might be responsible for the sticky behaviour of the porcine ZP towards PCV2.

It is also important to note that Bureau et al. (2005) reported that hyaluronidase is able to remove PCV2 from contaminated embryos. In fact, those authors demonstrated that when trypsin was used as proteolytic enzyme in washing buffers, more virions remained bound to the zona pellucida than when hyaluronidase was used.

It's well known that sperm head surface is involved in acrosome reaction which results in the release of digestive enzymes (including hyaluronidase) required for zona penetration (Yanagimachi et al. 1994; Flesch & Gadella 2000). Therefore we can hypothesize that, during the fertilization process, membrane capacitation modifications first and acrosome reaction subsequently, with hyaluronidase and other enzymes release, could have induced, at least in part, the detach of PCV2 from sperm head plasma membrane. More recently Blomqvist et al. (2011) showed that Single Layer Centrifugation (SLC) on Androcoll-P, followed by swim-up, is effective in removing PCV2 virus from boar semen. Moreover, Bucci et al. (2013) demonstrated that SLC using Androcoll-P induces some capacitation-related changes in boar sperm membrane, possibly as a consequence of the complete removal of seminal plasma and the partial removal of cholesterol from sperm membrane due to Androcoll centrifugation.

Basing on the above mentioned observations, we tested the grade of spermatozoa infection after 2 h at 39°C in capacitation medium and acrosome reaction induction by adding A23187 as previously described by Bucci et al. (2012). Spermatozoa subjected to *in vitro* induction of capacitation and acrosome reaction showed a reduction of PCV2 amount (unpublished data). These results could explain, at least in part, the inability of sperm cell to produce infected blastocysts.

In conclusion, the present study shows that PCV2 attaches to the surface of spermatozoa but it seems not to have the ability to carry the virus into the oocyte at fertilization and to produce infected embryos.

Conflict of interest statement

None of the authors of this manuscript have any conflict of interest to declare.

Acknowledgements

The work was supported by a Bologna University grant (Ricerca Fondamentale Orientata).

We thank P. Cordioli (Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia- Romagna) for providing PCV2 isolates.

REFERENCES

Bielanski A, Larochelle R, Algire J, Magar R. 2004. Distribution of PCV-2 DNA in the reproductive tract, oocytes and embryos of PCV-2 antibody-positive pigs. *Veterinary Record* **155**, 597–598.

Bielanski A, Loewen KS, Del Campo RM, Sirard MA, Willadsen S. 1993. Isolation of bovine herpesvirus-1 (BHV-1) and bovine viral diarrhea virus (BVDV) in association with the *in vitro* production of bovine embryos. *Theriogenology* **40**, 531-538.

Blomqvist G, Persson M, Wallgren M, Wallgren P, Morrell JM. 2011. Removal of virus from boar semen spiked with porcine circovirus type 2. *Animal Reproduction Science* **126**, 108–114.

Boquest AC, Grupen CG, Harrison SJ, 2002. Production of cloned pigs from cultured fetal fibroblast cells. *Biology of Reproduction* **66**, 1283–1287.

Brackett BG, Oliphant G. 1975. Capacitation of rabbit spermatozoa *in vitro*. *Biology of Reproduction* **12**, 260 –274.

Bucci D, Galeati G, Tamanini C, Vallorani C, Rodriguez-Gil JE, Spinaci M. 2012. Effect of sex sorting on CTC staining, actin cytoskeleton and tyrosine phosphorylation in bull and boar spermatozoa. *Theriogenology* **77**, 1206-1216.

Bucci D, Spinaci M, Morrell J, Vallorani C, Tamanini C, Guidetti R, Galeati G. 2013. Effects of single layer centrifugation with Androcoll-P on boar sperm. *Animal Reproduction Science* **138**, 276-281.

Bureau M, Dea S, Sirard MA. 2005. Evaluation of virus decontamination techniques for porcine embryos produced *in vitro*. *Theriogenology* **63**, 2343-2355.

Chae C. 2005. A review of porcine circovirus 2-associated syndromes and diseases. *Veterinary Journal* **169**, 326–336.

Flesch FM, Gadella BM. 2000. Dynamics of the mammalian sperm plasma membrane in the process of fertilization. *Biochimica et Biophysica Acta* **1469**, 197–235.

Forni M, Zannoni A, Tamanini C, Bacci ML. 2003. Opposite regulation of clusterin and LH receptor in the swine corpus luteum during luteolysis. *Reproduction Nutrition and Development* **43**, 517–25.

Funahashi H, Cantley TC, Day BN. 1997. Synchronization of meiosis in porcine oocyte by exposure to dibutyryl cyclic adenosine monophosphate improves developmental competence following *in vitro* fertilization. *Biology of Reproduction* **57**, 49–53.

Harris JD, Hibler DW, Fontenot GK, Hsu KT, Yurewicz EC, Sacco AG. 1994. Cloning and characterization of zona pellucida genes and cDNAs from a variety of mammalian species: the ZPA, ZPB and ZPC gene families. *DNA Sequence* **4**, 361–393.

Hesse R, Kerrigan M, Rowland RR. 2008. Evidence for recombination between PCV2a and PCV2b in the field. *Virus Research* **132**, 201–207.

Kim J, Han DU, Choi C, Chae C. 2001. Differentiation of porcine circovirus (PCV)-1 and PCV-2 in boar semen using a multiplex nested polymerase chain reaction. *Journal of Virological Methods* **98**, 25–31.

Ladekjaer-Mikkelsen AS, Nielsen J, Storgaard T, Bøtner A, Allan G, McNeilly F. 2001. Transplacental infection with PCV-2 associated with reproductive failure in a gilt. *Veterinary Record* **148**, 759–760.

392

393 Misinzo G, Delputte PL, Meerts P, Lefebvre DJ, Nauwynck HJ. 2006. Porcine
394 circovirus 2 uses heparan sulfate and chondroitin sulfate B glycosaminoglycans as
395 receptors for its attachment to host cells. *Journal of Virology* **80**, 3487-3494.

396

397 O'Connor B, Gauvreau H, West K, Bogdan J, Ayroud M, Clark EG, Konoby C, Allan
398 G, Ellis JA. 2001. Multiple porcine circovirus 2-associated abortions and reproductive
399 failure in a multisite swine production unit. *Canadian Veterinary Journal* **42**, 551–553.

400

401 Ouardani M, Wilson L, Jetté R, Montpetit C, Dea S. 2000. Multiplex PCR for detection
402 and typing of porcine circoviruses. *Journal of Clinic Microbiology* **38**, 1707-1714.

403

404 Pal N, Huang YW, Madson DM, Kuster C, Meng XJ, Halbur PG, Opriessing T. 2008.
405 Development and validation of a duplex realtime PCR assay for the simultaneous
406 detection and quantification of porcine circovirus type 2 and an internal control on
407 porcine semen samples. *Journal of Virological Methods* **149**, 217–225.

408

409 Petters RM, Wells KD. 1993. Culture of pig embryos. *Journal Reproduction Fertility* **48**,
410 61–73.

411

412 Sanchez RE, Nauwynck HJ, McNeilly F, Allan GM, Pensaert MB. 2001. Porcine
413 circovirus 2 infection in swine fetuses inoculated at different stages of gestation.
414 *Veterinary Microbiology* **83**, 169–176.

415

Sarli G, Morandi F, Panarese S, Bacci B, Ferrara D, Bianco C, Fusaro L, Bacci ML, Galeati G, Dottori M, Bonilauri P, Lelli D, Leotti G, Vila T, Joisel F, Allan G, Benazzi C, Ostanello F. 2012. Reproduction in porcine circovirus type 2 (PCV2) seropositive gilts inseminated with PCV2b spiked semen. *Acta Veterinaria Scandinavica* **54**, 51.

Segalés J, Allan GM, Domingo M. 2005. Porcine circovirus diseases. *Animal Health Research Reviews* **6**, 119–142.

Segalés J, Rosell C, Domingo M. 2004. Pathological findings associated with naturally acquired porcine circovirus type 2 associated disease. *Veterinary Microbiology* **98**, 137–149.

Spinaci M, Merlo B, Zannoni A, Iacono E, De Ambrogi M, Turba ME, Zambelli D. 2007. In vitro production of cat blastocysts of predetermined sex using flow cytometrically sorted semen. *Theriogenology* **67**, 872-877.

Stringfellow DA. 1998. Recommendations for the sanitary handling of in-vivo-derived embryos. In: Stringfellow DA, Seidel SM (eds), *Manual of the International Embryo Transfer Society*, pp. 79–84. Int. Embryo Transfer Soc. Savoy, IL.

Yanagimachi R, Knobil E, Neill JD. 1994. Mammalian Fertilization. In: Knobil E, Neill JD (eds), *The Physiology of Reproduction* (2nd edn), pp. 189–317. Raven Press Ltd, New York.

Table 1 Blastocysts obtained by IVF with infected (INF) or non-infected spermatozoa (CTR).

Groups	No. oocytes	No. blastocysts (%)
CTR	250	36 (14.4)
INF	250	33 (13.2)

Table 2 Evaluation of PCV2a or PCV2b presence in blastocysts obtained by IVF.

Groups	No. blastocysts examined	No. blastocysts positive	
		PCV2a	PCV2b
CTR	15	4	0
INF	15	2	1

Table 3 PCV2a and PCV2b positivity in the six pools of immature oocytes, cumulus cells and follicular fluid.

Pool n°	PCV2a positivity			PCV2b positivity		
	Oocytes	Cumulus cells	Follicular fluids	Oocytes	Cumulus cells	Follicular fluids
1	-	-	-	-	-	+
2	-	-	-	-	-	-
3	-	-	+	-	-	-
4	-	+	-	+	+	-
5	+	-	-	-	+	+
6	-	-	-	-	-	+

Figures captions

Figure 1 Representative image of agarose gels of nested PCR for PCV2a and PCV2b products (**A**, **B**) and β -actin (**C**) of blastocysts. **A**: CTR blastocysts; **B**: blastocysts obtained after IVF with PCV2b experimentally spiked semen. Lanes 1-5; 10-14: blastocysts; Lane 7: PCV2a positive control; Lane 8: PCV2b positive control; Lane 6: negative control of PCR; MWM: DNA molecular weight marker.

Figure 2 Representative image of agarose gels of nested PCR for PCV2a and PCV2b products of different pools (n=6; Lanes 1-6) of: oocytes (**A**); cumulus cells (**B**); follicular fluid (**C**). Lane 7: DNA molecular weight marker. Lane 8: PCV2a and PCV2b positive control.

Figure 3 Representative image of agarose gel of PCV2 PCR of infected spermatozoa (Lane 4) and washed infected spermatozoa (Lanes 1-3). Lane 5: negative control of PCR; Lane 6: PCV2b positive control; MWM: molecular weight marker.





Figure 1



476

477 **Figure 2**

478




Figure 3