

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Determination of lipid and phenolic fraction in two hazelnut (*Corylus avellana* L.) cultivars grown in Poland

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Ciemniewska-Zytkiewicz, H., Verardo, V., Pasini, F., Bryś, J., Koczoń, P., Caboni, M.F. (2015). Determination of lipid and phenolic fraction in two hazelnut (*Corylus avellana* L.) cultivars grown in Poland. *FOOD CHEMISTRY*, 168(1 February 2015), 615-622 [10.1016/j.foodchem.2014.07.107].

Availability:

This version is available at: <https://hdl.handle.net/11585/552099> since: 2016-07-14

Published:

DOI: <http://doi.org/10.1016/j.foodchem.2014.07.107>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Ciemniewska-Zytkiewicz, Hanna; Verardo, Vito; Pasini, Federica; Bryś, Joanna; Koczoń, Piotr; Caboni, Maria Fiorenza

Determination of lipid and phenolic fraction in two hazelnut (*Corylus avellana* L.) cultivars grown in Poland

FOOD CHEMISTRY 2015 vol. 168 pages 615-622

The final published version is available online at:

<https://doi.org/10.1016/j.foodchem.2014.07.107>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Manuscript Number: FOODCHEM-D-13-04870R2

Title: Determination of lipid and phenolic fraction in two hazelnut (*Corylus avellana* L.) cultivars grown in Poland

Article Type: Research Article (max 7,500 words)

Keywords: hazelnuts, tocopherols, sterols, phenolics, phospholipids

Corresponding Author: Dr. Vito Verardo, Ph.D.

Corresponding Author's Institution: Università di Bologna

First Author: Hanna Ciemniewska-Żytkiewicz, PhD student

Order of Authors: Hanna Ciemniewska-Żytkiewicz, PhD student; Vito Verardo, Ph.D.; Federica Pasini, PhD; Joanna Bryś; Piotr Koczoń; Maria Fiorenza Caboni

Abstract: The fatty acid, tocopherol, sterol, phospholipid and phenolic compositions of Polish hazelnuts (Kataloński and Webba Cenny) were examined. Particularly, free + esterified and bound tocopherol, sterol and phenolic compounds were determined.

The major fatty acids found in hazelnuts were oleic and linoleic acids. α -tocopherol was the most abundant tocopherol accounting for 90-92% of the total content. Bound tocopherols represented 45.5 and 21.7% of total tocopherols in Kataloński and Webba Cenny cultivar, respectively. Total free + esterified sterols were between 62.0 and 75.7% of total sterols and β -sitosterol was the first sterol in the two samples. Phosphatidylcholine was the most common phospholipid, accounting for 72.2% for Kataloński and 67.5% Webba Cenny, respectively. The most abundant fatty acids in the phospholipid fraction were oleic equally with palmitic acids.

Twelve free and six bound phenolic compounds were identified and quantified in hazelnut kernel, instead nine free and six bound phenolic compounds were determined in hard shell.

Highlights

- free and bound tocopherol, sterol and phenolic compounds were determined
- Twelve free and six bound phenolic compounds were identified in kernel
- nine free and six bound phenolic compounds were identified in hard shell
- Bound tocopherols represented 45.5 and 21.7% of total tocopherols

**Determination of lipid and phenolic fraction in two hazelnut (*Corylus avellana* L.)
cultivars grown in Poland**

Running title: *Determination of lipid and phenolic compounds in Polish hazelnuts*

Hanna Ciemniewska-Żytkiewicz^{a*}, Vito Verardo^{b**}, Federica Pasini^c, Joanna Bryś^a, Piotr
Koczoń^a, Maria Fiorenza Caboni^{b,c}

^aDepartment of Chemistry, Faculty of Food Sciences, Warsaw University of Life Sciences,
Nowoursynowska St. 166, 02-787 Warsaw, Poland

^bInterdepartmental Centre for Industrial Agri-Food Research (CIRI Agroalimentare),
University of Bologna, piazza Goidanich 60, I-47521, Cesena (FC), Italy

^cDepartment of Agricultural and Food Sciences, University of Bologna, piazza Goidanich 60,
I-47521, Cesena (FC), Italy

* *Corresponding author*. Tel: +48 22 59 37 607, fax: +48 22 59 37 635

***Corresponding author*. Tel: +39 0547 338117, fax: +39 0547 382348

Email address: hanna_ciemniewska_zytkiewicz@sggw.pl (H. Ciemniewska-Żytkiewicz),
vito.verardo@unibo.it (V. Verardo)

Abstract

The fatty acid, tocopherol, sterol, phospholipid and phenolic compositions of Polish hazelnuts (*Kataloński* and *Webba Cenny*) were examined. Particularly, free + esterified and bound tocopherol, sterol and phenolic compounds were determined.

The major fatty acids found in hazelnuts were oleic and linoleic acids. α -tocopherol was the most abundant tocopherol accounting for 90-92% of the total content. Bound tocopherols represented 45.5 and 21.7% of total tocopherols in *Kataloński* and *Webba Cenny* cultivar, respectively. Total free + esterified sterols were between 62.0 and 75.7% of total sterols and β -sitosterol was the first sterol in the two samples. Phosphatidylcholine was the most common phospholipid, accounting for 72.2% for *Kataloński* and 67.5% *Webba Cenny*, respectively. The most abundant fatty acids in the phospholipid fraction were oleic equally with palmitic acids.

Twelve free and six bound phenolic compounds were identified and quantified in hazelnut kernel, instead nine free and six bound phenolic compounds were determined in hard shell.

Keywords: hazelnuts, tocopherols, sterols, phenolics, phospholipids

1. Introduction

Hazelnuts (*Corylus avellana* L.) are the second most popular nuts worldwide just after almonds with a global production average at nearly one million tonnes (MT) annually (Contini, Baccelloni, Massantini & Anelli, 2008; Schmitzer, Slatnar, Veberic, Stampar & Solar, 2011). The world's hazelnut production is mainly covered by two main market players; Turkey produces about 430 000 MT/year and Italy about 130 000 MT/year in unshelled basis (FAOSTAT, 2011). However, there are also smaller but significant producers, such as, USA, Azerbaijan, Georgia, China, Iran, Spain, France, Kirgizstan, Poland and Croatia, listed here in descending order of production abundance.

Poland has no long standing tradition of producing large-fruited hazelnuts, but there has been a constant increase in output since beginning production in the 1980s. In accordance with the FAO database, production output came very recently to over 3000 MT/year in unshelled basis (FAOSTAT, 2011). The cultivation area is mainly composed of scattered and small orchards, located mostly in Lublin region (south-eastern part of Poland) and also in south and central Poland.

The lipid fraction forming the major part of hazelnuts (~60%), is composed of non polar (98.8%) and polar (1.2%) constituents (Alasalvar, Shahidi, Liyanapathirana & Ohshima, 2003a; Alasalvar, Amaral, Satir & Shahidi, 2009a). Triacylglycerols are the major nonpolar lipid class, representing nearly 100% of the total nonpolar lipids in hazelnut oil (Alasalvar, Shahidi, Liyanapathirana & Ohshima, 2003b). Minor lipidic compounds are the sterols and tocopherols: total sterol content varies from about 120 to 250 mg/100 g of hazelnut oil (Alasalvar et al. 2009a; Amaral, Casal, Citova, Santos, Seabra & Oliveira, 2006a; Crews et al. 2005). The differences occur among cultivars, though, sitosterol is the major sterol. Hazelnuts are an excellent source of tocols ranging from 11 to 45 mg/100 g of oil (Alasalvar et al. 2009a; Amaral, Casal, Seabra & Oliveira, 2006b; Crews et al. 2005). The major tocopherol is

α -tocopherol, accompanied by γ - and β -tocopherols. Hazelnut oil has been reported to have the highest α -tocopherol level among tree nut oils (Kornsteriner, Wagner & Elmadfa, et al. 2006).

Hazelnuts are an abundant source of several vitamins and minerals. They are a source of fibre, which has a nutritional function for humans, but also contain an array of phytochemicals including phenolics.

The content of phenolics in hazelnuts serves as a significant criterion in evaluating hazelnut quality (Alasalvar, Karamać, Kosińska, Rybarczyk, Shahidi & Amarowicz, 2009b; Contini et al., 2008). Five phenolic acids have been identified and quantified (both, free and esterified forms) by Alasalvar, Karamać, Amarowicz & Shahidi (2006) and Shahidi, Alasalvar & Liyanapathirana (2007) in the hazelnut kernel and by-products. Recently, Jakopic, Mikulic Petkovsek, Likožar, Solar, Stampar & Veberic (2011) detected nine flavan-3-ols, two benzoic acids derivatives, three flavonols and phloretin glycoside, as free phenolic compounds in hazelnuts.

The phenolics in fruits have been quantified as a freely extractable and the nonextractable polyphenols (Arranz, Saura-Calixto, Shaha & Kroon, 2009). At present, only the extractable phenolics in free and esterified forms of hazelnuts and its by-products have been thoroughly studied (Shahidi, Alasalvar & Liyana-Pathirana, 2007), whereas significant amounts of bioactive bound polyphenols that remain in the residues from free phenolic extraction have not been studied qualitatively yet. However, Yang (2009) took into account the concentration of total, free and bound phenolics in kernels of nine types of tree nuts, including hazelnuts. Nevertheless, the identification of the single bound phenolic compounds have not been studied so far.

The principal aim of this research was to investigate the content of lipidic (including fatty acid composition, tocopherol, sterol and phospholipid analysis), and phenolic compounds of

Polish hazelnut varieties *Kataloński* and *Webba Cenny* commonly cultivated in Poland. Analytical data on total, free and bound tocopherols, as well as sterols and phenolics collected with up-to-date instrumental methods were compared and related to available literature data.

2. Materials and Methods

2.1. Chemicals

All the solvents and reagents were purchased from Merck (Darmstadt, Germany). The standard compounds were supplied by Sigma-Aldrich (Saint Louis, MO, USA).

2.2. Samples

The investigated hazelnuts (*Corylus avellana* L.), *Kataloński* and *Webba Cenny* varieties were procured at the orchard located in the south of Poland (Jankowice, Pszczyna 50°0'5"N 18°59' 18"E) in 2012. The plants (both varieties) were grown in open field with the same agronomical conditions. Preliminary sun-dried nuts (3 days at 20–25 °C) were stored unshelled at 10°C for one month until they were analysed (final moisture content <7% water in d.m.). The hazelnuts *Kataloński* and *Webba Cenny* listed in *The Polish National List of Fruit Plant Varieties 2013* are the varieties of which seed material is eligible for production and marketing in Poland and is allowed to be distributed on the European Union territory. Hazelnuts were manually cracked and shelled before chopping. The fibrous skin, particularly distinctive for *Kataloński* variety, was removed by hand and a brown skin (pellicle) was left for analysis.

2.3. Moisture content

Approximately 5 g of the sample were dried in a drying oven for 8 hours in 105°C until constant weight according to AOAC (1995), following with moisture calculations.

2.4. Total lipid extraction (TL)

The hazelnut oil was extracted using the procedure described by Boselli, Velazco, Caboni & Lercker (2001). Approximately 10 g of the sample was homogenised with 100 mL of a chloroform/methanol solution (1/1 v/v) in a glass bottle with a screw-cap. The bottle was kept at 60°C for 20 min before adding an additional 100 mL of chloroform. After 3 min of homogenisation, the content of the mixture was filtered through the filter paper. The filtrate was mixed thoroughly with 70 mL of 1 M KCl solution and left overnight at 4°C in order to phase separation. The organic phase was collected and dried with a rotary evaporator at 40°C, dissolved in 5 mL *n*-hexane/isopropanol solution (4/1, v/v) and stored at -18°C until it was analysed.

2.5. Isolation of total unsaponifiable compounds by hot saponification (HS)

Hot saponification was performed as described by Caligiani, Bonzanini, Palla, Cirlini & Bruni (2010) with minor modifications. Briefly, approximately 10 g of hazelnuts were finely chopped using the manual chopper (Kitchen Mate, UK), the 0.5 mL of dihydrocholesterol (c = 2.0 mg/mL) was added and saponification was carried out by boiling and stirring for 1 h with 100 mL of 1.0 N potassium hydroxide in ethanol-water solution (4/1 v/v). After cooling, 100 mL of distilled water was added, and the sample transferred to a separating funnel and extracted four times with 50 mL of diethyl ether. The ether extracts were pooled into a separating funnel and washed four times with 50 mL of distilled water. The organic phase was dried with anhydrous sodium sulphate, filtered, dried and the residue was weighed.

2.6. Total fatty acid analysis

The fatty acid composition of hazelnut oil samples was determined from TL extract as FAMES by capillary gas chromatography analysis after alkaline treatment as described by

Christie (1982). The chromatographic conditions were the same as reported by Verardo, Gómez-Caravaca, Gori, Losi, & Caboni (2013).

2.7. Tocopherol analysis

Due to free and total tocopherols determination, approximately 300 mg of fat from TL and 5 mg of fat from HS extract were dissolved in 1 mL and 7 mL of *n*-hexane, respectively. The solutions were filtered through a 0.45 µm nylon filter. The tocopherols were determined by HPLC (Agilent 1200 series, Palo Alto, CA, USA) equipped with a fluorimeter detector (Agilent, Palo Alto, CA, USA). The excitation wavelength was 290 nm and the emission one was 325 nm. The column used was a Luna Hilic Phenomenex column (250 mm x 4.6 mm i.d., 5 µm particle size) in isocratic conditions as reported by Gómez-Caravaca, Verardo & Caboni (2010). Calibration curve was constructed with α -tocopherol standard solution and it was used for quantification.

2.8. Free + esterified and bound phytosterol (PS) analysis

To determine the free + esterified phytosterols, approximately 0.5 mL of dihydrocholesterol (c = 2.0 mg/mL) was added to 250 mg of fat from TL extract and saponification was conducted at room temperature. After saponification, the organic fraction was washed with 10 mL diethyl ether/water (1/1 v/v), and further the unsaponifiable matter was extracted in sequence; two times with 10 mL diethyl ether, and washed two times with 10 mL 0.5 N aqueous KOH and again two times with 10 mL of distilled water. The diethyl ether solvent was removed under vacuum and the unsaponifiable matter was used for the free + esterified sterol determination.

To determine the total sterols, 5 mg of fat from HS extract was used for analysis.

The unsaponifiable matter from TL and HS extracts were silylated according to Sweeley, Bentley, Makita & Wells (2002), and they were analysed using a GC/MS (GCMS-QP2010 Plus, Shimadzu, Tokyo, Japan) in the chromatographic conditions reported by Cardenia, Rodriguez-Estrada, Baldacci, Savioli & Lercker (2012).

Data were filed and processed by the software GCMSsolution ver. 2.50 SU1 from Shimadzu. PS identification was achieved by comparing peak mass spectra with peaks of standard mixture and with GC-MS data reported by Pelillo, Iafelice, Marconi & Caboni (2003). Quantification of identified phytosterols was performed in relation to dihydrocholesterol used as internal standard.

2.9. Phospholipid determination

To determine the phospholipids in hazelnut, approximately 100 mg of fat from TL extract were weighed and dissolved in 1 mL of 88/12 (v/v) chloroform/methanol system and used for the HPLC analysis.

The quantitation of the phospholipid classes was performed using HPLC-ELSD. The chromatographic method used for the separation of the polar lipids extracted from hazelnut oil by Verardo, Gomez-Caravaca, Gori, Losi & Caboni (2013) was carried out. Phospholipid separation was performed on an Agilent liquid chromatography HP 1200 Series (Agilent Technologies, Palo Alto, California, USA). The detector was an evaporative light scattering detector (ELSD; PL-ELS1000, Polymer Laboratories, Church Stretton, Shropshire, UK). The control of the HPLC system was accomplished by the software Agilent ChemStation (Agilent Technologies, Santa Clara, CA, USA) whilst chromatograms registration and data processing were assessed by ClarityLite (ver. 2.4.0.190, DataApex, Praha, The Czech Republic). The separation was achieved using a silica column, 150 mm \times 3 mm with 3 μ m particle diameter

(Phenomenex, Torrance, CA, USA). The calibration curves were prepared separately for each phospholipid identified.

2.10. Determination of phospholipid fatty acids

About 20 mg of fat from TL extract were dried under nitrogen, dissolved in 0.5 mL of chloroform and loaded on a Silica gel 60 TLC plate 20 x 20 cm (Merck KGaA, Darmstadt, Germany). The mobile phase was 100 mL of a mixture *n*-hexane/diethyl ether 3/2 (v/v). Phospholipid band was visualised under UV light (254 nm) by spraying the TLC plate with a 0.02% (m/v) ethanolic solution of 2,7-dichlorofluorescein (sodium salt) and then scraped off and collected. Phospholipids were extracted three times with chloroform (3 x 1 mL). Organic extracts were pooled, dried under nitrogen and, to convert fatty acids to the corresponding methyl esters (FAMES), the method of Christie (1982) was carried out. FAMES were determined by GC-FID according to Verardo et al. (2013).

2.11. Extraction and determination of free phenolic compounds

The free phenolic compounds were extracted using the optimised protocol described by Verardo, Bendini, Cerretani, Malaguti, Cozzolino & Caboni (2009). Briefly, 4 g of the chopped hazelnuts kernel and shell were extracted in an ultrasonicator using 40 mL of ethanol/water solution (4/1 v/v) at 40°C for 10 min. After centrifugation at 3,500 rpm for 15 min, the supernatant was collected and the residue was re-extracted under the same conditions. The free phenolic composition of the extracts was determined with RP-HPLC–DAD–MS, as previously described by Bocalandro et al. (2012).

2.12. Extraction and determination of bound phenolic compounds

Both, the kernel and shell residues of free phenolics ethanol extraction were taken under alkaline hydrolysis as reported by Bonoli, Verardo, Marconi & Caboni (2004). Briefly, the residue from free phenolic extraction was digested with 100 mL of 2 N NaOH in distilled water at room temperature for 20 h with no light access and stirring under nitrogen gas. The mixture was then brought to pH 2-3 by adding 37% hydrochloric acid in a cooling ice bath and extracted with 100 mL of *n*-hexane to remove the lipid fraction. The final solution was extracted three times with 100 mL of ethyl acetate/diethyl ether (1/1 v/v). The organic fractions were pooled and evaporated to dryness. The phenolic compounds were reconstituted with 4 mL of 1/1 methanol/water (v/v) and stored at -18°C until it was analysed. The polyphenol composition of the extracts was determined with RP-HPLC–DAD–MS, as previously described by Bocalandro et al (2012).

2.13. Statistical analysis

Relative standard deviation was obtained, where appropriate, for all data collected. One-way analysis of variance, ANOVA (Tukey's honest significant difference multiple comparison) was evaluated using Statistica 8 software (2006, StatSoft, Tulsa, OK, USA). *p*-values lower than 0.05 were considered statistically significant. All chemical analyses were carried out in triplicate (n=3) for each sample, and the analytical data were used for statistical comparisons.

3. Results and Discussion

3.1. Fatty acid composition

The fatty acid composition of studied hazelnuts is given in Table 1. The predominant fatty acid was oleic acid (C18:1 ω9), followed by linoleic acid (C18:2 ω6), with the mean values being in close agreement with the ranges reported elsewhere (Amaral et al., 2006a; Alasalvar et al., 2003b; Bacchetta et al., 2013). Based on the analysis conducted, statistically significant

differences ($p<0.05$) concerning content of major fatty acids in *Kataloński* and *Webba Cenny* were found.

Kataloński reported higher percentages of linoleic acid and, consequently, PUFA content as compared to *Webba Cenny*. *Webba Cenny* however showed the higher content of oleic acid, MUFA and SFA than *Kataloński*. Several authors (Alasalvar et al., 2003a) demonstrated that the content of oleic acid in various hazelnut cultivars is inversely correlated with the content of linoleic acid. This is reasonably consistent with the results herein reported for *Kataloński* and *Webba Cenny* cultivars.

Other major fatty acids were palmitic and stearic acids that represented more than 4 and 1% of total fatty acids, respectively.

SFA content for *Kataloński* and *Webba Cenny* varieties in present study were less than 6 and 7%, respectively. The average SFA content of 75 cultivars from Spain, Italy, Greece, Slovenia, France and Portugal studied by Bacchetta et al. (2013) accounted for 8.43%, total SFA value in the Portuguese cultivars ranged from 7.5 to 10% (Amaral et al., 2006a), and Turkish cultivar made up with 7.85% SFA of the total fatty acids of hazelnut oil (Alasalvar et al., 2003b). The data obtained in this work can be due to the different weather conditions in Poland; that are colder and rainy if compared with the Mediterranean climate of Turkey, Italy or Spain (Gantner, 2005). However, according to Piskornik & Korfel (1993), nuts obtained in these conditions contain considerably less saturated fatty acids (SFA).

3.2. Free (+ esterified), bound and total tocopherol and sterol content

Firstly, all tocopherol and sterol values presented in literature on hazelnuts so far have been reported on the oil basis and expressed in mg/kg, µg/g or mg/100g of oil (Matthäus & Özcan, 2012; Kornsteriner et al., 2006). However, the total values of tocopherols and sterols

expressed on whole-nut basis have not been published yet, excepting one report on total α -tocopherol content presented by Alasalvar et al. (2003a).

In the present study, the hot saponification was preliminary tested using 2.2 N KOH solution, as reported by Caligiani et al. (2010), however, the tocopherols, especially β - and γ -homologues were degraded during the saponification process. The similar KOH concentration for direct saponification (11% w/v solution \approx 2 N) was previously used by Katsanidis & Addis (1999) on animal origin matrix and they concluded that the investigated method is only useful for the analysis of the α -homologues. Because of that, lower KOH concentrations were tested (data not showed) and the data underlined that a solution of KOH 1 N reported better results for this matrix.

After establishing the procedure, free, bound and total tocopherols were determined. Results are shown in Table 2 and were expressed as dry base weight (moisture was 3.96 and 3.34 % w/w for *Kataloński* and *Webba Cenny* cultivars, respectively). Among the tocopherols identified, α -tocopherol was most abundant accounting for 90-92% of the total tocopherol content, followed by γ - (4.3-7.0%) and β -tocopherol (2.4-4.1%).

The extraction method (TL and HS) influenced significantly ($p < 0.05$) all tocopherol recovery and total amounts of α -, γ - and β -homologues. Herein, total tocopherol content was 236.2 and 205.8 mg/kg of d.w. nut for *Kataloński* and *Webba Cenny* cultivars, respectively. The α , γ and β tocopherols obtained directly from hot saponification of the matrix were on average 1.5, 1.3 and 2.0 fold higher, respectively, than free forms obtained on the oil basis. These data confirmed that bound tocopherols are contained in hazelnuts and the direct saponification of the sample is needed to determine their content.

The free α -tocopherol concentration also expressed as mg/kg d.w. nut amounted to 204.3 and 186.5 for *Kataloński* and *Webba Cenny* cultivars, respectively. These results were in the same order of magnitude as the data reported by Alasalvar et al. (2003a) where α -tocopherol was

289 determined with concentration of 24 mg/100 g of nut. In order to compare the hazelnuts
 290 cultivated in Poland with wider range of literature, the results of free tocopherols were
 291 recalculated and expressed on the oil basis. As can be seen in Supplemental Figure S1,
 292 *Kataloński* cultivar had significantly ($p<0.05$) higher concentration of α - (298.6 $\mu\text{g/g}$ oil), γ -
 293 (20.7 $\mu\text{g/g}$ oil) and total tocopherols (327.1 $\mu\text{g/g}$ oil) than *Webba Cenny* (283.6, 18.6, 309.6
 294 $\mu\text{g/g}$ oil, respectively). In case of β -tocopherol (*Kataloński* 7.8 $\mu\text{g/g}$ oil, *Webba Cenny* 8.4
 295 $\mu\text{g/g}$ oil), there was no statistically significant differences between cultivars ($p<0.05$). In
 296 general, results of free tocopherol contents are in agreement with available data (Kornsteriner
 297 et al., 2006; Alasalvar et al., 2003b; Savage, McNeil & Dutta, 1997).

298 Bound tocopherols represented the 45.5 and 21.7% of total tocopherols in *Kataloński* and
 299 *Webba Cenny* cultivar, respectively. As reported for the free form, α -tocopherol was the first
 300 bound tocopherol and accounted for the 81.0 and 88.2% of total bound tocopherols in
 301 *Kataloński* and *Webba Cenny* samples, respectively. Bound γ -tocopherol was the second
 302 bound tocopherol and its content was 15.4 and 3.1 mg/kg of nut for *Kataloński* and *Webba*
 303 *Cenny* samples representing the 14.3 and 6.9% of total bound tocopherols, respectively.
 304 Bound β -tocopherol accounted for the 4.7 and 4.9% of total bound tocopherol content.
 305

306 Seven sterols and three stanols were identified. Table 3 presents their free + esterified, bound
 307 and total form content.

308 Total content (free + esterified + bound) of phytosterols found in *Kataloński* and *Webba*
 309 *Cenny* samples was 1522.2 and 1303.2 mg/kg d.w. nut, respectively. These results are in the
 310 same order of magnitude as those of Nyström, Schär & Lampi (2012).

311 No statistical differences were reported in free + esterified phytosterols content in both
 312 samples; total free + esterified sterol compounds were the 62.0 and 75.7% of total sterols in
 313 *Kataloński* and *Webba Cenny* samples, respectively ($p<0.05$). Sitosterol was the first free

sterol in the two samples tested and accounted for the 84.0% of total free + esterified sterols. The other free + esterified sterols, in decreasing order of abundance, were campesterol (53.5 and 50.0 mg/kg d.w. for *Kataloński* and *Webba Cenny* samples), and Δ^5 -avenasterol (36.7 and 39.5 mg/kg d.w. for *Kataloński* and *Webba Cenny* samples). Other major compounds were the saturated sterols, namely sitostanol and campestanol.

Other minor free + esterified sterol compounds that were determined were Δ^7 -avenasterol, stigmasterol, chlerosterol, fucosterol and cholesterol.

Bound sterols in *Kataloński* and *Webba Cenny* samples were the 38 and 24% of total sterols, respectively. As reported for the free sterols, sitosterol, campesterol and Δ^5 -avenasterol were the first, second and third bound sterols in all samples. The sum of sitosterol, campesterol and Δ^5 -avenasterol was 92 and 85.5% of total bound sterols in *Kataloński* and *Webba Cenny*, respectively.

In order to provide comparable results, the concentration of free + esterified sterols was recalculated and expressed on the oil basis (Supplemental Figure S2). The most abundant phytosterols, namely, sitosterol, campesterol, and Δ^5 -avenasterol found in *Kataloński* and *Webba Cenny* oils accounted for 1894.5 and 1689.7 mg/kg, 128.9 and 101.5 mg/kg, 88.4 and 80.2 mg/kg, respectively. Obtained sterol results for both cultivars are in the wide agreement with those presented in literature. As an example, Matthä us & Özcan (2012) determined that hazelnut cultivars oil grown in Turkey contained 1222.2–4947.3 mg/g sitosterol, 81.1–445.9 mg/g campesterol, and 60.1–143.1 mg/g Δ^5 -avenasterol, whereas Savage et al. (1997) reported 1416–1693 mg/g sitosterol, 78–114 mg/g campesterol, and 33–170 mg/g Δ^5 -avenasterol.

3.3. Phospholipid (PL) analysis

The phospholipid content and the distribution of individual phospholipid species were determined in the two hazelnut cultivars (Table 4). Three phospholipids (PC – phosphatidylcholine, PE – phosphatidylethanolamine; PI – phosphatidylinositol) were fully separated and identified according to Parcerisa, Codony, Boatella & Rafecas (1999) and Alasalvar et al. (2003b). The *Kataloński* cultivar was significantly higher in total phospholipid content than the *Webba Cenny* ($p < 0.05$). The PC fraction was the most common phospholipid, accounting for 72.2% for *Kataloński* and 67.5% *Webba Cenny*, and followed by PE and PI, respectively. These results are in the close agreement with the data of Mirliakbari & Shahidi (2008), that reported the contents of PC and PI in the amount of 4.8 and 0.8 mg/g of oil, respectively. Different results were reported by Parcerisa et al. (1999), where the total PL content were accounted as ten fold lower (1.09 mg/g of oil). The occurred discrepancy could be coupled with the applied response factor for the PL calibration curve; herein the calibration curve was prepared separately for PC, PE and PI, whereas Parcerisa et al. (1999) assumed the PC response factors for PE and PI calculation. Additionally, the same authors (Parcerisa et al., 1999) analysed different cultivars that were obtained in different harvest time, agronomical management, soil, and climate conditions.

Nevertheless, limited information regarding phospholipids in hazelnuts is available in the literature.

To determine the phospholipid fatty acids and to obtain the purified phospholipid fraction from the total lipid extract, a TLC separation was carried out. This technique permits the isolation of the different lipid classes (Montealegre, Verardo, Gómez-Caravaca, García-Ruiz, Marina & Caboni, 2012).

The most abundant fatty acids in the PL fraction were oleic equally with palmitic acid, followed by stearic, palmitoleic and linoleic acids, whereas *Kataloński* cultivar was approximately three fold richer in the linoleic acid than *Webba Cenny* (Table 5). Fatty acid

profiles detected for PL fraction of both cultivars were consistently different and richer in SFA than that observed for total lipids (triglycerides + polar fats) according to Parcerisa et al. (1999) that reported the considerable higher concentration of SFA in PC than in TAG class.

3.4. Determination of free and bound phenolic compounds in kernel and shell

Phenolic compounds in raw shelled hazelnuts were analysed. Two different extraction methods were carried out to obtain free and bound phenolics from *Kataloński* and *Webba Cenny* cultivars grown in Poland.

The extracted compounds were identified by analysing UV and MS data and quantified by DAD detection. The quantification was performed by comparison with calibration curves of proanthocyanidin B-2 (0.03–1.00 g/L), taxifolin (0.06–1.00 g/L), (+)-catechin (0.10–1.00 g/L), gallic acid (0.12–1.20 g/L), quercetin (0.10–1.0 g/L), phloretin (0.05–1.00 g/L) and syringic acid (0.10–1.0 g/L) dissolved in methanol and filtered with a 0.20 µm filter.

Twelve free phenolic compounds were identified and quantified in hazelnut kernel. Briefly, two phenolic acids (gallic and protocatechuic acids) and ten flavonoids (catechin and epicatechin, 2 procyanidin dimmers, 2 procyanidin trimers, 3 flavonols and 1 chalcone) were determined. These compounds were previously identified in hazelnut kernel by several authors (Shahidi et al., 2007; Jakopic et al., 2011; Schmitzer et al., 2011).

Table 6 shows all the identified compounds determined in the hazelnut samples. *Kataloński* cultivar showed the highest content of total free phenolic compounds (31.7 µg/g d.w. nut) and its content was about two times higher than for *Webba Cenny*.

Phenolic acids accounted for the 38 and 32% of total free phenolic compounds in *Kataloński* and *Webba Cenny* kernel, respectively. Gallic acid was the first phenolic acid in the two samples and its concentration ranged from 4.1 µg/g in *Webba Cenny* to 11.1 µg/g in *Kataloński*, where the highest concentration of this phenolic compound was reported. These

results accorded to other authors (Alasalvar, Pelvan & Amarowicz, 2010; Pelvan, Alasalvar & Uzman, 2012).

Flavan-3-ols were the first free phenolic compounds in the both samples; in fact catechin and their oligomers represented the 48 and 89% of total free phenolics, respectively. Procyanidin dimmers were contained in higher quantities and ranged from 9.7 to 11.8 $\mu\text{g/g}$ d.w. nut in the analysed samples. These results were in the same order of magnitude as those reported in literature (Jakopic et al., 2011; Schmitzer et al. 2011).

Three flavonols, namely myricetin-3-O-rhamnoside, quercetin-pentoside and quercetin-3-O-rhamnoside were identified and quantified. According to Jackopic et al. (2011) quercetin-rhamnoside was the first flavonol, and herein it is two times higher in *Kataloński* than in *Webba Cenny*.

Six bound phenolic compounds were identified in the hazelnut kernels (Table 6). All of them are phenolic acids.

Kataloński and *Webba Cenny* samples showed a similar total content of phenolic compounds ranged between 17.4 and 19.7 $\mu\text{g/g}$ d.w.

Compound with principal fragment ions at m/z 179 and 139 m/z was identified as caffeic acid derivative and it was the first bound phenolic compound in the two studied cultivars. It accounted for 40.3 and 71.9% of total bound phenolics in *Kataloński* and *Webba Cenny* samples, respectively.

The other compounds that were identified in both samples, according to the following characteristic ions, were: $[\text{M}-\text{H}]^-$ 153, corresponding to protocatechuic acid; $[\text{M}-\text{H}]^-$ 167, vanillic acid; $[\text{M}-\text{H}]^-$ 223, sinapic acid; $[\text{M}-\text{H}]^-$ 163, *p*-coumaric acid; $[\text{M}-\text{H}]^-$ 191, quinic acid. Their content in two cultivar tested was reported in Table 6.

HPLC-DAD-ESI-MS was used to determine the free and phenolic component present in hard shell hazelnuts. Up-to-date, there are no sufficient data available on free phenolics contained

413 in hazelnut hard shell. Shahidi et al. (2007) reported free phenolic compounds in hazelnut
414 hard shell by HPLC-DAD method for the first time, but there were tentatively identified and
415 quantified merely five phenolic acids. The other authors determined the free phenolic fraction
416 in the hazelnut shell by spectrophotometrical methods (Stévigny, Rolle, Valentini & Zeppa,
417 2007; Contini et al., 2008).

418 Nine free phenolic compounds were identified in *Kataloński* and *Webba Cenny* samples used
419 MS detector. Briefly, five phenolic acids, three flavonoids and a chalcone compound were
420 determined in the hazelnut samples (Table 6). Total free phenolic content in *Kataloński*
421 samples was two fold higher than in *Webba Cenny*.

422 According to Shahidi and co-workers (2007), gallic acid was the first phenolic acid in hard
423 shell; its content was 2.7 and 0.8 µg/g d.w. nut in *Kataloński* and *Webba Cenny*, respectively.
424 The second and third phenolic acids were protocatechuic and vanillic acids respectively.
425 Moreover, sinapic and *p*-coumaric acids were, also, identified, but their content was lower
426 than LOQ. Quercetin-pentoside and catechin were the principal flavonoids in the hard shell
427 but myricetin-3-O-rhamnoside and phloretin-2-O-glucoside were also identified and
428 quantified in both tested cultivars.

429 As far as we know, six bound phenolic compounds were identified and quantified in hazelnut
430 shell for the first time. Total bound phenolic content in hard shell was about 7-8 times higher
431 than free phenolic content.

432 Four phenolic acids and two flavonoids were identified (Table 6). Phenolic acids were the
433 73.1 and 71.3% of total bound phenolic compounds in *Kataloński* and *Webba Cenny*,
434 respectively. Protocatechuic and vanillic acids were the first phenolic acids and their content
435 was higher than 30 µg/g d.w. nut. Gallic acid and its quinic derivative were also quantified.

The two representative flavonoids were taxifolin and catechin. Taxifolin was not previously described in hazelnut but it is a component of other woody structure, such as pine bark (Bocalandro et al. 2012).

4. Conclusions

To sum up, different extraction methods were used for examining lipids and phenolic antioxidants of two hazelnut cultivars grown in Poland. The results confirmed that direct saponification of hazelnut is necessary to determine the total sterol and tocopherol content, because these components in hazelnuts are present in both free and bound forms.

HPLC-DAD-ESI-MS was used to identify and quantify the free and bound phenolic compounds in hazelnut kernels and in the hard shells. Hazelnut shells represent a rich source of natural phenolic antioxidants.

Acknowledgements

The authors wish to thank Mr. Patrick MacNeil who assisted in the proof-reading of the manuscript.

References

- Alasalvar, C., Amaral, J., Satir, G., Shahidi, F. (2009a). Lipid characteristics and essential minerals of native Turkish hazelnut varieties (*Corylus avellana* L.). *Food Chemistry*, 113, 919–25.
- Alasalvar, C., Karamać, M., Amarowicz, R., Shahidi, F. (2006). Antioxidant and antiradical activities in extracts of hazelnut kernel (*Corylus avellana* L.) and hazelnut green leafy cover. *Journal of Agricultural and Food Chemistry*, 54, 4826–4832.

460 Alasalvar, C., Karamać, M., Kosińska, A., Rybarczyk, A., Shahidi, F., Amarowicz, R.
 461 (2009b). Antioxidant activity of hazelnut skin phenolics. *Journal of Agricultural and Food*
 462 *Chemistry*, 57, 4645–4650.

463 Alasalvar, C., Pelvan, E., Amarowicz, R. (2010). Effects of roasting on taste-active
 464 compounds of turkish hazelnut varieties (*Corylus avellana* L.). *Journal of Agricultural and*
 465 *Food Chemistry*, 58, 8674–8679.

466 Alasalvar, C., Shahidi, F., Liyanapathirana, C., Ohshima, T. (2003a). Turkish Tombul
 467 hazelnut (*Corylus avellana* L.). 1. Compositional characteristics. *Journal of Agricultural*
 468 *and Food Chemistry*, 51, 3790–3796.

469 Alasalvar, C., Shahidi, F., Ohshima, T., Wanasundara, U., Yurttas, H.C., Liyanapathirana,
 470 C.M., Rodrigues, F.B. (2003b). Turkish tombul hazelnut (*Corylus avellana* L.). 2. Lipid
 471 characteristics and oxidative stability. *Journal of Agricultural and Food Chemistry*, 51,
 472 3797–3805.

473 Amaral, J., Casal, S., Citova, I., Santos, A., Seabra, R., Oliveira, B. (2006a). Characterization
 474 of several hazelnut (*Corylus avellana* L.) cultivars based in chemical, fatty acid and sterol
 475 composition. *European Food Research and Technology*, 222, 274–80.

476 Amaral, J., Casal, S., Seabra, R., Oliveira, B. (2006b). Effects of roasting on hazelnut lipids.
 477 *Journal of Agricultural and Food Chemistry*, 54, 1315–1321.

478 AOAC. 1995. Official Methods of Analysis, Association of Official Analytical Chemists,
 479 Washington, DC, USA.

480 Arranz, S., Saura-Calixto, F., Shaha, S., Kroon, P. A. (2009). High contents of nonextractable
 481 polyphenols in fruits suggest that polyphenol contents of plant foods have been
 482 underestimated. *Journal of Agricultural and Food Chemistry*, 57, 7298–7303.

483 Bacchetta, L., Aramini, M., Zini, A., Di Giammatteo, V., Spera, D., Drogoudi, P., Rovira, M.,
 484 Silva, A. P., Solar, A., Botta, R. (2013). Fatty acids and alpha-tocopherol composition in

485 hazelnut (*Corylus avellana* L.): a chemometric approach to emphasize the quality of
 486 European germplasm. *Euphytica*, 191, 57–73.

487 Bocalandro, C., Sanhueza, V., Gómez-Caravaca, A.M., Gonzalez-Alvarez, J., Fernandez, K.,
 488 Roeckel, M., Rodriguez-Estrada, M.T. (2012). Comparison of the composition of *Pinus*
 489 *radiata* bark extracts obtained at bench- and pilot-scales. *Industrial Crops and Products*,
 490 38, 21–26.

491 Bonoli, M., Verardo, V., Marconi, E., Caboni, M.F. (2004). Antioxidant Phenols in barley
 492 (*Hordeum vulgare* L.) flour: comparative spectrophotometric study among extraction
 493 methods of free and bound phenolic compounds. *Journal of Agricultural and Food*
 494 *Chemistry*, 52, 5195–5200.

495 Boselli, E., Velazco, V., Caboni, M. F., Lercker, G. (2001). Pressurized liquid extraction of
 496 lipids for the determination of oxysterols in egg-containing food. *Journal of*
 497 *Chromatography A*, 917, 239–244.

498 Caligiani, A., Bonzanini, F., Palla, G., Cirlini, M., Bruni, R. (2010). Characterization of a
 499 potential nutraceutical ingredient: pomegrenate (*Punica granatum* L.) seed oil saponifiable
 500 fraction. *Plant Foods for Human Nutrition*, 65, 277–283.

501 Cardenia, V., Rodriguez-Estrada, M.T., Baldacci, E., Savioli, S., Lercker, G. (2012). Analysis
 502 of cholesterol oxidation products by fast gas chromatography/mass spectrometry. *Journal*
 503 *of Separation Science*, 35, 424–430.

504 Christie, W.W. (1982). A simple procedure for rapid transmethylation of glycerolipids and
 505 cholesteryl esters. *Journal of Lipid Research*, 23, 1072–1075.

506 Contini, M., Baccelloni, S., Massantini, R., Anelli, G. (2008). Extraction of natural
 507 antioxidants from hazelnut (*Corylus avellana* L.) shell and skin wastes by long maceration
 508 at room temperature. *Food Chemistry*, 110, 659–669.

509 Crews, C., Hough, P., Godward, J., Brereton, P., Lees, M., Guiet, S., Winkelmann, W. (2005).
 510 Study of the main constituents of some authentic hazelnut oils. *Journal of Agricultural and*
 511 *Food Chemistry*, 53, 4843–4852.

512 FAOSTAT (2011), <http://faostat.fao.org/>, (access date: 04.09.2013).

513 Gantner, M. (2005). Susceptibility of large-fruited hazel cultivars grown in Poland to major
 514 pest and their crop productivity. *Acta Horticulturae*, 686, 377-384.

515 Gómez-Caravaca, A.M., Verardo, V., Caboni, M.F. (2010). Chromatographic techniques for
 516 the determination of alkyl-phenols, tocopherols and other minor polar compounds in raw
 517 and roasted cold pressed cashew nut oils. *Journal of Chromatography A*, 1217, 7411–7417.

518 Jakopic, J., Mikulic Petkovsek, M., Likozar, A.; Solar, A., Stampar, F., Veberic, R. (2011).
 519 HPLC–MS identification of phenols in hazelnut (*Corylus avellana* L.) kernels. *Food*
 520 *Chemistry*, 124, 1100–1106.

521 Katsanidis, E., Addis, P.B. (1999). Novel HPLC analysis of tocopherols, tocotrienols and
 522 cholesterol in tissue. *Free Radical Biology and Medicine*, 27, 1137–1140.

523 Kornsteriner, M., Wagner, K.H., Elmadfa, I. (2006). Tocopherols and Total phenolics in 10
 524 different nut types. *Food Chemistry*, 98, 381–387.

525 Matthäus, B., Özcan, M.M. (2012). The comparison of properties of the oil and kernels of
 526 various hazelnuts from Germany and Turkey. *European Journal of Lipid Science and*
 527 *Technology*, 114, 801–806.

528 Miraliakbari, H., Shahidi, F. (2008). Lipid class compositions, tocopherols and sterols of tree
 529 nut oils extracted with different solvents. *Journal of Food Lipids*, 15, 81–96.

530 Montealegre, C., Verardo, V., Gómez-Caravaca, A.M, García-Ruiz, C., Marina, M.L.,
 531 Caboni, M.F. (2012). Molecular characterization of phospholipids by high-performance
 532 liquid chromatography combined with an evaporative light scattering detector, high-
 533 performance liquid chromatography combined with mass spectrometry, and gas

534 chromatography combined with a flame ionization detector in different oat varieties
 535 *Journal of Agricultural and Food Chemistry*, 60, 10963–10969.

536 Nyström, L., Schär, A., Lampi, A.M. (2012). Steryl glycosides and acylated steryl glycosides
 537 in plant foods reflect unique sterol patterns. *European Journal of Lipid Science and*
 538 *Technology*, 114, 656–669.

539 Parcerisa, J., Codony, R., Boatella, J., Rafecas, M. (1999). Triacylglycerol and phospholipid
 540 composition of hazelnut (*Corylus avellana* L.) lipid fraction during fruit development.
 541 *Journal of Agricultural and Food Chemistry*, 47, 1410-1415.

542 Pelillo, M., Iafelice, G., Marconi, E., Caboni, M.F. (2003). Identification of plant sterols in
 543 hexaploid and tetraploid wheats using gas chromatography with mass spectrometry. *Rapid*
 544 *Communications in Mass Spectrometry*, 17, 2245-2252.

545 Pelvan, E., Alasalvar, C., Uzman, S. (2012). Effects of roasting on the antioxidant status and
 546 phenolic profiles of commercial turkish hazelnut varieties (*Corylus avellana* L.). *Journal*
 547 *of Agricultural and Food Chemistry*, 60, 1218–1223.

548 Piskornik, Z., Korfel, J. (1993). Jakościowo-ilościowa analiza kwasów tłuszczowych w oleju
 549 z orzechów laskowych pochodzących z Włoch i z Polski. Conference „O lepszą jakość
 550 produktów ogrodnich”, Kraków, 257-261.

551 Savage, G. P., McNeil, D. L., Dutta, P. C. (1997). Lipid composition and oxidative stability of
 552 oils in hazelnuts (*Corylus avellana* L.) grown in New Zealand. *Journal of the American*
 553 *Chemical Society*, 74, 755–759.

554 Schmitzer, V., Slatnar, A., Veberic, R., Stampar, F., Solar, A. (2011). Roasting affects
 555 phenolic composition and antioxidative activity of hazelnuts (*Corylus avellana* L.).
 556 *Journal of Food Science*, 76, 14–18.

- Shahidi, F., Alasalvar, C., Liyana-Pathirana, C. M. (2007). Antioxidant phytochemicals in hazelnut kernel (*Corylus avellana* L.) and hazelnut byproducts. *Journal of Agricultural and Food Chemistry*, 55, 1212–1220.
- Stévigny, C., Rolle, L., Valentini, N., Zeppa, G. (2007). Optimization of extraction of phenolic content from hazelnut shell using response surface methodology. *Journal of the Science of Food and Agriculture*, 87, 2817–2822.
- Sweeley, C.C., Bentley, R., Makita, M., Wells, W. (2002). Gas–liquid chromatography of trimethylsilyl derivatives of sugars and related substances. *Journal of the American Chemical Society*, 85, 2497–2507.
- Verardo, V., Bendini, A., Cerretani, L., Malaguti, D., Cozzolino, E., Caboni, M.F. (2009). Capillary gas chromatography analysis of lipid composition and evaluation of phenolic compounds by micellar electrokinetic chromatography in italian walnuts (*Juglans regia* L.): irrigation and fertilization influence. *Journal of Food Quality*, 32, 262–281.
- Verardo, V., Gómez-Caravaca, A.M., Gori, A., Losi, G., Caboni, M.F. (2013). Bioactive lipids in the butter production chain from Parmigiano Reggiano cheese area. *Journal of Science and Food Agriculture*, 93, 3625-3633.
- Yang, J. (2009). Brazil nuts and associated health benefits: A review. *LWT - Food Science and Technology*, 42, 1573–1580.

2 Data expressed as means \pm standard deviation ($n=3$). The different lower case letters (a-b) in the same row indicate
3 significantly different values ($p < 0.05$)

5 **Table 2.** Tocopherol content in the two hazelnut varieties cultivated in Poland (mg/kg d.w. nut)

Tocopherols		Variety	
		<i>Kataloński</i>	<i>Webba Cenny</i>
α-tocopherol	free	117.3 ± 2.4 a	147.1 ± 4.9 b
	bound	87.0 ± 2.9 a	39.4 ± 1.4 b
	total	204.3 ± 23.3 a	186.5 ± 16.2 a
γ-tocopherol	free	7.9 ± 0.4 a	10.2 ± 0.00 b
	bound	15.4 ± 0.5 a	3.1 ± 0.5 a
	total	23.3 ± 0.2 a	13.3 ± 0.5 b
β-tocopherol	free	3.5 ± 0.0 a	3.8 ± 0.2 a
	bound	5.1 ± 0.2 a	2.2 ± 0.5 b
	total	8.6 ± 0.2 a	6.0 ± 0.3 b
Total	free	128.7 ± 2.8 a	161.1 ± 5.1 b
	bound	107.5 ± 6.9 a	44.7 ± 5.4 b
	total	236.2 ± 23.7 a	205.8 ± 15.4 a

6 Data expressed as means ± standard deviation (*n*=3). The different lower case letters (a-b) in the same row indicate
7 significantly different values (*p* < 0.05)

8

9 **Table 3.** Sterols content in two hazelnut varieties cultivated in Poland (mg/kg d.w. nut)

Sterols		Variety	
		<i>Kataloński</i>	<i>Webba Cenny</i>
Cholesterol	free + esterified	2.4 ± 0.2 a	1.6 ± 0.6 a
	bound	6.7 ± 0.4 b	1.8 ± 0.2 a
	total	9.1 ± 0.5 b	3.4 ± 0.3 a
Campesterol	free + esterified	53.5 ± 1.0 b	50.0 ± 0.6 a
	bound	35.8 ± 2.1 b	17.3 ± 1.5 b
	total	89.3 ± 1.3 b	67.3 ± 1.0 a
Campestanol	free + esterified	11.0 ± 1.0 b	8.0 ± 0.7 a
	bound	2.7 ± 0.6 a	6.1 ± 1.5 b
	total	13.7 ± 0.8 a	14.1 ± 1.0 a
Stigmasterol	free + esterified	6.1 ± 0.5 a	6.0 ± 0.1 a
	bound	3.2 ± 0.3 b	1.2 ± 0.0 a
	total	9.3 ± 0.4 b	7.2 ± 0.5 a
Stigmastanol	free + esterified	1.2 ± 0.3 a	0.9 ± 0.1 a
	bound	2.3 ± 0.5 b	1.2 ± 0.1 a
	total	3.5 ± 0.3 b	2.1 ± 0.1 a
Chlerosterol	free + esterified	5.7 ± 0.3 b	4.3 ± 0.7 a
	bound	7.1 ± 1.2 b	2.8 ± 0.6 a
	total	12.8 ± 0.5 b	7.1 ± 0.3 a
Sitosterol	free + esterified	789.6 ± 31.2 a	832.7 ± 36.2 a
	bound	472.1 ± 23.4 b	242.1 ± 18.4 a
	total	1261.7 ± 48.5 b	1074.8 ± 29.2 a
Sitostanol	free + esterified	28.2 ± 1.7 a	32.7 ± 0.2 b
	bound	15.9 ± 0.8 a	29.1 ± 0.9 b
	total	44.1 ± 0.9 a	61.8 ± 1.2 b
Δ5-avenasterol	free + esterified	36.7 ± 2.6 a	39.5 ± 1.7 a
	bound	24.7 ± 2.5 b	11.3 ± 2.0 a
	total	61.4 ± 2.7 b	50.8 ± 2.5 a
Fucoesterol	free + esterified	3.4 ± 0.5 a	4.2 ± 0.6 a
	bound	1.1 ± 0.1	1.3 ± 0.3 a
	total	4.5 ± 0.4 a	5.5 ± 0.5 a
Δ7-avenasterol	free + esterified	6.1 ± 0.1 a	6.8 ± 0.4 a
	bound	6.7 ± 0.3 b	3.8 ± 0.5 a
	total	12.8 ± 0.6 b	9.0 ± 0.4 a
Total	free + esterified	943.9 ± 47.0 a	986.7 ± 31.6 a
	bound	578.3 ± 33.4 b	316.5 ± 30.8 a
	total	1522.2 ± 41.6 b	1303.2 ± 36.3 a

10 Data expressed as means ± standard deviation (n=3). The different lower case letters (a-b) in the same row indicate
11 significantly different values ($p < 0.05$)

Table 4. Phospholipid content (mg/g of oil) and percentage of each phospholipid (on total phospholipid content) in Polish hazelnut varieties.

Phospholipids	Variety			
	<i>Kataloński</i>		<i>Webba Cenny</i>	
	mg/g of oil	%	mg/g of oil	%
PC	7.9 ± 0.0 a	72.2	5.3 ± 0.1 b	67.5
PE	2.1 ± 0.0 a	19.6	1.8 ± 0.0 b	23.2
PI	0.9 ± 0.0 a	8.2	0.7 ± 0.0 b	9.3
Total	10.9 ± 0.1 a		7.9 ± 0.1 b	

PC - Phosphatidylcholine, PE - Phosphatidylethanolamine; PI - Phosphadidylinositol.

Data expressed as means ± standard deviation ($n=3$). The different lower case letters (a-b) in the same row indicate significantly different values ($p < 0.05$)

Table 5. Phospholipid fatty acid composition (% area of total methyl esters) of hazelnut varieties grown in Poland.

Fatty acids	Variety	
	<i>Kataloński</i>	<i>Webba Cenny</i>
C16:0	32.3 ± 0.1 a	34.7 ± 0.6 b
C16:1	11.0 ± 1.4 a	12.5 ± 0.8 a
C18:0	14.6 ± 1.9 a	14.9 ± 0.5 a
C18:1 ω9	32.4 ± 0.4 a	34.5 ± 0.8 b
C18:2 ω6	9.7 ± 0.2 a	3.4 ± 0.5 b
Σ SFA	46.8 ± 1.9 a	49.6 ± 1.0 a
Σ MUFA	43.4 ± 1.7 a	47.0 ± 1.5 a
Σ PUFA	9.7 ± 0.2 a	3.4 ± 0.5 b

Data expressed as means ± standard deviation (*n*=3). The different lower case letters (a-b) in the same row indicate significantly different values (*p* < 0.05)

26 **Table 6.** Free and bound phenolic compounds ($\mu\text{g/g}$ d.w. nut or shell) isolated in kernel and hard shell of analysed hazelnuts

Phenolic compounds	[M-H] ⁻ (m/z)	Fragments (m/z)	Free phenolics				Bound phenolics			
			Kernel		Hard shell		Kernel		Hard shell	
			<i>Kataloński</i>	<i>Webba Cenny</i>	<i>Kataloński</i>	<i>Webba Cenny</i>	<i>Kataloński</i>	<i>Webba Cenny</i>	<i>Kataloński</i>	<i>Webba Cenny</i>
Gallic acid ¹	169	125	11.1 ± 0.1 b	4.1 ± 0.0 c	2.7 ± 0.0 d	0.8 ± 0.0 e	nd	nd	16.9 ± 0.1 a	13.9 ± 0.0 b
Caffeic acid derivative ²	179	135	nd	nd	nd	nd	7.9 ± 0.0 b	12.5 ± 0.0 a	nd	nd
Protocatechuic acid ¹	153	109	1.1 ± 0.1 d	1.1 ± 0.1 d	0.5 ± 0.0 e	1.1 ± 0.0 d	2.5 ± 0.0 c	4.8 ± 0.0 b	33.0 ± 0.2 a	33.6 ± 0.1 a
Procyanidin dimer ³	577	451 289	10.5 ± 0.1 a	5.3 ± 0.0 b	nd	nd	nd	nd	nd	nd
Procyanidin trimer ³	865	577	0.8 ± 0.0 a	1.2 ± 0.1 a	nd	nd	nd	nd	nd	nd
Catechin ⁴	289	245	0.5 ± 0.0 d	1.0 ± 0.0 c	0.8 ± 0.0 c	0.3 ± 0.0 d	nd	nd	16.0 ± 0.1 b	20.5 ± 0.0 a
Vanillic acid ¹	167	123	nd	nd	0.4 ± 0.0 e	0.2 ± 0.0 e	2.9 ± 0.0 d	3.5 ± 0.0 c	34.8 ± 0.1 b	36.0 ± 0.0 a
Procyanidin trimer ³	865	577	2.2 ± 0.1 b	3.0 ± 0.5 a	nd	nd	nd	nd	nd	nd
Taxifolin ⁵	303	-	nd	nd	nd	nd	nd	nd	21.5 ± 0.1 a	22.1 ± 0.0 a
Procyanidin B2 ³	577	451 289	1.3 ± 0.0 b	4.5 ± 0.2 a	nd	nd	nd	nd	nd	nd
Galloylquinic acid ¹	343	191	nd	nd	nd	nd	nd	nd	17.4 ± 0.0 b	22.3 ± 0.0 a
Sinapic acid ²	223	-	nd	nd	< LOQ	< LOQ	4.0 ± 0.1 b	6.9 ± 0.0 a	nd	nd
p-coumaric acid ²	163	119	nd	nd	< LOQ	< LOQ	5.1 ± 0.0 a	2.1 ± 0.2 b	nd	nd
Quinic acid ¹	191	-	nd	nd	nd	nd	5.1 ± 0.0	< LOQ	nd	nd
Epicatechin ⁴	289	245	0.1 ± 0.0 a	0.1 ± 0.0 a	nd	nd	nd	nd	nd	nd

Myricetin-3-O-rhamnoside ⁶	463	317	0.8 ± 0.1 a	0.5 ± 0.1 b	0.2 ± 0.0 c	< LOQ	nd	nd	nd	nd
Quercetin-pentoside ₆	433	301	1.0 ± 0.0 b	2.9 ± 0.0 a	1.0 ± 0.0 b	0.2 ± 0.0 c	nd	nd	nd	nd
Quercetin-3-O-rhamnoside ⁶	447	301	2.8 ± 0.0 b	4.9 ± 0.1 a	nd	nd	nd	nd	nd	nd
Phloretin-2-O-glucoside ⁷	435	273	0.3 ± 0.0 b	0.9 ± 0.0 a	0.1 ± 0.0 c	0.1 ± 0.0 c	nd	nd	nd	nd
Total			31.7 ± 0.2 c	16.7 ± 0.4 d	5.6 ± 0.2 e	2.5 ± 0.1 f	19.7 ± 0.2 d	17.4 ± 0.2 d	139.6 ± 0.6 b	148.5 ± 0.2 a

27 Data expressed as means ± standard deviation (*n*=3). nd: not detected. The different lower case letters (a-b) in the same row indicate significantly different values (*p* < 0.05)

28 LOQ: limit of quantification

29 1= quantified with gallic acid at 280 nm; 2 = quantified with syringic acid at 280 nm; 3 = quantified with procyanidin B2 at 280 nm; 4 = quantified with

30 catechin at 280 nm; 5 = quantified with taxifolin at 280 nm; 6 = quantified with quercetin at 350 nm; 7 = quantified with phloretin at 280 nm

Supplementary Material

[Click here to download Supplementary Material: Fig1.pdf](#)

Supplementary Material

[Click here to download Supplementary Material: Fig2.pdf](#)