

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Impact of bisphenol A (BPA) on early embryo development in the marine mussel *Mytilus galloprovincialis*:
Effects on gene transcription

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Balbi, T., Franzellitti, S., Fabbri, R., Montagna, M., Fabbri, E., Canesi, L. (2016). Impact of bisphenol A (BPA) on early embryo development in the marine mussel *Mytilus galloprovincialis*: Effects on gene transcription. ENVIRONMENTAL POLLUTION, 218, 996-1004 [10.1016/j.envpol.2016.08.050].

Availability:

This version is available at: <https://hdl.handle.net/11585/566458> since: 2021-11-30

Published:

DOI: <http://doi.org/10.1016/j.envpol.2016.08.050>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Balbi, Teresa; FRANZELLITTI, SILVIA; Fabbri, Rita; Montagna, Michele; FABBRI, ELENA; Canesi, Laura: Impact of bisphenol A (BPA) on early embryo development in the marine mussel *Mytilus galloprovincialis*: Effects on gene transcription. ENVIRONMENTAL POLLUTION 218. 0269-7491

DOI: 10.1016/j.envpol.2016.08.050

The final published version is available online at:

<http://dx.doi.org/10.1016/j.envpol.2016.08.050>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Impact of bisphenol A (BPA) on early embryo development in the marine mussel *Mytilus galloprovincialis*: Effects on gene transcription[☆]

Teresa Balbi ^{a,1}, Silvia Franzellitti ^{b,1}, Rita Fabbri ^a, Michele Montagna ^a, Elena Fabbri ^{b,*}, Laura Canesi ^a

^a Department of Earth, Environment and Life Sciences (DISTAV), University of Genoa, Corso Europa 26, 16132, Genova, Italy

^b Department of Biological, Geological and Environmental Sciences, University of Bologna, Campus of Ravenna, via S. Alberto 163, 48123 Ravenna, Italy

ARTICLE INFO

Keywords:

Bisphenol
Marine mussel
Larvae
Gene transcription
Shell formation
Endocrine disruptors

ABSTRACT

Bisphenol A (BPA), a monomer used in plastic manufacturing, is weakly estrogenic and a potential endocrine disruptor in mammals. Although it degrades quickly, it is pseudo persistent in the environment because of continual inputs, with reported concentrations in aquatic environments between 0.0005 and 12 µg/L. BPA represents a potential concern for aquatic ecosystems, as shown by its reproductive and developmental effects in aquatic vertebrates.

In invertebrates, endocrine related effects of BPA were observed in different species and experimental conditions, with often conflicting results, indicating that the sensitivity to this compound can vary considerably among related taxa. In the marine mussel *Mytilus galloprovincialis* BPA was recently shown to affect early development at environmental concentrations. In this work, the possible effects of BPA on mussel embryos were investigated at the molecular level by evaluating transcription of 13 genes, selected on the basis of their biological functions in adult mussels. Gene expression was first evaluated in trocophorae and D veligers (24 and 48 h post fertilization) grown in physiological conditions, in comparison with unfertilized eggs. Basal expressions showed a general up regulation during development, with distinct transcript levels in trocophorae and D veligers. Exposure of fertilized eggs to BPA (10 µg/L) induced a general upregulation at 24 h pf, followed by down regulation at 48 h pf. *Mytilus* Estrogen Receptors, serotonin receptor and genes involved in biomineralization (Carbonic Anhydrase and Extracellular Matrix Protein) were the most affected by BPA exposure. At 48 h pf, changes in gene expression were associated with irregularities in shell formation, as shown by scanning electron microscopy (SEM), indicating that the formation of the first shelled embryo, a key step in mussel development, represents a sensitive target for BPA. Similar results were obtained with the natural estrogen 17β estradiol. The results demonstrate that BPA and E₂ can affect *Mytilus* early development through dysregulation of gene transcription.

1. Introduction

Aquatic environments are the ultimate reservoirs for many environmental anthropogenic chemicals, including those that can interfere with the functions of natural hormones. These compounds, defined as endocrine disrupting chemicals (EDCs) can adversely impact reproduction and development in humans and

wildlife (Zoeller et al., 2012). Fish and other aquatic organisms often have the greatest exposures to such chemicals during critical periods of development or even their entire life cycles.

Bisphenol A (BPA), generally considered one of the most common EDCs (Rubin, 2011; Vandenberg et al., 2013), is a monomer used in the manufacture of polycarbonate plastics and in a variety of plastic products, including food and beverage packaging, flameretardants, adhesives, building materials, electronic components, and paper coatings (EFSA FIP, 2015). Due to its widespread applications, BPA has become one of the highest production volume chemicals (5.5 million tons per year) in the last decades (Alonso Magdalena et al., 2015; Flint et al., 2012). Although it degrades quickly, it is pseudo persistent in the environment because of

[☆] This paper has been recommended for acceptance by Maria Cristina Fossi.

^{*} Corresponding author.

E-mail address: elena.fabbri@unibo.it (E. Fabbri).

¹ Equally contributed to the work.

continual inputs, leading to reported concentrations in aquatic environments between 0.0005 and 12 µg/L (Flint et al., 2012). Available data underline that at environmental concentrations BPA may represent a potential concern for aquatic ecosystems (Tisler et al., 2016). In aquatic vertebrates, BPA causes developmental and reproductive effects, as well disturbances in immune function and metabolism (Canesi and Fabbri, 2015). In invertebrates, endocrine related effects of BPA have been observed in different species and experimental conditions, with often conflicting results. Some invertebrates appear to be quite sensitive to BPA (freshwater and marine mollusks, insect larvae, marine copepods in particular), with effects documented at environmentally relevant concentrations (reviewed in Canesi and Fabbri, 2015; Flint et al., 2012).

Research on the impact of BPA in model aquatic organisms has major regulatory implications towards the ecosystem health. In this light, studies involving exposure during early life stages in invertebrates, which can be highly sensitive to environmental perturbations, would greatly help identifying potential adverse effects of BPA as an EDC, in analogy with what reported in vertebrates (Hamlin and Guillette, 2011; Santangeli et al., 2016; Tse et al., 2013).

We have recently shown that in the marine mussel *Mytilus galloprovincialis* exposure of fertilized eggs to BPA at environmentally relevant concentrations affects the formation of fully developed D larvae at 48 h post fertilization (pf) (Fabbri et al., 2014). Larval development in mussels chronologically unfolds following a mechanical remodeling of the form and behavior, which requires controlled mechanisms engaged in tissue distribution and organ system differentiation (Ackerman et al., 1994). In *Mytilus*, after external oocyte fertilization embryos undergo fast cell divisions that lead to their segmentation, and differentiate into motile forms (trochophore, 24 h pf), with successive development of various shelled forms (D veligers) from 48 h pf (Marin et al., 2008). Although a number of genetic regulators during late stages of larval development have been recently identified (Bassim et al., 2014), no information on the molecular mechanisms underlying the physiological processes involved in the early critical stages, such as the formation of the first shelled embryo, is available.

In this work, the effects of BPA on gene transcription in early embryo development of *M. galloprovincialis* were investigated. Fertilized eggs were exposed to BPA 1 µg/L (<EC₅₀, LOEC 10×) and 10 µg/L (>EC₅₀, LOEC 100×) as previously described (Fabbri et al., 2014), and the differential expression of 13 gene products related to known biological functions in adult mussels was assessed. These include neuroendocrine signaling, antioxidant defense, biomineralization, stress response, apoptosis, autophagy, and immune response. Transcriptional profiles were evaluated in trochophore and D veligers grown under physiological conditions or in the presence of BPA at environmentally realistic concentrations (Corrales et al., 2015). Morphology of control and BPA exposed embryos was evaluated by scanning electron microscopy (SEM). The effects of BPA on embryo development, gene expression and morphology were compared with those of the natural estrogen 17β estradiol (E₂).

2. Methods

2.1. Animal holding and larval rearing

Sexually mature mussels (*M. galloprovincialis* Lam.), purchased from an aquaculture farm in the Ligurian Sea (La Spezia, Italy) between November 2014 and March 2014, were transferred to the laboratory and acclimatized in static tanks containing aerated artificial sea water (ASTM, 2004), pH 7.9–8.1, 36 ppt salinity (1 L/animal), at 16 ± 1 °C.

Gamete collection and fertilization were performed following

the procedure described by Fabbri et al. (2014). Details are reported in Fig. S1 (Supporting Information Appendix 1) with the description of the 96 microwell embryotoxicity assay performed to assess the effects of E₂. Unfertilized eggs (about 24,000 eggs/mL) obtained from at least 6 female individuals were collected by centrifugation at 400×g for 10 min at 4 °C, and the resulting pellet was frozen in liquid nitrogen. Eggs were fertilized with an egg:sperm ratio 1:10 in polystyrene 6 well plates and a final 6 mL volume. After 30 min, fertilization success (n. fertilized eggs/n. total eggs × 100) was verified by microscopical observation (>85%). At 30 min pf, BPA or E₂ were added to fertilize eggs in each well from 1 mg/L concentrated stock solutions prepared in ethanol, and suitably diluted to reach the final nominal concentrations of 1 and 10 µg/L for BPA, and 10 µg/L for E₂. Concentrations were chosen on the basis of the dose response curve obtained in the *Mytilus* 48 h embryotoxicity test for BPA (Fabbri et al., 2014) and E₂ (present study; Fig. S1, Supporting Information Appendix 1). Control wells (negative controls) contained only ASW. Four replicates for each experimental condition were made. In all cases, ethanol final concentration did not significantly affect the biological endpoints analysed (data not shown). Concentrations of BPA and E₂ in test solutions were checked by LC/MS (Table S1, Supporting Information Appendix 2).

2.2. RNA extraction and cDNA preparation

At 24 and 48 h pf larvae were collected by a nylon mesh (40 µm pore filter) and washed with ASW. Three wells for each condition were pooled in order to obtain approximately 7000 embryos/replicate. The larval suspension was centrifuged at 800×g for 10 min at 4 °C. Larval pellets and unfertilized eggs were lysed in 1 mL of the TRI Reagent (Sigma Aldrich, Milan, Italy) and total RNA was further extracted using the DirectZol kit (Zymo Research, Freiburg, Germany) following manufacturer's instructions. RNA concentration and quality were verified using the Qubit RNA assay (Thermo Fisher, Milan, Italy) and electrophoresis using a 1.2% agarose gel under denaturing conditions. First strand cDNA for each sample was synthesized from 1 µg total RNA according to Balbi et al. (2014).

2.3. Real time quantitative PCR (qPCR) analysis

Primers pairs employed for qPCR analysis were as reported in previous studies or were designed with Primer Express (Thermo Fisher, Milan, Italy) using nucleotide sequences retrieved from the GeneBank database (<https://www.ncbi.nlm.nih.gov/genbank/>) for *Mytilus galloprovincialis* (Table S2, Supporting Information Appendix 2). qPCR reactions were performed in duplicate in a final volume of 10 µL containing 5 µL iTaq universal master mix with ROX

(BioRad Laboratories, Milan, Italy), 2 µL diluted cDNA, and 0.2 µM specific primers (Table S2, Supporting Information Appendix 2). A control lacking cDNA template (no template) and a minus reverse transcriptase (no RT) control were included in the qPCR analysis to determine the specificity of target cDNA amplification. Amplifications were performed in a StepOne real time PCR system apparatus (Thermo Fisher, Milan, Italy) using a standard "fast mode" thermal protocol. For each target mRNA, melting curves and agarose gel electrophoresis were utilized to verify the specificity of the amplified products and the absence of artifacts. The amplification efficiency of each primer pair was calculated using a dilution series of cDNA (Table S2, Supporting Information Appendix 2). A preliminary stability analysis of 6 established candidate reference transcripts (Cubero Leon et al., 2012; Franzellitti and Fabbri, 2013) was performed to achieve a robust normalization of qPCR data in mussel larval stages. Detailed methods and results are reported in Supporting Information Appendix 3. *HEL* and *EF α1* were selected

as the best performing combination of reference gene products for data normalization. Calculations of relative expression of target mRNAs was performed by a comparative C_T method (Schmittgen and Livak, 2008) using the StepOne software tool (Thermo Fisher, Milan, Italy). Data were reported as relative expression (fold change or \log_2 transformed fold changes according to the data ranges) with respect to unfertilized eggs (basal gene expression across larval development) or control samples within each life stage (BPA or E_2 treatments).

2.4. Scanning electron microscopy (SEM)

At different times post fertilization, depending on the experiment, control and treated embryos grown in 96 microwells were fixed in 3% glutaraldehyde in ASW. After fixation, samples from 6 microwells were pooled, placed onto Whatman 22 μ m filters, dehydrated in an ascending series of ethanol washes (50%–80%–90%–100%) and air dried. Then samples were sputter coated with gold, and observed at 20 kV with a Vega3 Tescan scanning electron microscope.

2.5. Statistics

Statistical comparisons between unfertilized eggs vs trocophora/D veligers, or controls vs BPA/ E_2 treated larvae (within each developmental stage) were performed using the non parametric one way ANOVA (Kruskal Wallis test) followed by the Mann Whitney U test ($P < 0.05$), after deviations from parametric ANOVA assumptions being verified (Normality: Shapiro Wilk's test; equal variance: F test). Data analyses were performed using GraphPad Prism 5 software (GraphPad Inc.).

Additional statistical approaches were utilized for analysis of qPCR data to evaluate:

- i) Basal gene expression across larval development. Ordination analysis (carried out by Principal Coordinates, PCO) and clustering of basal transcript expressions across the larval stages were performed using the PRIMER v6 software (PRIMER E Ltd, UK), which employed \log_2 transformed fold change variations of the target transcripts to calculate similarity matrices based on the Euclidean distance (999 permutations). The similarity matrices were also used to perform a canonical analysis of principal coordinates ordination (CAP) to attribute the specific transcripts responsible for condition differences (i.e. differences between larval stages).
- ii) Effects of BPA or E_2 treatments on target transcript expressions in different larval stages. Relative expression data from BPA or E_2 treatments were submitted to permutation multivariate analysis of variance (PERMANOVA) using the PERMANOVA+ add on in PRIMER v6 (Anderson et al., 2008). Log transformed copy number variations of the target transcripts were used to calculate similarity matrices based on the Euclidean distance (999 permutations). Factors considered were "developmental stage" and "treatment" (BPA or E_2). Pseudo F values in the PERMANOVA main tests were evaluated in terms of significance (Anderson et al., 2008). When the main test revealed statistical differences ($P < 0.05$), PERMANOVA pairwise comparisons were carried out (using the Euclidean distance matrix and 999 permutations). Distance based redundancy linear modeling (DISTLM) followed by a redundancy analysis (dbRDA) in PRIMER with PERMANOVA+ extension was also performed on BPA or E_2 treatments to examine the relationship between the multivariate dataset (i.e. the suite of transcripts assayed and their expression levels) and the predictor variables (life stage and BPA/ E_2 treatment). Numerical metric for life stage progression

was indicated by the post fertilization time (0: unfertilized egg; 24 h: trocophora; 48 h: D veliger). Treatment was indicated by the nominal concentrations of exposure (BPA: 0, 1, and 10 μ g/L; E_2 : 0 and 10 μ g/L). DISTLM used the BEST selection procedure and adjusted R^2 selection criteria.

3. Results

3.1. Transcriptional profiles during early larval development

Transcription of selected genes related to neuroendocrine signaling (estrogen receptors *MeER1*, *MeER2*, serotonin receptor, *5 HTR*), antioxidant defense (superoxide dismutase, *SOD*, catalase, *CAT*), biomineralization (extrapallial protein, *EP*, carbonic anhydrase, *CA*), detoxification response (glutathione *S* transferase, *GST* π , ABC transporter p glycoprotein, *ABCB*), stress response (70 kDa heat shock proteins, *HSP70*), apoptosis (*p53*), autophagy (serine/threonine protein kinase mTOR, *mTor*), immune response (Toll like receptor, *TLR i*), was evaluated at both 24 and 48 h pf in embryos grown under normal physiological conditions. Data on relative expression of each transcript across larval development (fold changes with respect to unfertilized eggs) are reported in Fig. 1A. To help comparing overall transcript levels among datasets, fold change variations (\log_2 transformed) of target transcripts were subjected to a cluster analysis and to an ordination analysis using a Principal Coordinate Ordination (PCO) extraction procedure (Fig. 1B). Basal transcript expressions showed substantial time course variations across developmental stages with general and significant up regulation (Fig. 1A). The highest mRNA levels were reported for *MeERs*, *5 HTR*, *CA* and *p53* (Fig. 1A). On the contrary, down regulation was observed for *GST* π in both larval stages (Fig. 1A). The PCO analysis (Fig. 1B) detected two principal coordinates (PCO1 and PCO2) explaining about 97% of total variation (89.8% and 5.8%, respectively). Samples from unfertilized eggs formed one group with evident separation from those belonging to trocophora and D veliger stages, which also formed distinct clusters. The canonical analysis of principal coordinates ordination (CAP analysis) attributed much of the differences between these latter two stages to *5 HTR*, *EP*, *CA*, *MeER1*, *p53*, *CAT*, *SOD*, *ABCB* profiles (Fig. 1C).

3.2. Effects of BPA on gene transcription

Fertilized eggs were exposed to BPA 1 μ g/L ($<EC_{50}$, LOEC 10 \times), and 10 μ g/L ($>EC_{50}$, LOEC 100 \times), as previously described (Fabbri et al., 2014), and gene transcription was evaluated at different times pf (24 h and 48 h) in comparison with untreated samples (Fig. 2). Distinct changes in gene transcription at different stages and at the tested BPA concentrations were observed. In general, a small though significant up regulation of most transcripts (from +35% to +55%, depending on the transcript and the developmental stage), was observed in larvae grown under 1 μ g/L BPA treatment (Fig. 2A). Significant down regulation was observed for *SOD* and *EP* in trocophorae, and for *SOD* and *CA* in D veligers (from 30% to 60%) (Fig. 2A). More consistent effects were observed in larvae grown under 10 μ g/L treatment. As shown in Fig. 2B, all transcripts were significantly up regulated in trocophorae, with the exception of *HSP70*, which showed down regulation (Fig. 2B). A maximum 3.5 fold increase was observed for the *EP* transcript. In contrast, in D veligers 10 out of 13 transcripts were significantly down regulated, with mRNA levels ranging from 30% to 54% with respect to controls (Fig. 2B). Only *SOD* and *CA* were significantly up regulated, with *CA* showing up to 4.2 fold increase with respect to controls (Fig. 2B).

Distance based linear model (DISTLM) analysis revealed that

Fig. 1. Transcriptional profiles of selected gene products during early larval development in mussels. (A) Gene transcription was evaluated by qPCR in embryos of *M. galloprovincialis* grown under physiological conditions at 24 h (trochophora) and 48 h (D-veliger) post fertilization. Data, reported as \log_2 -transformed relative expressions with respect to unfertilized eggs, represent the mean \pm SD (N = 12). Expression of all transcripts was significantly changed at both 24 and 48 h pf with respect to unfertilized eggs (1 way non parametric ANOVA followed by the Mann Whitney U test; $P < 0.05$). Further statistical differences were evaluated between trochophorae and D veligers ($^*P < 0.05$, $^{**}P < 0.01$, $^{***}P < 0.001$ according to the Mann Whitney U test). (B) Principal coordinates ordination (PCO) bi plot of mRNA expression levels with super imposed cluster analysis by condition (Euclidean distance matrix; 999 permutations). (C) Canonical analysis of principal coordinates ordination (CAP analysis) illustrating transcripts whose expression patterns are likely to discriminate the differences between life stages. Transcripts to be displayed in the CAP bi plot were filtered according to their Spearman correlation's coefficients ($r > 0.6$ with at least CAP1 or CAP2).

expression profiles were strongly dependent on embryo development (time from pf) that explained about 91% total variation (Fig. 2C). Nevertheless, BPA treatment accounted for about 5% of total variation, which mostly explained the observed changes in transcript expressions at 48 h pf and at 10 µg/L BPA treatment (Fig. 2C). Results from PERMANOVA and permutation *t* test analyses demonstrated that the effects of BPA were statistically significant in both larval stages and at both concentrations tested ($P < 0.05$; Table S3, Supporting Information Appendix 2). PERMANOVA analysis also showed a significant interaction between the two factors (developmental stage and BPA treatment; $P < 0.05$; Table S3, Supporting Information Appendix 2).

3.3. SEM analysis

Larval development was observed by Scanning electron microscopy (SEM) first in larvae grown under physiological conditions, and representative images are reported in Fig. 3(a–f). Fig. 3a and b show a trocophora at 24 h pf, and the detail of the ciliated epithelium, respectively. During the trocophora stage, the onset of shell mineralization occurs (generally at 20 h pf). The process is initiated by a specialized group of ectodermal cells, the shell field,

that are responsible for production of the first outer organic shell layer, the periostracum (Weiss et al., 2002). The saddle shaped shell field is well visible in Fig. 3c. From about 32 h pf, first shelled embryos could be observed (Fig. 3d); at this stage, most shells were extremely fragile, and damaged by the sample preparation procedure for SEM observations. Micrometric spherules of amorphous calcium carbonate (ACC) could be observed both at the surface and margins of the valvae (Fig. 3e) typical of initial shell formation (Cartwright et al., 2012; Weiss et al., 2002). Finally, at 48 h pf the D veliger stage is reached (Fig. 3f), with shells showing straight hinge, symmetric valvae, and uniform surface, indicating that formation of the calcified shell occurred simultaneously in the whole shell field. In samples exposed to BPA (10 µg/L), no appreciable morphological changes were observed at 24 h pf (Trochophora) (not shown). On the other hand, clear changes were observed in the morphology of D larvae at 48 h pf (Fig. 3g–i). Common shell alterations were rough surface, thinner valvae, convex hinge, asymmetry and/or gap between valvae; moreover, many samples showed the externalized velum (ciliated swimming and feeding organ), a feature typical of the pre veliger stage (Fig. 3i and inset to Fig. 3i).




Fig. 2. Effects of BPA on gene transcription in *Mytilus* embryos at 24 h and 48 h post fertilization times. Expression levels of target transcripts were evaluated by qPCR following 1 µg/L (A) and 10 µg/L (B) BPA treatment. Data are reported as mean \pm SD of the relative expression with respect to untreated samples within each life stage (N = 4). *P < 0.05, **P < 0.01 according to the Mann-Whitney U test. (C) DISTLM modeling with distance-based redundancy analysis (dbRDA) to explore the amount of the variation in gene transcription to be attributed to BPA treatment (Euclidean Distance resemblance matrix, 999 permutations).

3.4. Effects of 17 β estradiol on mussel embryo development and gene transcription

The effects of BPA were compared with those of the natural estrogen 17 β estradiol (E_2). Fig. S1a (Supporting Information Appendix 1) shows results from the 48 h embryotoxicity assay performed utilizing E_2 in the same concentration range as that previously utilized for BPA (Fabbri et al., 2014). E_2 induced a

dose dependent decrease in normal larval development that was significant from 0.1 µg/L, with a calculated EC_{50} value of 6.904 µg/L (95% confidence interval of 3.284–14.511 µg/L). Optical microscopy observations indicated that, as previously observed with BPA (Fabbri et al., 2014), E_2 mainly induced a delay in development, with a prevalent increase in the percentage of trocophorae/not fully developed D veligers, and a variable percentage of malformed embryos at different concentrations. At the highest concentration




Fig. 3. Scanning electron microscopy (SEM) of *Mytilus* embryos. a–f: representative images of different stages of larvae grown under physiological conditions: a) trocophora at 24 h pf, and b) detail of the ciliated epithelium; c) trocophora with super-imposed margins of the saddle-shaped shell field; d) early D-veliger at 36 h pf, showing the presence of amorphous calcium carbonate (ACC) spherules (e); f) normal D-veliger at 48 h pf, with straight hinge, symmetric valves and uniform surface. g–i: representative images of the effects of BPA (10 µg/L) of shell morphology in D-veligers at 48 h pf, showing asymmetric valves and rough surface (g), convex hinge and thinner valves (h), externalized velum (i).

tested (1000 µg/L) more than 90% embryos were at the pre veliger stage or showed gross shell malformations (not shown). At 10 µg/L E_2 , SEM observations showed that most common shell alterations induced by E_2 were hinge malformations, asymmetric and thinned valves and rough surface (Fig. S1b–d, Supporting Information Appendix 1).

The effects of E_2 (10 µg/L) on gene transcription are illustrated in Fig. 4A. The results show that most transcripts were significantly down regulated both at 24 h and 48 h pf, to a different extent depending on the type of gene and developmental stage. In particular, at 24 h pf E_2 induced a general decrease in gene transcription, except for 5 *HTR*, *ABCB*, *EP* and *CA*. Dramatic down regulation ($\leq 70\%$ with respect to controls) was observed for *MeER1*, *CAT*, *p53*, and *mTor*, and almost complete inhibition resulted for *TLR i* transcription. At 48 h pf, a significant down regulation was observed for all transcripts except for *HSP70* and *mTor*, whose mRNA levels were unchanged, and for *ABCB*, that was up regulated.

DISTLM analysis revealed that E_2 treatment accounted for 75% of total variation, and affected both trocophora and D veliger stages (Fig. 4B). Results from PERMANOVA and permutation *t* test analyses demonstrated that the effects of E_2 were significant in both larvae stages ($P < 0.05$; Table S4, Supporting Information Appendix 2). PERMANOVA analysis also showed a significant interaction between the two factors (developmental stage and E_2 treatment; $P < 0.05$; Table S4, Supporting Information Appendix 2).

4. Discussion

We have previously shown that BPA significantly affected 48 h embryo development in *M. galloprovincialis*, with a LOEC and EC_{50} of 0.1 µg/L and 3.68 µg/L, respectively (Fabbri et al., 2014). Since these values fall within the concentration range of BPA found in coastal environments (Flint et al., 2012), these data underlined the sensitivity of mussel embryos towards exposure to environmental




Fig. 4. Effects of E_2 ($10 \mu\text{g/L}$) on gene transcription in *Mytilus* embryos at 24 h and 48 h post fertilization times. (A) Expression levels of target transcripts evaluated by qPCR. Data are reported as the mean \pm SD of the relative expression with respect to untreated samples within each life stage ($N = 4$). * $P < 0.05$, ** $P < 0.01$ according to the Mann-Whitney U test. (B) DISTLM modeling with distance-based redundancy analysis (dbRDA) to explore the amount of the variation in gene transcription is attributable to E_2 treatment (Euclidean Distance resemblance matrix, 999 permutations).

concentrations of BPA, and supported the hypothesis that this compound could act as an EDC in marine bivalves. The results here obtained demonstrated that BPA affects gene transcriptions in early mussel embryos, with effects depending on the BPA concentration, the target transcript and the developmental stage. These data represent the first indication of molecular mechanisms for BPA impacts on developmental stages in a marine bivalve.

Basal transcription of 13 genes involved in different biological processes was first evaluated in larvae grown under physiological conditions at both 24 and 48 h pf, and relative expression was quantified with respect to unfertilized eggs. Genes were selected on the basis of their recognized physiological functions in *Mytilus* spp. and/or of their sensitivity to different environmental chemicals, including BPA (i.e. neuroendocrine signaling, antioxidant defenses, biomineralization, detoxification processes, stress response, apoptosis, autophagy, immunity). These pathways are reported to provide potent defenses in embryos against common stressors, including changes in temperature, hypoxia, and pollutants, to permit optimal development despite relevant environmental challenges (Hamdoun and Epel, 2007). To the best of our knowledge, the results represent the first data on basal mRNA expressions in early development of *M. galloprovincialis*. All transcripts were significantly up regulated in both trocophora and D veliger stages with respect to unfertilized eggs, except for GST π , which resulted significantly down regulated in both trocophore and D veligers compared to eggs, thus suggesting that early embryos rely on maternal GST, if any, for Phase II biotransformation pathways. PCO

and cluster analyses disclosed a further distinct clustering between trocophore and D veliger samples, consistent with the progressive differentiation of different processes. Main contribution to the observed data ordination was provided by the increased expressions of 5 HTR, EP, CA, MeER1, p53, CAT, SOD, ABCB in D veligers compared to trocophorae, as suggested by CAP analysis. If on one hand the results obtained for 5 HTR confirm the early and progressive development of serotonergic transmission in *Mytilus* spp. reported by previous immuno histochemical investigations (Voronezhskaya et al., 2008), on the other hand increased expressions of SOD, and to a lesser extent, of other transcripts with cytoprotective functions (i.e. ABCB, CAT, p53), indicate a reinforcement of defenses in the first D shelled embryo stage.

Although a number of genes involved in biomineralization are being described in different bivalve species (Bassim et al., 2015; Vendrami et al., 2016), those involved in the initial formation of a first calcified shell (prodissoconch I), a key step in early development, have not been identified yet. Carbonic anhydrase (CA) is known to regulate matrix mineralization by generating an acidic environment (Clark et al., 2010). *Mytilus* EP (Extrapallial Protein), an acidic calcium binding protein, regulates the production of different polymorphs of calcium carbonate (Yin et al., 2009). Both EP and CA showed the highest expression in D veligers, supporting the results from SEM analyses, showing that the formation of a complete normal shell occurred only at 48 h pf, and confirming the key physiological role of both transcripts in initial biomineralization (Medakovic, 2000; Yin et al., 2009, 2005).

Exposure of fertilized eggs to 1 µg/L BPA resulted in small but significant up regulation of most transcripts. However, downregulation was observed for *SOD*, and for *EP* and *CA* in trocophore and D veligers, respectively. More evident changes were observed at 10 µg/L. In these conditions, distinct trends in gene transcription were observed at different post fertilization periods, with a general up regulation in the trocophore and down-regulation in D veligers for all genes tested, except for *CA* and *SOD*, whose mRNA levels were further increased at 48 h pf and at 10 µg/L BPA treatment. Interestingly, transcription of both *EP* and *CA* was strongly affected by BPA exposure at both developmental stages.

The effects of BPA on mussel embryos were compared with those of the natural estrogen 17β estradiol (*E*₂). *E*₂ significantly affected larval development in the 48 h embryotoxicity tests (*EC*₅₀

6.904 µg/L), with effects similar to those of BPA (Fabbri et al., 2014). Therefore, gene transcription was evaluated after exposure to 10 µg/L *E*₂. At 24 h pf, *E*₂ induced a general decrease in gene transcription, in particular of genes involved in apoptosis, autophagy and immunity, showing a distinct effect with respect to BPA. At 48 h pf, *E*₂, like BPA, induced down regulation of most genes, including those involved in biomineralization. An exception to this trend was represented by *ABCB*, whose transcription was unaffected in trocophore, and significantly increased in D veligers. At the 1.0 µg/L concentration, the main morphological effects induced by BPA (Fabbri et al., 2014) and *E*₂ (this work) were represented by the presence of immature, not fully developed D shell larvae at 48 h pf. Similar results were obtained when BPA or *E*₂ were added at 24 h pf, instead of at 30 min post oocyte fertilization (not shown), indicating that the transition from the trocophore stage to the first D shelled larva represents a critical developmental step affected by both compounds. Accordingly, a common effect of BPA and *E*₂ observed at 48 h pf was the general down regulation of gene expression and alterations in shell formation, as indicated by SEM analysis.

Overall, the results demonstrate that both the estrogenic chemical BPA and the natural estrogen *E*₂ affect *Mytilus* early development through dysregulation of gene transcription. In the marine gastropod *Haliotis diversicolor*, BPA affected different stages of larval development, with an *EC*₅₀ of 1.02 µg/L at 96 h (completion of metamorphosis) (Liu et al., 2011). Furthermore, proteomic analysis indicated that both chemicals interfered with different physiological pathways, including energy and substance metabolism, cell signaling, formation of cytoskeleton and cilium, immune and stress responses at the same time, leading to the failure of metamorphosis (Liu et al., 2011).

The results of the present work underline that in *Mytilus* *MeERs* and 5 *HTR* represent significant targets for both BPA and *E*₂ during early embryo development. Similar results on *MeER* expression were observed in the hepatopancreas of adult mussels exposed to BPA (Canesi et al., 2011, 2007, 2006). Taken together the results support a common neuroendocrine related effect of these compounds in mussels. However, from an accurate analysis on the literature on the effects of steroids and EDCs Scott (2013) concluded that there is no indisputable bioassay evidence that vertebrate sex steroids (or their agonist/antagonists) have endocrine or reproductive roles in mollusks. Estrogen receptors (ERs) generally show constitutive transcriptional expressions in mollusks, and are not activated by either natural estrogens or estrogenic compounds, this rising a debate on their physiological function (Bridgham et al., 2014). In this respect, molluscan ERs seem to resemble mammalian orphan estrogen related receptors (ERRs), that play multiple and overlapping roles in cell growth, differentiation and metabolism (Huss et al., 2015). However, cloning and phylogenetic analysis of estrogen like receptors in *Mytilus edulis* and *M. galloprovincialis*

revealed the *Me/MgER2* is an ER, whereas the *Me/MgER1* is an ERR (Nagasawa et al., 2015). Our results show that in physiological conditions both transcripts encoding the *MeER1* and *MeER2* isoforms were among the most up regulated across different developmental stages, while these receptors were down regulated by exposure to both BPA and *E*₂ in D veligers. These data indicate that *Mytilus* ERs may play a role in early development and represent significant targets for estrogenic chemicals. Similar results were obtained for the 5 *HTR* transcript, encoding a putative type 1 serotonin receptor (Cubero Leon et al., 2010; Franzellitti and Fabbri, 2013), indicating that BPA and *E*₂ have a significant impact on early development of the serotonergic signaling in mussels (Kreiling et al., 2001; Voronezhskaya et al., 2008).

In addition, the results underline how estrogenic compounds significantly affected transcription of *EP* and *CA* involved in mussel biomineralization, with down regulation of both transcripts observed at 48 h pf. These data were supported by SEM observations, which clearly showed similar irregularities in shell formation in both BPA and *E*₂ treated embryos at 48 h pf. These effects are in line with the effects of estrogenic chemicals on bone metabolism observed in vertebrate systems (Agas et al., 2013). Overall, these represent the first data on the effects of EDCs on early biomineralization processes in marine invertebrates. The effects of BPA on the shell structure are under investigation. Calcifying larvae of marine species are particularly vulnerable to abiotic stressors (Przeslawski et al., 2015). Interestingly, the shell malformations

induced by BPA and *E*₂ were similar to those observed in *M. galloprovincialis* embryos by inhibiting of the synthesis of chitin, a key biological component that forms the framework for other macromolecules that guide initial shell deposition (Schonitzer and Weiss, 2007; Weiss and Schonitzer, 2006), as well as by exposure to high pCO₂ or low pH (Kadar et al., 2010; Kurihara et al., 2008). In this light, chitin synthesis may represent an important target for the effects of both chemical pollutants and altered environmental parameters in the developing embryo. Studies on the effects of individual chemicals that represent potential EDCs on early life stages of sensitive marine species may contribute to the development of future multiple stressor studies in a global change scenario.

5. Conclusions

The results of the present study show that environmental levels of BPA and *E*₂ may adversely affect physiological processes involved in early embryo development of marine mussels through dysregulation of transcription for genes involved in different key regulatory pathways. In particular, the results address the transition from the trocophore stage to the first D shelled larva as a very sensitive target for estrogenic compounds. The results indicate that, although physiological processes in early embryos are well buffered for environmental challenges (Hamdoun and Epel, 2007), anthropogenic stressors can overwhelm this intrinsic robustness and compromise developmental processes.

Acknowledgement

The research has been funded by the Italian Ministry of University and Research (RFO2014) to E.F. and to L.C.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.envpol.2016.08.050>.

References

- Ackerman, J.D., Sim, B., Nichols, S.J., Claudi, R., 1994. A review of the early life history of zebra mussels (*Dreissena polymorpha*): comparisons with marine bivalves. *Can. J. Zool.* 72, 1169–1179. <http://dx.doi.org/10.1139/z94-157>.
- Agas, D., Sabbieti, M.G., Marchetti, L., 2013. Endocrine disruptors and bone metabolism. *Arch. Toxicol.* 87, 735–751. <http://dx.doi.org/10.1007/s00204-012-0988-y>.
- Alonso-Magdalena, P., Quesada, L., Nadal, A., 2015. Prenatal exposure to BPA and offspring outcomes: the diabetogenic behavior of BPA. *Dose-Response* 13, 1–8. <http://dx.doi.org/10.1177/1559325815590395>.
- Anderson, M., Gorley, R., Clarke, K., 2008. PERMANOVA+ for Primer: Guide to Software and Statistical Methods.
- ASTM, 2004. Standard Guide for Conducting Static Acute Toxicity Tests Starting with Embryos of Four Species of Saltwater Bivalve Molluscs. <http://dx.doi.org/10.1520/E0724-98>.
- Balbi, T., Smerilli, A., Fabbri, R., Ciacci, C., Montagna, M., Grasselli, E., Brunelli, A., Pojana, G., Marcomini, A., Gallo, G., Canesi, L., 2014. Co-exposure to n- TiO_2 and Cd^{2+} results in interactive effects on biomarker responses but not in increased toxicity in the marine bivalve *Mytilus galloprovincialis*. *Sci. Total Environ.* 493, 355–364. <http://dx.doi.org/10.1016/j.scitotenv.2014.05.146>.
- Bassim, S., Chapman, R.W., Tanguy, A., Moraga, D., Tremblay, R., 2015. Predicting growth and mortality of bivalve larvae using gene expression and supervised machine learning. *Comp. Biochem. Physiol. Part D. Genom. Proteom.* 16, 59–72. <http://dx.doi.org/10.1016/j.cbd.2015.07.004>.
- Bassim, S., Tanguy, A., Genard, B., Moraga, D., Tremblay, R., 2014. Identification of *Mytilus edulis* genetic regulators during early development. *Gene* 551, 65–78. <http://dx.doi.org/10.1016/j.gene.2014.08.042>.
- Bridgham, J.T., Keay, J., Ortlund, E.A., Thornton, J.W., 2014. Vestigialization of an allosteric switch: genetic and structural mechanisms for the evolution of constitutive activity in a steroid hormone receptor. *PLoS Genet.* 10 <http://dx.doi.org/10.1371/journal.pgen.1004058>.
- Canesi, L., Borghi, C., Ciacci, C., Fabbri, R., Vergani, L., Gallo, G., 2007. Bisphenol-A alters gene expression and functional parameters in molluscan hepatopancreas. *Mol. Cell. Endocrinol.* 276, 36–44. <http://dx.doi.org/10.1016/j.mce.2007.06.002>.
- Canesi, L., Ciacci, C., Iorullo, L.C., Bett, M., Guarnieri, T., Tavorali, S., Gallo, G., 2006. Immunomodulation by 17 β -estradiol in bivalve hemocytes. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 291, R664–R673. <http://dx.doi.org/10.1152/ajpregu.00139.2006>.
- Canesi, L., Fabbri, E., 2015. Environmental effects of BPA: focus on aquatic species. *Dose. Response* 13. <http://dx.doi.org/10.1177/1559325815598304>.
- Canesi, L., Negri, A., Barmo, C., Banni, M., Gallo, G., Viarengo, A., Dondero, F., 2011. The organophosphate Chlorpyrifos interferes with the responses to 17 β -estradiol in the digestive gland of the marine mussel *Mytilus galloprovincialis*. *PLoS One* 6. <http://dx.doi.org/10.1371/journal.pone.0019803>.
- Cartwright, J.H.E., Checa, A.G., Gale, J.D., Gebauer, D., Sainz-Díaz, C.L., 2012. Calcium carbonate polymorphism and its role in biomineralization: how many amorphous calcium carbonates are there? *Angew. Chem. Int. Ed. Engl.* 51, 11960–11970. <http://dx.doi.org/10.1002/anie.201203125>.
- Clark, M.S., Thorne, M.A., Vieira, F.A., Cardoso, J.C., Power, D.M., Peck, L.S., 2010. Insights into shell deposition in the Antarctic bivalve *Laternula elliptica*: gene discovery in the mantle transcriptome using 454 pyrosequencing. *BMC Genom.* 11, 362.
- Corrales, J., Kristofco, L.A., Steele, W.B., Yates, B.S., Breed, C.S., Williams, E.S., Brooks, B.W., 2015. Global assessment of bisphenol A in the environment: review and analysis of its occurrence and bioaccumulation. *Dose. Response* 13. <http://dx.doi.org/10.1177/1559325815598308>.
- Cubero-Leon, E., Ciocan, C.M., Hill, E.M., Osada, M., Kishida, M., Itoh, N., Kondo, R., Minier, C., Rotchell, J.M., 2010. Estrogens disrupt serotonin receptor and cyclooxygenase mRNA expression in the gonads of mussels (*Mytilus edulis*). *Aquat. Toxicol.* 98, 178–187. <http://dx.doi.org/10.1016/j.aquatox.2010.02.007>.
- Cubero-Leon, E., Ciocan, C.M., Minier, C., Rotchell, J.M., 2012. Reference gene selection for qPCR in mussel, *Mytilus edulis*, during gametogenesis and exogenous estrogen exposure. *Environ. Sci. Pollut. Res.* 19, 2728–2733. <http://dx.doi.org/10.1007/s11356-012-0772-9>.
- EFSA FIP, 2015. Scientific opinion on the risks to public health related to the presence of bisphenol A (BPA) in foodstuffs. *EFSA J.* 13, 3978. <http://dx.doi.org/10.2903/j.efsa.2015.3978>.
- Fabbri, R., Montagna, M., Balbi, T., Raffo, E., Palumbo, F., Canesi, L., 2014. Adaptation of the bivalve embryotoxicity assay for the high throughput screening of emerging contaminants in *Mytilus galloprovincialis*. *Mar. Environ. Res.* 99, 1–8. <http://dx.doi.org/10.1016/j.marenvres.2014.05.007>.
- Flint, S., Markle, T., Thompson, S., Wallace, E., 2012. Bisphenol A exposure, effects, and policy: a wildlife perspective. *J. Environ. Manage.* 104, 19–34. <http://dx.doi.org/10.1016/j.jenvman.2012.03.021>.
- Franzellitti, S., Fabbri, E., 2013. Cyclic-amp mediated regulation of ABCB mRNA expression in mussel haemocytes. *PLoS One* 8, e61634. <http://dx.doi.org/10.1371/journal.pone.0061634>.
- Hamdoun, A., Epel, D., 2007. Embryo stability and vulnerability in an always changing world. *Proc. Natl. Acad. Sci. U. S. A.* 104, 1745–1750. <http://dx.doi.org/10.1073/pnas.0610181104>.
- Hamlin, H.J., Guillette, L.J., 2011. Embryos as targets of endocrine disrupting contaminants in wildlife. *Birth Defects Res. C. Embryo Today* 93, 19–33. <http://dx.doi.org/10.1002/bdrc.20202>.
- Huss, J.M., Garbacz, W.G., Xie, W., 2015. Constitutive activities of estrogen-related receptors: transcriptional regulation of metabolism by the ERR pathways in health and disease. *Biochim. Biophys. Acta* 1852, 1912–1927. <http://dx.doi.org/10.1016/j.bbdis.2015.06.016>.
- Kadar, E., Simmance, F., Martin, O., Voulvoulis, N., Widdicombe, S., Mitov, S., Lead, J.R., Readman, J.W., 2010. The influence of engineered Fe_2O_3 nanoparticles and soluble (FeCl_3) iron on the developmental toxicity caused by CO_2 -induced seawater acidification. *Environ. Pollut.* 158, 3490–3497. <http://dx.doi.org/10.1016/j.envpol.2010.03.025>.
- Kreiling, J.A., Jessen-Eller, K., Miller, J., Seegal, R.F., Reinisch, C.L., 2001. Early development of the serotonergic and dopaminergic nervous system in *Spisula solidissima* (surf clam) larvae. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* 130, 341–351. [http://dx.doi.org/10.1016/S1095-6433\(01\)00394-4](http://dx.doi.org/10.1016/S1095-6433(01)00394-4).
- Kurihara, H., Asai, T., Kato, S., Ishimatsu, A., 2008. Effects of elevated pCO_2 on early development in the mussel *Mytilus galloprovincialis*. *Aquat. Biol.* 4, 225–233. <http://dx.doi.org/10.3354/ab00109>.
- Liu, Y., Tam, N.F.Y., Guan, Y., Yasojima, M., Zhou, J., Gao, B., 2011. Acute toxicity of nonylphenols and bisphenol A to the embryonic development of the abalone *Haliotis diversicolor* supertexta. *Ecotoxicology* 20, 1233–1245.
- Marin, F., Luquet, G., Marie, B., Medakovic, D., 2008. Molluscan shell proteins: primary structure, origin, and evolution. *Curr. Top. Dev. Biol.* 80, 209–276. [http://dx.doi.org/10.1016/S0070-2153\(07\)80006-8](http://dx.doi.org/10.1016/S0070-2153(07)80006-8).
- Medakovic, D., 2000. Carbonic anhydrase activity and biomineralization process in embryos, larvae and adult blue mussels *Mytilus edulis* L. *Helgol. Mar. Res.* 54, 1–6. <http://dx.doi.org/10.1007/s101520050030>.
- Nagasawa, K., Treen, N., Kondo, R., Otoki, Y., Itoh, N., Rotchell, J.M., Osada, M., 2015. Molecular characterization of an estrogen receptor and estrogen-related receptor and their autoregulatory capabilities in two *Mytilus* species. *Gene* 564, 153–159. <http://dx.doi.org/10.1016/j.gene.2015.03.073>.
- Przeslawski, R., Byrne, M., Mellin, C., 2015. A review and meta-analysis of the effects of multiple abiotic stressors on marine embryos and larvae. *Glob. Chang. Biol.* 21, 2122–2140. <http://dx.doi.org/10.1111/gcb.12833>.
- Rubin, B.S., 2011. Bisphenol A: an endocrine disruptor with widespread exposure and multiple effects. *J. Steroid Biochem. Mol. Biol.* 127, 27–34. <http://dx.doi.org/10.1016/j.jsbmb.2011.05.002>.
- Santangel, S., Maradonna, F., Gioacchini, G., Cobellis, G., Piccinetti, C.C., Dalla Valle, L., Carnevali, O., 2016. BPA-induced deregulation of epigenetic patterns: effects on female zebrafish reproduction. *Sci. Rep.* 6, 21982. <http://dx.doi.org/10.1038/srep21982>.
- Schmittgen, T.D., Livak, K.J., 2008. Analyzing real-time PCR data by the comparative C_T method. *Nat. Protoc.* 3, 1101–1108. <http://dx.doi.org/10.1038/nprot.2008.73>.
- Schönitzer, V., Weiss, L.M., 2007. The structure of mollusc larval shells formed in the presence of the chitin synthase inhibitor Nikkomycin Z. *BMC Struct. Biol.* 7, 71. <http://dx.doi.org/10.1186/1472-6807-7-71>.
- Scott, A.P., 2013. Do mollusks use vertebrate sex steroids as reproductive hormones? II. Critical review of the evidence that steroids have biological effects. *Steroids* 78, 268–281. <http://dx.doi.org/10.1016/j.steroids.2012.11.006>.
- Tisler, T., Krel, A., Gerzeli, U., Erjavec, B., Dolenc, M.S., Pintar, A., 2016. Hazard identification and risk characterization of bisphenols A, F and AF to aquatic organisms. *Environ. Pollut.* 212, 472–479. <http://dx.doi.org/10.1016/j.envpol.2016.02.045>.
- Tse, W.K.F., Yeung, B.H.Y., Wan, H.T., Wong, C.K.C., 2013. Early embryogenesis in zebrafish is affected by bisphenol A exposure. *Biol. Open* 2, 466–471. <http://dx.doi.org/10.1242/bio.20134283>.
- Vandenberg, L.N., Hunt, P.A., Myers, J.P., Vom Saal, F.S., 2013. Human exposures to bisphenol A: mismatches between data and assumptions. *Rev. Environ. Health* 28, 37–58. <http://dx.doi.org/10.1515/reveh-2012-0034>.
- Vendrami, D.L.J., Shah, A., Telesca, L., Hoffman, J.L., 2016. Mining the transcriptomes of four commercially important shellfish species for single nucleotide polymorphisms within biomineralization genes. *Mar. Genom.* 27, 17–23. <http://dx.doi.org/10.1016/j.margen.2015.12.009>.
- Voronezhskaya, E.E., Nezhlin, L.P., Odintsova, N.A., Plummer, J.T., Croll, R.P., 2008. Neuronal development in larval mussel *Mytilus trossulus* (Mollusca: Bivalvia). *Zoomorphology* 127, 97–110. <http://dx.doi.org/10.1007/s00435-007-0055-z>.
- Weiss, I.M., Schönitzer, V., 2006. The distribution of chitin in larval shells of the bivalve mollusk *Mytilus galloprovincialis*. *J. Struct. Biol.* 153, 264–277. <http://dx.doi.org/10.1016/j.jsb.2005.11.006>.
- Weiss, I.M., Tüross, N., Addadi, L., Weiner, S., 2002. Mollusc larval shell formation: amorphous calcium carbonate is a precursor phase for aragonite. *J. Exp. Zool.* 293, 478–491. <http://dx.doi.org/10.1002/jez.90004>.
- Yin, H., Ji, B., Dobson, P.S., Mosbahi, K., Glidle, A., Gadegaard, N., Freer, A., Cooper, J.M., Cusack, M., 2009. Screening of biomineralization using microfluidics. *Anal. Chem.* 81, 473–478. <http://dx.doi.org/10.1021/ac801980b>.
- Yin, Y., Huang, J., Paine, M.L., Reinhold, V.N., Chastee, N.D., 2005. Structural characterization of the major extrapallial fluid protein of the mollusc *Mytilus edulis*: implications for function. *Biochemistry* 44, 10720–10731. <http://dx.doi.org/10.1021/bi0505565>.
- Zoeller, R.T., Brown, T.R., Doan, L.L., Gore, A.C., Skakkebaek, N.E., Soto, A.M., Woodruff, T.J., Vom Saal, F.S., 2012. Endocrine-disrupting chemicals and public health protection: a statement of principles from the Endocrine Society. *Endocrinology* 153, 4097–4110. <http://dx.doi.org/10.1210/en.2012-1422>.