

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Nuclear Inositide Signaling via Phospholipase C

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Ratti, S., Mongiorgi, S., Ramazzotti, G., Follo, M.Y., Mariani, G.A., Suh, P., et al. (2017). Nuclear Inositide Signaling via Phospholipase C. JOURNAL OF CELLULAR BIOCHEMISTRY, 118(8), 1969-1978 [10.1002/jcb.25894].

Availability:

This version is available at: <https://hdl.handle.net/11585/597672> since: 2020-01-14

Published:

DOI: <http://doi.org/10.1002/jcb.25894>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the accepted manuscript of:

Ratti S, Mongiorgi S, Ramazzotti G, Follo MY, Mariani GA, Suh PG, McCubrey JA, Cocco L, Manzoli L. Nuclear Inositide Signaling Via Phospholipase C. J Cell Biochem. 2017 Aug;118(8):1969-1978.

Available at <https://doi.org/10.1002/jcb.25894>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Nuclear Inositide Signaling via Phospholipase C[†]

**Stefano Ratti¹, Sara Mongiorgi¹, Giulia Ramazzotti¹, Matilde Y. Follo¹, Giulia A. Mariani¹,
Pann-Ghill Suh², James A. McCubrey³, Lucio Cocco^{1*} and Lucia Manzoli¹**

¹ Cellular Signalling Laboratory, Department of Biomedical and Neuromotor Sciences, University
of Bologna, via Irnerio 48, 40126 Bologna, Italy

² School of Life Sciences, Ulsan National Institute of Science and Technology, Ulsan, Republic of
Korea

³ Department of Microbiology & Immunology, Brody School of Medicine, East Carolina
University, Greenville, NC 27834, USA

* Correspondence to: Lucio Cocco, Cellular Signalling Laboratory, Department of Biomedical and
Neuromotor Sciences, University of Bologna, via Irnerio 48, 40126 Bologna, Italy. E-mail:
lucio.cocco@unibo.it

[†]This article has been accepted for publication and undergone full peer review but has not been
through the copyediting, typesetting, pagination and proofreading process, which may lead to
differences between this version and the Version of Record. Please cite this article as doi:
[10.1002/jcb.25894]

ABSTRACT: the existence of an independent nuclear inositide pathway distinct from the cytoplasmic one has been demonstrated in different physiological systems and in diseases. In this prospect we analyze the role of PI-PLC β 1 nuclear isoform in relation to the cell cycle regulation, the cell differentiation and different physiopathological pathways focusing on the importance of the nuclear localization from both molecular and clinical point of view. PI-PLC β 1 is essential for G1/S transition through DAG and Cyclin D3 and plays also a central role in G2/M progression through Cyclin B1 and PKC α . In the differentiation process of C2C12 cells PI-PLC β 1 increases in both myogenic differentiation and osteogenic differentiation. PI-PLC β 1 and Cyclin D3 reduction has been observed in Myotonic Dystrophy (DM) suggesting a pivotal role of these enzymes in DM physiopathology. PI-PLC β 1 is also involved in adipogenesis through a double phase mechanism. Moreover PI-PLC β 1 plays a key role in the normal hematopoietic differentiation where it seems to decrease in erythroid differentiation and increase in myeloid differentiation. In Myelodysplastic Syndromes (MDS) PI-PLC β 1 has a genetic and epigenetic relevance and it is related to MDS patients' risk of Acute Myeloid Leukemia (AML) evolution. In MDS patients PI-PLC β 1 seems to be also a therapeutic predictive outcome marker. In the central nervous system PI-PLC β 1 seems to be involved in different pathways in both brain cortex development and synaptic plasticity related to different diseases. Another PI-PLC isozyme that could be related to nuclear activities is PI-PLC ζ that is involved in infertility processes. This article is protected by copyright. All rights reserved

KEY WORDS: SIGNALING, INOSITIDES, NUCLEUS, PHOSPHOLIPASE C, PATHOPHYSIOLOGY

NUCLEAR INOSITIDE SIGNALING

During the last 30 years nuclear inositide signaling has been investigated in many different systems and diseases. The eukaryotic outer nuclear membrane is made of protein and phospholipids and comes from the endoplasmic reticulum. For this reason cell fractionation showed nuclei with a high amount of lipids, due to the incomplete removal of the nuclear double layer. After the total removal of the nuclear envelope by detergents it was interesting to observe that still a significant quantity of lipids was present inside the nucleus. Even though nuclear Phosphatidylcholine (PC) is the main phospholipid, Phosphatidylinositols (PIs), present in small amount (Tribble et al., 2016), have specific role in intranuclear signaling therefore in nuclear function regulation and behave differently from their counterparts in the cytoplasm. The autonomous nuclear PI cycle in eukaryotic cells is involved in different regulation processes, from cell proliferation to differentiation and many others (Irvine, 2003). In this prospect we will try to describe the state of the art of nuclear inositide signaling via Phospholipase C (PLC), the pathophysiological involvement of PI-PLCs, the fields remained undiscovered and the possible researches related to this intriguing and complicated world.

HISTORICAL BACKGROUND

In 1987, for the first time, using Friend cell highly purified nuclei (washed free of nuclear membrane by Triton) Cocco et al. suggested that Phosphatidylinositol-4-phosphate (PI4P) and Phosphatidylinositol-4,5-bisphosphate (PI(4,5)P₂) were specifically present inside the nucleus. This work proposed that these PIs could have an important role in signal transduction parallel to the more documented cytoplasmic one (Cocco et al., 1987). Later, the stimulation with Insulin-like growth factor-1 (IGF-1) of Swiss 3T3 fibroblasts demonstrated the IGF-1 regulation of a distinct nuclear PI metabolism (Cocco et al., 1988). Moreover the treatment of Swiss 3T3 fibroblasts with IGF-1 determined a decrease in nuclear PI4P and PI(4,5)P₂ with a consequential increase in Diacylglycerol (DAG) and therefore a translocation of protein kinase C (PKC) inside the nucleus. On the contrary Swiss 3T3 fibroblast treated with bombesin showed activation only of the classical cytoplasmic pathway (Divecha et al., 1991, Martelli et al., 1991). The presence of these enzymes at the nuclear level suggests their essential role in nuclear signaling that, otherwise, would not be possible to be realized at the cytoplasmic level only. Moreover for being able to identify a certain amount of these PIs inside the nucleus their presence should be stable or, at least, long enough to be localized and of course to perform a certain nuclear activity. For these reasons these data opened a new field of research in nuclear inositide signaling via PLC, independent from the cytoplasmic one. Since the beginning of these experiments up to now there have been lots of enlightenments on this theme, but many aspects remain still unclear and critical.

NUCLEAR INOSITIDE SUBCELLULAR LOCALIZATION AND FUNCTIONS

If it is known that PIs are also present within the nucleus, their precise topography is still under investigation. Different studies focused on nuclear compartments in order to clarify PI nuclear geography. The main difficulty is that some PIs have different possible routes of synthesis and also removal that are localized in different regions inside and outside the nucleus (Irvine, 2003). For example PI(4,5)P₂ could be synthesized by PIP-5-kinase (PIP5K) or by PIP-4-kinase (PIP4K) starting from different substrates, respectively PI-4-phosphate (PI-4P) and PI-5P (Bulley et al., 2015). PIP5K and PIP4K have both different isoforms with different localization from the plasma membrane to

the nucleus itself: PIP5K α (plasma membrane), PIP5K β (both plasma membrane and perinuclear region), and PIP5K γ (in focal adhesions and adherent junctions in epithelial cells); PIP4K2A (plasma membrane and cytoplasm), PIP4K2B (nucleus) and PIP4K2C (endomembrane compartment) (Bultsma, et al. 2010). Considering our geographical focus the main question here is which class of these two PIPKs is responsible for the PI(4,5)P₂ synthesis inside the nucleus. Several experiments evidenced that PIP5K was accountable of almost 90% of PI(4,5)P₂ production within the nucleus, while PIP4K was responsible for only a little amount (Balla & Balla, 2006). These data suggest that there are several possible pathways for the same PI synthesis, related to its localization and function. Moreover PI4-kinase is mainly present in nuclear peripheral matrix, instead PI-PLC, DAG kinase, and PI4P5-kinase seem to be present in the internal one (Payraastre et al., 1992). On rat liver cells the isolation of nuclei with Triton X-100 at a concentration of 0,04%, in order to remove completely the bilayer nuclear envelope, showed that there were still several nuclear pools of different PIs such as PI, PI(4,5)P₂, phosphatidic acid (PA), DAG, and phosphatidylinositol 3,4,5-triphosphate (PI(3,4,5)P₃) (Vann et al., 1997). Moreover it has been evidenced that PI(4,5)P₂ is related to nuclear speckles. These speckles are interchromatin granule clusters where different PIs and other related enzymes are localized regulating chromatin remodeling, nuclear functions and RNA splicing (Boronenkov et al., 1998, Osborne et al., 2001).

The complexity of this topic is related to the difficulties of reproducible nuclear purification techniques and the variables in the metabolism and localization of these enzymes. In any case, these data suggest that PIs are somehow and at some point present inside specific nuclear regions and can therefore perform nuclear functions. Their temporary or stable permanence within the nucleus should then be useful for some nuclear activities that, instead, would not be possible in the cytoplasm (Tribble et al., 2016).

Indeed nuclear PIs are involved in many activities and can control several fundamental nuclear processes. As we saw above, their localization in specific nuclear regions named speckles strongly suggests their role in transcription regulation and in chromatin remodeling (Boronenkov et al., 1998). Proteins can bind PIs through Plant Homeo Domains (PHD) fingers making the interaction with other nuclear proteins, ligands, histones and chromatin possible. PIs can modify histone N-tails characteristics regulating the activity of several enzymes involved in chromatin remodeling, leading to different chromatin architecture and therefore gene transcription. Moreover they are able to modulate gene transcription interacting with transcription complexes and nuclear proteins. PIs can also directly affect RNA maturation affecting several gene expression and therefore PIs can have pivotal roles in different physiopathological processes (Shah et al., 2013). Between all the PIs that are present in the nucleus of eukaryotic cells we will mainly focus on the PI-PLCs and its related enzymes trying to analyze their physiopathological role.

NUCLEAR PI-PLC FEATURES

PLCs are a class of 13 isozymes, divided in six families: PI-PLC- β (1-4), - γ (1-2), - δ (1, 3 and 4), - ϵ , - ζ and η (1-2) (Gresset et al., 2012). These different enzymes share conserved structural features and can all hydrolyze PI(4,5)P₂ in response to several ligand bonds with cell surface receptors. This process produces DAG and inositol-1,4,5-trisphosphate (IP₃). IP₃ is responsible for the release of Ca²⁺ that, together with DAG, leads to the activation of PKC. This molecular cascade is essential for transmembrane signal transductions that result in cellular communication, proliferation and differentiation. Beyond these routes, PLCs are also localized within the nucleus and are independent from the cytoplasmic pathways. As seen before, we now know that several PIs are

localized inside the nucleus itself, but the specific mechanism of activation is still uncertain. Cytoplasmic PLC activation is related to G-protein-coupled receptor (GPCR) and receptor tyrosine kinase (RTK), but the nuclear activation apparatus is not completely defined. It is possible that IGF-I dependent signal, through a negative feedback by PKC α , modulates nuclear PI-PLC activation (Xu, et al., 2001). Therefore it seems that the nuclear route has different initial triggers and different mechanism compared to the cytoplasmic ones. Between the different PLC isozymes we will focus on PI-PLC β 1, the most investigated PI-PLC within the nucleus (Martelli et al., 1992).

PI-PLC β 1 gene is located on the short arm of chromosome 20 (20p12) and produces two splicing variants: PI-PLC β 1a (150 kDa) and PI-PLC β 1b (140 kDa) that differentiate one another for the C-terminal sequence and their localization (Bahk et al., 1998). Specifically, both of them have a Nuclear Localization Sequence (NLS), but PLC β 1a also has a Nuclear Export Sequence (NES), and for this reason it localizes also in the cytoplasm (Follo et al., 2006). On the other hand, PI-PLC β 1b is mainly located inside the nucleus (Martelli et al., 2005). In order to clarify the mechanism of PI-PLC β 1 nuclear import, PLC β 1a isoform was mutated at the C-terminal sequence. This experiment resulted in a significant reduction in the nuclear localization of the enzyme confirming that the C-terminal sequence is involved in the nuclear import (Kim, et al., 1996).

NUCLEAR INOSITIDE SIGNALLING IN CELL CYCLE REGULATION

We will now analyze the role of this enzyme in cell cycle regulation even if we have to consider the difficulties of synchronizing or arresting in a specific cell cycle phase our cell population. Moreover, working with lipids makes always hard to obtain highly purified nuclei in order to be sure to observe the nuclear PI polls.

Nuclear PI-PLC β 1: Nuclear PI-PLC β 1 is involved in cell cycle progression and can regulate the levels of different cyclins such as cyclin D3, cyclin E and cyclin B1. Cells overexpressing PI-PLC β 1 show an increase of Cyclin D3 levels and a higher percentage of cells in S phase, compared to the controls in K562 human erythroleukemia cell line (Poli et al., 2016a). Moreover PLC β 1 can regulate Cyclin E that plays a pivotal role in S-phase progression (Piazzi et al., 2015). PI-PLC β 1 can therefore modulate G1/S cell cycle transition through these cyclins. Another class of nuclear PIs that very recently resulted to be essential during this cell cycle phase in K562 cells are the Diacylglycerol Kinases (DGKs) and particularly the DGK α (Poli et al., 2016b). Furthermore it is interesting to observe that PI-PLC β 1 can be also involved in G2/M cell cycle progression. In 2005 Lukinovic-Skudar et al. described two peaks of PLC during both G1/S transition and G2/M progression (Lukinovic-Skudar et al., 2005). We will now analyze the nuclear PI-PLC β 1 mainly related PIs.

Nuclear DAG: as it occurs in the cytoplasm, also in the nucleus PI-PLC activity is responsible for DAG and IP₃ synthesis. IP₃ should perform his activity as a Ca²⁺ mobilizer inside the nucleus, too and nuclear DAG shows level changes during the cell cycle. As before, it seems that DAG follows nuclear PI-PLC oscillations observed in G1/S and G2/M phases (Lukinovic-Skudar et al., 2005). These data gave more strength to the previous experiments performed on HL60 cell lines (Sun et al., 1997) and U937 cell lines (Deacon et al., 2002) underlining the pivotal role of PLC β 1 and its related enzymes in cell cycle regulation. It has been demonstrated a peak in nuclear DAG production followed by an increase of PKC α and cyclin B1 in G2/M K562 synchronized cells. Indeed the inhibition of PLC activity leads to a reduction in nuclear IP₃ synthesis with a consequential decrease in nuclear translocation of PKC α and Cyclin B1 that are essential in G2/M phase (Poli et

al., 2014). As previously shown the role of PI-PLC is central for the PI cascade in cell cycle regulation and interaction with Cyclins. The probable consequence of DAG generation within the nucleus is an activation of nuclear PKC that leads to several cellular processes.

Nuclear PKC: PKC can be present inside the nucleus in different isoforms (Irvine, 2003). Both its two isoforms PKC α and PKC β II can translocate inside the nucleus after DAG stimulation. As for other PIs, this translocation should be long enough to be observed and to be able to play some role in nuclear pathways. In Swiss 3T3 cell line PKC α can translocate inside the nuclei as a result of a probable nuclear DAG stimuli (Martelli et al., 1991) and also in U937 cell line the translocation of PKC β II inside the nuclei was showed after DAG increase (Deacon et al., 2002). The intriguing data is that PKC can translocate inside the nuclei also after other stimuli such as IGF-1, Platelet derived growth factor or Bryostatins (Irvine, 2003). In addition to these data the same stimuli can determine the translocation in the nucleus of both PKC α and PKC β II isoforms. Another interesting aspect is that a PI-PLC β 1 stable overexpression is related with a decrease of PKC α levels, and this decrease is possibly related to PKC α hyper and stable activation induced by DAG continuous production (Poli et al., 2014). In G2/M-synchronized MEL cells the change in levels of PI-PLC β 1 altered nuclear PKC α accumulation with consecutive alteration in the mitotic process (Fiume et al., 2009). PKC has indeed different targets, like the Lamin B1, strongly linked to the cellular cycle. As we saw above, there is a PI cascade that starting from nuclear PI-PLC β 1, through DAG and IP $_3$ or and Ca $^{2+}$, activates PKC α /PKC β II and other molecules controlling the cell cycle. It is possible that DAG and so PKC regulate cell cycle and cell proliferation in relation to different signals and cell population.

NUCLEAR PI(4,5)P $_2$: PI(4,5)P $_2$ is not only a substrate for PI-PLC, but this PI seems to be essential in cell cycle progression and other nuclear functions. As reported before, PI(4,5)P $_2$ is largely produced by PIP5K that can interact with Retinoblastoma protein (pRb) which, in turn, controls the progression to S phase in the cell cycle (Divecha et al., 2002). The link between PIP5K and pRb can determine PIP5K activation and then PI(4,5)P $_2$ synthesis. For this reason PI(4,5)P $_2$ plays an important role in cell cycle progression and in the nuclear PI-PLC cascade. Moreover PI(4,5)P $_2$ could have structural and also regulatory function in RNA splicing, indeed this class of PIs resides in the interchromatin granular structures interacting dynamically with different chromatin structures during the cell cycle (Osborne et al., 2001) (*Figure 1*).

NUCLEAR PI-PLC β 1 IN CELL DIFFERENTIATION

Nuclear PI-PLC β 1 plays a crucial role in cell differentiation of different cell systems. We will focus on the hematopoietic, osteogenic, myogenic and adipogenic systems in order to concentrate on the different function and variations of this pivotal PI in cell differentiation. Indeed PI-PLC β 1 seems to have reverse modulation in distinctive cell systems (*Figure 2*).

Hematopoietic differentiation: PI-PLC β 1 has shown to be involved in the hematopoietic system performing pivotal roles in both erythroid and myeloid differentiation. Interestingly this nuclear PI has opposite behaviour in these two subsystems: in the erythroid differentiation PI-PLC β 1 seems to decrease, whereas in the myeloid differentiation PI-PLC β 1 appears to increase (Follo et al., 2015). More specifically in murine erythroleukemia cells (MEL) and in normal CD34+ cells with

erythropoietin stimulation towards erythroid lineage PI-PLC β 1 has shown to decrease during the erythroid differentiation process. Furthermore PI-PLC β 1 has presented the same trend also in Myelodysplastic Syndromes (MDS), responding to erythropoiesis stimulating agents (Follo et al., 2012). These data suggest that PI-PLC β 1 plays an inhibiting role in erythroid differentiation that is countered by PI-PLC γ 1, another PI-PLC isozyme that increases in erythroid differentiation. The negative role of PI-PLC β 1 in erythroid differentiation was first observed inside the nucleus with an increase of PI(4,5)P₂ and a decrease of both PI-PLC β 1 and DAG in differentiating MEL cells (Cocco et al., 1987).

In myeloid differentiation the role of nuclear PI-PLC β 1, studied using a MDS experimental model, seems to be opposite to the one just observed in erythroid differentiation. Indeed MDS cells treated with azacitidine, a hypomethylating agent used for the induction of a normal myeloid differentiation, determined a decrease of PI-PLC β 1 promoter methylation and thus an increase of nuclear PI-PLC β 1. The potential positive role of PI-PLC β 1 in myeloid differentiation could be correlated with the recruitment of specific myeloid transcription factors, such as Myeloid zinc finger-1 (MZF-1), during PI-PLC β 1 expression increase (Cocco et al., 2016). Observing also the clinical aspects of MDS patients treated with azacitidine it is very interesting to point that the increase of PI-PLC β 1 expression is not only related to a favorable clinical response, but also to the durable response to the treatment. Indeed the reduction in PI-PLC β 1 expression can be associated with a loss of response even in patients that had, at first, a positive response to azacitidine (Fili et al., 2013). These data about the variation in PI-PLC β 1 expression could be useful to divide MDS patients that could undergo to azacitidine treatment into two groups: the ones who will probably show a favorable long term response to the drug and the ones who would be refractory to the treatment and thus could avoid its intrinsic risks (Cocco et al., 2015). Afterwards we will see MDS clinical aspects and possibilities related to PI-PLC β 1 and other PIs more in detail.

Osteogenic differentiation: nuclear PI-PLC β 1 seems to be involved also in osteogenic differentiation in relation to different osteogenic molecules. The osteogenic differentiation is stimulated by several signaling pathways, mainly controlled by bone morphogenic proteins (BMPs), Osterix (Osx/Sp7) and runt-related transcription factor 2 (runx2).

C2C12 cells, a mouse skeletal muscle cell line, can differentiate into osteoblasts after BMP-2 stimulation (Katagiri et al., 1994). Very interestingly, during the osteogenesis induced by BMP-2 in C2C12 cells it has been shown an increase of PI-PLC β 1 expression. In addition the down-regulation of PI-PLC β 1 gene expression results in an inhibition of the osteogenic differentiation process induced in C2C12 by BMP-2, as demonstrated by Alkaline phosphatase (ALP) reduction. Instead PI-PLC β 1 overexpression determines an increase of osteogenic differentiation as demonstrated from the rise of ALP and Osx/Sp7 (Ramazzotti et al., 2016). Indeed ALP and Bone gamma-carboxyglutamic acid containing protein (Bglap or Osteocalcin) are osteoblast marker genes that increase in response to the action of the main bone morphogenic proteins that control the osteoblastic differentiation.

PI-PLC β 1 is also related to miR-124, another molecule that is very relevant during the control of osteogenic differentiation. miR-124 can promote the myogenic differentiation and inhibit the osteogenic differentiation by regulating Osx/sp7. Specifically the inhibition of miR-214 in C2C12 cells leads to an increase of PI-PLC β 1 protein level and furthers the osteogenesis (Feng et al., 2011). Another very intriguing aspect of the relation between nuclear PIs and osteogenic differentiation is that, even without BMP-2, the overexpression of PI-PLC β 1 still determines a certain level of osteoblast differentiation, albeit reduced. This could signify the capability of PI-

PLC β 1, per se, to induce osteogenic differentiation through possible downstream molecules.

Myogenic differentiation: There are different pathways related to the myogenic differentiation that can be activated by several events and that can lead to various cellular processes. The two main families of the transcription factors related to myogenic differentiation are the Myogenic Regulatory Factors (MRFs) and the Myocyte Enhancer Factor-2 (MEF2) (Ramazzotti et al., 2016). C2C12 cells under low serum conditions can differentiate into myotubes. These cells synthesize muscle specific proteins like MyoD and Myogenin (MyoG) that are essential for the myoblast alignment, elongation and fusion that allows myotube formation. During the myotube formation nuclear PI-PLC β 1 increased in C2C12 cells induced to differentiate. This PI-PLC β 1 increase was associated with a Cyclin D3 level increase, too (Cocco et al., 2016). MyoD stimulates Cyclin D3 expression that can thus binds the unphosphorylated pRb responsible for the myoblast cell cycle recession (Ramazzotti et al., 2016). Moreover Cyclin D3 promoter has a binding sequence for PI-PLC β 1, and this bond could be responsible for Cyclin D3 activation in relation to PI-PLC β 1 activity (Faenza et al., 2012). More specifically the transcription factor c-jun needs to bond to Cyclin D3 promoter in order to determine its activation during PI-PLC β 1 signaling cascade (Ramazzotti et al., 2016). This possible pathway starts with PI-PLC β 1 catalytic activity that leads to IP₃ synthesis, which is phosphorylated by Inositol Polyphosphate Multikinase (IPMK) with the production of higher phosphorylated inositol species such as IP₄, IP₅ and IP₆. IP₅ can induce β -catenin nuclear translocation, activating c-jun, thus Cyclin D3 promoter. β -catenin target genes are involved in several cellular processes like proliferation, differentiation and cancerogenic stimuli. This pathway could explain the key role of nuclear PI-PLC β 1 in myogenic differentiation. It is worthwhile to remember that also PI-PLC γ 1 expression increases during the myogenic differentiation. Interestingly, even if Cyclin D3 is essential in myogenic differentiation it seems not to be involved during osteogenesis. Indeed Cyclin D3 levels do not increase during osteogenesis neither after PI-PLC β 1 overexpression, for this reason it seems to promote myogenic, but not osteogenic differentiation (Ramazzotti et al., 2016). We will analyze later in this prospect the clinical correlation between PI-PLC β 1 and myogenic differentiation in the Myotonic dystrophies.

Adipogenic differentiation: nuclear PI-PLC β 1 and cyclin D3 could be involved in adipocyte differentiation, too. During the differentiation of 3T3-L1 adipocytes there is an up-regulation of the nuclear PI-PLC β 1. Interestingly there have been observed two different phases of this differentiation process. In the first phase, after 5 minutes of incubation with the differentiation media, the pathway is regulated by protein kinase RNA-like endoplasmic reticulum kinase (pERK) and PKC α . In this phase no further nuclear PI-PLC β 1 recruiting is needed, probably because the nuclear PI-PLC β 1 levels are already sufficient at this stage. In the second phase, after 2 days of differentiation, there is the necessity of PI-PLC β 1 nuclear translocation, independently from pERK and PKC α control. Interestingly the overexpression of PLC mutants, lacking the ERK phosphorylation site or the nuclear localization sequence showed that both PI-PLC β 1 phases are essential for an effective differentiation. Moreover the inhibition of PI-PLC β 1 avoided the overexpression of the complex formed by Cyclin D3 and Cyclin-dependent-kinase-4(CDK4), suggesting a role of PI-PLC β 1 in the cell cycle control during adipogenic differentiation (O'Carroll et al., 2009).

NUCLEAR PI-PLCs IN DISEASES

PI-PLCs have been studied as pivotal enzymes, clinical molecular targets and clinical prognostic/diagnostic factors in many physiopathological processes. The different PI-PLC isozymes are related to several pathologies in many human tissues and the understanding of these mechanisms could open new path for developing innovative therapeutic strategies. In this prospect we will focus on the diseases characterized by alterations of nuclear PI-PLCs.

Myelodysplastic Syndromes(MDS): MDS are a heterogeneous group of pathologies characterized by hematopoietic stem cell alteration that lead to anemia, neutropenia, bleeding and infections. Moreover 30% of MDS patients can evolve into acute myeloid leukemia (AML). The IPSS (International Prognostic Scoring Systems) and the WPSS (WHO classification-based Scoring System) classifications, mainly based on the blast number and karyotype, are used to divide MDS patients into two major groups: high and low risk of AML progression. Recently, the evidence of a clinical correlation between the presence of a PI-PLC β 1 monoallelic gene deletion and the progression of MDS to AML opened new perspectives of research and treatments (Follo et al., 2008). The gene encoding for PI-PLC β 1 has been mapped on chromosome 20p12. High-risk MDS patients should undergo allogenic hematopoietic stem cell transplantation (allo-HSCT), but for those who are not candidates, hypomethylating agents (HMAs), such Azacitidine (AZA), are now the first therapeutic choice. HMAs can also be used in low risk MDS patients. AZA is both a hypomethylating and a direct cytotoxic agent for abnormal hematopoietic cells. Patients affected by MDS with a high risk of AML evolution show a reduction in the expression of the nuclear PI-PLC β 1 variant that can also be related to the presence of a mono-allelic deletion of the gene. As PI-PLC β 1b can physiologically regulate the progression through the G1 phase of the cell cycle, the drastic reduction of nuclear PI-PLC β 1 expression could alter the normal cell cycle in patients with MDS. These data could suggest the analysis of PI-PLC β 1 gene expression as a prognostic marker for MDS progression to AML. Nuclear PLC β 1 in MDS is also epigenetically relevant (Jhanwar et al., 2015). In fact, several studies have shown that PLC β 1 is a molecular target for AZA (Follo et al., 2009). As a matter of fact, high-risk and low-risk patients that respond to the drug have shown an early increase of nuclear PI-PLC β 1 expression, a reduction of PI-PLC β 1 promoter methylation, an induction of normal myeloid differentiation and a better prognosis. Moreover, as described above, the increase of PI-PLC β 1 expression and the reduction in PI-PLC β 1 promoter methylation are not only related to a favourable clinical response, but also to a durable response to the treatment. This effect resulted particularly interesting, because, as AZA treatment needed several cycles in order to observe a clinical response, the molecular response could specifically and rapidly predict the future patients' outcome during demethylating therapies. In this way it would be possible to personalize MDS patients' therapies in order to differentiate the ones who would benefit of the treatment from the ones who would be refractory to the treatment and thus could avoid it (Cocco et al., 2016). Furthermore several studies showed that in MDS high-risk patients there is an inverse correlation between PI-PLC β 1 and PI3K expression. PI3K can phosphorylate Protein Kinase B (PKB) also known as Akt (Toker and Marmiroli, 2014). When there is a constitutive activation of Akt, through different phosphorylation, there is a decrease in MDS cell apoptosis that can be reverted after PI-PLC β 1 increase due to the demethylating therapies. Indeed PI-PLC β 1 increase could reduce the level of p-Akt inducing a higher level of apoptosis (Follo et al., 2007). This mechanism is still partially undiscovered, but a possible explication could be that as PI(4,5)P₂ is a common substrate of both PI-PLC β 1 and PI3Ks, the alterations in PI-PLC β 1 levels could cause PI(4,5)P₂ hydrolysis, reducing its availability for PI3K. This could eventually inhibit Akt activation (Follo et al., 2015). All these mechanisms are extremely interesting for the potentiality of nuclear PI-PLC β 1 to become a prognosis stratification marker and a treatment predictive outcome marker in patients

with MDS.

Nuclear inositides have been studied also in AML that can be a direct evolution of MDS. As we saw earlier in this prospect PIP4K2A catalyzes the phosphorylation of PI5P, synthesizing PI(4,5)P₂. The role of PIP4K2A has been examined in human MLL-AF9 and in murine AML cells, and further investigated also in human AML samples (Jude et al., 2015). AML cells were sensible to PIP4K2A depletion determining a cyclin-dependent kinase inhibitor CDKN1A (p21) accumulation, with thus an activation of the apoptosis. This mechanism seems to be related to the activation of mammalian Target of Rapamycin(mTOR), too. Considering this pathway it is then possible that PIP4K2A could also regulate mTOR. In addition it has been observed significant increase of PI5P following PIP4K2A depletion (Jude et al., 2015). PI5P has been related to TP53 and Akt activation and to cellular reactive oxygen species (ROS) accumulation in normal hematopoietic cells (Jones et al., 2013). Moreover PI5P might increase in response to several stress stimuli, mimicking cell stress induction. This mechanism could induce AML cells to apoptosis or perhaps increase their susceptibility to other critical processes, such as DNA damage. It is still not clear why normal hematopoietic cells and AML cells have different response to PIP4K2A depletion, but these differences could be used for developing new therapeutic strategies against AML (Jude et al., 2015).

Myotonic dystrophies (DM): DM are autosomal dominant neuromuscular degenerative disorders, characterized by a widely variable clinical picture and a slow progressive course, whose onset can occur at any age. The clinical picture is characterized by loss of muscle mass, myotonia, cataracts, defects in the cardiac conduction system, endocrine abnormalities and cognitive deficits in congenital cases. Moreover there is an anticipation phenomenon thus the disease onset tends to occur at an increasingly younger age from generation to generation in the same family.

These diseases are classified into DM type I (DM1) and DM type II (DM2). The majority of patients suffer of DM1, but both DM are caused by DNA tandem repeats that result in aberrant RNA accumulation in the nucleus that causes alteration in RNA-binding protein localization. In DM1 the mutated gene is called myotonic dystrophy protein kinase (DMPK) and encodes a myosin kinase expressed in skeletal muscles. This gene is located in the long arm of chromosome 19 (19q13.3). In the 3'UTR of this locus there is an unstable CTG triplet that can be repeated from 50 to 4000 times when in the normal population the range varies from 5 to 37 times. In DM2 there is a defect in zinc finger protein 9 (ZNF9) gene on chromosome 3 (3q21). The alteration is caused by CCTG expansion from 75 to over 11000, but in this case there seems to be no difference in the severity or the early onset of the disease (Cho and Tapscott, 2007). CUG triplet repeat RNA-binding protein 1 (CUGBP1) plays a central role in alternative splicing of specific target genes and can interact with Initiation Factor 2(eIF2) and Cyclin D3 inducing normal myogenic differentiation. This process can be altered in DM, indeed CUGBP1-eIF2 interactions are reduced in myotonic dystrophy differentiating cells with an impaired muscle differentiation (Faenza et al., 2012). Nevertheless, in DM1, ectopic expression of Cyclin D3 helps the increase of CUGBP1-eIF2 complex, improving the myogenic differentiation marker expression (Salisbury et al., 2008). Moreover, during the myogenesis in DM cells there is a decrease of PI-PLCβ1 expression that could be related to the reduction in Cyclin D3 transcription and induction as described above in the role of PI-PLCβ1 and Cyclin D3 in myogenic differentiation. Interestingly the normalization of PI-PLCβ1 expression in DM1 and DM2 myoblasts can cause a Cyclin D3 expression increase, determining a partially restored phenotype of the myotubes (Faenza et al., 2012). All these data suggest that both PI-PLCβ1 and Cyclin D3 could be investigated for future possible molecular therapies in order to induce a correct skeletal muscle differentiation in DM.

Brain disorders: PI-PLC isozymes are highly expressed in different brain structures. Specific neurotransmitters (NTs) and hormones trigger PI-PLC pathways, determining various activities in many brain regions (Yang et al., 2016). PI-PLC β 1 is present at high concentration in hippocampus, amygdala, lateral septum, olfactory bulb and cerebral cortex (Fukaya et al., 2008). Indeed it has been related to cortical development and synaptic plasticity (Spires et al., 2005). This PI has been associated with epilepsy and to schizophrenia in PI-PLC β 1 knocked out mice (Yang et al., 2016). Losses of PI-PLC β 1 were also observed in patients with epileptic encephalopathies, schizophrenia and bipolar disorders (Kurian et al., 2010, Lo Vasco et al., 2012, Lo Vasco et al., 2013). Also PI-PLC γ 1 has been correlated to neuronal cell migration and synaptic plasticity and its activity has been suggest to be of some relevance in epilepsy, Huntington disease, depression, bipolar disorders and Alzheimer disease (Yang et al., 2016). Moreover PI-PLC β 1 seems to be central in the control of endocannabinoid neuronal excitability in most of the brain areas, indeed in the postsynaptic sites, and it's pivotal in the synthesis of DAG which hydrolysis catalyzed by DAG-Lipase α (DAGL α) leads to 2-arachidonyl-glycerol (2-AG), the most represented CB1 cannabinoid receptor endogenous agonist. Released from the postsynaptic neuron, 2-AG can activate CB1 cannabinoid receptors on the presynaptic sites repressing other NT release (Montaña et al., 2012). Moreover PI-PLC β 1 can respond to two separate signals: depolarization and receptor activation, maintaining the brain inhibitory circuits through 2-AG (del Caño et al., 2014). Interestingly the pathway related to PI-PLC β 1/DAGL α /2-AG seems to be at neuronal nuclear level. In adult rat brain, double immunofluorescence staining and confocal laser scanning showed an overlapping pattern of both PI-PLC β 1 and DAGL α with nuclear speckles marker SC-35 and NeuN/Fox3. In addition PI-PLC β 1 was also highly expressed in neuronal nuclear regions full of PI(4,5)P $_2$. These data were enforced by the fact that the co-localization with Nuclear Pore Complex and Lamin B1 was instead not found (Montaña et al., 2012). There are also evidences of the relation between 2-AG and nuclear Peroxisome proliferator-activated receptor gamma (PPAR γ) as a receptor or as a prostaglandin precursor (del Caño et al., 2014). All these intriguing data prompt to better investigate the neuronal nuclear localization and mechanisms of PI-PLC β 1 related molecules in different pathophysiological brain functions. Furthermore, considering nuclear PI-PLC β 1 pivotal role in cell cycle regulation and the controversial and complicate role of cell cycle in several central nervous system insults and pathologies, it would be very fascinating to study nuclear PI-PLC β 1 functions during diverse brain injuries forms (del Caño et al., 2014).

Infertility: PI-PLC ζ is a specific sperm protein that has been related to nuclear infertility mechanisms. PI-PLC ζ activation and nuclear translocation mechanisms remain unknown, but at nuclear level this PI has been specifically connected with molecular oocyte activation. This PI-PLC is essential in inducing Ca $^{2+}$ release via IP $_3$ pathway (Saunders et al., 2002). Ca $^{2+}$ oscillations within the oocyte have been related to different processes responsible for its activation such as the cortical granule exocytosis, the release of meiotic arrest, the gene expression regulation, the recruitment of maternal mRNA, the pronuclear formation and the initiating of embryogenesis. The specific mechanism of these reactions is still unclear, but the central role of Ca $^{2+}$ is fundamental for the beginning of embryogenesis (Amdani et al., 2016). The principal cause of infertility after Intra Cytoplasmatic Sperm Injection (ICSI) is considered to be the alteration of oocyte activation mechanisms of which sperm defects seem to be the main responsible (Amdani et al., 2016). From one side, cRNA PI-PLC ζ and recombinant protein microinjections in mice and cattle oocyte determine Ca $^{2+}$ oscillations and oocyte activation (Saunders et al., 2002). From the other side

infertile man sperm with alteration in PI-PLC ζ quantity and quality fail to induce Ca²⁺ oscillations and therefore oocyte activation (Amdani et al., 2016) . For these reasons it is possible that PI-PLC ζ could be pivotal in fertility processes and could be considered a prognostic/diagnostic molecular marker in order to identify male patients that could benefit of assisted reproductive technology. Moreover, even if Ca²⁺ ionophores are the main agent for artificial oocyte activation, PI-PLC ζ could be a safer potential therapeutic agent (Amdani et al., 2016).

CONCLUSIONS

In this prospect we tried to highlight our understanding of nuclear inositide signaling in different cell systems and pathologies. This topic is at the same time very testing and intriguing due to the technical difficulties in focusing on the nuclear localization of such complicate pathways and to the importance of these enzymes in several pathophysiological mechanisms. We analyzed the PI nuclear signaling through PI-PLCs mainly focusing on PI-PLC β 1, the most studied nuclear PI-PLC, and its related molecules like PI(4,5)P₂, DAG, PKCs, cyclin D3 and many others. We started from the beginning of this lipid nuclear study in the late 80s up to the most recent molecular and clinical implications and possibilities. PI-PLC β 1 and its associated molecules have been implicated to be essential in the cell cycle during both G1/S transition and G2/M progression. Moreover PI-PLC β 1 is involved as pivotal enzyme in the normal mesenchymal cell differentiation in osteogenesis, myogenesis and adipogenesis. During the myogenic differentiation of C2C12 cells it has been shown an increase of both nuclear PI-PLC β 1 and Cyclin D3. On the contrary, PI-PLC β 1 and Cyclin D3 reduction has been observed in Myotonic Dystrophy (DM), suggesting a pivotal role of these enzymes in DM physiopathology. Similarly PI-PLC β 1 expression increases in the normal osteogenesis induced by BMP-2 in C2C12 cells and the down-regulation of PI-PLC β 1 gene expression determines an inhibition of the osteogenic differentiation as demonstrated by ALP reduction. PI-PLC β 1 seems to increase also in adipogenesis and its inhibition seems to stop the overexpression of the Cyclin D3-CDK4 complex. This central role has been also demonstrated in the hematopoietic differentiation with probably opposite behaviour between erythroid and myeloid differentiation. Very interestingly the nuclear inositide signaling via PI-PLC β 1 helped to create a bridge between molecular and clinical analyses, giving the opportunity to explore new potential therapeutic strategies and markers. Indeed there have been many progresses especially in understanding MDS pathological mechanism in AML evolution and in response to the therapies. Indeed nuclear PI-PLC β 1 could be used both as a prognostic marker in AML evolution and as a therapy predictive marker in MDS patients. PI-PLC β 1 has been investigated in the central nervous system where it seems to be involved in different processes and pathways in both brain cortex development and synaptic plasticity related to different disease. Nuclear PI-PLC β 1 is also involved in neurotransmitters regulation, suggesting a pivotal role in different brain mechanisms. Another PI-PLC isozyme that could be related to nuclear activities is PI-PLC ζ that is involved in oocyte activation and infertility processes where it could be considered as a molecular marker in order to identify male patients that could benefit of assisted reproductive technology. There are certainly many questions that remain to be clarified in relation to nuclear PI cycle and functions, but the importance of these themes and the related clinical prospects stimulate to persist on this field of investigation.

Acknowledgements: this work was supported by grant from Intesa San Paolo to LC, grants from Italian MIUR-PRIN 2015 to LM and MYF, grant #111104 from East Carolina University to JAM, grant from National Research foundation of Korea to PGS.

REFERENCES

- Amdani SN, Yeste M, Jones C, Coward K. (2016). Phospholipase C zeta (PLC ζ) and male infertility: Clinical update and topical developments. *Adv Biol Regul.* 61:58-67
- Bahk Y. Y., Song H., Baek S. H., Park B. Y., Kim H., Ryu S. H., & Suh P. G. (1998). Localization of two forms of phospholipase C- β 1, a and b, in C6Bu-1 cells. *Biochimica et Biophysica Acta - Lipids and Lipid Metabolism*, 1389(1): 76–80.
- Balla A., & Balla T. (2006). Phosphatidylinositol 4-kinases: old enzymes with emerging functions. *Trends in Cell Biology*, 16(7): 351–361.
- Boronenkov IV, Loijens JC, Umeda M, Anderson RA. (1998). Phosphoinositide signaling pathways in nuclei are associated with nuclear speckles containing pre-mRNA processing factors. *Mol Biol Cell* 9:3547–3560.
- Bulley S. J., Clarke J. H., Droubi A., Giudici M. L., & Irvine R. F. (2015). Exploring phosphatidylinositol 5-phosphate 4-kinase function. *Advances in Biological Regulation*, 57: 193–202.
- Bultsma Y., Keune W.-J., & Divecha N. (2010). PIP4K β interacts with and modulates nuclear localization of the high-activity PtdIns5P-4-kinase isoform PIP4K α . *The Biochemical Journal*, 430(2): 223–35.
- Cho D. H., & Tapscott S. J. (2007). Myotonic dystrophy: Emerging mechanisms for DM1 and DM2. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease*, 1772(2): 195–204.
- Cocco L., Finelli C., Mongiorgi S., Clissa C., Russo D., Bosi C., ... Follo M. Y. (2015). An increased expression of PI-PLC 1 is associated with myeloid differentiation and a longer response to azacitidine in myelodysplastic syndromes. *Journal of Leukocyte Biology*, 98(5): 769–780.
- Cocco L., Gilmour R. S., Ognibene A., Letcher A. J., Manzoli F. A., & Irvine R. F. (1987). Synthesis of polyphosphoinositides in nuclei of Friend cells. Evidence for polyphosphoinositide metabolism inside the nucleus which changes with cell differentiation. *The Biochemical Journal*, 248(3): 765–70.
- Cocco L., Manzoli L., Faenza I., Ramazzotti G., Yang Y. R., McCubrey J. A., ... Follo M. Y. (2016). Modulation of nuclear PI-PLC β 1 during cell differentiation. *Advances in Biological Regulation*, 60: 1–5.
- Cocco L., Martelli A. M., Gilmour R. S., Ognibene A., Manzoli F. A., & Irvine R. F. (1988). Rapid changes in phospholipid metabolism in the nuclei of Swiss 3T3 cells induced by treatment of the cells with insulin-like growth factor I. *Biochemical and Biophysical Research Communications*, 154(3): 1266–72.
- Deacon E. M., Pettitt T. R., Webb P., Cross T., Chahal H., Wakelam M. J. O., & Lord J. M. (2002). Generation of diacylglycerol molecular species through the cell cycle: a role for 1-stearoyl, 2-arachidonyl glycerol in the activation of nuclear protein kinase C- β 1 at G2/M. *Journal of Cell Science*, 115(Pt 5): 983–989.
- Del Caño, G., Montaña, M., Aretxabala, X., González-Burguera, I., López de Jesús, M., Barrondo, S., & Sallés, J. (2014). Nuclear phospholipase C- β 1 and diacylglycerol LIPASE- α in brain cortical neurons. *Advances in Biological Regulation*, 54: 12–23.
- Divecha N., Banfic H., Irvine R. F., Banfić H., & Irvine R. F. (1991). The polyphosphoinositide cycle exists in the nuclei of Swiss 3T3 cells under the control of a receptor (for IGF-I) in the plasma membrane, and stimulation of the cycle increases nuclear diacylglycerol and apparently induces translocation of protein kinase. *Embo J*, 10(11): 3207–3214.
- Divecha N., Roefs M., Los A., Halstead J., Bannister A., & D'Santos C. (2002). Type IPIPKs interact with and are regulated by the retinoblastoma susceptibility gene product-pRB. *Curr. Biol.*, 12(7): 582–587.
- Faenza I., Blalock W., Bavelloni a., Schoser B., Fiume R., Pacella S., ... Cocco L. (2012). A role for PLC 1 in myotonic dystrophies type 1 and 2. *The FASEB Journal*, 26(7): 3042–3048.
- Feng Y., Cao J.-H., Li X.-Y., & Zhao S.-H. (2011). Inhibition of miR-214 expression represses proliferation and differentiation of C2C12 myoblasts. *Cell Biochemistry and Function*, 29(5): 378–83.
- Fili C., Malagol, M., Follo M. Y., Finelli C., Iacobucci I., Martinelli G., ... Russo D. (2013). Prospective phase II study on 5-days azacitidine for treatment of symptomatic and/or erythropoietin unresponsive patients with low/INT-1-risk myelodysplastic syndromes. *Clinical Cancer Research*, 19(12): 3297–3308.

- Fiume R., Ramazzotti G., Teti G., Chiarini F., Faenza I., Mazzotti G., ... Cocco L. (2009). Involvement of nuclear PLC β 1 in lamin B1 phosphorylation and G2/M cell cycle progression. *FASEB Journal : Official Publication of the Federation of American Societies for Experimental Biology*, 23(3): 957–66.
- Follo, M. Y., Bosi, C., Finelli, C., Fiume, R., Faenza, I., Ramazzotti, G., ... Cocco, L. (2006). Real-time PCR as a tool for quantitative analysis of PI-PLC β 1 gene expression in myelodysplastic syndrome. *International Journal of Molecular Medicine*, 18(2): 267–271.
- Follo, M. Y., Finelli, C., Clissa, C., Mongiorgi, S., Bosi, C., Martinelli, G., ... Cocco, L. (2008). Phosphoinositide-Phospholipase C 1 Mono-Allelic Deletion Is Associated With Myelodysplastic Syndromes Evolution Into Acute Myeloid Leukemia. *Journal of Clinical Oncology*, 27(5): 782–790.
- Follo, M. Y., Finelli, C., Mongiorgi, S., Clissa, C., Bosi, C., Testoni, N., ... Cocco, L. (2009). Reduction of phosphoinositide-phospholipase C β 1 methylation predicts the responsiveness to azacitidine in high-risk MDS. *Proceedings of the National Academy of Sciences of the United States of America*, 106(39): 16811–16816.
- Follo, M. Y., Manzoli, L., Poli, A., McCubrey, J. A., & Cocco, L. (2015). PLC and PI3K/Akt/mTOR signalling in disease and cancer. *Advances in Biological Regulation*, 57: 10–16.
- Follo, M. Y., Mongiorgi, S., Bosi, C., Cappellini, A., Finelli, C., Chiarini, F., ... Martelli, A. M. (2007). The Akt/mammalian target of rapamycin signal transduction pathway is activated in high-risk myelodysplastic syndromes and influences cell survival and proliferation. *Cancer Research*, 67(9): 4287–4294.
- Follo, M. Y., Mongiorgi, S., Clissa, C., Paolini, S., Martinelli, G., Martelli, a M., ... Cocco, L. (2012). Activation of nuclear inositide signalling pathways during erythropoietin therapy in low-risk MDS patients. *Leukemia*, 26(12): 2474–82.
- Fukaya M., Uchigashima M., Nomura S., Hasegawa Y., Kikuchi H., & Watanabe M. (2008). Predominant expression of phospholipase C β 1 in telencephalic principal neurons and cerebellar interneurons, and its close association with related signaling molecules in somatodendritic neuronal elements. *The European Journal of Neuroscience*, 28(9): 1744–59.
- Gresset A., Sondek J., & Harden T. K. (2012). The phospholipase C isozymes and their regulation. *Sub-Cellular Biochemistry*, 58: 61–94.
- Irvine R. F. (2003). Nuclear lipid signalling. *Nature Reviews Molecular Cell Biology*, 4(5): 349–361.
- Jhanwar SC (2015). Genetic and epigenetic pathways in myelodysplastic syndromes: A brief overview. *Adv Biol Regul.* 58:28-37
- Jones, D. R., Foulger, R., Keune, W.-J., Bultsma, Y., & Divecha, N. (2013). PtdIns5P is an oxidative stress-induced second messenger that regulates PKB activation. *FASEB Journal : Official Publication of the Federation of American Societies for Experimental Biology*, 27(4): 1644–56.
- Jude J. G., Spencer G. J., Huang X., Somerville T. D. D., Jones D. R., Divecha N., & Somervaille T. C. P. (2015). A targeted knockdown screen of genes coding for phosphoinositide modulators identifies PIP4K2A as required for acute myeloid leukemia cell proliferation and survival. *Oncogene*, 34(10): 1253–62.
- Katagiri T., Yamaguchi A., Komaki M., Abe E., Takahashi N., Ikeda T., ... Suda T. (1994). Bone morphogenetic protein-2 converts the differentiation pathway of C2C12 myoblasts into the osteoblast lineage. *Journal of Cell Biology*, 127(6 1): 1755–1766.
- Kim C. G., Park D., & Rhee S. G. (1996). The Role of carboxyl-terminal basic amino acids in G q α -dependent activation, particulate association, and nuclear localization of phospholipase C- β 1. *The Journal of Biological Chemistry*, 271(35): 21187–21192.
- Kurian M. A., Meyer E., Vassallo G., Morgan N. V., Prakash N., Pasha S., ... Maher E. R. (2010). Phospholipase C β 1 deficiency is associated with early-onset epileptic encephalopathy. *Brain*, 133(10): 2964–2970.
- Lo Vasco V. R., Longo L., & Polonia P. (2013). Phosphoinositide-specific Phospholipase C β 1 gene deletion in bipolar disorder affected patient. *Journal of Cell Communication and Signaling*, 7(1): 25–29.
- Lo Vasco V., Cardinale G., & Polonia P. (2012). Deletion of PLCB1 gene in schizophrenia-affected patients. *Journal of Cellular and Molecular Medicine*, 16(4): 844–851.

- Lukinovic-Skudar, V., Donlagic, L., Banfic, H., & Visnjic, D. (2005). Nuclear phospholipase C-beta1b activation during G2/M and late G1 phase in nocodazole-synchronized HL-60 cells. *Biochimica et Biophysica Acta*, 1733(2-3): 148-56.
- Martelli AM, Fiume R, Faenza I, Tabellini G, Evangelista C, Bortul R, Follo MY, Falà F, Cocco L. (2005). Nuclear phosphoinositide specific phospholipase C (PI-PLC)-beta 1: a central intermediary in nuclear lipid-dependent signal transduction. *Histol Histopathol* 20(4):1251-60
- Martelli A. M., Gilmour R. S., Bertagnolo V., Neri L. M., Manzoli L., & Cocco L. (1992). Nuclear localization and signalling activity of phosphoinositidase C beta in Swiss 3T3 cells. *Nature*, 358(6383): 242-5.
- Martelli A. M., Neri L. M., Gilmour R. S., Barker P. J., Huskisson N. S., Manzoli F. A., & Cocco L. (1991). Temporal changes in intracellular distribution of protein kinase C in Swiss 3T3 cells during mitogenic stimulation with insulin-like growth factor I and bombesin: translocation to the nucleus follows rapid changes in nuclear polyphosphoinositides. *Biochemical and Biophysical Research Communications*, 177(1): 480-7.
- Montaña M., García del Caño G., López de Jesús M., González-Burguera I., Echeazarra L., Barrondo S., & Sallés J. (2012). Cellular neurochemical characterization and subcellular localization of phospholipase C β 1 in rat brain. *Neuroscience*, 222: 239-68.
- O'Carroll S. J., Mitchell M. D., Faenza I., Cocco L., & Gilmour R. S. (2009). Nuclear PLCbeta1 is required for 3T3-L1 adipocyte differentiation and regulates expression of the cyclin D3-cdk4 complex. *Cellular Signalling*, 21(6): 926-35. Retrieved from
- Osborne S. L., Thomas C. L., Gschmeissner S., & Schiavo G. (2001). Nuclear PtdIns(4,5)P2 assembles in a mitotically regulated particle involved in pre-mRNA splicing. *Journal of Cell Science*, 114(Pt 13): 2501-11.
- Payraastre, B., Nievers, M., Boonstra, J., Breton, M., Verkleij, A. J., & Van Bergen en Henegouwen, P. M. (1992). A differential location of phosphoinositide kinases, diacylglycerol kinase, and phospholipase C in the nuclear matrix. *The Journal of Biological Chemistry*, 267(8): 5078-84.
- Piazzì M., Blalock W. L., Bavelloni A., Faenza I., Raffini M., Tagliavini F., ... Cocco L. (2015). PI-PLC β 1b affects Akt activation, cyclin E expression, and caspase cleavage, promoting cell survival in pro-B-lymphoblastic cells exposed to oxidative stress. *FASEB Journal : Official Publication of the Federation of American Societies for Experimental Biology*, 29(4): 1383-94.
- Poli A., Billi A. M., Mongiorgi S., Ratti S., Mccubrey J. A., Suh P. G., ... Ramazzotti, G. (2016a). Nuclear Phosphatidylinositol Signaling: Focus on Phosphatidylinositol Phosphate Kinases and Phospholipases C. *Journal of Cellular Physiology*, 231(8): 1645-1655.
- Poli A., Fiume R., Baldanzi G., Capello D., Ratti S., Gesi M., ... Follo M. Y. (2016b). Nuclear Localization of Diacylglycerol Kinase Alpha in K562 Cells Is Involved in Cell Cycle Progression. *Journal of Cellular Physiology*, 10.1002/jcp.25642
- Poli A., Ramazzotti G., Matteucci A., Manzoli L., Lonetti A., Suh P.-G., ... Cocco L. (2014). A novel DAG-dependent mechanism links PKC α and Cyclin B1 regulating cell cycle progression. *Oncotarget*, 5(22): 11526-11540.
- Ramazzotti G., Faenza I., Fiume R., Billi A. M., Manzoli L., Mongiorgi S., ... Follo M. Y. (2016). PLC- β 1 and cell differentiation: An insight into myogenesis and osteogenesis. *Advances in Biological Regulation*, 5-9. 10.1016/j.jbior.2016.10.005
- Salisbury E., Sakai K., Schoser B., Huichalaf C., Schneider-Gold C., Nguyen H., ... Timchenko L. T. (2008). Ectopic expression of cyclin D3 corrects differentiation of DM1 myoblasts through activation of RNA CUG-binding protein, CUGBP1. *Experimental Cell Research*, 314(11-12): 2266-2278.
- Saunders C. M., Larman M. G., Parrington J., Cox L. J., Royse J., Blayney L. M., ... Lai F. A. (2002). PLC zeta: a sperm-specific trigger of Ca(2+) oscillations in eggs and embryo development. *Development (Cambridge, England)*, 129(15): 3533-44.
- Shah Z. H., Jones D. R., Sommer L., Foulger R., Bultsma Y., D'Santos C., & Divecha N. (2013). Nuclear phosphoinositides and their impact on nuclear functions. *FEBS Journal*, 280(24): 6295-6310.

Spires T. L., Molnár Z., Kind P. C., Cordery P. M., Upton A. L., Blakemore C., & Hannan A. J. (2005). Activity-dependent regulation of synapse and dendritic spine morphology in developing barrel cortex requires phospholipase C- β 1 signalling. *Cerebral Cortex*, 15(4): 385–393.

Sun B., Murray N. R., & Fields A. P. (1997). A role for nuclear phosphatidylinositol-specific phospholipase C in the G2/M phase transition. *J.Biol.Chem.*, 272(0021–9258):26313–26317.

Toker A., & Marmiroli S. (2014). Signaling specificity in the Akt pathway in biology and disease. *Advances in Biological Regulation*, 55: 28–38.

Tribble EK, Ivanova PT, Grabon A, Alb JG Jr, Faenza I, Cocco L, Brown HA, Bankaitis VA(2016).

Quantitative profiling of the endonuclear glycerophospholipidome of murine embryonicfibroblasts. *Journal of Lipid Research*, 57(8):1492-506

Vann L. R., Wooding F. B., Irvine R. F., & Divecha N. (1997). Metabolism and possible compartmentalization of inositol lipids in isolated rat-liver nuclei. *The Biochemical Journal*, 327 (Pt 2):569-76.

Xu A., Wang Y., Xu L. Y., & Gilmour R. S. (2001). Protein Kinase C α -mediated Negative Feedback Regulation Is Responsible for the Termination of Insulin-like Growth Factor I- induced Activation of Nuclear Phospholipase C β 1 in Swiss 3T3 Cells. *The Journal of Biological Chemistry*. 276(18):14980-6

Yang YR, Kang DS, Lee C, Seok H, Folio MY, Cocco L, Suh PG.(2016) Primary phospholipase C and brain disorders. *Adv Biol Regul*. 61:80-5

Ledends to figures:

Figure 1: in this figure we can observe the nuclear PI-PLC β 1 pathway related to the cell cycle. PI-PLC β 1 is essential for G1/S transition through DAG, Cyclin D3-Cdk4 complex and plays a central role in G2/M progression through Cyclin B1 and PKC α . Indeed DAG attracts Cyclin B1 and PKC α from the cytoplasm into the nucleus. We can also see that PIP5K is pivotal for PI(4,5)P2 production within the nucleus, while PIP4K was responsible for only a little amount.

Figure 2: PI-PLC β 1 level changes during cell differentiation. PI-PLC β 1 increases during the normal osteogenic and myogenic differentiation in C2C12 cells. PI-PLC β 1 also increases in adipogenic differentiation in 3T3-L1 cells. During erythroid differentiation there is a decrease in PI-PLC β 1 level demonstrated in MEL, CD34+ and MDS cells. PI-PLC β 1 seems to increase in MDS cells and also in MDS patients samples during the myeloid differentiation induced by Azacitidine.




Figure 1




Figure 2