



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Occurrence of *Salmonella enterica* subsp. *enterica* in bivalve molluscs and associations with *Escherichia coli* in molluscs and faecal coliforms in seawater

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Rubini, S., Galletti, G., D'Incau, M., Govoni, G., Boschetti, L., Berardelli, C., et al. (2018). Occurrence of *Salmonella enterica* subsp. *enterica* in bivalve molluscs and associations with *Escherichia coli* in molluscs and faecal coliforms in seawater. *FOOD CONTROL*, 84, 429-435 [10.1016/j.foodcont.2017.08.035].

Availability:

This version is available at: <https://hdl.handle.net/11585/614905> since: 2021-01-20

Published:

DOI: <http://doi.org/10.1016/j.foodcont.2017.08.035>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Silva Rubini, Giorgio Galletti, Mario D'Incau, Guido Govoni, Lari Boschetti, Chiara Berardelli, Stefania Barbieri, Giuseppe Merialdi, Andrea Formaglio, Enrica Guidi, Mauro Bergamini, Silvia Piva, Andrea Serraino, Federica Giacometti. *Occurrence of Salmonella enterica subsp. enterica in bivalve molluscs and associations with Escherichia coli in molluscs and faecal coliforms in seawater*. Food Control 84 (2018) 429e435.

The final published version is available online at:

<http://dx.doi.org/10.1016/j.foodcont.2017.08.035>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26

Occurrence of *Salmonella* spp. in bivalve molluscs and associations with *Escherichia coli* in molluscs and faecal coliforms in seawater

Silva Rubini^a, Giorgio Galletti^a, Mario D’Incau^a, Guido Govoni^b, Lari Boschetti^c, Chiara Berardelli^c, Stefania Barbieri^d, Giuseppe Meriardi^a, Andrea Formaglio^d, Enrica Guidi^d, Mauro Bergamini^d, Silvia Piva^e, Andrea Serraino^e, Federica Giacometti^{e*}

^aExperimental Institute for Zooprophyllaxis in Lombardy and Emilia Romagna, Via Bianchi 9, 25124 Brescia, Italy

^bInternational Committee of the Red Cross (ICRC), 19 Avenue de la Paix 1202, Geneva, Switzerland

^cPublic Health Unit of Ferrara, U.O.D. Igiene degli alimenti di Origine Animale, Via Cassoli 30, 44121 Ferrara, Italy

^dDepartment of Medical Sciences, Branch of Public Health Medicine, University of Ferrara, Via Fossato di Mortara 64/b, 44121 Ferrara, Italy

^eDepartment of Veterinary Medical Sciences, University of Bologna, Via Tolara di Sopra 50, 40064 Ozzano dell’Emilia, Bologna, Italy

(*) **Corresponding author:** Federica Giacometti, Tel.: +39 51 2097320; fax: +39 51 2097346. E-mail address: federica.giacometti3@unibo.it (F. Giacometti).

Keywords: bivalve molluscs, seawaters, *Salmonella* spp., *Escherichia coli*, faecal coliforms

ABSTRACT

The objectives of this study were to present data on the presence of *Salmonella* spp. and on the enumeration of *Escherichia coli* and faecal coliforms respectively in different species of bivalve

27 molluscs and seawater and to conduct a retrospective evaluation to assess the capacity of *E. coli* in
28 molluscs and faecal coliforms and *Salmonella* spp. in sea and brackish water to predict the presence
29 of *Salmonella* spp. in bivalve molluscs, and therefore, the risk of exposure for consumers. Data were
30 collected from 4972 seawater samples and 5785 live bivalve molluscs samples (2877 *Ruditapes*
31 *philippinarum*, 2177 *Mytilus galloprovincialis*, 256 *Chamela gallina* and 475 *C. gigas* and *O. edulis*)
32 collected in the molluscs production area of Ferrara, Northern Italy, from 1997 to 2015. An overall
33 *Salmonella* spp. occurrence of 2.2% was reported in water and molluscs, with percentages varying
34 depending on the type of sample and on the classification areas. All the 237 *Salmonella* strains were
35 identified as *Salmonella enterica* and a total of 53 different serovars were observed. Significant
36 associations between the fecal indicators and presence of *Salmonella* spp. were observed both
37 applying EU and USA criteria, but, it should be noted that the EU approach seems to be more stringent
38 achieving the goal of identifying the most critical batches (94 out of the 100) whereas, following the
39 USA approach, a not negligible and higher number of batches compliant for faecal coliforms but
40 contaminated by *Salmonella* spp. has to be mentioned. In any case, the faecal indicators *E. coli* in
41 molluscs and faecal coliforms in seawaters reflect only in part the presence of *Salmonella* spp. in
42 molluscs and the consequent potential risk for consumers. Microbiological evaluation of seawaters
43 seems to have a minor impact into the prediction of *Salmonella* spp. presence in molluscs.

44

45 **1. Introduction**

46 In Italy bivalve molluscs such as Manila clams (*Ruditapes philippinarum*) and mussels (*Mytilus*
47 *galloprovincialis*) represent products of great economic importance whereas striped clams (*Chamela*
48 *gallina*) and oysters (*Crassostrea gigas* and *Ostrea edulis*) are of less economic importance. Overall
49 they are widely distributed in the food trade and, specifically for *R. philippinarum*, Italy is the second
50 producer after China (Turolla, 2008).

51 In relation to the sanitary control of shellfish produced and sold for human consumption, two main
52 **control systems** are in place in the European Community (EC) and United States of America (USA):

53 these are the principal systems used worldwide and countries that trade with EU and/or USA will use
54 either one, or a hybrid of the two systems (Gosling, 2015). In Europe, according to Regulation
55 853/2004, areas in which bivalve molluscs are cultivated in marine or brackish water must be
56 previously classified by veterinary authorities as production area A, B or C depending on the content
57 of *Escherichia coli* in the soft parts and flesh and intravalvular liquid of harvested bivalves, with an
58 upper limit of respectively 230, 4600 and 46000 MPN *E. coli*/100 g sample material is for class A, B
59 and C areas. Bivalve molluscs from class A area can be placed directly on the market provided that
60 they comply with microbiological criteria requirements, whereas molluscs from class B area must be
61 purified by resuspension at class A area, or heat-treated before distribution, and, finally molluscs
62 from class C area, a resuspension at class A area over a long period of time or a sufficient heat
63 treatment is needed.

64 In relation to food safety criteria laid down in EC Regulation 2073/2005 concerning bivalve to be
65 placed on the market, the absence of *Salmonella* spp. in 25 g of flesh and an upper limit of 230 MPN
66 *E. coli*/100 g sample material are mandatory; according to EC Regulation 2285/2015, from 1st January
67 2017, 20% of the samples may contain *E. coli* between 230 and 700 MPN/100 g sample material
68 while the remaining 80% of the samples must be below 230 MPN/100 g sample material. In the USA,
69 the official controls system is based on the National Shellfish Sanitation Program (NSSP), that is a
70 federal/state cooperative programme recognized by Food and Drug Administration (FDA) and the
71 Interstate Shellfish Conference that is substantially different from the EU one; for example, the
72 microbiological monitoring is based on water testing and considers either total coliforms or faecal
73 coliforms, not *E. coli*, and to achieve 'approved' status the geometric mean faecal coliforms count
74 per water sample from an area must not exceed 14 Most probable Number (MPN)/100 ml, and the
75 ninetieth percentile must not exceed 43 MPN/100 ml (NSSP, 2017).

76 It is well known that live bivalve molluscs, being suspension feeders that gain nourishment by
77 pumping large volumes of water from the environment through their gills, actively filter and retain
78 particles from their surrounding water, including free living or particle bound bacteria, viruses and

79 parasites. This highlights the role of bivalve molluscs as vehicle for several hazards that could result
80 in potential health risks for consumers, particularly if live bivalve molluscs are eaten raw or lightly
81 cooked. With specific regard to bacteria, the microbiota found in bivalve molluscs include indigenous
82 bacteria that naturally occur in marine or estuarine environments (mostly *Vibrio* spp.), non-
83 indigenous bacteria, usually enteric bacteria, derived from faecal contamination (mostly *Salmonella*
84 spp., *E. coli*, *Shigella* spp. and rarely *Campylobacter* spp. and *Yersinia enterocolitica*), and bacteria
85 from contamination during food preparation and processing by the distribution industry or consumers
86 (*Bacillus cereus*, *Staphylococcus aureus* and *Clostridium perfringens*) (Anacleto, Pedro, Leonor, Rosa,
87 & Marques, 2013).

88 *Salmonella* is the second most common cause of human gastroenteritis (EFSA & ECDC, 2016): the
89 risks of foodborne illness associated with *Salmonella* in molluscs are low compared to viruses and
90 *Vibrio* spp. (Iwamoto, Ayers, Mahon, & Swerdlow, 2010; NACMCF, 1992) but there is a
91 considerable amount of epidemiological data regarding the presence of *Salmonella* in seafood and
92 related illness. The EU summary report on zoonoses, zoonotic agents and food-borne outbreaks in
93 2015 reports that, for the causative agent of foodborne outbreaks by food vehicle, in “fish, shellfish,
94 molluscs, crustaceans and products thereof”, histamine was the leading cause of strong-evidence
95 outbreaks (52.5%) followed by calicivirus including Norwalk-like virus (norovirus) (25%) and
96 *Salmonella* (12.5%) (EFSA & ECDC, 2016).

97 Even if *Salmonella* spp. are not natural inhabitants of the aquatic environment, several *Salmonella*
98 serovars are widely distributed in water (sea, estuarine, river) and in a variety of seafood, with the
99 highest prevalence in molluscs, shrimp, clams, and various fish species (Novoslavskij et al., 2016).

100 *Salmonella* spp. and, in general, faecally derived enteric pathogens, are introduced into the aquatic
101 environment via anthropogenic activities like inappropriate disposal of human wastes, agricultural
102 runoffs or sewage discharges (Malham et al., 2014) as well as wildlife (Obiri-Danso & Jones, 2000),
103 and, given the reported ability to survive long periods in different aquatic environments, these
104 microorganisms could pass into new hosts (Amagliani, Brandi, & Schiavano, 2012). Yet the use of

105 faecal bacterial indicators in bivalve monitoring was based on their relation to the probable
106 occurrence of human faecal pathogens: it has been well remarked that no correlation was between
107 faecal microorganisms and both *Vibrio* spp. and viruses for humans, but, the true evidence of this
108 correlation in relation to *Salmonella* spp. is still a matter of debate.

109 The current publication presents data on the presence of *Salmonella* spp. and on the enumeration of
110 *E. coli* in different species of bivalve molluscs, namely Manila clams (*R. philippinarum*), mussels
111 (*M. galloprovincialis*), striped clams (*C. gallina*) and oysters (*C. gigas* and *O. edulis*), and on faecal
112 coliform levels in water samples from seashore and inland channels of the Ferrara province, Emilia-
113 Romagna Region, Northern Italy with a continuous sampling history from 1997 to 2015. In addition,
114 a retrospective evaluation was conducted to assess: i) the capacity of the faecal indicator *E. coli* to
115 predict the presence of *Salmonella* spp. in bivalve molluscs, and therefore, the risk of exposure for
116 consumers; ii) the efficacy of sea and brackish water analysis for faecal coliforms and *Salmonella*
117 spp. to predict the presence of *Salmonella* spp. in live bivalve molluscs iii) the comparison of the
118 regulations currently in force in USA versus Europe.

119

120 **2. Material and methods**

121 *2.1. Sample and data collection*

122 Microbiological records for *E. coli*, faecal coliforms and *Salmonella* spp. analyses of live bivalve
123 molluscs and water were collated from official monitoring performed by the regional Veterinary
124 Authority from 1997 to 2015 and from a shellfish monitoring program that has been carried out since
125 1997 in the province of Ferrara, Emilia Romagna region, Italy. A total of 10757 samples were
126 collected, respectively 4972 seawater samples (of which 1237 in class A area and 3735 in class B
127 area) and 5785 live bivalve molluscs samples; the examined bivalve molluscs comprise 2877 Manila
128 clams (*R. philippinarum*, all in class B area), 2177 mussels (*M. galloprovincialis*, 969 and 1208 in
129 class A and B areas), 256 striped clams (*C. gallina*, all in class A area) and 475 oysters (*C. gigas* and
130 *O. edulis*, respectively 62 and 413 in class A and B areas). A total of 4815 paired samples of bivalve

131 mollusc and seawater were collected. Each mollusc sample comprised at least 10 live individuals
132 (commercial size or adult product) that were analyzed for *Salmonella* spp. detection and *E. coli*
133 enumeration. Seawater samples were collected in from 0.2 m below the water surface using 1.2 L
134 polypropylene or glass bottle for *Salmonella* spp. detection and for faecal coliforms count. All
135 samples were transported under chilled conditions to the laboratory and processed within 24 h after
136 collection.

137

138 2.2. Sampling site

139 All the samplings were performed in the north-western area of the Adriatic Sea facing the Po river
140 delta, the major Italian river which, from spring to estuary, flows through the Po Valley (Pianura
141 Padana) for a total of 652 Km. The Po Valley is a high populated area with abundant large animals
142 intensive farms. The Po River, near to its end in the Adriatic sea, creates a wide delta with a surface
143 area of 31 Km² and an average depth of 1.5 m; its hydrographic network is for the most part artificially
144 regulated, and, as a consequence, freshwater flows are partially independent of rain events. More than
145 one third of the lagoon surface is exploited for clam farming, with an annual production that reached
146 a maximum of 87000 t y⁻¹ in 2011 (Bison, 2012).

147 The areas of molluscs production have been divided into 5 sub-areas (see Figure 1): i) long-line: the
148 marine class A area used to breed mostly mussels and secondary oysters; ii) Lupini: the coastal marine
149 area including seawaters between 1 and 2 nautical miles that is classified as class A area in which
150 natural banks of striped clams are present and harvested; iii) B-Out: the narrow sea coastal area and
151 inland waters classified as class B area; iv) B-In: class B area that includes the inner channels directly
152 connected to the sea, together with internal waters; v) Sacca: the class B area included between the
153 Po river and the marine coastline. All these last three sub-areas were used to breed mussels, Manila
154 clams and oysters.

155

156 2.3. Microbiological analyses

157 All samples were analyzed at the Experimental Institute for Zooprohylaxis of Lombardy and Emilia
158 Romagna in Ferrara. For live bivalve molluscs, the isolation of *Salmonella* spp. was performed using
159 the official International Organization for Standardization (ISO) cultural methods, UNI EN ISO
160 6579:1993 still 2002, the ISO 6579:2002 since 2003 with its technical corrigendum and amendments,
161 whereas, *E. coli* was enumerated using a 5 tube 3 dilutions MPN procedure based on ISO 16649-3.
162 For water analysis, the isolation of *Salmonella* spp. was carried out filtering a sample volume of 1000
163 mL through 0.45 µm-pore size membrane filters according to APAT CNR IRSA 7080 procedure
164 (APAT, 2003); the enumeration of faecal coliform and *E. coli* from 1998 to 2014 were performed by
165 a five tube three dilutions MPN methods according respectively to APAT CNR IRSA 7020 and 7030
166 procedures (APAT, 2003), whereas from August 2014, a membrane filtering method was applied
167 (APAT, 2003).

168 Cultures displaying a reaction typical of *Salmonella* (an alkaline slant and acid butt, with or without
169 production of H₂S) were confirmed by biochemical tests using miniaturized galleries, e.g. API-20E
170 strip (bioMérieux, Marcy l'Etoile, France) or MID (Microgen Bioproduct Ltd, Camberley, United
171 Kingdom) (Anacleto et al. 2013; APAT, 2003). The serotyping of *Salmonella* strains was performed
172 using commercial antisera (BBL Becton Dickinson, Franklin Lakes, USA; Statens Serum Institut,
173 Copenhagen, Denmark); following the White-Kauffman-Le Minor serotyping scheme (Baudart,
174 Lemarchand, & Brisabois, 2000).

175

176 2.4. Statistical analysis

177 Fisher's exact test were used to study the association between the occurrence of *Salmonella* spp. in
178 molluscs and: i) *E. coli* levels in molluscs; ii) faecal coliform levels in paired seawater; iii) presence
179 of *Salmonella* spp. in paired seawater.

180 To perform the statistical analysis the *E. coli* counts in molluscs were divided based on the *E. coli*
181 contamination level laid down into EC Regulation 2073/2005 as food safety criteria applicable for
182 products from class A areas, namely collected for direct human consumption and placed on the

183 market, irrespective of the fact that samples were collected from class A or B areas. Therefore, two
184 groups were considered using 230 MPN/100g as cutoff. Similarly results of fecal coliforms count in
185 water were divided in three categories identified according to the classification approach used in USA
186 with standards set for categories ranging from waters with low contamination levels, ≤ 14 , between
187 14 and ≤ 43 and > 43 MPN/100 ml water. Data on faecal coliforms was available since 1999.

188

189 3. Results

190 A total of 237 out of 10757 samples (2.2%) were positive for *Salmonella* spp., respectively 137
191 seawater samples (2.8%) and 100 bivalve molluscs (1.7%): the presence of *Salmonella* spp. ranged
192 from 0 to 3.4% depending on the type of sample and on the classification areas (Table 1).

193 The overall *Salmonella* spp. prevalence differs considerably through the years and between water
194 class areas and species of molluscs: water prevalences were of 0 to 4.2% and of 0 to 13.8%
195 respectively for class A and B areas; in molluscs, *R. philippinarum* (collected only in class B area)
196 reported higher prevalences of 0 to 8.2% followed by *M. galloprovincialis* (0 to 4.3% and of 0 to
197 2.4% respectively for class A and B areas), *C. gallina* (collected only in class A area) with prevalences
198 of 0 to 4.3% and lastly *C. gigas* and *O. edulis* with prevalences of 0 to 2.1%.

199 All the 237 *Salmonella* strains were identified as *Salmonella enterica* and a total of 53 different
200 serovars were observed (Figure 2), respectively 43 and 32 serovars from the 137 and 100 seawater
201 and molluscs positive samples. *Salmonella* ser. Typhimurium resulted the dominant serotype
202 (26.9%), followed by its monophasic variant 4,[5],12:i:- (11.8%), *Salmonella* Derby (6.3%),
203 *Salmonella* Newport (5.5%), and *Salmonella* Thompson (4.6%).

204 A significant association was observed between *E. coli* levels in molluscs and presence of *Salmonella*
205 spp. in molluscs samples, both from class A and B areas. In more details, an association was observed
206 in *R. philippinarum* from class B area (no *R. philippinarum* were on class A area) and in *M.*
207 *galloprovincialis* collected only from class A area, not for class B area (table 2).

208 A significant association was observed between faecal coliforms levels in seawater and presence of
209 *Salmonella* spp. both in total molluscs and *R. philippinarum* samples (table 3).

210 A significant association between presence of *Salmonella* spp. in molluscs and paired seawater was
211 observed only for *R. philippinarum* samples (see table 4), where a co-presence of *Salmonella* spp. in
212 13 samples was observed. In 5 samples the same serovar was isolated in the two different matrices,
213 respectively *S. enterica* serovars Typhimurium (3 samples), its monophasic variant 4,[5],12:i:- (1
214 sample) and *Salmonella* Thompson (1 sample).

215

216 **4. Discussion**

217 Only safe food should be placed on the market (Regulation EC 178/2002). With regards to bivalve
218 molluscs, the climate change, the pollution and the anthropogenic factors as well as the food
219 distributions and consumers behaviour changes (undercooked or raw seafood consumption, mainly
220 bivalve molluscs) have created a cunning environment in which each element may have a potential
221 impact on food safety. In this context, it appears evident that the aquatic environment and its quality
222 and/or safety is a critical point for live bivalve molluscs that, being excellent bio-samplers, reflect the
223 quality of the surrounding water in a given location. This study, reporting data on the presence of
224 *Salmonella* spp. and on the enumeration of *E. coli* in different species of bivalve molluscs and of fecal
225 coliform in water samples, with a continuous sampling history over 19-year-period (1997 through
226 2015), is useful to present the real scenario in which bivalve molluscs are bred in the molluscs
227 production area of Ferrara, leading us to perform some considerations.

228 An overall occurrence of *Salmonella* spp. of 2.2% was reported in water and molluscs collected in
229 the considered production area: this rate could be considered low, but, at the same time poses a not
230 negligible threat for the assessment of the risks of faecal pollutions in the aquatic environment. The
231 significance of *Salmonella* spp. as human pathogen and as a leading cause of food-borne illness is
232 well known, but, the observed occurrence suggests to reconsider the role and the ecology of this
233 allochthonous pathogen at the base of the marine and estuarine environment and ecosystem; in fact,

234 although allochthonous organisms are historically viewed as transient member of the microbial
235 community of coastal waters, the coastal environment may serve as an important niche for these
236 microorganisms to persist, exchange genetic material, and grow (Bienfang et al., 2011).

237 The observed *Salmonella* spp. occurrence in seawater and bivalve molluscs is in agreement with
238 available data in literature, in which the prevalence rates considerably vary depending on climate
239 conditions, the season and area of samplings, the type of considered seafood and similar or different
240 environmental conditions (estuarine or seawater, after rainfall, low solar radiation) (Amagliani et al.,
241 2012; Novoslavskij et al., 2016); in Italy, *Salmonella* spp. prevalence rates from 0 to 3.1% were
242 reported (Normanno, Parisi, Addante, Quaglia, & Dambrosio, 2006; Mazzette, Virgilio, Piras,
243 Tempesta & Serra, 2010; Serracca et al., 2010; Prato et al., 2013; Fusco, Aprea, Galiero, Guarino, &
244 Viscardi, 2011; Carraro et al., 2015; EFSA & ECDC, 2016). In our 19-years period of monitoring in
245 live bivalve molluscs, from class A areas (products that can be placed directly on the market) only 6
246 out of the 1287 (0.5%) samples collected resulted *Salmonella* spp. positive, showing a good insight
247 the official monitoring usefulness. In the other hand, a total of 94 samples resulted *Salmonella* spp.
248 positive among the 4498 (2%) molluscs collected from class B areas (molluscs that may be placed on
249 the market only after treatment in a purification centre or after relaying); no food safety considerations
250 could be gathered from these data, but, again, the EU classification areas approach achieves the goal
251 of identifying the most critical batches (94 out of the 100).

252 Lastly, it should be noted that 87 out of total 100 *Salmonella* spp. positive molluscs were *R.*
253 *philippinarum* and, therefore, an higher *Salmonella* spp. prevalence in Manila clams than the other
254 bivalve molluscs species considered in this study has to be reported, according with the studies of
255 Anacleto et al. (2013), that reported an occurrence of 17% and 25%, respectively for *V. pullastra* and
256 *R. philippinarum*, and of Carraro et al. (2015) that observed the presence of *Salmonella* spp. in only
257 one sample of Manila clam among the 540 samples analyzed.

258 In literature, other studies investigated the correlation between the presence of *Salmonella* spp. and
259 the bacterial indicator of faecally pollution *E. coli* in bivalve molluscs, even if with contrasting results,

260 but, as far as we known, no other studies were on the comparison of bacterial indicators of faecal
261 pollution *E. coli* in molluscs and faecal coliforms in water (representative of EU and USA monitoring
262 approaches) in relation to the power of prediction of the presence of *Salmonella* spp. in bivalve
263 molluscs.

264 Literature reports that bacterial indicators of faecal pollution in water provide an adequate indicator
265 of *Salmonella* spp. presence in waters (Efstratiou, Mavridou, & Richardson, 2009; Ferguson, Coote,
266 Ashbolt, & Stevenson, 1996; Morinigo, Castro, Balebona, Munoz, & Borrego, 1992) and in bivalve
267 molluscs, particularly in *R. philippinarum*, or in other bivalve species like oysters (Anacleto et al.,
268 2013), but, on several occasions or no *Salmonella* spp. was detected in samples with high indicator
269 counts or *Salmonella* spp. was detected in samples with low indicator counts (Dionisio, Joao, Ferreira,
270 Fidalgo, & Garc, 2000; Efstratiou, Mavridou, & Richardson, 2009; Hood, Ness, & Blake, 1983;
271 Mannas, Mimouni, Chaouqy, Hamadi, & Martinez-urtaza, 2014; Morigligo, Cornax, Oz, Romero, &
272 Borrego, 1990). In this context, despite our monitoring was not drawn with the specific aim of testing
273 these associations, and therefore, it could be biased by several factors, our study results is a useful
274 and robust tool for assessing the *E. coli*, faecal coliform and *Salmonella* spp. parameters all together
275 in a long period of time and in a specific area and considering both water versus molluscs and mollusc
276 versus mollusc and some considerations could be performed from our findings.

277 A significant associations between fecal indicators and presence of *Salmonella* spp. was observed
278 both applying EU and USA criteria prevalently for *R. philippinarum*, whereas, the weight carried for
279 the other observed associations resulted of less importance or partially affected by *R. philippinarum*
280 results. Considering that the *R. philippinarum* data we analyzed belongs exclusively from class B
281 area, it seems evident that or this fact may has biased our findings or *R. philippinarum* really has a
282 different behavior with respect with other bivalve species.

283 It should be noted that, following EU regulations and its official monitoring system, only two batches
284 (0.2%) with *E. coli* levels ≤ 230 MPN/100 g in molluscs from class A area showed the presence of
285 *Salmonella* spp. in molluscs, whereas, following the NSSP, a total of 58 batches (1.6%) with faecal

286 coliforms ≤ 14 MPN/100 ml in seawaters, were contaminated by *Salmonella* spp. Shellfish from
287 “approved” waters, namely waters in which the MPN faecal coliforms median does not exceed the
288 considered value of ≤ 14 MPN/100 ml water, can be sold directly on the market without prior
289 treatment (Gosling, 2015). In this specific case, a not negligible and higher number of batches
290 compliant for the faecal indicators but contaminated by *Salmonella* spp. has to be mentioned. This
291 scenario is not representative of the current situation for two reasons: i) the sampling plan was not
292 designed to assess the microbiological quality of waters according to NSSP; ii) the statistical
293 evaluation could not be performed calculating the geometric mean of fecal coliforms enumeration
294 but only singular values were considered. On the other hand calculating the gemetric median of all
295 data collected in this study on fecal coliform contamination of water a level of 12 MPN/100 ml water
296 was obtained, finding that could be considered compliant with NSSP. Despite our attempts, in one
297 hand, this estimation should be considered anyway a forced parameter which remains an important
298 data gap having some implications in the final outputs of this association, but in the other, it should
299 be noted that, in general, concentrations of faecal coliform bacteria under steady-state conditions are
300 approximately 10 to 100 times higher in bivalves than the surrounding seawater (Strubbia, Lyons, &
301 Lee, 2016; Bernard, 1989). This demonstrates the fact that the occurrence of faecal bacteria in waters
302 is highly variable and depending on several factors including pollution sources as well as
303 environmental conditions that are not easily identifiable and testable. This observation should be
304 taken into account in relation to the different approaches used for faecal indicators, that in the EU are
305 measured in shellfish flesh while in the USA in shellfish-growing waters.

306

307 **5. Conclusion**

308 This study has reported the presence of *Salmonella* spp. in seawaters and in bivalve molluscs collected
309 in the Ferrara area from 1997 to 2015 and has demonstrated that its presence varied by bivalve species
310 considered, classification areas in which molluscs were collected and sampling occasion. The faecal
311 indicators *E. coli* in molluscs and faecal coliforms in seawaters reflect only in part the presence of

312 *Salmonella* spp. in molluscs and the consequent potential risk for consumers. Microbiological
313 evaluation of seawaters seems to have a minor impact into the prediction of *Salmonella* spp. presence
314 in molluscs.

315

316 **References**

317 178/2002/EC, R. (2002). Regulation (EC) No. 178/2002 of the European Parliament and of the
318 Council laying down the general principles and requirements of food law, establishing the
319 European Food Safety Authority and laying down procedures in matters of food safety. In E.
320 Union (Ed.), 178/2002. Brussels, Belgium: The European parliament and the council of the
321 European Union.

322 853/2004/EC, R. (2004). Regulation (EC) No. 853/2004 of the European Parliament and of the
323 Council laying down specific hygiene rules for food of animal origin. In E. Union (Ed.),
324 853/2004. Brussels, Belgium: The European parliament and the council of the European Union.

325 2073/2005/EC, R. (2005). Commission Regulation (EC) No. 2073/2005 on microbiological criteria
326 for foodstuffs. In E. Union (Ed.), 2073/2005. Brussels, Belgium: The European parliament and
327 the council of the European Union.

328 2285/2015/ EC, Commission Regulation No. 2285/2015 amending Annex II to Regulation (EC) No
329 854/2004 of the European Parliament and of the Council laying down specific rules for the
330 organisation of official controls on products of animal origin intended for human consumption
331 as regards certain requirements for live bivalve molluscs, echinoderms, tunicates and marine
332 gastropods and Annex I to Regulation (EC) No 2073/2005 on microbiological criteria for
333 foodstuffs. In E. Union (Ed.), 2285/2015. Brussels, Belgium: The European parliament and the
334 council of the European Union.

335 Amagliani, G., Brandi, G., & Schiavano, G. F. (2012). Incidence and role of *Salmonella* in seafood
336 safety. *Food Research International*, 45, 780-8.

337 Anacleto, P., Pedro, S., Nunes, M. L., Rosa, R., & Marques, A. (2013). Microbiological composition
338 of native and exotic clams from Tagus estuary: Effect of season and environmental parameters.
339 *Marine pollution bulletin*, 74(1), 116-124.

340 APAT CNR IRSA. 2003. APAT Manuali e Linee Guida 29/2003. Metodi analitici per le acque-
341 Volume 3, 7000 - Metodi per la determinazione di microorganismi indicatori di
342 inquinamento e di patogeni - APAT CNR IRSA 7020 Coliformi fecali, pag. 875.
343 <http://www.irsa.cnr.it/Docs/Capitoli/1000.pdf>

344 APAT CNR IRSA. 2003. APAT Manuali e Linee Guida 29/2003. Metodi analitici per le acque-
345 Volume 3, 7000 - Metodi per la determinazione di microorganismi indicatori di
346 inquinamento e di patogeni - APAT CNR IRSA 7030 *Escherichia coli.*, pag. 883.
347 <http://www.irsa.cnr.it/Docs/Capitoli/1000.pdf>

348 APAT CNR IRSA. 2003. APAT Manuali e Linee Guida 29/2003. Metodi analitici per le acque-
349 Volume 3, 7000 - Metodi per la determinazione di microorganismi indicatori di
350 inquinamento e di patogeni - APAT CNR IRSA 7080 *Salmonella* spp., pag. 927.
351 <http://www.irsa.cnr.it/Docs/Capitoli/1000.pdf>

352 Baudart, J., Lemarchand, K., Brisabois, A., & Lebaron, P. (2000). Diversity of *Salmonella* strains
353 isolated from the aquatic environment as determined by serotyping and amplification of the
354 ribosomal DNA spacer regions. *Applied and Environmental Microbiology*, 66(4), 1544-1552.

355 Bernard, F. R. (1989). Uptake and elimination of coliform bacteria by four marine bivalve molluscs.
356 *Canadian Journal of Fisheries and Aquatic Science*, 46, 1592-1599.

357 Bienfang, P. K., DeFelice, S. V., Laws, E. A., Brand, L. E., Bidigare, R. R., Christensen, S., et al.
358 (2010). Prominent human health impacts from several marine microbes: history, ecology, and
359 public health implications. *International Journal of Microbiology*, 2011.

360 Bison, G.O. (2012). Gli obiettivi per il futuro: fidelizzare il consumatore al brand “Vongola di Goro”
361 e commercializzare confezioni sottovuoto con sugo pronto a fianco. *Il Pesce* 2 (Mar/Apr):52
362 <http://www.pubblicitaitalia.com/ilpesce/2012/2/11664.html>

363 Carraro, V., Sanna, C., Brandas, V., Sanna, A., Pinna, A., & Coroneo, V. (2015). Hygiene and health
364 risks associated with the consumption of edible lamellibranch molluscs. *International Journal*
365 *of Food Microbiology*, 201, 52-57.

366 Catalao Dionisio, L. P., Joao, M., Ferreiro, V. S., Fidalgo, M. L., Garcia Rosado, M. E., & Borrego,
367 J.J. (2000). Occurrence of *Salmonella* spp. in estuarine and coastal waters of Portugal. *Antonie*
368 *Van Leeuwenhoek*, 78, 99-106.

369 Efstratiou, M. A., Mavridou, A., & Richardson, C. (2009). Prediction of *Salmonella* in seawater by
370 total and faecal coliforms and Enterococci. *Marine Pollution Bulletin*, 58(2), 201-205.

371 EFSA & ECDC. (2016). The European Union summary report on trends and sources of zoonoses,
372 zoonotic agents and food-borne outbreaks in 2015. *EFSA Journal*, 14(12), 4634. European
373 Food Safety Authority and European Centre for Disease Prevention and Control (EFSA and
374 ECDC).

375 Ferguson, C. M., Coote, B. G., Ashbolt, N. J., & Stevenson, I. M. (1996). Relationships between
376 indicators, pathogens and water quality in an estuarine system. *Water Research*, 30, 2045-2054.

377 Fusco, G., Aprea, G., Galiero, G., Guarino, A., & Viscardi, M. (2013). *Escherichia coli*, *Salmonella*
378 spp., Hepatitis A Virus and Norovirus in bivalve molluscs in Southern Italy. *Veterinaria*
379 *italiana*, 49(1), 55-58.

380 Geldreich, E. (1970). Applying bacteriological parameters to recreational water quality. *Journal of*
381 *the American Water Works Association*, 62, 113-120.

382 Gosling, E. (2015). *Marine bivalve molluscs*. (2nd ed.). Chichester, UK: Wiley- Blackwell.

383 Hood, M. A., Ness G. E., & Blake NJ. (1983). Relationship among fecal coliforms, *Escherichia coli*,
384 and *Salmonella* spp. in shellfish. *Applied and Environmental Microbiology*, 45(1), 122-126.

385 ISO. (1993). ISO 6579:1993. Microbiology - General guidance on methods for the detection of
386 *Salmonella*. Geneva, Switzerland: International Organization for Standardization (ISO).

387 ISO. (2002). ISO ISO 6579:2002. Microbiology of food and animal feeding stuffs - Horizontal
388 method for the detection of *Salmonella* spp. Geneva, Switzerland: International Organization
389 for Standardization (ISO).

390 ISO. (2005). ISO/TS 16649-3:2005. Microbiology of food and animal feeding stuffs - Horizontal
391 method for the enumeration of beta-glucuronidase-positive *Escherichia coli* - Part 3: Most
392 probable number technique using 5-bromo-4-chloro-3-indolyl-beta-D-glucuronide. Geneva,
393 Switzerland: International Organization for Standardization (ISO).

394 Iwamoto, M., Ayers, T., Mahon, B. E., & Swerdlow D. L. (2010). Epidemiology of seafood-
395 associated infections in the United States. *Clinical Microbiological Review*, 23, 399-410.

396 Malham, S. K., Rajko-Nenow, P., Howlett, E., Tuson, K. E., Perkins, T. L., Pallett, D. W., et al.
397 (2014). The interaction of human microbial pathogens, particulate material and nutrients in
398 estuarine environments and their impacts on recreational and shellfish waters. *Environmental*
399 *Science: Processes & Impacts*, 16 (9), 2145-2155.

400 Mannas, H., Mimouni, R., Chaouqy, N., Hamadi, F., & Martinez-Urtaza, J. (2014). Occurrence of
401 *Vibrio* and *Salmonella* species in mussels (*Mytilus galloprovincialis*) collected along the
402 Moroccan Atlantic coast. *SpringerPlus*, 3(1), 265.

403 Martinez-Manzanares, E., Moriñigo, M. A., Castro, D., Balebona, M. C., Muñoz, M. A. & Borrego
404 J. J. (1992). Relationship between indicators of fecal pollution in shellfish-growing water and
405 the occurrence of human pathogenic microorganisms in shellfish. *Journal of Food Protection*,
406 55(8), 609-614.

407 Mazzette, R., Virgilio, S., Piras, A., Tempesta, A., Serra, S., Sferlazzo, G., et al. (2010). Valutazione
408 dell'efficacia dei sistemi di depurazione nei confronti di *E. coli*, *Salmonella* spp. e *Vibrio* spp.
409 in mitili allevati nel Golfo di Olbia. *Il Pesce*, 5, 165-172.

410 Moriñigo, M. A., Còrnax, R., Munòz, M. A., Romero, P., & Borrego, J. J. (1990). Relationships
411 between *Salmonella* spp. and indicator microorganisms in polluted natural waters. *Water*
412 *Research*, 24(1), 117e120.

413 Moriñigo, M. A., Martínez-Manzanares, E., Muñoz, M. A., Balebona, M. C., & Borrego, J. J. (1993).
414 Reliability of several microorganisms to indicate the presence of *Salmonella* in natural waters.
415 *Water Science and Technology*, 27(3-4), 471-474.

416 NACMCF. (1992). Microbiological criteria for raw molluscan shellfish. *Journal of Food Protection*,
417 55, 463–480. National Advisory Committee on Microbiological Criteria for Foods.

418 Normanno, G., Parisi, A., Addante, N., Quaglia, N. C., Dambrosio, A., Montagna, C., & Chiocco, D.
419 (2006). *Vibrio parahaemolyticus*, *Vibrio vulnificus* and microorganisms of fecal origin in
420 mussels (*Mytilus galloprovincialis*) sold in the Puglia region (Italy). *International Journal of*
421 *Food Microbiology*, 106(2), 219-222.

422 Novoslavskij, A., Terentjeva, M., Eizenberga, I., Valciņa, O., Bartkevičs, V. & Bērziņš, A. (2016).
423 Major foodborne pathogens in fish and fish products: a review. *Annals of Microbiology*, 66(1),
424 1-15.

425 NSSP. (2017). Guide for the Control of Molluscan Shellfish: 2015 Revision (Updated February
426 2017).
427 [https://www.fda.gov/downloads/Food/GuidanceRegulation/FederalStateFoodPrograms/UCM](https://www.fda.gov/downloads/Food/GuidanceRegulation/FederalStateFoodPrograms/UCM505093.pdf)
428 [505093.pdf](https://www.fda.gov/downloads/Food/GuidanceRegulation/FederalStateFoodPrograms/UCM505093.pdf)

429 Obiri-Danso, K., & Jones, K. (2000). Intertidal sediments as reservoirs for hippurate negative
430 campylobacters, salmonellae and faecal indicators in three EU recognised bathing waters in
431 North West England. *Water Research*, 34, 519-527.

432 Prato, R., Martinelli, D., Tafuri, S., Barbuti, G., Quarto, M., Germinario, C. A., & Chironna, M.
433 (2013). Safety of shellfish and epidemiological pattern of enterically transmitted diseases in
434 Italy. *International Journal of Food Microbiology*, 162(2), 125-128.

435 Serracca, L., Verani, M., Battistini, R., Rossini, I., Carducci, A., & Ercolini, C. (2010). Evaluation of
436 Adenovirus and *E. coli* as indicators for human enteric viruses presence in mussels produced in
437 La Spezia Gulf (Italy). *Letters in Applied Microbiology*, 50(5), 462-467.

- 438 Strubbia, S., Lyons, B. P., & Lee, R. J. (2016). Geographical and temporal variation of E. coli and
439 norovirus in mussels. *Marine pollution bulletin*, 107(1), 66-70.
- 440 Turolla, E. (2008). La venericoltura in Italia. En A. Lovatelli, A. Farías & I. Uriarte (eds). FAO
441 Actas de Pesca y Acuicultura. No. 12. Roma, FAO. pp. 177–188.
- 442 <http://www.fao.org/3/a-i0444s/i0444s13.pdf>
- 443 Viaroli, P., Marinov, D., Bodini, A., Giordani, G., Galbiati, L., Somma, F., et al. (2007). Analysis
444 of clam farming scenarios in the Sacca di Goro lagoon. *Transitional Waters Monographs*, 1,
445 71-92.