



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE  
DELLA RICERCA

## Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

An ISH technique for the early detection of *Enterospora nucleophila*, an intranuclear microsporidian causing emaciative disease in gilthead sea bream

This is the submitted version (pre peer-review, preprint) of the following publication:

*Published Version:*

Hossam Eldin N., Caffara M., Fioravanti M.L., Sitjà-Bobadilla A., Palenzuela O. (2017). An ISH technique for the early detection of *Enterospora nucleophila*, an intranuclear microsporidian causing emaciative disease in gilthead sea bream.

*Availability:*

This version is available at: <https://hdl.handle.net/11585/622814> since: 2019-01-14

*Published:*

DOI: <http://doi.org/>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

**An ISH technique for the early detection of *Enterospora nucleophila*, an intranuclear microsporidian causing emaciative disease in gilthead sea bream****Hossam Eldin N<sup>1,2</sup>**, Caffara M<sup>3</sup>, Fioravanti M<sup>3</sup>, Sitjà-Bobadilla A<sup>1</sup>, Palenzuela O<sup>1</sup><sup>1</sup>IATS-CSIC, Ribera De Cabanes, Spain, <sup>2</sup>National Institute of Oceanography and Fisheries (NIOF), Hurghada, Egypt, <sup>3</sup>Dept. Veterinary Medical Sciences, University of Bologna, Bologna, Italy

*Enterospora nucleophila* is a microsporidian parasite causing serious emaciative disease in cultured gilthead sea bream (GSB), *Sparus aurata*, and closely related to *Enterocytozoon hepatopenaei* from shrimp. The parasite can be found within the nuclei of enterocytes and rodlet cells of the fish intestinal epithelium, and occasionally in the cytoplasm of phagocytes in more advanced infections. However, spores are the only stage that can be unmistakably identified but even this may require thorough histopathological examination and use of chitin fluorescent stains for a proper confirmatory diagnosis, due to their minute size and intranuclear location. In the absence of spores, the infection can be suspected from a remarkable hypercellularity and the presence of altered nuclei in the epithelial layer. This results in a very poor correlation of disease signs with diagnosis of *E. nucleophila* infection, and to current uncertainty about its real impact in GSB culture.

This work describes the development and application of an *in situ* hybridization (ISH) technique as a powerful tool to overcome current diagnostic limitations for this species, and to decipher basic data on the infection and disease onset. We designed DIG-labelled oligonucleotide probes targeting unique regions of the (+) strand of *E. nucleophila* rDNA gene, and we developed an ISH protocol that results in good staining of infected host cells prior to the development of spores or other conspicuous stages.

On clinically infected samples, numerous ISH-positive cells are present which are not stained with fluorescent whiteners that bind to parasite spores. In contrast, poor staining of spores was obtained with the DNA probes, due to their limited penetration in these stages and the low number of target gene copies that they harbour. These aspects might be improved using specific steps (e.g., chitinase or more aggressive permeabilization strategies) but they are not necessary in a clinical diagnostic context.

Due to the relevance of *E. nucleophila* infections it is essential to decipher unknown aspects of the parasite's biology and course of infection using proper diagnostic tools. Their cryptic nature makes this a challenging task for microsporidians like *E. nucleophila*, which can benefit from the method hereby presented.

Funding of presentation:

This study was partially supported by EU H2020 program through the ParaFishControl project (634429).