

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Rational Degenerations of M-Curves, Totally Positive Grassmannians and KP2-Solitons

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Simonetta Abenda, P.G.G. (2018). Rational Degenerations of M-Curves, Totally Positive Grassmannians and KP2-Solitons. COMMUNICATIONS IN MATHEMATICAL PHYSICS, 361(3), 1029-1081 [10.1007/s00220-018-3123-y].

Availability:

This version is available at: <https://hdl.handle.net/11585/638436> since: 2021-06-29

Published:

DOI: <http://doi.org/10.1007/s00220-018-3123-y>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Abenda, S., Grinevich, P.G. Rational Degenerations of M-Curves, Totally Positive Grassmannians and KP2-Solitons. Commun. Math. Phys. 361, 1029–1081 (2018).

The final published version is available online at: <https://doi.org/10.1007/s00220-018-3123-y>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

RATIONAL DEGENERATIONS OF M-CURVES, TOTALLY POSITIVE GRASSMANNIANS AND KP2-SOLITONS.

SIMONETTA ABENDA AND PETR G. GRINEVICH

ABSTRACT. We establish a new connection between the theory of totally positive Grassmannians and the theory of M-curves using the finite-gap theory for solitons of the KP equation. Here and in the following KP equation denotes the Kadomtsev-Petviashvili 2 equation (see [\(1\)](#)), which is the first flow from the KP hierarchy. We also assume that all KP times are **real**. We associate to any point of the real totally positive Grassmannian $Gr^{TP}(N, M)$ a reducible curve which is a rational degeneration of an M-curve of minimal genus $g = N(M - N)$, and we reconstruct the real algebraic-geometric data á la Krichever for the underlying real bounded multiline KP soliton solutions. From this construction it follows that these multiline solitons can be explicitly obtained by degenerating regular real finite-gap solutions corresponding to smooth M-curves. In our approach we rule the addition of each new rational component to the spectral curve via an elementary Darboux transformation which corresponds to a section of a specific projection $Gr^{TP}(r + 1, M - N + r + 1) \mapsto Gr^{TP}(r, M - N + r)$.

CONTENTS

1. Introduction	2
1.1. Main results and plan of the paper	4
2. Multi-soliton KP solutions	6
2.1. The heat hierarchy and the dressing transformation	7
2.2. Real finite-gap KP solutions and M-curves	8
3. Algebraic-geometric approach to real bounded regular multiline KP solitons	10
3.1. Algebraic-geometric construction for soliton data in $Gr^{TP}(1, M)$	11
3.2. The algebraic-geometrical setting for soliton data in $Gr^{TP}(N, M)$	14
4. The reducible M-curve and the effective divisor for soliton data in $Gr^{TP}(N, M)$	20
5. Proof of Theorem 3	22
5.1. The algebraic part	23
5.2. The analytic part	28
6. Analytic properties of the effective divisor	33
6.1. Zero divisor for the vacuum wave function	33
6.2. Zero divisor for the normalized Darboux transformed wave function	34
7. $\Gamma(\xi)$ and the vacuum divisor for soliton data in $Gr^{TP}(2, 4)$	36
7.1. $\Gamma(\xi)$ and its desingularization for generic soliton data in $Gr^{TP}(2, 4)$	36
7.2. The leading order coefficients and vectors of the vacuum wavefunction	37
7.3. The vacuum divisor for soliton data in $Gr^{TP}(2, 4)$	38
Appendix A. Points in $Gr^{TP}(N, M)$ and totally positive matrices in classical sense	39
Appendix B. Lemmas for the proof of Theorem 6	42

Key words and phrases. Total positivity; Grassmannians; KP finite-gap theory; real solitons; M-curves.

This work has been partially supported by PRIN “Teorie geometriche e analitiche dei sistemi Hamiltoniani in dimensioni finite e infinite”, by the project “EQUATIONS”, by the Russian Foundation for Basic Research, grant 17-01-00366, by the program “Fundamental problems of nonlinear dynamics”, Presidium of RAS .

1. INTRODUCTION

Since the seminal works of Schoenberg, Gantmacher and Krein [50, 19], totally positive matrices have appeared in connection with problems from different areas of pure and applied mathematics, including small vibrations of mechanical systems, statistical mechanics, approximation theory, combinatorics, graph theory (for more details see [38, 47]). A generalization of total positivity for generic reductive groups and their flag varieties was introduced by Lusztig in [36]. In [37] this construction was extended to the partial flag manifolds, including Grassmannians associated with arbitrary reductive groups. A parametrization of the totally non-negative part of a flag variety was obtained by Marsh and Rietsch in [40]. The development of criteria to test total positivity is also deeply related to the foundation of the cluster algebra theory of Fomin and Zelevinsky [17, 18]. Relevant recent applications include the development of Poisson geometry in cluster algebras [20], and the computation of scattering amplitudes for on-shell diagrams for the planar limit of $\mathcal{N} = 4$ super Yang Mills theory [4, 5]. In recent years, total positivity has proven to be an efficient tool to investigate the asymptotic properties of multi-line solitons for the KP equation [6, 7, 8, 10, 11, 14, 28, 29, 30] which we study in this paper in the finite-gap setting.

The topological classification of real algebraic curves is also deeply connected with the theory of integrable systems and statistical models. In particular M-curves [24], i.e. real algebraic curves with maximal number of components, naturally arise in the theory of integrable systems, such as the real finite-gap theory of the KP equation [16] the theory of finite-gap at one energy two-dimensional Schrödinger operators at the energies below the ground state [52, 53], quantum cluster systems [21] and in statistical models such as dimer models [27]. Let us point out that in these papers the M-curves (Harnack's curves) arise as the spectral curves for two-dimensional models with double-periodic boundary conditions or their quasiperiodic generalizations, and, generically, they are smooth. An investigation of the relative positions of the branches of real algebraic curves of given degree (and similarly for algebraic surfaces) is the first part of the Hilbert's 16th problem. The term M-curve was first introduced by Petrovsky [46] ("M" means "maximal"). Additional information about this topic can be found in the review papers [44, 54].

In this paper we establish a connection of different nature between the theory of totally positive Grassmannians, and rational degenerations of M curves, in the framework of the solitonic limit of real finite gap solutions for the Kadomtsev-Petviashvili 2 (KP) equation

$$(1) \quad \partial_x (-4\partial_t u + 6u\partial_x u + \partial_x^3 u) + 3\partial_y^2 u = 0,$$

where ∂_z denotes the usual partial derivative with respect to the variable z .

We recall that **a relevant question is:** can all **real regular** KP multisoliton solutions be obtained by degenerating **real regular** finite-gap KP solutions. The importance of this problem was pointed out by S.P. Novikov. In this paper we provide a positive answer to the above question for real bounded regular multiline KP solitons, associated to the principal cell $Gr^{\text{TP}}(N, M)$.

We remark that, even if the same soliton solution can be obtained from different degenerate algebraic-geometric data, Baker-Akhiezer functions are still an important tool of integration in such limiting case [15]. For instance, in the case of the open Toda lattice, such approach has been used by Krichever and Vaninsky [34] to construct a Baker-Akhiezer function on a reducible

singular curve which is the limit of the hyperelliptic spectral curve associated to the periodic Toda lattice.

Multiline KP solitons can be obtained from the Wronskian method [56, 41, 42, 25] starting from N independent solutions of the heat hierarchy depending on M phases and are naturally identified with a certain finite-dimensional reduction of the Sato Grassmannian [49]. The KP multiline soliton solutions are real bounded and regular for all (x, y, t) if and only if they are parametrized by points in the totally nonnegative part of the Grassmannian, $Gr^{\text{TNN}}(N, M)$ [29]. The asymptotic behavior in space-time and the tropicalization for this class of solutions has been thoroughly investigated in [10, 11, 28, 29, 30] using the combinatorial classification of Postnikov [48] of $Gr^{\text{TNN}}(N, M)$. According to Krichever scheme of KP finite gap theory [31, 32], this family of solutions also originates from regular quasi-periodic solutions in the solitonic limit.

In this paper we show that real bounded regular multiline KP soliton solutions originate from **regular real quasi-periodic finite-gap** solutions in the solitonic limit.

Regular real quasi-periodic solutions are parametrized by degree g non-special divisors on genus g Riemann surfaces which possess an antiholomorphic involution of decomposing type which fixes the maximum number $g + 1$ of ovals [16]*. In such a case, there exists a homological basis of cycles such that the α cycles correspond to g (finite) ovals and the essential singularity of the wavefunction is in the remaining (infinite) oval. Finally, for a fixed spectral curve, such solutions are parametrized by degree g non-special divisors of the Baker–Akhiezer function with exactly one pole in each finite oval. In the solitonic limit a certain number of cycles shrinks and the non-singular curve degenerates to a reducible curve of rational type.

Let us fix the M phases, $\kappa_1 < \dots < \kappa_M$. Then generic regular bounded multi-line KP solitons are parametrized by points in $Gr^{\text{TP}}(N, M) \subset Gr^{\text{TNN}}(N, M)$, i.e. points in the Grassmannian with all Plücker coordinates strictly positive [48]. These solutions then depend on $N(M - N)$ parameters (the dimension of the Grassmannian). We show that for any compact subset in $Gr^{\text{TP}}(N, M)$ we may fix a spectral curve, which is a rational degeneration of regular M-curves of genus $N(M - N)$, and the solutions are parametrized by $N(M - N)$ point divisors on it.

The starting point of the construction is the following observation: for any soliton data in $Gr^{\text{TP}}(N, M)$, the Sato dressed wave function is defined on \mathbb{CP}^1 , which we denote Γ_0 , with $M + 1$ marked points (the phases $\kappa_1, \dots, \kappa_M$ and the essential singularity P_0). On Γ_0 , the normalized Sato dressed wave function is a Baker–Akhiezer function for the given soliton data with a real N -point divisor. To obtain a degree $N(M - N)$ divisor, in our construction we attach additional components to Γ_0 in such a way that the resulting reducible curve possesses the $N(M - N) + 1$ real components (ovals), and we extend the Baker–Akhiezer function to these new components so that the degree of the divisor matches the number of finite ovals and the reality constraints of Dubrovin–Natanzon theorem. To make effective the relation between Sato’s and finite-gap approaches for this family of KP soliton solutions in $Gr^{\text{TP}}(N, M)$, we use total positivity in classical sense. Our construction gives a new relation between the theory of integrable systems, the theory of algebraic curves and total positivity. In [2], we modify our approach and extend the present construction to all positroid cells in the stratification of $Gr^{\text{TNN}}(N, M)$ characterized in [48].

A relevant open question is the characterization of the regular M-curves whose rational limit has been constructed here and in [2]. Another important open question is also the thorough investigation of the following problem: start from a given reducible curve and a divisor compatible with the reality and regularity conditions in [16] and identify the soliton data in $Gr^{\text{TNN}}(N, M)$.

*In the following, with a slight abuse of notation, we call regular M-curves the Riemann surfaces satisfying Dubrovin and Natanzon’s hypotheses.

In [11], one of us has identified the $(M - 1)$ -dimensional variety of soliton data in $Gr^{TP}(N, M)$ associated to a specific rational degeneration of a hyperelliptic curve and proved that the vacuum KP divisor for such soliton data coincides with the Toda divisor found in [34].

In a forthcoming paper we also plan to investigate explicitly the relations between our construction and the characterization of the asymptotic behaviour of the multi-line solitons in [11, 30], i.e. to characterize the asymptotics of the zero divisor $\mathcal{D}(x, y, t, \vec{0})$ on $\Gamma(\xi)$. Finally it would be relevant to investigate possible connections between KP soliton theory and statistical models such as dimers in the disk [35] or with field theoretical models in the framework of the planar limit of $\mathcal{N} = 4$ -SYM [4, 5] through their common combinatorial characterization in $Gr^{TNN}(N, M)$.

1.1. Main results and plan of the paper. The main result of this paper is that, for soliton solutions associated with the principal cell $Gr^{TP}(N, M)$, we can always fix algebraic-geometric data on *reducible* curves, generating these solutions, with additional requirements that these curves are rational degenerations of some family of regular M-curves and that the divisor points satisfy the reality and regularity conditions of [16].

The first nontrivial case we study is $Gr^{TP}(1, M)$ (see also [1]). In this case the curve Γ is obtained by attaching a second copy of \mathbb{CP}^1 to Γ_0 at the phases $\kappa_1, \dots, \kappa_M$, and it is the rational degeneration of a hyperelliptic M-curve of genus $M - 1$. The divisor has degree $M - 1$ and exactly one point in each real finite oval. We do the construction of the KP wavefunction on Γ in two steps. We first extend the Sato vacuum wavefunction to a degree $M - 1$ meromorphic function, $\Psi(P, \vec{t})$ on $\Gamma \setminus \{P_0\}$ with exactly one simple pole in each finite oval (see [13]) and such that at the Darboux marked point Q_1 belonging to the infinite oval, $\Psi(Q_1, \vec{t})$ coincides with the generator of the dressing transformation for the KP solution (see [14]). Then, after normalization, the dressing of Ψ , $\tilde{\Psi}(P, \vec{t})$ has a degree $M - 1$ pole divisor with exactly one simple pole in each finite oval and its restriction to Γ_0 is the normalized Sato dressed wavefunction (see Lemma [1]).

For soliton data in $Gr^{TP}(N, M)$, we use a similar scheme: we first glue $(N + 1)$ copies of \mathbb{CP}^1 , $\Gamma = \Gamma_0 \sqcup \Gamma_1 \sqcup \dots \sqcup \Gamma_N$ in pairs at some real ordered points creating $N(M - N) + 1$ ovals. We then extend Sato vacuum (zero potential) wavefunction on $\Gamma_0 \setminus \{P_0\}$ to a meromorphic vacuum wavefunction on $\Gamma \setminus \{P_0\}$ with a degree $N(M - N)$ divisor on $\Gamma \setminus \Gamma_0$ such that exactly one divisor point belongs to each finite oval and, at each marked Darboux point Q_r , $r \in [N]$, we require that Ψ coincides for all time with one of the N generators of the dressing transformation associated to the given soliton data. Then, after normalization, the dressing of Ψ , $\tilde{\Psi}(P, \vec{t})$ has a degree $N(M - N)$ pole divisor with exactly one simple pole in each finite oval and its restriction to Γ_0 is the normalized Sato dressed wavefunction (see Theorem [4]). The advantage of this multistep procedure is that it is easier to control indirectly the position of the effective KP divisor imposing conditions on the vacuum wavefunction than to directly extend the Baker Akhiezer function to $\Gamma \setminus \{P_0\}$.

In the general case $Gr^{TP}(N, M)$, the main technical problem is to coherently glue more copies of \mathbb{CP}^1 in pairs creating the necessary number of ovals, in such a way to control the position of the effective divisor.

To solve this problem we use the following two-step construction:

- (1) We provide an algebraic model of the vacuum wave function using total positivity to control the values of the vacuum wavefunction at all marked points.

[†]Here and in the following, unless differently specified, \vec{t} means the whole sequence of times $\vec{t} = (x, y, t, t_4, t_5, \dots)$ associated to the KP hierarchy.

- (2) We introduce a family of curves depending on a parameter $\xi \gg 1$, and we show that, for any point in $Gr^{\text{TP}}(N, M)$ there exists ξ_0 such that for any $\xi > \xi_0$ the analytic extension of the Sato vacuum wave function to the curve $\Gamma(\xi)$ coincides with the algebraic model.

The topological type of the resulting reducible rational curve $\Gamma = \Gamma(\xi)$ is independent of $\xi \gg 1$ and it is the same for all points in $Gr^{\text{TP}}(N, M)$. In particular the real part of Γ , $\Gamma_{\mathbb{R}}$ has $N(M - N) + 1$ real ovals, and Γ is the rational degeneration of a regular M-curve of minimal genus $g = N(M - N)$. In section 7 we construct explicitly the reducible nodal curve associated to generic points in $Gr^{\text{TP}}(2, 4)$ and the underlying regular M-curve of genus 4, which is a 3-sheeted covering of the Riemann sphere.

More precisely, the leading order behavior of the **vacuum** wavefunction $\Psi(P, \vec{t})$ at all marked points of $\Gamma(\xi)$, for $\xi \gg 1$, is ruled explicitly by an algebraic recursion (see Theorems 5 and 6) associated to a conveniently normalized totally positive band matrix \hat{A} representing the soliton datum in $Gr^{\text{TP}}(N, M)$. The exact behavior of $\Psi(P, \vec{t})$ at the marked points on $\Gamma(\xi)$ is ruled via an upper triangular matrix $\hat{A}(\xi)$ explicitly computed in the proof of Theorem 6.

Thanks to the recursive relations in Theorem 5, we control the sign of Ψ at the double points and, therefore, the position of the vacuum divisor in the ovals of $\Gamma(\xi)$. In section 7 we explicitly compute such vacuum divisor for the algebraic curve associated to generic soliton data in $Gr^{\text{TP}}(2, 4)$. The characterization of the vacuum wave-function is contained in Theorem 3 where we prove that $\Psi(P, \vec{t})$

- (1) coincides with Sato's vacuum wavefunction on Γ_0 ;
- (2) is meromorphic of degree $N(M - N)$ on $\Gamma \setminus \{P_0\}$;
- (3) possesses $M - N$ simple poles on each Γ_r , $r \in [N]$;
- (4) possesses exactly one simple pole in each finite oval of Γ ;
- (5) coincides with an explicit basis of heat hierarchy solution $f_{\xi}^{(r)}(\vec{t})$, which generate the Darboux transformation, at the marked Darboux point $Q_r \in \Gamma_r$, for any $r \in [N]$ and for all \vec{t} .

Properties (4) and (5) are sufficient to guarantee the complete control of the position of the effective divisor of the normalized KP wavefunction $\tilde{\Psi}(P, \vec{t}) = \tilde{\Psi}_{\xi}(P, \vec{t})$, which is obtained applying the dressing (Darboux) transformation to the vacuum wavefunction: the effective divisor $\mathcal{D} = \mathcal{D}(\xi)$ satisfies both the conditions imposed by Dubrovin–Natanzon theorem and the constraints imposed by Sato dressing (see Theorem 4). More precisely, we prove that $\tilde{\Psi}(P, \vec{t})$ has the following properties:

- (1) its restriction to Γ_0 coincides with Sato's KP dressed wavefunction;
- (2) it is meromorphic of degree $N(M - N)$ on $\Gamma \setminus \{P_0\}$;
- (3) it possesses N poles on Γ_0 ;
- (4) it possesses $M - N - 1$ simple poles on each Γ_r , $r \in [N]$;
- (5) it possesses exactly one simple pole in each finite oval of Γ .

We also provide explicit estimates for the position of the divisor \mathcal{D} in Theorem 7 in the given local coordinates. We stress that such local coordinates are associated to a totally positive basis in Fomin–Zelevinsky sense [17] and, in this sense, we establish a natural correspondence between points in $Gr^{\text{TP}}(N, M)$ and the algebraic–geometric data associated to the corresponding soliton solutions.

Plan of the paper:

- In section 2, we recall some known facts about regular finite gap and multi-soliton solutions of the KP equation.

- In section [3](#), we first associate the rational degeneration of a genus $(M-1)$ hyperelliptic M-curve and construct the effective divisor for soliton data in $Gr^{\text{TP}}(1, M)$. Then we present the main ideas of the algebraic–geometric construction for soliton data in $Gr^{\text{TP}}(N, M)$.
- Section [4](#) contains the principal statements. In Theorem [4](#), we extend the Baker–Akhiezer function for generic soliton data in $Gr^{\text{TP}}(N, M)$ to the curve $\Gamma(\xi)$. The proof follows immediately from Theorem [3](#), in which we extend Sato’s vacuum wavefunction to $\Gamma(\xi)$ so that it takes the desired values at all marked points (double points and Darboux points).
- The proof of Theorem [3](#) is carried out in detail in Section [5](#). The first part of the proof is algebraic and fixes the leading order behavior of the vacuum wavefunction at the double points and at the Darboux points (Lemma [2](#) and Theorem [5](#)). The matrix associated to such leading order behavior is in band form and we characterize its properties in Appendix [A](#). The second part of the proof (Theorem [6](#)) is analytic and contains the explicit construction of the vacuum wavefunction. Some Lemmas necessary to its proof are in Appendix [B](#).
- In section [6](#) we characterize the pole divisor \mathcal{D} and the zero divisor $\mathcal{D}(\vec{t})$ of the KP–eigenfunction $\tilde{\Psi}(P, \vec{t})$.
- In section [7](#) we apply our construction to soliton data in $Gr^{\text{TP}}(2, 4)$. In particular we explicitly construct the curve $\Gamma(\xi)$ and show that it is the rational degeneration of an M-curve of genus 4. We also construct the vacuum divisor and check the algebraic identities for the vacuum wavefunction in such case.

Notation: We use the following notations throughout the paper:

- (1) N and M are positive integers such that $N < M$;
- (2) For $s \in \mathbb{N}$ let $[s] = \{1, 2, \dots, s\}$; if $s, j \in \mathbb{N}$, $s < j$, then $[s, j] = \{s, s+1, s+2, \dots, j-1, j\}$;
- (3) For a given matrix A we denote by $A_{[j_1, \dots, j_q]}^{[i_1, \dots, i_p]}$ the $p \times q$ submatrix of A formed by the elements $A_{j_l}^{i_m}$, $m \in [p]$, $l \in [q]$;
- (4) If $p = q$, $\Delta_{[j_1, \dots, j_p]}^{[i_1, \dots, i_p]}(A)$ denotes the determinant of the submatrix $A_{[j_1, \dots, j_p]}^{[i_1, \dots, i_p]}$.
- (5) For a given matrix A , $\Delta_{[j_1, \dots, j_n]}(A)$ denotes the determinant of the $n \times n$ matrix, combined from the last n rows of the columns j_1, \dots, j_n ;
- (6) $\vec{t} = (t_1, t_2, t_3, \dots)$, where $t_1 = x$, $t_2 = y$, $t_3 = t$;
- (7) $\theta(\zeta, \vec{t}) = \sum_{n=1}^{\infty} \zeta^n t_n$,
- (8) We denote the real phases $\kappa_1 < \kappa_2 < \dots < \kappa_M$ and $\theta_j \equiv \theta(\kappa_j, \vec{t})$.

2. MULTI-SOLITON KP SOLUTIONS

The KP equation [\(1\)](#) was originally introduced by Kadomtsev and Petviashvili [\[26\]](#) to study the stability of the Korteweg de Vries equation under weak transverse perturbation in the y direction. It has remarkable properties coming from the fact that it is the first non trivial flow [\[56\]](#) of the so-called KP Hierarchy (see the monographs [\[13, 15, 25, 43, 45\]](#)). The family of regular real bounded KP multi-line soliton solutions may be characterized with different approaches: the Wronskian method, a special reduction of the Sato Grassmannian and in the solitonic limit of real finite-gap theory. Below we briefly recall how these solutions may be obtained via these different approaches.

2.1. The heat hierarchy and the dressing transformation. Let $A = (A_j^i)$, be an $N \times M$ real matrix and fix $\kappa_1 < \dots < \kappa_M$. In the following $\vec{t} = (x, y, t, t_4, t_5, \dots)$ indicates an infinite number of real times unless specified differently. Following [41], let us consider N linear independent solutions

$$(2) \quad f^{(i)}(\vec{t}) = \sum_{j=1}^M A_j^i e^{\theta_j}, \quad i \in [N],$$

to the heat hierarchy[‡]

$$(3) \quad \partial_y f = \partial_x^2 f, \quad \partial_{t_l} f = \partial_x^l f, \quad l = 2, 3, \dots,$$

and define their Wronskian

$$(4) \quad \tau(\vec{t}) = Wr(f^{(1)}, \dots, f^{(N)}) \equiv \sum_I \Delta_I(A) \prod_{\substack{i_1 < i_2 \\ i_1, i_2 \in I}} (\kappa_{i_2} - \kappa_{i_1}) e^{i \sum \theta_i}$$

where the sum is over all N -element ordered subsets I in $[M]$, *i.e.* $I = \{1 \leq i_1 < i_2 < \dots < i_N < M\}$ and $\Delta_I(A)$ are the maximal minors of the matrix A , *i.e.* the Plücker coordinates for the corresponding point in the finite dimensional Grassmannian $Gr(N, M)$.

The multi-line soliton solution to the KP equation (1) is defined by the following formula:

$$(5) \quad u(\vec{t}) = 2\partial_x^2 \log(\tau(\vec{t})).$$

For the full KP hierarchy it is rather easy to check that the condition $\Delta_I(A) \geq 0$ for all I is necessary and sufficient to have (5) regular and real bounded for all times. A very nontrivial result of [29] states that this condition is necessary and sufficient also in the case of the first flow from the KP hierarchy. In such case, the equivalence class of A , $[A]$ is a point in the totally non-negative Grassmannian [48]

$$Gr^{\text{TNN}}(N, M) = GL_N^+ \backslash Mat_{N,M}^{\text{TNN}},$$

where $Mat_{N,M}^{\text{TNN}}$ is the set of real $N \times M$ matrices of maximal rank N with nonnegative maximal minors $\Delta_I(A)$ and GL_N^+ is the group of $N \times N$ matrices with positive determinants.

Since left multiplication by $N \times N$ matrices with positive determinants preserves the KP multisoliton solution $u(\vec{t})$ in (5), there is a natural bijection between KP regular real bounded multi-line solitons (5) and points in $Gr^{\text{TNN}}(N, M)$.

Let us recall the construction of the wave function for the multi-line soliton solutions. According to Sato theory [49] all KP soliton solutions may be obtained from the dressing (inverse gauge) transformation of the vacuum eigenfunction $\Psi^{(0)}(\zeta, \vec{t}) = \exp(\theta(\zeta, \vec{t}))$, which solves

$$\partial_x \Psi^{(0)}(\zeta, \vec{t}) = \zeta \Psi^{(0)}(\zeta, \vec{t}), \quad \partial_{t_l} \Psi^{(0)}(\zeta, \vec{t}) = \zeta^l \Psi^{(0)}(\zeta, \vec{t}), \quad l \geq 2,$$

via the dressing (*i.e.* gauge) operator

$$W(\vec{t}) = 1 - \sum_{j=1}^{\infty} \chi_j(\vec{t}) \partial_x^{-N},$$

under the condition that W satisfies Sato equations

$$\partial_{t_n} W = B_n W - W \partial_x^n, \quad n \geq 1,$$

[‡]We remark that the class of solutions to (3) that we consider in this paper is not the general one.

where $B_n = (W\partial_x^n W^{-1})_+$ is the differential part of the operator $W\partial_x^n W^{-1}$. Then

$$L = W\partial_x W^{-1} = \partial_x + \frac{u(\vec{t})}{2}\partial_x^{-1} + \dots, \quad u(\vec{t}) = 2\partial_x \chi_1(\vec{t}),$$

and

$$\hat{\Psi}^{(0)}(\zeta; \vec{t}) = W\Psi^{(0)}(\zeta; \vec{t})$$

are respectively the KP-Lax operator, the KP-potential (KP solution) and the KP-eigenfunction, i.e.

$$L\hat{\Psi}^{(0)}(\zeta; \vec{t}) = \zeta\hat{\Psi}^{(0)}(\zeta; \vec{t}), \quad \partial_{t_l}\hat{\Psi}^{(0)}(\zeta; \vec{t}) = B_l\hat{\Psi}^{(0)}(\zeta; \vec{t}), \quad l \geq 2,$$

where $B_l = (W\partial_x^l W^{-1})_+ = (L^l)_+$.

The dressing transformation associated to the line solitons (5) corresponds to the following choice of the dressing operator

$$W = 1 - w_1(\vec{t})\partial_x^{-1} - \dots - w_N(\vec{t})\partial_x^{-N},$$

where $w_1(\vec{t}), \dots, w_N(\vec{t})$ are uniquely defined as solutions to the following linear system of equations

$$(6) \quad \partial_x^N f^{(i)} = w_1\partial_x^{N-1}f^{(i)} + \dots + w_N f^{(i)}, \quad i \in [N],$$

and, in such case, $w_1(\vec{t}) = \partial_x \tau / \tau$ and $u(\vec{t}) = 2\partial_x w_1(\vec{t}) = 2\partial_x^2 \log(\tau)$. Moreover

$$(7) \quad D^{(N)} \equiv W\partial_x^N = \partial_x^N - \partial_x^{N-1}w_1(\vec{t}) - \dots - w_N(\vec{t}).$$

We observe that w_1, \dots, w_N is the solution to the linear system (6) if and only if

$$(8) \quad D^{(N)} f^{(i)} \equiv W\partial_x^N f^{(i)} = 0, \quad i \in [N].$$

Moreover, if the above identity holds, then

$$\partial_{t_l}(D^{(N)} f^{(i)}) = 0, \quad \forall l \in \mathbb{N},$$

that is, by construction, the N -th order Darboux transformation is associated with the N eigenfunctions $f^{(1)}(\vec{t}), \dots, f^{(N)}(\vec{t})$, of the KP Lax Pair with zero potential for the infinite multiplicity eigenvalue.

The KP-eigenfunction associated to this class of solutions is

$$\hat{\Psi}^{(0)}(\zeta; \vec{t}) = W\Psi^{(0)}(\zeta; \vec{t}) = (1 - w_1(\vec{t})\zeta^{-1} - \dots - w_N(\vec{t})\zeta^{-N}) e^{\theta(\zeta, \vec{t})},$$

or, equivalently,

$$(9) \quad D^{(N)}\Psi^{(0)}(\zeta; \vec{t}) \equiv W\partial_x^N \Psi^{(0)}(\zeta; \vec{t}) = (\zeta^N - \zeta^{N-1}w_1(\vec{t}) - \dots - w_N(\vec{t})) \Psi^{(0)}(\zeta; \vec{t}) = \zeta^N \hat{\Psi}^{(0)}(\zeta; \vec{t}).$$

2.2. Real finite-gap KP solutions and M-curves. The general method to construct periodic and quasi-periodic solutions to the KP equation is due to Krichever [31, 32]: let Γ be a smooth algebraic curve of genus g with a marked point P_0 and let ζ^{-1} be a local parameter in Γ in a neighborhood of P_0 such that $\zeta^{-1}(P_0) = 0$. The triple $(\Gamma, P_0, \zeta^{-1})$ defines a family of exact solutions to (1) parametrized by degree g non-special divisors \mathcal{D} defined on $\Gamma \setminus \{P_0\}$.

The finite gap solutions of (1) are constructed starting from the commutation representation §56

$$(10) \quad [-\partial_y + B_2, -\partial_t + B_3] = 0,$$

[§]The representation of KP as commutation of operators is also known in literature as Zakharov-Shabat equation or zero-curvature condition.




FIGURE 1. Left: a regular M-curve, $g = 2$, 3 real ovals (painted grey), involution σ is reflection, orthogonal to the Π plane. Right: degeneration of a genus 2 M-curve, 3 real ovals (painted grey).

where

$$B_2 \equiv (L^2)_+ = \partial_x^2 + u, \quad B_3 = (L^3)_+ = \partial_x^3 + \frac{3}{4}(u\partial_x + \partial_x u) + \tilde{u},$$

and $\partial_x \tilde{u} = \frac{3}{4} \partial_y u$. Then, the Baker-Akhiezer function $\tilde{\Psi}(P, \vec{t})$ meromorphic on $\Gamma \setminus \{P_0\}$, with poles at the points of the divisor \mathcal{D} and essential singularity at P_0 of the form

$$\tilde{\Psi}(\zeta, \vec{t}) = e^{\zeta x + \zeta^2 y + \zeta^3 t + \dots} (1 - \chi_1(\vec{t})\zeta^{-1} - \dots - \chi_N(\vec{t})\zeta^{-N} - \dots)$$

is an eigenfunction of the following linear differential operators

$$\partial_y \tilde{\Psi} = B_2 \tilde{\Psi}, \quad \partial_t \tilde{\Psi} = B_3 \tilde{\Psi},$$

and in such case, imposing compatibility condition (10), $u(\vec{t}) = 2\partial_x \chi_1(\vec{t})$ satisfies the KP equation.

The divisor of poles \mathcal{D} does not depend on the times \vec{t} . In contrast to it, the divisor of zeroes $\mathcal{D}(\vec{t})$ depends on all times. The Abel transform of $\mathcal{D}(\vec{t})$ is a linear function of times \vec{t} , therefore such transform linearizes the KP hierarchy. \mathcal{D} is an effective divisor, therefore $\Psi(\zeta, \vec{0}) \equiv 1$, and at the point $\vec{t} = \vec{0}$ the divisor of zeroes coincides with the divisor of poles, $\mathcal{D}(\vec{0}) = \mathcal{D}$. The last identity justifies the use of the same letter for both divisors.

After fixing a canonical basis of cycles $a_1, \dots, a_g, b_1, \dots, b_g$ and a basis of normalized holomorphic differentials $\omega_1, \dots, \omega_g$ on Γ , that is $\oint_{a_j} \omega_k = 2\pi i \delta_{jk}$, $\oint_{b_j} \omega_k = B_{kj}$, $j, k \in [g]$, the KP solution takes the form

$$(11) \quad u(x, y, t) = 2\partial_x^2 \log \theta(xU^{(1)} + yU^{(2)} + tU^{(3)} + z_0) + c_1,$$

where θ is the Riemann theta function and $U^{(k)}$, $k \in [3]$ are vectors of the b -periods of the following normalized meromorphic differentials, holomorphic on $\Gamma \setminus \{P_0\}$ and with principal parts $\hat{\omega}^{(k)} = d(\zeta^k) + \dots$, $k \in [3]$, at P_0 (see [31, 16]).

The necessary and sufficient conditions for the smoothness and realness of the solution (11) associated with smooth curve Γ of genus g have been proven by Dubrovin and Natanzon (see [16] and references therein): Γ must be an M-curve, that is it possesses an antiholomorphic involution[¶]

$$\sigma : \Gamma \rightarrow \Gamma, \quad \sigma^2 = 1, \quad \sigma(P_0) = P_0, \quad \sigma^*(\zeta) = \bar{\zeta},$$

such that the set of fixed points of σ consists of $g+1$ ovals (the maximum number of ovals [23]), $\Omega_0, \Omega_1, \dots, \Omega_g$. These ovals are called “fixed” or “real”. The set of real ovals divides Γ into two connected components. Each of these components is homeomorphic to a sphere with $g+1$ holes. In Figure 1[left] we show an example.

[¶]Here $\bar{\cdot}$ denotes complex conjugation.

On such smooth M-curve of genus g it is possible to fix a basis of cycles such that the essential singularity P_0 belongs to one oval Ω_0 (which is called “infinite” oval), and the remaining g fixed ovals Ω_j , $j \in [g]$ coincide with the a_j -cycles of this basis:

$$\sigma(a_j) = a_j, \quad \sigma(b_j) = -b_j, \quad j \in [g].$$

We call the ovals Ω_j , $j \in [g]$ “finite”. Finally, **in order to have regular real quasi-periodic solutions it is necessary and sufficient that each finite oval contains exactly one pole divisor point** [16].

The proof that this condition is sufficient is rather simple. At $\vec{t} = \vec{0}$ each finite oval contains exactly one zero divisor point. During real times evolution, each point of the zero divisor moves in the corresponding oval, and no point of the zero divisor can touch the oval Ω_0 . Taking into account that the singularities of the solution arise when a divisor point touches the point P_0 , we see that for real times solutions are regular. It is easy to check that this argument is valid for degenerate M-curves if the distance of the pole divisor from the intersection of the finite ovals and the infinite ovals is positively bounded from below.

3. ALGEBRAIC-GEOMETRIC APPROACH TO REAL BOUNDED REGULAR MULTILINE KP SOLITONS

Soliton solutions of KP correspond to algebraic-geometric data associated to rational curves obtained by shrinking some cycles to double points ([33], see also the book [15] and references therein). After applying the Darboux transformation ([7]) we obtain the wave function ([9]) defined on a Riemann sphere denoted by Γ_0 throughout the paper, whose effective divisor $\mathcal{D}^{(0)} = \{\gamma_l^{(0)}; l \in [M-1]\} \subset \Gamma_0 \setminus \{P_0\}$, consists of the N real simple poles of the dressed wavefunction ([9]) at time $\vec{t} = \vec{0} = (0, 0, \dots)$,

$$(12) \quad (\gamma_l^{(0)})^N - w_1(\vec{0})(\gamma_l^{(0)})^{N-1} - \dots - w_{N-1}(\vec{0})\gamma_l^{(0)} - w_N(\vec{0}) = 0, \quad l \in [N],$$

which satisfy $\kappa_1 \leq \gamma_1^{(0)} < \gamma_2^{(0)} < \dots < \gamma_N^{(0)} \leq \kappa_M$ [39], after convenient labeling.

The sufficient part of Dubrovin and Natanzon’s proof holds also when the algebraic M-curve is singular. Since the multi-soliton solutions associated to points in $Gr^{\text{TNN}}(N, M)$ are real bounded regular for all \vec{t} , it is natural to expect that they may be associated to algebraic-geometric data on **reducible** curves which are rational degenerations of regular M-curves. In the following we provide such construction. For example, in Figure 1[right] we show the rational degeneration of the smooth M-curve of Figure 1[left]. For what concerns the degenerate M-curve, we describe them in terms of their real parts represented as a collection of circles in the plane (with non-intersecting interiors) with marked points, where each marked point at one circle is connected to the corresponding marked point at another circle. For an example, see Figure 2[left]. Let us point out that this diagram is a pure topological representation, it does not respect the complex structure. To obtain a degenerate M-curve it is necessary and sufficient that this diagram can be drawn in the plane without intersection (see Figure 2).

In our text we present a solution of the following

Problem. Associate a rational degeneration of a smooth M-curve and a divisor satisfying the reality and regularity conditions, to a regular real bounded multiline soliton solution of KP represented by the following data:

- (1) The fixed numbers $M, N, M > N$.
- (2) A set of M real ordered phases $\kappa_1 < \kappa_2 < \dots < \kappa_M$.
- (3) A point $[A]$ in the totally positive Grassmannian $Gr^{\text{TP}}(N, M)$.

Remark 1. *Of course, the solution to this problem is not unique, and our construction provides an infinite number of curves depending on a parameter.*

To be more precise, in our text we associate the following algebraic-geometrical objects to a given soliton datum:

- (1) A reducible curve Γ which is the rational degeneration of a smooth M-curve of genus $N(M - N)$ equal to the dimension of the totally positive Grassmannian. Γ has exactly $N(M - N) + 1$ ovals. The curve Γ_0 in our approach is one of the irreducible components of Γ .
- (2) On Γ we construct a unique wave-function $\tilde{\Psi}$ such that its restriction on Γ_0 is the normalized Sato wave function.
- (3) On $\Gamma \setminus \{P_0\}$ the wave function $\tilde{\Psi}$ has effective divisor of degree $N(M - N)$ which coincides with (12) on the restriction to Γ_0 . The essential singularity $P_0 \in \Gamma_0$ lies in the infinite oval of Γ . Each finite oval of Γ contains exactly one divisor point.

In our paper we construct the wave function $\tilde{\Psi}$ in **three steps**:

- (1) We first extend the vacuum wave function $\Psi^{(0)} = e^{\theta(\zeta; \vec{t})}$ from the curve Γ_0 to the curve Γ as a meromorphic function Ψ in $\Gamma \setminus \{P_0\}$. The poles of this function are called the **vacuum divisor**. The function Ψ satisfies the heat hierarchy for all values of spectral parameter $\zeta \in \Gamma$.
- (2) We then apply the Darboux transformation (dressing) $D^{(N)}$ to Ψ . In the algebraic-geometric setting, the dressing corresponds to the following shift of the divisor: we add a N -th order pole to the point P_0 and N simple zeroes to some points Q_1, \dots, Q_N . The divisor of $D^{(N)}\Psi$ is non-effective.
- (3) We renormalize the dressed wave function:

$$\tilde{\Psi}(\zeta, \vec{t}) = \frac{D^{(N)}\Psi(\zeta, \vec{t})}{D^{(N)}\Psi(\zeta, \vec{0})}.$$

The divisor of $\tilde{\Psi}$ is the effective KP divisor.

Remark 2. *If we work with regular spectral curves, the shift of the divisor corresponds to a phase shift in the KP solution $u(\vec{t})$. In the case of degenerate curves, the Jacobian consists of several components, and some of them correspond to trivial solutions of KP, and other ones correspond to nontrivial solutions. By applying the Darboux transformation we jump from one component to another, therefore we generate nontrivial solutions from the trivial ones.*

Remark 3. *The advantage of our multi-step procedure is that it is easier to control indirectly the position of the dressed KP divisor via the position of vacuum divisor than directly extend the Sato divisor to Γ . Indeed, we are able to control the sign of the vacuum wave function at all double points and its value at the Darboux points through an algebraic lemma.*

3.1. Algebraic-geometric construction for soliton data in $Gr^{\text{TP}}(1, M)$. In the case of soliton data $[A] = [A_1, \dots, A_M] \in Gr^{\text{TP}}(1, M)$, the expected minimal degree of the divisor is $M - 1 = \dim(Gr^{\text{TP}}(1, M))$. Since Sato constraints produce a one point divisor on Γ_0 , we need to glue at least another copy of \mathbb{CP}^1 to Γ_0 creating double points and M ovals. Then, on such reducible curve, we must extend the wavefunction to a Baker–Akhiezer function with $M - 1$ poles with the following constraints: the poles are simple, the essential singularity $P_0 \in \Gamma_0$ belongs to one such oval and each other oval contains exactly one such pole.

To achieve such a goal, we make the ansatz that Γ is the rational degeneration of a hyperelliptic curve of genus $M - 1$ such that all branch points are real. Let v be the hyperelliptic involution




FIGURE 2. Left: The real part of degenerated M-curve from the previous example is represented as a pair of circles with 3 connecting lines. Right: Not a M-curve, genus = 2, the diagram is non-planar, 1 real oval

and $\Gamma_1 = v(\Gamma_0)$. Here σ is the standard complex conjugation of the Riemann spheres Γ_0, Γ_1 , $\sigma\Gamma_0 = \Gamma_0$, $\sigma\Gamma_1 = \Gamma_1$. The double points κ_j , $j \in [M]$, are fixed points of both involutions σ, v . Finally, let $Q_1 = v(P_0)$. By definition, $\Gamma = \Gamma_0 \sqcup \Gamma_1$ possesses M ovals, $\Omega_0, \Omega_{1,1}, \dots, \Omega_{1,M-1}$ such that $P_0, Q_1 \in \Omega_0$ and, for each $l \in [M-1]$, $\Omega_{1,l}$ is the topological circle formed by the connected union of the intervals $[\kappa_l, \kappa_{l+1}] \subset \Gamma_0$ and $[v(\kappa_l), v(\kappa_{l+1})] \subset \Gamma_1$ (see Figure 3).

We then extend the vacuum wavefunction $\Psi^{(0)}(P, \vec{t})$ to a meromorphic function $\Psi(P, \vec{t})$ on $P \in \Gamma \setminus \{P_0\}$ with an $(M-1)$ -point divisor in the intersection of Γ_1 with the finite ovals $\Omega_{1,l}$, $l \in [M-1]$. We also set the value of $\Psi(Q_1, \vec{t})$ to the heat hierarchy solution $f^{(1)}(\vec{t})$ so that the Darboux transformation

$$D^{(1)} = \partial_x - \frac{\partial_x f^{(1)}(\vec{t})}{f^{(1)}(\vec{t})}$$

creates a non-effective divisor with one zero at Q_1 for all \vec{t} . Then, after normalization, the effective KP divisor satisfies both the reality and regularity conditions in [16] and Sato's constraints.

Without loss of generality, let us choose the representative matrix $[A] \in Gr^{\text{TP}}(1, M)$ which satisfies $\sum_{l=1}^M A_l = 1$. Then, the vacuum wavefunction on $\Gamma = \Gamma_0 \sqcup \Gamma_1$ is defined as follows

$$(13) \quad \Psi(\zeta, \vec{t}) = \begin{cases} \Psi^{(0)}(\zeta; \vec{t}) = e^{\theta(\zeta; \vec{t})}, & \zeta \in \Gamma_0, \\ \Psi^{(1)}(\zeta; \vec{t}) = \frac{\sum_{l=1}^M A_l E_l(\vec{t}) \prod_{j \neq l}^M (\zeta - \kappa_j)}{\prod_{r=1}^{M-1} (\zeta - b_r)}, & \zeta \in \Gamma_1 \end{cases}$$

with

$$\theta \equiv \theta(\zeta; \vec{t}) = \sum_{i \geq 1} \zeta^i t_i, \quad E_l(\vec{t}) = e^{\theta(\kappa_l; \vec{t})}, \quad l \in [M],$$

where the vacuum divisor b_j consists of $M-1$ real simple poles on Γ_1 , uniquely defined by the gluing conditions:

$$\Psi^{(0)}(\kappa_l; \vec{t}) = \Psi^{(1)}(\kappa_l; \vec{t}), \quad l \in [M].$$

It is easy to check that

$$\kappa_1 < b_1 < \kappa_2 < b_2 < \kappa_3 < \dots < \kappa_{M-1} < b_{M-1} < \kappa_M.$$

Moreover, by definition,

$$(14) \quad \Psi^{(1)}(Q_1; \vec{t}) = \lim_{\zeta \rightarrow \infty} \Psi^{(1)}(\zeta; \vec{t}) = \sum_{l=1}^M a_l E_l(\vec{t}) = f^{(1)}(\vec{t}), \quad \forall \vec{t},$$




FIGURE 3. The rational degeneration of a hyperelliptic curve is associated to soliton data in $Gr^{TP}(1, M)$. Left: the vacuum divisor. Right: a possible effective divisor after the Darboux transformation.

is the heat hierarchy solution generating the Darboux transformation $D^{(1)}$ for the given soliton data. Then, it is immediate to verify that the KP wavefunction

$$(15) \quad \tilde{\Psi}(\zeta, \vec{t}) = \frac{D^{(1)}\Psi(\zeta, \vec{t})}{D^{(1)}\Psi(\zeta, \vec{0})} = \begin{cases} \tilde{\Psi}^{(0)}(\zeta; \vec{t}) = \frac{\zeta - \gamma_1^{(0)}(\vec{t})}{\zeta - \gamma_1^{(0)}} e^{\theta(\zeta; \vec{t})}, & \zeta \in \Gamma_0, \\ \tilde{\Psi}^{(1)}(\zeta; \vec{t}) = \frac{\sum_{1 \leq j < l \leq M} A_l A_j (\kappa_l - \kappa_j)^2 E_l(\vec{t}) E_j(\vec{t}) \prod_{s \neq j, l}^M (\zeta - \kappa_s)}{f^{(1)}(\vec{t}) \sum_{1 \leq j < l \leq M} A_l A_j (\kappa_l - \kappa_j)^2 \prod_{s \neq j, l}^M (\zeta - \kappa_s)}, & \zeta \in \Gamma_1, \end{cases}$$

has effective divisor $\mathcal{D}^{(1)} = \{\gamma_1^{(0)}; \gamma_1^{(1)}, \dots, \gamma_{M-2}^{(1)}\}$, which satisfies the required conditions. Indeed the following Lemma holds.

Lemma 1. [1] Let $\kappa_1 < \dots < \kappa_M$, $[A] \in Gr^{TP}(1, M)$ as above, and let $\tilde{\Psi}(\zeta, \vec{t})$ be as in (15). Then:

- (1) the restriction of the divisor $\mathcal{D}^{(1)}$ on Γ_0 consists of one pole: $\mathcal{D}^{(1)} \cap \Gamma_0 = \{\gamma_1^{(0)}\}$, where $\gamma_1^{(0)} = \frac{\sum_{j=1}^M A_j \kappa_j}{\sum_{j=1}^M A_j}$;
- (2) $\mathcal{D}^{(1)} \cap \Gamma_1 = \{\gamma_1^{(1)}, \dots, \gamma_{M-2}^{(1)}\}$, where $\gamma_r^{(1)}$, $r \in [M-2]$, are the real simple roots of $\sum_{1 \leq j < l \leq M} A_l A_j (\kappa_l - \kappa_j)^2 \prod_{s \neq j, l}^M (\zeta - \kappa_s) = 0$;
- (3) each finite oval contains exactly one pole and no pole is in the infinite oval, $\#(\mathcal{D}^{(1)} \cap \Omega_{1,l}) = 1$, for any $l \in [M-1]$, and $\mathcal{D}^{(1)} \cap \Omega_0 = \emptyset$.

Moreover, if γ_0 coincides with a double point, that is for some $\bar{l} \in [2, M-1]$, $\gamma_1^{(0)} = \kappa_{\bar{l}}$, then also $\gamma_{\bar{l}-1}^{(1)} = \kappa_{\bar{l}}$ and $\mathcal{D}^{(1)} \cap ([\kappa_{\bar{l}-1}, \kappa_{\bar{l}+1}] \setminus \{\kappa_{\bar{l}}\}) = \emptyset$.

We remark that the case in which $\gamma_0^{(1)}$ coincides with a double point is not generic and, for any $\epsilon > 0$ there exists $\vec{t}_0 = (x_0, y_0, t_0, 0, \dots)$ with $\|\vec{t}_0\| < \epsilon$ such that $\gamma_1^{(0)}(\vec{t}_0) \neq \kappa_{\bar{l}}$. Finally, $\mathcal{D}^{(1)} \cap \Omega_0 = \emptyset$ implies that, under our hypotheses, the pole divisor satisfies a stronger conditions than in [39], that is

$$\gamma_1^{(0)}, \gamma_l^{(1)} \in]\kappa_1, \kappa_M[, \quad \forall l \in [M-2].$$

If we start from the vacuum wavefunction as in (13) and apply the Darboux transformation $D^{(N)}$ associated to a generic point in $Gr^{TP}(N, M)$, we obtain an effective divisor which coincides with Sato's on Γ_0 and consists of further $M-1$ distinct poles on Γ_1 . In [1] one of us (S.A.) has characterized the $(N-1)$ -dimensional variety of soliton data in $Gr^{TP}(N, M)$ such that the Darboux transformation $D^{(N)}$ generates a non-effective divisor with a zero of order N in Q_1 , so that the effective divisor has $M-N-1$ real simple poles on Γ_1 and is compatible with Dubrovin and Natanzon conditions (see Remark 2). Such multi-line solitons are naturally related to the




FIGURE 4. Left: The real part of Γ for $M = 6$, $N = 3$. Right: The 10 real ovals of Γ for $M = 6$, $N = 3$.

finite Toda lattice (see [1] and references therein) and correspond to a well-defined immersion of $Gr^{TP}(1, M) \hookrightarrow Gr^{TP}(N, M)$, for any fixed $N \in [M - 1]$. In particular, in [1] it is remarked that the KP vacuum divisor b_1, \dots, b_{M-1} of the KP vacuum wavefunction (13) on Γ_1 coincides with the Toda divisor in [34] upon identification of the KP phases with the Toda spectrum and of the soliton datum $[A] \in Gr^{TP}(1, M)$ in the Toda IVP.

3.2. The algebraic-geometrical setting for soliton data in $Gr^{TP}(N, M)$. Now we present the main geometric construction of our paper.

Definition 1. The rational spectral curve. Assume that we have the following data

- (1) The fixed numbers M, N , $M > N$.
- (2) A set of M real ordered phases $\kappa_1 < \kappa_2 < \dots < \kappa_M$.

To such data we associate the curve Γ , obtained by gluing $N + 1$ copies of \mathbb{CP}^1 , $\Gamma_0, \Gamma_1, \dots, \Gamma_N$, in the following way:

Denote by ζ the local parameter on each copy of \mathbb{CP}^1 . We have the following marked points:

- (1) On Γ_0 we have M real ordered marked points $\kappa_1 < \kappa_2 < \dots < \kappa_M$ and the infinite point P_0 , $\zeta(P_0) = \infty$.
- (2) On each Γ_r , $r \in [N]$ we have $M - N + 1$ real negative ordered marked points in the local coordinate ζ

$$(16) \quad \lambda_1^{(r)} = 0 > \lambda_2^{(r)} > \dots > \lambda_{M-N+1}^{(r)},$$

$M - N$ real positive ordered marked points in the local coordinate ζ

$$(17) \quad 0 < \alpha_2^{(r)} < \alpha_3^{(r)} < \dots < \alpha_{M-N+1}^{(r)},$$

and the infinite point Q_r , $\zeta^{-1}(Q_r) = 0$.

To such data we associate the following set of gluing rules:

- (1) We glue $\lambda_l^{(1)} \in \Gamma_1$ to $\kappa_{N+l-1} \in \Gamma_0$, for $l \in [M - N + 1]$;
- (2) For any $r \in [2, N]$, we glue $\lambda_1^{(r)} \in \Gamma_r$ to $\kappa_{N-r+1} \in \Gamma_0$;
- (3) For any $r \in [2, N]$, and for all $l \in [2, M - N + 1]$, we glue $\lambda_l^{(r)} \in \Gamma_r$ to $\alpha_l^{(r-1)} \in \Gamma_{r-1}$.

Let us recall that we have the standard complex conjugation on each copy of \mathbb{CP}^1 , and, taking into account that we glue only real points, we obtain a complex conjugation on Γ . It is easy to check that we construct a connected reducible curve whose real part possesses the needed




FIGURE 5. Left: Ovals passing through double points in agreement with the planar diagram. Right: Oval passing the double points not in agreement with planar diagram.

number of ovals. In Proposition 1 we characterize the topological properties of Γ and we show the gluing rules and the ovals structure in Figure 4 for the case $N = 3$, $M = 6$.

Proposition 1. *Let $\Gamma = \Gamma_0 \sqcup \Gamma_1 \sqcup \dots \sqcup \Gamma_N$ be the connected reducible curve of Definition 1. Then the real part of Γ which we denote $\Gamma_{\mathbb{R}}$ possesses $1 + (M - N)N$ ovals and each oval is topologically equivalent to a circle. Each double point of Γ is a common point to exactly a pair of ovals. Let us denote Ω_0 the oval containing the infinity point $P_0 \in \Gamma_0$ (we call this oval infinite), and $\Omega_{r,n}$, $r \in [N]$, $n \in [M - N]$ be the remaining $(M - N) \times N$ (finite) ovals.*

Then $Q_r \in \Omega_0$, $r \in [N]$, $\Omega_{r,k}$ are defined by the following properties:

- (1) For $r = 1$, $n \in [M - N]$,

$$\begin{aligned} \Omega_{1,n} \cap \Gamma_0 &= [\kappa_{N+n-1}, \kappa_{N+n}], & \Omega_{1,n} \cap \Gamma_1 &= [\lambda_{n+1}^{(1)}, \lambda_n^{(1)}]; \\ \Omega_{1,n} \cap \Gamma_r &= \emptyset, & r &\in [2, N]; \end{aligned}$$

- (2) For $r \in [2, N]$

$$\begin{aligned} \Omega_{r,1} \cap \Gamma_0 &= [\kappa_{N-r+1}, \kappa_{N-r+2}], & \Omega_{r,1} \cap \Gamma_{r-1} &= [\lambda_1^{(r-1)}, \alpha_2^{(r-1)}], \\ \Omega_{r,1} \cap \Gamma_r &= [\lambda_2^{(r)}, \lambda_1^{(r)}], & \Omega_{r,1} \cap \Gamma_j &= \emptyset, \quad \forall j \in [N] \setminus \{0, r-1, r\}; \end{aligned}$$

- (3) For $r \in [2, N]$ and $n \in [2, M - N]$,

$$\begin{aligned} \Omega_{r,n} \cap \Gamma_{r-1} &= [\alpha_n^{(r-1)}, \alpha_{n+1}^{(r-1)}], & \Omega_{r,n} \cap \Gamma_r &= [\lambda_{n+1}^{(r)}, \lambda_n^{(r)}], \\ \Omega_{r,n} \cap \Gamma_j &= \emptyset, & \forall j &\in [N] \setminus \{r-1, r\}; \end{aligned}$$

Proof. The real ovals of Γ are defined as the union of the corresponding intervals. The structure of the ovals can be easily determined from the gluing law settled in Definition 1. \square

Remark 4. *Our curve can be obtained as a degeneration of a smooth compact Riemann surface of genus $N \times (M - N)$ with an antiholomorphic involution σ having $N \times (M - N) + 1$ fixed ovals. Therefore this smooth Riemann surface is an M-curve (see [44]), and it is natural to treat Γ as a reducible rational M-curve.*

Remark 5. *In our representation of the real part of our curve as a planar diagram, we use the following rule: the ovals pass the double points in such a way that they do not intersect each other on the diagram (see Fig. 5). Moreover, a perturbation is admissible if it respects the antiholomorphic involution σ , and if it preserves the number of ovals (see Fig. 6). For example, in the $Gr^{TP}(1, M)$ case, we assume, that after perturbation the double points split into pairs of **real** points. The perturbations splitting double points into pairs of complex conjugate point also respect the antiholomorphic involution, but they reduce the number of ovals and are forbidden.*

In Section 7 we represent rational degenerations of smooth M-curves corresponding to the soliton data in $Gr^{TP}(2, 4)$ as partial normalizations of reducible algebraic plane nodal curve whose irreducible components are rational (for necessary definitions see [3]). We also show that such reducible plane curve is the rational degeneration of a smooth M-curve of genus 4.




FIGURE 6. Left: A curve with double points and a pair of ovals passing through the double point. Middle: An admissible perturbation of the curve: the number of ovals is preserved. Right: This perturbation is not admissible since two ovals merge.

Let us now describe the analytic properties of the vacuum wave function. Let us introduce the following notation: the restriction of Ψ to the component Γ_r is denoted by $\Psi^{(r)}$; if $P \in \Gamma_r$ and $\zeta = \zeta(P)$, then

$$(18) \quad \Psi(\zeta, \vec{t}) \Big|_{\Gamma_r} = \Psi^{(r)}(\zeta, \vec{t}).$$

Definition 2. The vacuum wave function. Assume that we have the following data

- (1) The reducible M -curve Γ as in Definition 1.
- (2) A collection of $N \times (M - N)$ real divisor points $b_k^{(r)} \in \Gamma_r$, $r \in [N]$, $k \in [M - N]$ such that $\lambda_{k+1}^{(r)} < b_k^{(r)} < \lambda_k^{(r)}$.

To such data we associate the vacuum wave function $\Psi(P, \vec{t})$, where P is a point of the curve Γ , and \vec{t} is the vector of KP times (we may use the standard assumption that only a finite number of $t_j \neq 0$) with the following analytic properties:

- (1) $\Psi^{(0)}(\zeta, \vec{t})$ is the Sato vacuum KP wave function:

$$(19) \quad \Psi^{(0)}(\zeta, \vec{t}) = e^\theta,$$

where

$$(20) \quad \theta \equiv \theta(\zeta, \vec{t}) = \sum_{i \in [\infty]} \zeta^i t_i, \quad t_1 = x, \quad t_2 = y, \quad t_3 = t, \dots$$

- (2) For any fixed collection of KP times \vec{t} the function $\Psi^{(r)}(\zeta, \vec{t})$, $r > 0$ is meromorphic in ζ on Γ_r , and its divisor is exactly $b_1^{(r)}, \dots, b_{M-N}^{(r)}$.
- (3) **Matching rules at the double points:** at all double points of Γ obtained by gluing a pair of points $P_1 \in \Gamma_i$ and $P_2 \in \Gamma_j$ the wave function has the same value for all \vec{t} : $\Psi(P_1, \vec{t}) \equiv \Psi(P_2, \vec{t})$. More precisely,
 - (a) For $r = 1$ and $l \in [M - N + 1]$ the values of $\Psi^{(1)}$ at the point $\lambda_l^{(1)}$ is equal to the value of $\Psi^{(0)}$ at the point κ_{N+l-1} :

$$(21) \quad \Psi^{(1)}(\lambda_l^{(1)}, \vec{t}) = \Psi^{(0)}(\kappa_{N+l-1}, \vec{t}), \quad \forall \vec{t};$$

- (b) For $r \in [2, N]$ the value of $\Psi^{(r)}$ at the point $\lambda_1^{(r)}$ is equal to the value of $\Psi^{(0)}$ at the point κ_{N-r+1} :

$$(22) \quad \Psi^{(r)}(\lambda_1^{(r)}, \vec{t}) = \Psi^{(0)}(\kappa_{N-r+1}, \vec{t}), \quad \forall \vec{t};$$

- (c) For $r \in [2, N]$ and $l \in [2, M - N + 1]$ the value of $\Psi^{(r)}$ at the point $\lambda_l^{(r)}$, is equal to the value of $\Psi^{(r-1)}$ at the point $\alpha_l^{(r-1)}$:

$$(23) \quad \Psi^{(r)}(\lambda_l^{(r)}, \vec{t}) = \Psi^{(r-1)}(\alpha_l^{(r-1)}, \vec{t}), \quad \forall \vec{t}.$$

Proposition 2. *The properties (1)-(3) uniquely define the function $\Psi(P, \vec{t})$. Moreover, for all $P \in \Gamma \setminus \{P_0, b_j^{(r)}, r \in [N], j \in M - n\}$ the function $\Psi(P, \vec{t})$ is a smooth function in the variables \vec{t} . If, in addition, P is a real point, i.e. $\sigma P = P$, then $\Psi(P, \vec{t})$ is real for real \vec{t} .*

The proof of this statement is standard and we omit it.

Remark 6. *Properties (1), (2) and (3) in Definition 2 immediately impose that the vacuum wave function takes the following form:*

$$(24) \quad \Psi^{(r)}(\zeta, \vec{t}) = \sum_{l=1}^{M-N+1} \hat{B}_l^{(r)} \frac{\prod_{j \neq l} (\zeta - \lambda_j^{(r)})}{\prod_{k=1}^{M-N} (\zeta - b_k^{(r)})} V_l^{(r)}(\vec{t}), \quad r \in [1, N],$$

for some real coefficients $\hat{B}_l^{(r)}$, $l \in [M - N + 1]$, and $b_k^{(r)}$, $k \in [M - N]$ depending only on ξ , where

$$(25) \quad V_l^{(r)}(\vec{t}) = \begin{cases} e^{\theta_{N+l-1}(\vec{t})}, & l \in [M - N + 1], & r = 1 \\ e^{\theta_{N-r+1}(\vec{t})} & l = 1, & r \in [2, N] \\ \Psi^{(r-1)}(\alpha_l^{(r-1)}, \vec{t}) & l \in [2, M - N + 1], & r \in [2, N]. \end{cases}$$

Let us define $f_r(\vec{t}) = \Psi(Q_r, \vec{t})$, $r \in [N]$. Then by applying the Darboux transformation we obtain the dressed (unnormalized) KP wave function, associated to the set f_r of the heat hierarchy solutions. Then its normalization

$$(26) \quad \tilde{\Psi}(P, \vec{t}) = \frac{D^{(N)} \Psi(P, \vec{t})}{D^{(N)} \Psi(P, \vec{0})},$$

is the Baker-Akhiezer function for the KP multisoliton solution used in the Krichever construction in the case of reducible spectral curve. It also satisfy the Dubrovin-Natanzon reality and regularity constraints.

Our goal is to solve the inverse problem: **construct the spectral data from a given soliton datum, consisting of M ordered phases and a point $[A]$ of the totally positive Grassmannian $Gr^{TP}(N, M)$.** Such inverse problem is equivalent to construct a curve and a vacuum divisor such that the functions $\Psi(Q_r, \vec{t})$ form a basis of heat hierarchy solutions generating the Darboux transformation associated with the prescribed point $[A]$.

A solution of this problem is given in the next Section. Assume that we have succeeded in constructing such a curve and a vacuum divisor on it. Let us describe the action of the Darboux transformation in the algebraic-geometrical terms.

Theorem 1. *Assume that we fix the soliton data: a set of M real ordered phases $\mathcal{K} = \{\kappa_1 < \kappa_2 < \dots < \kappa_M\}$ and a point of the totally positive Grassmannian $[\hat{A}] \in Gr^{TP}(N, M)$. If, to such soliton data, we may associate a curve Γ as in Definition 1, a $N(M - N)$ vacuum divisor $b_j^{(r)}$ as in Definition 2 and a representative matrix \hat{A} , such that*

$$(27) \quad \Psi(Q_r, \vec{t}) = \sum_{j \in [M]} \hat{A}_j^{N-r+1} e^{\theta_j(\vec{t})}, \quad r \in [N],$$

then the Darboux transformation $D^{(N)}$ associated to the soliton data $(\mathcal{K}, [\hat{A}])$ generates the following **shift of divisor**: we add N simple zeroes at the points Q_r and N -th order pole at the point P_0 .




FIGURE 7. The KP effective divisor may be described as the following action of the Darboux (dressing) $D^{(N)}$ transformation on the effective vacuum divisor: poles are allowed to move inside the finite ovals in such a way that N poles lie in Γ_0 and $M - N - 1$ poles lie in each copy Γ_r , $r \in [N]$.

Proof. By construction, the Darboux transformed wave function is obtained from the vacuum wave function by applying the N -th order ordinary differential operator $D^{(N)}$ such that $\Psi(Q_r, \vec{t})$, $r \in [N]$, generate its kernel. Therefore

$$(28) \quad D^{(N)}\Psi(Q_r, \vec{t}) \equiv 0 \quad \text{for all } \vec{t}.$$

Property (28) exactly means that we add a simple zero at each Q_r to the divisor. Near the point P_0 we have

$$(29) \quad D^{(N)}\Psi_0(\zeta, \vec{t}) = (\zeta^N + O(\zeta^{N-1}))e^{\theta(\zeta, \vec{t})},$$

therefore an N -order pole is added at P_0 . Since the operator $D^{(N)}$ does not affect the poles $b_j^{(r)}$, $r \in [N]$, $j \in [M - N]$, of Ψ , the proof is complete. \square

Remark 7. We remark that the gluing conditions at all double points are preserved by the Darboux transformation for all \vec{t} .

Let us now characterize the effective divisor (see also Figure 7).

Theorem 2. Under the hypotheses of Theorem 1, the normalized wavefunction

$$(30) \quad \tilde{\Psi}(P, \vec{t}) = \frac{D^{(N)}\Psi(P, \vec{t})}{D^{(N)}\Psi(P, \vec{0})}, \quad P \in \Gamma \setminus \{P_0\}$$

has the following analytic properties:

- (1) **Characterization of the effective divisor:** $\tilde{\Psi}(\zeta, \vec{t})$ is meromorphic for $P \in \Gamma \setminus \{P_0\}$, $\zeta = \zeta(P)$, and it possesses an $N(M - N)$ divisor of real simple poles $\mathcal{D} = \{\gamma_s^{(0)}, \gamma_l^{(r)}\}$, $s \in [N]$, $r \in [N]$, $l \in [M - N - 1]$ independent on \vec{t} and with the following properties:
 - (a) The restriction of $\tilde{\Psi}(\zeta, \vec{t})$ to Γ_0 (we use again notations (18) and denote it $\tilde{\Psi}^{(0)}(\zeta, \vec{t})$) is the normalized Sato KP wave function:

$$(31) \quad \tilde{\Psi}^{(0)}(\zeta, \vec{t}) = \frac{D^{(N)}e^{\theta(\zeta, \vec{t})}}{D^{(N)}e^{\theta(\zeta, \vec{0})}},$$

where $\theta(\zeta, \vec{t})$ is as in (20) and $D^{(N)} = \partial_x^N - w_1(\vec{t})\partial_x^{N-1} - \dots - w_N(\vec{t})$ is the Darboux transformation associated to the soliton data $(\mathcal{K}, [A])$, with $\mathcal{K} = \{\kappa_1 < \dots < \kappa_M\}$.




FIGURE 8. Fig. 8: A small perturbation of a pair of zeroes at a double point.

In particular the divisor of poles restricted to Γ_0 is $\{\gamma_s^{(0)}, s \in [N]\}$ and is the zero set of the characteristic polynomial of $D^{(N)}$ at time $\vec{t} = \vec{0}$, i.e.

$$\zeta^N - w_1(\vec{0})\zeta^{N-1} - \dots - w_N(\vec{0}) = 0$$

- (b) For any fixed collection of KP times \vec{t} and $r \in [N]$ the restriction of $\tilde{\Psi}(\zeta, \vec{t})$ to Γ_r (we use again notations 18) and denote it $\tilde{\Psi}^{(r)}(\zeta, \vec{t})$ is meromorphic in ζ on Γ_r , and its divisor consists of $M - N - 1$ poles $\gamma_1^{(r)}, \dots, \gamma_{M-N-1}^{(r)}$;
- (c) There is exactly one divisor pole in each finite oval: $\#(\mathcal{D} \cap \Omega_{r,l}) = 1$, for all $r \in [N]$, $r \in [M - N]$. Here we use the counting rule, see Definition 3;
- (d) There is no pole in the infinite oval: $\mathcal{D} \cap \Omega_{r,l} = \emptyset$.
- (2) **Matching rules at the double points:** at all double points of Γ obtained by gluing a pair of points $P_1 \in \Gamma_{r_i}$ and $P_2 \in \Gamma_{r_j}$ the wave function has the same value for all \vec{t} : $\tilde{\Psi}(P_1, \vec{t}) \equiv \tilde{\Psi}(P_2, \vec{t})$. More precisely,
 - (a) For $r = 1$ and $l \in [M - N + 1]$ the value of $\tilde{\Psi}^{(1)}$ at the point $\lambda_l^{(1)}$ is equal to the value of $\tilde{\Psi}^{(0)}$ at the point κ_{N+l-1} :

$$(32) \quad \tilde{\Psi}^{(1)}(\lambda_l^{(1)}, \vec{t}) = \tilde{\Psi}^{(0)}(\kappa_{N+l-1}, \vec{t}), \quad \forall \vec{t};$$

- (b) For $r \in [2, N]$ the value of $\tilde{\Psi}^{(r)}$ at the point $\lambda_1^{(r)}$ is equal to the value of $\tilde{\Psi}^{(0)}$ at the point κ_{N-r+1} :

$$(33) \quad \tilde{\Psi}^{(r)}(\lambda_1^{(r)}, \vec{t}) = \tilde{\Psi}^{(0)}(\kappa_{N-r+1}, \vec{t}), \quad \forall \vec{t};$$

- (c) For $r \in [2, N]$ and $l \in [2, M - N + 1]$ the value of $\tilde{\Psi}^{(r)}$ at the point $\lambda_l^{(r)}$ is equal to the value of $\tilde{\Psi}^{(r-1)}$ at the point $\alpha_l^{(r-1)}$:

$$(34) \quad \tilde{\Psi}^{(r)}(\lambda_l^{(r)}, \vec{t}) = \tilde{\Psi}^{(r-1)}(\alpha_l^{(r-1)}, \vec{t}), \quad \forall \vec{t}.$$

Definition 3. (The counting rule) For \vec{t} fixed, we call the divisor $\mathcal{D}(\vec{t})$ of $\tilde{\Psi}(\zeta; \vec{t})$ generic, if no points of $\mathcal{D}(\vec{t})$ lie at the double points of Γ , otherwise we call it non generic. In the non generic case, we have at least a zero (resp. a pole) of $\tilde{\Psi}(P, \vec{t})$ at a double point $P = X$ belonging to a pair of finite ovals, that is $X \in \Gamma_{r_1} \cap \Gamma_{r_2}$, ($r_1 \neq r_2$). In such case, the function $\tilde{\Psi}(P, \vec{t})$ has simple zeroes (resp. simple poles) at X at both the components Γ_{r_1} and Γ_{r_2} , i.e. we have a collision of 2 divisor points $\gamma_{k_1}^{(r_1)} \in \Gamma_{r_1}$ and $\gamma_{k_2}^{(r_2)} \in \Gamma_{r_2}$. Then we use the following counting rule: if we have a pair of divisor points at a double point, then one of them is assigned to the first oval and the other is assigned to the second oval.

The counting rule has the following interpretation. If we have a divisor point at a double point, we may apply a generic small shift of \vec{t} , and we obtain a generic divisor with the property formulated in Theorem 2 (see Figure 8).

4. THE REDUCIBLE M-CURVE AND THE EFFECTIVE DIVISOR FOR SOLITON DATA IN $Gr^{TP}(N, M)$

In this section, in Theorem 4 we associate a spectral curve and an effective divisor to any given soliton datum $[\hat{A}] \in Gr^{TP}(N, M)$ represented by the normalized band matrix \hat{A} as in Definition 5. More precisely, for any fixed sufficiently large positive parameter ξ , we construct such algebraic-geometrical data on the spectral curve $\Gamma(\xi)$ as in Definition 4. The proof of Theorem 4 follows from Theorem 3, where we construct a vacuum wavefunction on $\Gamma(\xi)$ satisfying all the properties settled in Definition 2. We prove Theorem 3 in the next section: such proof is constructive and is divided into an algebraic and an analytic part. In particular, we explicitly relate the elementary Darboux transformations to a sequence of maps $Gr^{TP}(1, M - N + 1) \leftarrow Gr^{TP}(2, M - N + 2) \leftarrow \dots \leftarrow Gr^{TP}(r - 1, M - N + r - 1) \leftarrow Gr^{TP}(r, M - N + r) \leftarrow \dots \leftarrow Gr^{TP}(N, M)$, by constructing a specific upper triangular representative matrix $\hat{A}(\xi)$ for the soliton datum in $Gr^{TP}(N, M)$. Such matrix $\hat{A}(\xi)$ rules the exact behavior of the vacuum wavefunction at all marked points and it is obtained as a perturbation of the band matrix \hat{A} which rules the dominant behavior of the vacuum wavefunction at the same marked points. Finally, by construction, the two matrices represent the same point in $Gr^{TP}(N, M)$: $[\hat{A}(\xi)] = [\hat{A}]$.

Definition 4. *Specification of positions of the double points for the rational spectral curve.* Denote by $\Gamma(\xi)$, $\xi > 1$, the spectral curve from Definition 1 with the following collection of double points: on each Γ_r , $r \in [N]$

$$(35) \quad \lambda_1^{(r)} = 0, \quad \lambda_l^{(r)} = -\xi^{2(l-2)}, \quad l \in [2, M - N + 1],$$

$$(36) \quad \alpha_s^{(r)} = \xi^{2s-5}, \quad s \in [2, M - N + 1].$$

Remark 8. For soliton data in $Gr^{TP}(1, M)$, $M \geq 4$, we remark that $\Gamma(\xi)$ in Definition 4 is not equivalent as a complex variety to the rational degeneration of the hyperelliptic curve Γ used in section 3.1.

Definition 5. To any given element of the totally positive Grassmannian $Gr^{TP}(N, M)$ we associate the unique normalized band matrix:

$$(37) \quad \hat{A} = \begin{bmatrix} \hat{A}_1^1 & \hat{A}_2^1 & \hat{A}_3^1 & \hat{A}_4^1 & \dots & \hat{A}_{M-N+1}^1 & 0 & 0 & \dots & 0 & 0 & 0 \\ 0 & \hat{A}_2^2 & \hat{A}_3^2 & \hat{A}_4^2 & \dots & \hat{A}_{M-N+1}^2 & \hat{A}_{M-N+2}^2 & 0 & \dots & 0 & 0 & 0 \\ 0 & 0 & \hat{A}_3^3 & \hat{A}_4^3 & \dots & \hat{A}_{M-N+1}^3 & \hat{A}_{M-N+2}^3 & \hat{A}_{M-N+3}^3 & \dots & 0 & 0 & 0 \\ \vdots & & & & & \ddots & & & & & & \vdots \\ 0 & 0 & 0 & \dots & 0 & \hat{A}_{N-1}^{N-1} & \dots & \dots & \dots & \hat{A}_{M-2}^{N-1} & \hat{A}_{M-1}^{N-1} & 0 \\ 0 & 0 & 0 & 0 & \dots & 0 & \hat{A}_N^N & \dots & \dots & \hat{A}_{M-2}^N & \hat{A}_{M-1}^N & \hat{A}_M^N \end{bmatrix},$$

$$(38) \quad \sum_{j=r}^{M-N+r} A_j^r = 1, \quad r \in [N],$$

and the following set of heat hierarchy solutions:

$$(39) \quad f^{(N-r+1)}(\vec{t}) = \sum_{j=r}^{M-N+r} \hat{A}_j^r e^{\theta_j}, \quad r \in [N].$$

It is easy to check that all elements \hat{A}_j^i in this representation are positive.

Remark 9. In Appendix [A](#) we review a certain number of properties of the band matrix \hat{A} . In particular, there are two different natural normalizations for banded matrices:

- (1) Normalization [\(38\)](#) is convenient in the finite-gap approach, because it corresponds to the normalization of the wavefunction for effective divisors.
- (2) The normalization $A_i^i = 1$, $i \in [N]$, provides local affine coordinates $x_{r,s}$ on $Gr^{TP}(N, M)$ defined in Proposition [5](#) in Appendix [A](#).

The connection between these two normalizations is very simple:

$$(40) \quad \hat{A}_j^i = \frac{A_j^i}{A_i^i + A_{i+1}^i + \dots + A_{M-N+i}^i}.$$

Remark 10. The positive banded matrices arise in many applications, see, for example, [\[9\]](#), [\[12\]](#).

Theorem 3. (The vacuum wavefunction on $\Gamma(\xi)$) Let the soliton data $\mathcal{K} = \{\kappa_1 < \dots < \kappa_M\}$ and $[\hat{A}] \in Gr^{TP}(N, M)$ be fixed, with \hat{A} as in Definition [5](#). Let $f^{(r)}(\vec{t})$, $r \in [N]$, be the heat hierarchy solutions associated to \hat{A} as in [\(39\)](#). Let $\Gamma(\xi)$ denote the curve from Definition [4](#) for $\xi > 1$.

Then for any sufficiently big ξ there exists a unique collection of divisor points $b_k^{(r)} = b_k^{(r)}(\xi) \in]\lambda_{k+1}^{(r)}, \lambda_k^{(r)}[$ and positive coefficients $\epsilon_j^{(r)} = \epsilon_j^{(r)}(\xi)$, $r \in [2, N]$, $j \in [r-1]$ such that

$$(1) \quad (41) \quad \lim_{\zeta \rightarrow \infty} \Psi^{(1)}(\zeta, \vec{t}) = f^{(1)}(\vec{t}),$$

(2) For $r \in [2, N]$

$$(42) \quad \lim_{\zeta \rightarrow \infty} \Psi^{(r)}(\zeta, \vec{t}) = \frac{f^{(r)}(\vec{t}) + \sum_{j=1}^{r-1} \epsilon_j^{(r)} f^{(j)}(\vec{t})}{1 + \sum_{j=1}^{r-1} \epsilon_j^{(r)}}.$$

(3) At all double points the matching conditions of Definition [2](#) are fulfilled.

Moreover, if $\xi \rightarrow +\infty$, then all the parameters $\epsilon_j^{(r)}$ tend to 0.

Remark 11. The proof of Theorem [3](#) provides us with a small perturbation of the band matrix \hat{A} , $\hat{A}(\xi)$ which represents the same point in $Gr^{TP}(N, M)$ as \hat{A} : $[\hat{A}(\xi)] = [\hat{A}]$. Indeed the following set of solutions of the heat hierarchy is uniquely associated to [\(42\)](#)

$$(43) \quad f_\xi^{(r)}(\vec{t}) \equiv \frac{f^{(r)}(\vec{t}) + \sum_{j=1}^{r-1} \epsilon_j^{(r)} f^{(j)}(\vec{t})}{1 + \sum_{j=1}^{r-1} \epsilon_j^{(r)}} = \sum_{j=N-r+1}^M \hat{A}(\xi)_j^{N-r+1} e^{\theta_j}, \quad r \in [N].$$

Then matrix $\hat{A}(\xi)$ is upper triangular by construction and it is defined starting from \hat{A} in the following way: for each $r \in [N]$, the $(N-r+1)$ -th row of $\hat{A}(\xi)$ is the linear combinations of the last r rows of \hat{A} with the positive coefficients $\epsilon_k^{(r)}(\xi)$, that is

$$\hat{A}(\xi)_j^{N-r+1} = \hat{A}_j^{N-r+1} + \sum_{k=1}^{r-1} \epsilon_k^{(r)}(\xi) \hat{A}_j^{N-k+1}, \quad j \in [M].$$

The proof of this Theorem is provided in the section [5](#). We end this section with the Theorem which summarizes the properties of the effective divisor of the normalized KP wavefunction $\tilde{\Psi}$ which is obtained dressing the vacuum wavefunction in Theorem [3](#).

Theorem 4. (The effective divisor on Γ_ξ) Let $\xi \gg 1$, $1 \leq N < M$ be fixed and let $\Gamma = \Gamma(\xi)$ be as in Definition 1. Let the soliton data $\mathcal{K} = \{\kappa_1 < \dots < \kappa_M\}$ and $[\hat{A}] \in Gr^{TP}(N, M)$ be fixed and let $\Psi(\zeta, \vec{t})$ be the corresponding vacuum wavefunction on Γ satisfying Theorem 3. Let $D^{(N)} = \partial_x^N - w_1(\vec{t})\partial_x^{N-1} - \dots - w_N(\vec{t})$ be the Darboux (dressing) transformation associated to the soliton data. Then the effective divisor \mathcal{D}_ξ of the normalized dressed wavefunction

$$\tilde{\Psi}(\zeta, \vec{t}) = \frac{D^{(N)}\Psi(\zeta, \vec{t})}{D^{(N)}\Psi(\zeta, \vec{0})}$$

has degree $N(M - N)$ and satisfies the reality and regularity conditions.

More precisely let us denote $\mathcal{D}^{(0)} = \{\gamma_k^{(0)}, k \in [N]\}$ the set of solutions to (12)

$$(\gamma_l^{(0)})^N - w_1(\vec{0})(\gamma_l^{(0)})^{N-1} - \dots - w_{N-1}(\vec{0})(\gamma_l^{(0)}) - w_N(\vec{0}) = 0, \quad l \in [N]$$

and $\mathcal{D}_\xi^{(r)} = \{\gamma_l^{(r)}(\xi), l \in [M - N - 1]\}$ the set of solutions to

$$D^{(N)}\Psi^{(r)}(\zeta, \vec{0}) = 0, \quad r \in [N].$$

Then $\mathcal{D}_\xi = \mathcal{D}^{(0)} \cup \mathcal{D}_\xi^{(1)} \cup \dots \cup \mathcal{D}_\xi^{(N)}$ and it has the following properties:

- (1) $\mathcal{D}_\xi \cap \Gamma_0 = \mathcal{D}^{(0)}$ and all points $\gamma_k^{(0)}$ lying in Γ_0 are pairwise different;
- (2) For any fixed $r \in [N]$, $\mathcal{D}_\xi \cap \Gamma_r = \mathcal{D}_\xi^{(r)}$ and all points $\gamma_k^{(r)}(\xi)$ lying in Γ_r are pairwise different;
- (3) $\mathcal{D}_\xi \cap \Omega_0 = \emptyset$;
- (4) $\mathcal{D}_\xi \subset \bigcup_{r,j} \Omega_{r,j}$, that is each $\gamma_k^{(r)}(\xi)$ is real and lies in some finite oval;
- (5) Each finite oval $\Omega_{r,j}$ contains exactly one point of \mathcal{D} according to the counting rule from Definition 3.

The proof of Theorem 4 follows immediately from Theorems 1, 2 and 3.

Remark 12. Replacing the double points of the reduced curve Γ by thin handles, we pass from real bounded regular solitons to real bounded regular quasiperiodic finite-gap solutions to the KP equations. Moreover, in a neighborhood of the point $[A]$ we can work with a fixed curve Γ and we have a local bijection between the admissible divisors and an open subset in $Gr^{TP}(N, M)$, containing $[A]$.

5. PROOF OF THEOREM 3

The main ideas of the proof. The proof of the Theorem 3 may be divided in and algebraic and an analytic part.

In the first part of the proof we present a recursive algebraic model of the principal cell of $Gr^{TP}(N, M)$, providing a birational map of this cell to the positive octant of $\mathbb{R}^{N(M-N)}$. In fact this construction can be treated as a special case of the Whitney theorem [55]. We start defining natural projections: $\pi_N : Gr^{TP}(N, M) \rightarrow Gr^{TP}(N-1, M-1)$. If \hat{A} is the **band** matrix of Definition 5 representing the class $[\hat{A}] \in Gr^{TP}(N, M)$, then, removing the first row and the first column of \hat{A} we obtain a point in $Gr^{TP}(N-1, M-1)$, represented by a $(N-1) \times (M-1)$ band matrix with the same normalization of the initial \hat{A} . Therefore we have a chain of maps:

$$(44) \quad Gr^{TP}(N, M) \rightarrow \dots \rightarrow Gr^{TP}(r, M-N+r) \rightarrow Gr^{TP}(r-1, M-N+r+1) \rightarrow \dots \rightarrow Gr^{TP}(1, M-N+1),$$

which we use to settle a recursive construction of the totally positive band matrix \hat{A} starting from the last row and moving up (Theorem 5). The key point in such construction is Lemma 2

where, to the r -th row of \hat{A} , we associate a minimal set of $M - N$ vectors with non-negative entries with the following property: **the band matrix obtained by adding the $(r - 1)$ -th row is totally positive if and only if the row $r - 1$ is a linear combination with positive coefficients of these basic vectors and of the corresponding pivot vector.** The recursive procedure of Theorem 5 for the given band matrix \hat{A} then follows from the following two properties: 1) at any fixed step, the positive coefficients in Lemma 2 are uniquely defined by the entries of the r -th row of \hat{A} , and 2) the minimal set of vectors associated with the $(r - 1)$ -th row is computed starting from the set of vectors associated with the r -th row via an explicit transition matrix.

The second part of the proof uses the following idea: assume that we have a Riemann sphere with $M - N$ divisor points, $M - N + 1$ “input” points and $M - N$ “output” points. Then the value of the wave function at each output point is a linear combination of the values of the wave function at the input points. The coefficients of this transition matrix depend on the positions of the points and the divisor. **We show that the transition matrices arising in Theorem 5 admit arbitrary good approximation by transition matrices associated with Riemann spheres by choosing proper marked points.** Therefore we can construct a family of spectral curves with divisors such that the asymptotics of the wave function at double points for large ξ is ruled by the algebraic construction of the first part of the proof.

Moreover, this approximation is corrected at each step so that we obtain exact matching of the vacuum wavefunction at the double points and control its value at the Darboux points. In particular, we prove that $\Psi^{(r)}(\zeta, \vec{t})$ is positive if $\zeta > 0$ for all \vec{t} . Thus, the zeroes of the vacuum wave function are located at the required positions.

5.1. The algebraic part. In this section \mathbb{R}_+^n denotes the positive octant in the space \mathbb{R}^n : $x_j > 0$, $j \in [n]$. Let us also remark that the positive octant in \mathbb{R}^n is naturally isomorphic to a collection of $n + 1$ positive numbers $\{\hat{B}_1, \hat{B}_2, \dots, \hat{B}_{n+1}\}$ such that

$$\hat{B}_1 + \hat{B}_2 + \dots + \hat{B}_{n+1} = 1.$$

$$(45) \quad \hat{B}_1 = \frac{1}{1 + x_1 + \dots + x_n}, \quad \hat{B}_{j+1} = \frac{x_j}{1 + x_1 + \dots + x_n}, \quad j \in [n].$$

Let us start from $N = 1$. In the case of $Gr^{TP}(1, M)$ we have a very simple characterization: the matrix \hat{A} is totally positive if and only if it may be represented as:

$$(46) \quad \hat{A} = \hat{B}_1 \cdot [1, 0, 0, \dots, 0] + \hat{B}_2 \cdot [0, 1, 0, \dots, 0] + \dots + \hat{B}_M \cdot [0, 0, 0, \dots, 1],$$

where all \hat{B}_i are positive. We fix the unique representative in $[\hat{A}]$ by assuming:

$$(47) \quad \hat{B}_1 + \hat{B}_2 + \dots + \hat{B}_M = 1.$$

Formulas (45)-(47) define a birational map $\mathbb{R}_+^{(M-1)} \rightarrow Gr^{TP}(1, M)$.

In the next Lemma we express the first row of the band matrix \hat{A} as a linear combination with positive coefficients of a minimal set of $M - N + 1$ vectors whose non-negative entries depend only on the $(N - 1) \times (N - 1)$ minors of the band matrix obtained eliminating the first row and the first column of \hat{A} .

Lemma 2. (Principal Algebraic Lemma). *Assume that \hat{A} is an $N \times M$ matrix in banded form such that $\hat{A}_i^i > 0$, $\hat{A}_j^i = 0$, if and only if $j < i$ or $j > M - N + i$ and $\sum_{j=1}^M \hat{A}_j^i = 1$ for all $i \in [N]$. Assume also that \hat{A} has the following property: after removing the first row and the first column from \hat{A} we obtain a matrix $\hat{A}^{(0)}$ such that $[\hat{A}^{(0)}] \in Gr^{TP}(N - 1, M - 1)$.*

Then $[\hat{A}] \in Gr^{TP}(N, M)$ if and only if there exist $\hat{B}_n > 0$, $n \in [M - N + 1]$, such that $\sum_{n=1}^{M-N+1} \hat{B}_n = 1$ and the first line of \hat{A} can be represented in the following form

$$(48) \quad [\hat{A}_1^1, \hat{A}_2^1, \dots, \hat{A}_{M-N+1}^1, 0, \dots, 0] = \sum_{n=1}^{M-N+1} \hat{B}_n \hat{E}^n,$$

where \hat{E}^n denotes the following collection of vectors

$$(49) \quad \hat{E}^1 = [1, 0, 0, \dots, 0], \quad \hat{E}^n = [0, E_2^n, E_3^n, \dots, E_n^n, 0, \dots, 0], \quad n \in [2, M - N + 1],$$

$$E_j^n = \frac{\Delta_{[j, n+1, \dots, n+N-2]}}{\sum_{s=2}^n \Delta_{[s, n+1, \dots, n+N-2]}}, \quad j \in [2, n].$$

Moreover, in such case

$$(50) \quad \hat{B}_n = \begin{cases} \hat{A}_1^1, & n = 1, \\ \frac{\Delta_{[n, \dots, n+N-1]} (\sum_{s=2}^n \Delta_{[s, n+1, \dots, n+N-2]})}{\Delta_{[n, \dots, n+N-2]} \Delta_{[n+1, \dots, n+N-1]}}, & n \in [2, M - N + 1]. \end{cases}$$

Proof. To start with, let us present an alternative definition of the vectors \hat{E}^n . Consider the matrix $\hat{A}_{[2, 3, \dots, n+N-2]}^{[2, 3, \dots, N]}$ obtained from \hat{A} by taking the consequent columns 2, 3, \dots , $n+N-2$ and removing the first row. Using the same Gaussian elimination process as above we can transform $\hat{A}_{[2, 3, \dots, n+N-2]}^{[2, 3, \dots, N]}$ to the banded form. Denote by \mathcal{A}'^n the vector obtained from the first row of the banded form of $\hat{A}_{[2, 3, \dots, n+N-2]}^{[2, 3, \dots, N]}$ by adding one zero at the left-hand side and $M - N - n + 2$ zeroes at the right-hand side.

The Gaussian elimination does not affect $(n-1)$ -order minors, formed by the last $n-1$ rows and arbitrary columns. Therefore we have the following formulas:

$$(\mathcal{A}'^n)_j = \begin{cases} \frac{\Delta_{[j, n+1, \dots, n+N-2]}}{\Delta_{[n+1, \dots, n+N-2]}} & j \in [2, n], \\ 0 & j = 1 \text{ or } j > n+1, \end{cases}$$

$$\text{and } \hat{E}^n = (\mathcal{A}'^n) \cdot \left(\sum_{j=2}^n (\mathcal{A}'^n)_j \right)^{-1}.$$

Let us replace the first row of \hat{A} by \hat{E}^n , and denote the new matrix by $\tilde{A}^{(n)}$. Denote by $\tilde{A}_{[j, j+1, \dots, j+N-1]}^{(n)}$ the $N \times N$ submatrices of $\tilde{A}^{(n)}$ formed by N consecutive columns starting from the column j , $j \geq 2$. If $j \leq n-1$, the first row of $\tilde{A}_{[j, j+1, \dots, j+N-1]}^{(n)}$ is a linear combination of the other rows, and $\det(\tilde{A}_{[j, j+1, \dots, j+N-1]}^{(n)}) = 0$. If $j > n$, all elements of the first row are equal to 0 and $\det(\tilde{A}_{[j, j+1, \dots, j+N-1]}^{(n)}) = 0$. If $j = n$, the matrix $\tilde{A}_{[j, j+1, \dots, j+N-1]}^{(n)}$ is lower-triangular, therefore we have

$$\det(\tilde{A}_{[j, j+1, \dots, j+N-1]}^{(n)}) = \delta_j^n \cdot \frac{\Delta_{[n, n+1, \dots, n+N-2]} \Delta_{[n+1, n+2, \dots, n+N-1]}}{\left(\sum_{s=2}^n \Delta_{[s, n+1, \dots, n+N-2]} \right)},$$

where $n, j \in [2, M - N + 1]$, and δ_j^i denotes the standard Kronecker symbol. As a corollary we immediately obtain that the vectors \hat{E}^n , $n \in [0, M - N]$ are linearly independent, therefore any vector with zero elements in the positions $M - N + 2, M - N + 3, \dots, M$ can be uniquely represented as a linear combination of these vectors.

Denote by \tilde{A} the matrix, obtained from \hat{A} by replacing the first row with the vector, defined by the formula (48). Then

$$\Delta_{[n,n+1,\dots,n+N-1]}(\tilde{A}) = \hat{B}_n \cdot \frac{\Delta_{[n,n+1,\dots,n+N-2]} \Delta_{[n+1,n+2,\dots,n+N-1]}}{\left(\sum_{s=2}^n \Delta_{[s,n+1,\dots,n+N-2]}\right)},$$

where $n > 0$,

$$\Delta_{[1,2,\dots,N]}(\tilde{A}) = \Delta_{[1,2,\dots,N]}.$$

Therefore we have $\tilde{A} = \hat{A}$ if and only if \hat{B}_n are defined by (50) for $n > 1$, $\hat{B}_1 = \hat{A}_1^1$.

If all minors of the matrix \hat{A} are strictly positive, then all $\hat{B}_n > 0$. Conversely, if all $\hat{B}_n > 0$, then all N -order minors of \hat{A} formed by consequent columns are strictly positive, and applying Lemma 6 we obtain that all N -order minors are strictly positive. It completes the proof. \square

Corollary 1. *Assume that the matrix \hat{A} is the same as in the Principal Algebraic Lemma, and all $N \times N$ minors of \hat{A} are strictly positive. Denote by \check{A} the matrix, obtained from \hat{A} by removing the last s columns, $s < M - N$. If we apply to \check{A} the same procedure as in the Principal Algebraic Lemma, we obtain a collection of vectors \check{E}^n , $n = 1, 2, \dots, M - N - s + 1$, where \check{E}^n is obtained from \hat{E}^n by removing the last s zeroes.*

Proof. The proof follows directly from the following property of the collection \hat{E}^n : to define the element \hat{E}^n it is sufficient to know the first $N + n - 2$ columns of the matrix \hat{A} . \square

In Lemma 2 the bases \hat{E}^n are provided by explicit formulas. However, our construction of the vacuum wavefunction for a given spectral curve requires a recursive procedure for calculating the vectors \hat{E}^n , since such vectors rule the large ξ asymptotic behavior of the vacuum wave function at all double points and Darboux points. Next Theorem is the main result of this subsection and it provides such recursive algebraic construction.

In Theorem 5 to each row of \hat{A} we associate a unique collection of $M - N + 1$ normalized vectors $\hat{E}^{(r)[j]}$, $j \in [M - N + 1]$ with non-negative entries, such that this row is expressed as a linear combination of these vectors with positive coefficients $\hat{B}_l^{(r+1)} > 0$, $r \in [0, N - 1]$, $l \in [M - N + 1]$. For each row we organize such a collection of vectors as an $(M - N + 1) \times M$ matrix $\hat{E}^{(r)}$. We show that matrices $\hat{E}^{(r)}$ can be computed using a simple recursive procedure.

Theorem 5. (The recursive algebraic construction). *Let $N < M$ and let \hat{A} be the totally non-negative $N \times M$ band matrix as in Definition 5 representing a given point $[\hat{A}] \in Gr^{TP}(N, M)$.*

For any $r \in [0, N - 1]$ let us define a normalized $(M - N + 1) \times M$ matrix $\hat{E}^{(r)}$ with non-negative entries of the form:

$$\hat{E}^{(r)} = \begin{bmatrix} \hat{E}^{(r)[1]} \\ \hat{E}^{(r)[2]} \\ \vdots \\ \hat{E}^{(r)[M-N+1]} \end{bmatrix}, \quad \hat{E}^{(r)[j]} = [(\hat{E}^{(r)[j]})_1, (\hat{E}^{(r)[j]})_2, \dots, (\hat{E}^{(r)[j]})_M],$$

and normalization $\sum_{s=1}^M (\hat{E}^{(r)[j]})_s = 1$, by the following formulas:

- (1) For $r = 0$ the matrix $\hat{E}^{(0)}$ is defined by: $(\hat{E}^{(0)[l]})_j = \delta_j^{N+l-1}$.

(2) For $r \in [1, N-1]$ the matrix $\hat{E}^{(r)}$ is defined by: $(\hat{E}^{(r)[1]})_j = \delta_j^{N-r}$, and for $n \in [2, M-N+1]$

$$(51) \quad (\hat{E}^{(r)[n]})_j = \begin{cases} 0, & \text{if } j \in [N-r] \\ \frac{\Delta[j; N-r+n, N-r+n+1, \dots, N+n-2]}{\sum_{s=N-r+1}^{N-r+n-1} \Delta[s; N-r+n, N-r+n+1, \dots, N+n-2]}, & \text{if } j \in [N-r+1, N-r+n-1] \\ 0, & \text{if } j \in [N-r+n, M]. \end{cases}$$

For each $r \in [N]$ let us define a collection $\hat{B}_j^{(r)}$, $j \in [M-N+1]$ of positive coefficients as follows:

(1) For $r = 1$ the constants $\hat{B}_j^{(1)}$ are defined by: $\hat{B}_j^{(1)} = \hat{A}_{N+j-1}^N$, $j \in [M-N+1]$.

(2) For $r \in [2, N]$ the constants $\hat{B}_j^{(r)}$ are defined by:

$$(52) \quad \hat{B}_j^{(r)} = \begin{cases} \hat{A}_{N-r+1}^{N-r+1}, & \text{if } j = 1 \\ \frac{\Delta[N-r+j, \dots, N+j-1] (\sum_{s=N-r+2}^{N-r+j} \Delta[s; N-r+j+1, N-r+j+2, \dots, N+j-2])}{\Delta[N-r+j+1, \dots, N+j-1] \Delta[N-r+j, \dots, N+j-2]}, & \text{if } j \in [2, M-N+1] \end{cases}$$

Then the matrices $\hat{E}^{(r)}$ and the coefficients $\hat{B}_j^{(r)}$ have the following properties:

- (1) The constants $\hat{B}_l^{(r)}$ are normalized: $\sum_{s=1}^{M-N+1} \hat{B}_s^{(r)} = 1$.
- (2) For each $r \in [1, N]$ we have:

$$(53) \quad \hat{A}^{[N-r+1]} = \sum_{j=1}^{M-N+1} \hat{B}_j^{(r)} \hat{E}^{(r-1)[j]},$$

- (3) For each $r \in [1, N]$ we have:

$$(54) \quad \hat{E}^{(r)[2, M-N+1]} = \mathcal{B}^{(r)} \hat{E}^{(r-1)},$$

where $\mathcal{B}^{(r)}$ denotes the following $(M-N) \times (M-N+1)$ matrix:

$$(55) \quad \mathcal{B}^{(r)} = \begin{bmatrix} \frac{\hat{B}_1^{(r)}}{\hat{B}_1^{(r)}} & 0 & 0 & \dots & 0 & 0 \\ \frac{\hat{B}_1^{(r)}}{\hat{B}_1^{(r)} + \hat{B}_2^{(r)}} & \frac{\hat{B}_2^{(r)}}{\hat{B}_1^{(r)} + \hat{B}_2^{(r)}} & 0 & \dots & 0 & 0 \\ \frac{\hat{B}_1^{(r)}}{\hat{B}_1^{(r)} + \hat{B}_2^{(r)} + \hat{B}_3^{(r)}} & \frac{\hat{B}_2^{(r)}}{\hat{B}_1^{(r)} + \hat{B}_2^{(r)} + \hat{B}_3^{(r)}} & \frac{\hat{B}_3^{(r)}}{\hat{B}_1^{(r)} + \hat{B}_2^{(r)} + \hat{B}_3^{(r)}} & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & 0 & 0 \\ \frac{\hat{B}_1^{(r)}}{\sum_{l=1}^{M-N} \hat{B}_l^{(r)}} & \frac{\hat{B}_2^{(r)}}{\sum_{l=1}^{M-N} \hat{B}_l^{(r)}} & \frac{\hat{B}_3^{(r)}}{\sum_{l=1}^{M-N} \hat{B}_l^{(r)}} & \dots & \frac{\hat{B}_{M-N}^{(r)}}{\sum_{l=1}^{M-N} \hat{B}_l^{(r)}} & 0 \end{bmatrix}$$

Remark 13. For any fixed $r \in [N]$, the elements of the matrix $\hat{E}^{(r)}$ and the coefficients $\hat{B}_j^{(r)}$ are subtraction free rational expressions in the minors of \hat{A} formed with its last k rows for $k \leq r$, so they may be expressed as subtraction free rational expressions in the elements $x_{l,s}$, $l \in [r]$, $s \in [M-N]$ of the FZ-basis following Corollary [6](#). As a consequence, all the identities in the above Theorem may be expressed in invariant form, that is they are associated to the given point in the Grassmannian and not to the representative matrix \hat{A} .

The proof of the Theorem is based on the following two Lemmas:

Lemma 3. Let \hat{A} be a totally positive $N \times M$ matrix in banded as in Definition 5 and let $s \in [N-1]$ be fixed. Let us define

$$\hat{B}_j = \begin{cases} \hat{A}_s^s, & j = 1 \\ \frac{\Delta_{[s+j-1, \dots, N+j-1]} \cdot (\sum_{k=1}^{j-1} \Delta_{[s+k; s+j, \dots, N+j-2]})}{\Delta_{[s+j-1, \dots, N+j-2]} \Delta_{[s+j, \dots, N+j-1]}}, & j \in [2, M-N+1]. \end{cases}$$

Then

$$\sum_{j=1}^k \hat{B}_j = \frac{\sum_{j=1}^k \Delta_{[s+j-1; s+k, \dots, N+k-1]}}{\Delta_{[s+k, \dots, N+k-1]}}, \quad k \in [M-N+1].$$

In particular

$$\sum_{j=1}^{M-N+1} \hat{B}_j = \frac{\sum_{j=1}^{M-N+1} \Delta_{[s+j-1; M-N+s+1, \dots, M]}}{\Delta_{[M-N+s+1, \dots, M]}} = \sum_{j=s}^{M-N+s} \hat{A}_j^s \equiv 1.$$

Proof. The proof is by induction in l using the minors identity

$$\Delta_{[s+j-1; s+k-1, \dots, N+k-2]} \Delta_{[s+k, \dots, N+k-1]} + \Delta_{[s+k-1, \dots, N+k-1]} \Delta_{[s+j-1; s+k, \dots, N+k-2]} = \Delta_{[s+j-1; s+k, \dots, N+k-1]} \Delta_{[s+k-1, \dots, N+k-2]}$$

where $s < s+j-1 < s+k-1 < N+k-1$.

Indeed for $l = 1$ we just have

$$\hat{B}_1 \equiv \hat{A}_s^s = \frac{\Delta_{[s \dots N]}}{\Delta_{[s+1, \dots, N]}}.$$

Suppose the identity holds for $l = k-1$, then for $l = k$, using the minors identity, we immediately get

$$\begin{aligned} \sum_{l=1}^k \hat{B}_l &= \frac{\Delta_{[s+k-1, \dots, N+k-1]} \sum_{j=1}^{k-1} \Delta_{[s+j; s+k, \dots, N+k-2]}}{\Delta_{[s+k-1, \dots, N+k-2]} \Delta_{[s+k, \dots, N+k-1]}} + \frac{\sum_{j=1}^{k-1} \Delta_{[s+j-1; s+k-1, \dots, N+k-2]}}{\Delta_{[s+k-1, \dots, N+k-2]}} \\ &= \frac{\Delta_{[s; s+k, \dots, N+k-1]} + \Delta_{[s+k-1, \dots, N+k-1]}}{\Delta_{[s+k, \dots, N+k-1]}} + \frac{\sum_{j=2}^{k-1} \Delta_{[s+j-1; s+k, \dots, N+k-1]}}{\Delta_{[s+k, \dots, N+k-1]}} \end{aligned}$$

During this calculation we used the following formula: due to the banded structure of \hat{A} and $k \in [2, M-N+1]$: $\frac{\Delta_{[s; s+k-1, \dots, N+k-2]}}{\Delta_{[s+k-1, \dots, N+k-2]}} = \frac{\Delta_{[s; s+k, \dots, N+k-1]}}{\Delta_{[s+k, \dots, N+k-1]}} = \hat{A}_s^s$. \square

Lemma 4. Let \hat{A} be a totally positive $N \times M$ matrix in banded form as in Definition 5. Let $r \in [N-1]$, $k \in [2, M-N+1]$ and $j \in [N-r+1, N-r+k-1]$. Then we have the following identity

$$(56) \quad \sum_{n=1}^{k-1} \frac{\Delta_{[N-r+n, \dots, N+n-1]} \cdot \Delta_{[j; N-r+n+1, \dots, N+n-2]}}{\Delta_{[N-r+n+1, \dots, N+n-1]} \cdot \Delta_{[N-r+n, \dots, N+n-2]}} = \frac{\Delta_{[j; N-r+k, \dots, N+k-2]}}{\Delta_{[N-r+k, \dots, N+k-2]}}$$

Proof. The proof is again by induction. For $k = 2$ we have $j = N-r+1$, and the identity is trivial. Let $k > 2$, and suppose that for all $2 \leq k' \leq k-2$ the identity has been proven. Then for $j \in [N-r, N-r+k-2]$ we can write

$$\begin{aligned} \sum_{n=1}^{k-1} &\frac{\Delta_{[N-r+n, \dots, N+n-1]} \cdot \Delta_{[j; N-r+n+1, \dots, N+n-2]}}{\Delta_{[N-r+n+1, \dots, N+n-1]} \cdot \Delta_{[N-r+n, \dots, N+n-2]}} = \\ &= \sum_{n=1}^{k-2} \frac{\Delta_{[N-r+n, \dots, N+n-1]} \cdot \Delta_{[j; N-r+n+1, \dots, N+n-2]}}{\Delta_{[N-r+n+1, \dots, N+n-1]} \cdot \Delta_{[N-r+n, \dots, N+n-2]}} + \frac{\Delta_{[N-r+k-1, \dots, N+k-2]} \cdot \Delta_{[j; N-r+k, \dots, N+k-3]}}{\Delta_{[N-r+k, \dots, N+k-2]} \cdot \Delta_{[N-r+k-1, \dots, N+k-3]}} = \\ &= \frac{\Delta_{[j; N-r+k-1, \dots, N+k-3]}}{\Delta_{[N-r+k-1, \dots, N+k-3]}} + \frac{\Delta_{[N-r+k-1, \dots, N+k-2]} \cdot \Delta_{[j; N-r+k, \dots, N+k-3]}}{\Delta_{[N-r+k, \dots, N+k-2]} \cdot \Delta_{[N-r+k-1, \dots, N+k-3]}} = \\ &= \frac{\Delta_{[j; N-r+k-1, \dots, N+k-3]} \cdot \Delta_{[N-r+k, \dots, N+k-2]} + \Delta_{[N-r+k-1, \dots, N+k-2]} \cdot \Delta_{[j; N-r+k, \dots, N+k-3]}}{\Delta_{[N-r+k, \dots, N+k-2]} \cdot \Delta_{[N-r+k-1, \dots, N+k-3]}} = \end{aligned}$$

Applying the minor identity to the numerator, we obtain

$$\begin{aligned} \sum_{n=1}^{k-1} \frac{\Delta_{[N-r+n, \dots, N+n-1]} \cdot \Delta_{[j; N-r+n+1, \dots, N+n-2]}}{\Delta_{[N-r+n+1, \dots, N+n-1]} \cdot \Delta_{[N-r+n, \dots, N+n-2]}} &= \frac{\Delta_{[j; N-r+k, \dots, N+k-2]} \cdot \Delta_{[N-r+k-1, \dots, N+k-3]}}{\Delta_{[N-r+k, \dots, N+k-2]} \cdot \Delta_{[N-r+k-1, \dots, N+k-3]}} = \\ &= \frac{\Delta_{[j; N-r+k, \dots, N+k-2]}}{\Delta_{[N-r+k, \dots, N+k-2]}}. \end{aligned}$$

Assume now that $j = N - r + k - 1$. Then we have only one nonzero term in our sum:

$$\frac{\Delta_{[N-r+k-1, \dots, N+k-2]} \cdot \Delta_{[N-r+k-1; N-r+k, \dots, N+k-3]}}{\Delta_{[N-r+k, \dots, N+k-2]} \cdot \Delta_{[N-r+k-1, \dots, N+k-3]}} = \frac{\Delta_{[N-r+k-1; N-r+k, \dots, N+k-2]}}{\Delta_{[N-r+k, \dots, N+k-2]}}.$$

□

Now we are ready to prove Theorem 5.

Proof. The first item follows immediately from Lemma 3.

The second statement is exactly the Principal Algebraic Lemma, applied to the matrix, obtained from \hat{A} by removing the first $N - r$ rows and the first $N - r$ columns. Applying the formula (49) we immediately notice, indeed, that the vector $\hat{E}^{(r-1)[l]}$ is obtained from \hat{E}^l by adding $N - r$ zeroes from the left.

The last statement follows immediately from the Lemma 4. □

5.2. The analytic part. We do the proof recursively starting from the last line.

Let us introduce the following notation.

$$(57) \quad \psi_n^{(r)}(\vec{t}) = \sum_{j=1}^M \hat{E}_j^{(r)[n]} e^{\theta_j(\vec{t})} = \frac{\sum_{j=N-r+1}^{N-r+n-1} \Delta_{[j; N-r+n, N-r+n+1, \dots, N+n-2]} e^{\theta_j(\vec{t})}}{\sum_{s=N-r+1}^{N-r+n-1} \Delta_{[s; N-r+n, N-r+n+1, \dots, N+n-2]}},$$

where $\hat{E}_j^{(r)[n]}$ are as in Theorem 5. In Proposition 3 we compute the vacuum divisor points on the component Γ_1 , and we check that, at leading order in ξ , the value of the vacuum wave function at each double point coincides with that prescribed by Theorem 5 i.e. we check that $\psi_n^{(0)}(\vec{t})$ and $\psi_n^{(1)}(\vec{t})$ respectively are the leading order behavior at $\lambda_n^{(1)}$ and $\alpha_n^{(1)}$ when $\xi \gg 1$.

Proposition 3. Let $\xi \gg 1$ be fixed, and let $\Gamma = \Gamma(\xi)$ be as in Definition 4, with $\lambda_l^{(1)}$, $l \in [M - N + 1]$, $\alpha_s^{(1)}$, $s \in [2, M - N + 1]$, as in (35) and (36). Let

$$\Psi^{(1)}(\zeta, \vec{t}) = \sum_{l=1}^{M-N+1} \hat{B}_l^{(1)} \frac{\prod_{k \neq l}^{M-N+1} (\zeta - \lambda_k^{(1)})}{\prod_{k=1}^{M-N} (\zeta - b_k^{(1)})} e^{\theta_{N+l-1}(\vec{t})},$$

where $\hat{B}_l^{(1)}$, $b_k^{(1)}$ are unknown parameters. Then the requirements:

- (1) $\lim_{\zeta \rightarrow \infty} \Psi^{(1)}(\zeta, \vec{t}) = \sum_{j=N}^M \hat{A}_j^N e^{\theta_j(\vec{t})}$,
- (2) $\Psi^{(1)}(\lambda_l^{(1)}, \vec{t}) = e^{\theta_{N+l-1}(\vec{t})} > 0$, $l \in [M - N + 1]$

uniquely define the coefficients $\hat{B}_l^{(1)} = \hat{A}_{N+l-1}^N$, $l \in [M - N + 1]$ and the divisor points $b_k^{(1)} = b_k^{(1)}(\xi) \in]\lambda_{k+1}^{(1)}, \lambda_k^{(1)}[$, $k \in [M - N]$. Moreover $\Psi^{(1)}$ has the following properties:

- (1) $\Psi^{(1)}(\zeta, \vec{t}) > 0$, $\forall \zeta > 0$, and for all **real** \vec{t} . In particular, all $\alpha_n^{(1)} > 0$, therefore $\Psi^{(1)}(\alpha_n^{(1)}, \vec{t}) > 0$ for all n and for all real \vec{t} . Moreover, we have the following estimate: for $\alpha_n^{(1)}$ as in (36), $n \in [2, M - N + 1]$,
- (58)

$$\Psi^{(1)}(\alpha_n^{(1)}, \vec{t}) = \sum_{j=1}^M E_j^{(1)[n]}(\xi) e^{\theta_j(\vec{t})} = \left(\psi_n^{(1)}(\vec{t}) + \frac{\sum_{j=N+n-1}^M \hat{A}_j^N e^{\theta_j(\vec{t})} \xi^{-2(j-n-N+1)-1}}{\sum_{s=N}^{N+n-2} \hat{A}_s^N} \right) (1 + O(\xi^{-1})),$$

where

$$\psi_n^{(1)}(\vec{t}) = \sum_{j=N}^{N+n-2} \hat{A}_j^N e^{\theta_j(\vec{t})},$$

and for all $n \in [2, M - N + 1]$ the coefficients $E_j^{(1)[n]}(\xi)$ defined by (58) have the following properties:

- (a) $E_j^{(1)[n]}(\xi) \equiv 0$, if $j \in [N - 1]$;
 - (b) $E_j^{(1)[n]}(\xi)$ are rational functions in ξ for all $j \in [N, M]$;
 - (c) $\lim_{\xi \rightarrow \infty} E_j^{(1)[n]}(\xi) = \hat{E}_j^{(1)[n]}$, where $\hat{E}^{(1)}$ is as in Theorem 5 for $r = 1$.
- (2) The elementary symmetric functions in the $b_k^{(1)}(\xi)$,

$$\Pi_s^{(1)}(\xi) = \sum_{1 < j_1 < j_2 < \dots < j_s \leq M-N} \left(\prod_{l=1}^s b_{j_l}^{(1)} \right), \quad s \in [M - N]$$

are rational functions in ξ with coefficients depending only on \hat{A}_{N+l}^N , $l \in [0, M - N]$;

- (3) For $\xi \gg 1$, we have the following explicit asymptotic estimates for the divisor points $b_k^{(1)}$ on Γ_1 :

$$(59) \quad b_k^{(1)} = - \frac{\sum_{l=0}^{k-1} \hat{A}_{N+l}^N \xi^{2(k-1)} (1 + O(\xi^{-1}))}{\sum_{l=0}^k \hat{A}_{N+l}^N}, \quad k \in [M - N];$$

Proof. Requirement (1) is clearly equivalent to $\hat{B}_l^{(1)} = \hat{A}_{N+l-1}^N$, $l \in [M - N + 1]$. Requirement (2) is equivalent to:

$$\hat{A}_{N+l-1}^N \frac{\prod_{k \neq j}^{M-N+1} (\lambda_j^{(1)} - \lambda_k^{(1)})}{\prod_{k=1}^{M-N} (\lambda_j^{(1)} - b_k^{(1)})} = \delta_l^j, \quad j, l \in [M - N + 1].$$

Then, using Lemma 7 in Appendix B with $c_l = \hat{A}_{N+l-1}^N$, $l \in [M - N + 1]$, the coefficients $b_k^{(1)}$, $k \in [M - N]$ are uniquely defined and satisfy the required asymptotics (59). If $\zeta > 0$, then $\Psi^{(1)}(\zeta, \vec{t})$ is a finite sum of positive terms. Finally the asymptotic behavior of $\Psi^{(1)}(\alpha_n^{(1)}, \vec{t})$, $n \in [2, M - N + 1]$, easily follows from Lemma 7. \square

Let us now present the construction and the properties of the vacuum wavefunction on each Γ_r , for $r \in [2, N]$.

Theorem 6. Let $\xi > 1$ fixed and sufficiently big, $\lambda_l^{(r)}$, $l \in [M - N + 1]$, $\alpha_s^{(r)}$, $s \in [2, M - N + 1]$, $r \in [N]$, as in (35) and (36), $f^{(i)}(\vec{t})$, $i \in [N]$ as in (39) and let $\Psi^{(1)}(\zeta, \vec{t})$ satisfy Proposition 3.

For $r \in [2, N]$ let $\Psi^{(r)}(\zeta, \vec{t})$ as in (24) with $V_l^{(r)}(\vec{t})$ as in (25), $l \in [M - N + 1]$. Then for $r \in [2, N]$ properties

$$(60) \quad \lim_{\zeta \rightarrow \infty} \Psi^{(r)}(\zeta, \vec{t}) = \frac{f^{(r)}(\vec{t}) + \sum_{j=1}^{r-1} \epsilon_j^{(r)} f^{(j)}(\vec{t})}{1 + \sum_{j=1}^{r-1} \epsilon_j^{(r)}},$$

$$(61) \quad \Psi^{(r)}(\lambda_n^{(r)}, \vec{t}) = \begin{cases} e^{\theta_{N-r+1}(\vec{t})}, & n = 1 \\ \Psi^{(r-1)}(\alpha_n^{(r-1)}, \vec{t}), & n \in [2, M - N + 1], \end{cases}$$

uniquely define the coefficients $\hat{B}_l^{(r)} \equiv \frac{B_l^{(r)}}{1 + \sum_{k=i}^{r-1} \epsilon_k^{(r)}}$, the parameters $\epsilon_j^{(r)}$ and the divisor points

$b_k^{(r)} \in]\lambda_{k+1}^{(r)}, \lambda_k^{(r)}[$. Moreover

- (1) $B_l^{(r)} = B_l^{(r)}(\xi)$, $\epsilon_j^{(r)} = \epsilon_j^{(r)}(\xi)$ and the elementary symmetric functions in the divisor points $b_k^{(r)} = b_k^{(r)}(\xi)$,

$$\Pi_s^{(r)}(\xi) = \sum_{1 < j_1 < j_2 < \dots < j_s \leq M-N} \left(\prod_{l=1}^s b_{j_l}^{(r)} \right), \quad s \in [M - N]$$

are all rational function in ξ with coefficients depending only on $x_{l,s}$, $l \in [r]$, $s \in [M - N]$;

- (2) For $\xi \gg 1$, $B_l^{(r)} > 0$, $\epsilon_j^{(r)} > 0$ and:

$$(62) \quad B_l^{(r)} = \begin{cases} \hat{A}_{N-r+1}^{N-r+1}, & l = 1, \\ \hat{B}_l^{(r)} \cdot (1 + O(\xi^{-1})), & l \in [2, M - N + 1], \end{cases}$$

where $\hat{B}_l^{(r)}$ are defined by (52);

$$(63) \quad \epsilon_j^{(r)} = \eta_j^{(r)} \xi^{-j} (1 + O(\xi^{-1})), \quad j \in [1, r - 1],$$

where the positive constants $\eta_j^{(r)}$ are as in (88) and may be explicitly computed using (84) in Lemma 8 and (91) in Lemma 9;

- (3) For $\xi \gg 1$, the poles have the following asymptotics:

$$(64) \quad b_k^{(r)} = -\xi^{2(k-1)} \left(\frac{\sum_{l=1}^k \hat{B}_l^{(r)}}{\sum_{l=1}^{k+1} \hat{B}_l^{(r)}} \right) (1 + O(\xi^{-1})), \quad k \in [M - N];$$

- (4) $\Psi^{(r)}(\zeta, \vec{t}) > 0$, for all $\zeta > 0$ and for all \vec{t} , $r \in [2, N]$. In particular, all $\alpha_n^{(r)} > 0$, therefore $\Psi^{(r)}(\alpha_n^{(r)}, \vec{t}) > 0$ for all n and for all \vec{t} . Moreover, for any $n \in [2, M - N + 1]$ and for all \vec{t} ,

$$(65) \quad \Psi^{(r)}(\alpha_n^{(r)}, \vec{t}) = \sum_{j=1}^M E_j^{(r)[n]}(\xi) e^{\theta_j(\vec{t})} = \left\{ \psi_n^{(r)}(\vec{t}) + \frac{\sum_{j=N-r+n}^{N+n-2} \sigma_{n,j}^{(r)} e^{\theta_j} \xi^{-j+N-r+n-1}}{\sum_{s=N-r+1}^{N-r+n-1} \Delta_{[s; N-r+n, N-r+n+1, \dots, N+n-2]}} + \right. \\ \left. + \frac{\sum_{j=N+n-1}^M \Delta_{[N-r+n, N-r+n+1, \dots, N+n-2; j]} e^{\theta_j} \xi^{-r-2(j-N-n+1)}}{\sum_{s=N-r+1}^{N-r+n-1} \Delta_{[s; N-r+n, N-r+n+1, \dots, N+n-2]}} \right\} \times (1 + O(\xi^{-1})),$$

where the constants $\sigma_{n,j}^{(r)} > 0$ are recursively computed using Lemmas 9 and 10 and depend only on the affine coordinates $x_{l,k}$ on $Gr^{TNN}(N, M)$, $l \in [r]$, $s \in [M - N]$, defined in Proposition 5 in Appendix A. Moreover, for all $n \in [2, M - N + 1]$:

- (a) $E_j^{(r)[n]}(\xi) \equiv 0$ if $j \in [N - r]$;

- (b) $E_j^{(r)[n]}(\xi)$ are rational functions in ξ , for $j \in [N - r + 1, M]$
(c) $\lim_{\xi \rightarrow \infty} E_j^{(r)[n]}(\xi) = \hat{E}_j^{(r)[n]}$, where $\hat{E}^{(r)}$ is as in Theorem 5.

Proof. of Theorem 6. The proof goes through several steps and by induction.

Step 1: Direct proof for $r = 2$. Let $\Psi^{(1)}(\zeta, \vec{t})$ be as in Proposition 3 and let

$$\Psi^{(2)}(\zeta, \vec{t}) = \frac{\hat{B}_1^{(2)} \prod_{k \neq 1}^{M-N+1} (\zeta - \lambda_k^{(2)})}{\prod_{k=1}^{M-N} (\zeta - b_k^{(2)})} e^{\theta_{N-1}(\vec{t})} + \sum_{j=2}^{M-N+1} \frac{\hat{B}_j^{(2)} \prod_{k \neq j}^{M-N+1} (\zeta - \lambda_k^{(2)})}{\prod_{k=1}^{M-N} (\zeta - b_k^{(2)})} \Psi^{(1)}(\alpha_j^{(1)}, \vec{t}).$$

with all of the $\hat{B}_l^{(2)} = \frac{B_l^{(2)}}{1 + \epsilon_1^{(2)}}$, for $l \in [M - N + 1]$, $\epsilon_1^{(2)}$ and $b_k^{(2)}$ to be determined. Then:

a) We have

$$(66) \quad \lim_{\zeta \rightarrow \infty} \Psi^{(2)}(\zeta, \vec{t}) = \frac{f^{(2)}(\vec{t}) + \epsilon_1^{(2)} f^{(1)}(\vec{t})}{1 + \epsilon_1^{(2)}}$$

if and only if $B_1^{(2)} = \hat{A}_{N-1}^{N-1}$ and the remaining coefficients satisfy the linear system

$$\sum_{l=2}^{M-N+1} B_l^{(2)} E_s^{(1)[l]}(\xi) = \hat{A}_s^N \epsilon_1^{(2)} + \hat{A}_s^{N-1}, \quad s \in [N, M],$$

where the coefficients $E_s^{(1)[l]}(\xi)$ are the rational functions in ξ defined in (58). Then it is straightforward to check both the uniqueness of the solution of the above system for almost all $\xi > 1$ and the regularity properties in ξ of the coefficients $B_l^{(2)}$ and $\epsilon_1^{(2)}$. Moreover, using Lemma 8 for $r = 2$ with $\sigma_{n,l}^{(1)} = \hat{A}_{N+l-1}^N$, $l \in [1, M - N + 1]$, $n \in [2, M - N + 1]$, (58), the Principal Algebraic Lemma 2 and Lemma 10, we immediately get the required estimates for the coefficients ($l \in [2, M - N + 1]$),

$$B_l^{(2)} = \frac{\Delta_{[N+l-2, N+l-1]} \left(\sum_{j=1}^{l-1} \hat{A}_{N+j-1}^N \right)}{\hat{A}_{N+l-2}^N \hat{A}_{N+l-1}^N} (1 + O(\xi^{-1})), \quad \epsilon_1^{(2)} = \frac{\hat{A}_{M-1}^{N-1}}{\xi \hat{A}_{M-1}^N} (1 + O(\xi^{-1})).$$

Thanks to the positivity property of the matrix \hat{A} we immediately have $B_j^{(2)} > 0$, $j \in [M - N + 1]$ and $\epsilon_1^{(2)} > 0$ for all $\xi \gg 1$. Moreover, inserting $\vec{t} = \vec{0}$ in (66), we conclude $\sum_{j=1}^{M-N+1} \hat{B}_j^{(2)}(\xi) \equiv 1$. Finally all of the quantities may be expressed in invariant form using the FZ-basis $x_{l,k}$, $l = 1, 2$, $k \in [M - N]$.

b) The set of conditions

$$\Psi^{(2)}(\lambda_1^{(2)}, \vec{t}) = e^{\theta_{N-1}(\vec{t})}, \quad \Psi^{(2)}(\lambda_n^{(2)}, \vec{t}) = \Psi^{(1)}(\alpha_n^{(1)}, \vec{t}), \quad n \in [2, M - N + 1],$$

is equivalent to

$$\hat{B}_n^{(N-1)} \prod_{k \neq n}^{M-N+1} (\lambda_n - \lambda_k) = \prod_{k=1}^{M-N} (\lambda_n - b_k^{(2)}), \quad n \in [1, M - N + 1].$$

Using Lemmas 3 and 7 in the case $r = 2$, with $c_j = \hat{B}_j^{(2)}$, we immediately obtain the required conditions for the regularity in ξ , the position and the leading order expansion of the poles

($k \in [M - N]$),

$$b_k^{(2)} = -\xi^{2(k-1)} \frac{\hat{A}_{N+k}^N \left(\sum_{l=1}^k \Delta_{[N-2+l, N-1+k]} \right)}{\hat{A}_{N+k-1}^N \left(\sum_{l=1}^{k+1} \Delta_{[N-2+l, N+k]} \right)} (1 + O(\xi^{-1})).$$

c) If $\zeta > 0$, then $\Psi^{(2)}(\zeta, \vec{t})$ is a finite sum of positive terms. Finally, using Lemmas 9, 10 and Theorem 5, we get that the coefficients $E_j^{(2)[n]}(\xi)$ have the required regularity properties in ξ and that they satisfy the asymptotic expansion (65)

$$\begin{aligned} \Psi^{(2)}(\alpha_n^{(2)}, \vec{t}) &= \sum_{j=1}^M E_j^{(2)[n]}(\xi) e^{\theta_j(\vec{t})} = \left(\sum_{j=N-1}^{N+n-3} \Delta_{[j, N+n-2]} e^{\theta_j} + \frac{(\hat{A}_{N+n-2}^N)^2 \Delta_{[N+n-3, N+n-1]} e^{\theta_j}}{\hat{A}_{N+n-3}^N \hat{A}_{N+n-1}^N} \frac{e^{\theta_j}}{\xi} + \right. \\ &\quad \left. + \sum_{j=N+n-1}^M \frac{\Delta_{[N+n-2, j]} e^{\theta_j}}{\xi^{2(j-N-n+2)}} \right) \times \frac{(1+O(\xi^{-1}))}{\sum_{k=N-1}^{N+n-3} \Delta_{[k, N+n-2]}}; \end{aligned}$$

Step 2: The induction procedure. Let $r \in [3, N]$ and let us suppose we proved the Theorem for $i = 2, \dots, r-1$, and let us prove it for $i = r$. Let us denote $\hat{B}_l^{(r)} = \frac{B_l^{(r)}}{1 + \sum_{j=1}^{r-1} \epsilon_j^{(r)}}$, for $l \in [M - N + 1]$ and define

$$\Psi^{(r)}(\zeta, \vec{t}) = \frac{B_1^{(r)} e^{\theta_{N-r+1}(\vec{t})} \prod_{j \neq 1}^{M-N+1} (\zeta - \lambda_j^{(r)})}{(1 + \sum_{j=1}^{r-1} \epsilon_j^{(r)}) \prod_{k=1}^{M-N} (\zeta - b_k^{(r)})} + \sum_{n=2}^{M-N+1} \frac{B_n^{(r)} \Psi^{(r-1)}(\alpha_n^{(r-1)}, \vec{t}) \prod_{j \neq n}^{M-N+1} (\zeta - \lambda_j^{(r)})}{(1 + \sum_{j=1}^{r-1} \epsilon_j^{(r)}) \prod_{k=1}^{M-N} (\zeta - b_k^{(r)})},$$

with coefficients $B_n^{(r)}$, $\epsilon_j^{(r)}$ and $b_k^{(r)}$ to be determined.

a) We have

$$(67) \quad \lim_{\zeta \rightarrow \infty} \Psi^{(r)}(\zeta, \vec{t}) = \frac{f^{(r)}(\vec{t}) + \sum_{j=1}^{r-1} \epsilon_j^{(r)} f^{(j)}(\vec{t})}{1 + \sum_{j=1}^{r-1} \epsilon_j^{(r)}}$$

if and only if $B_1^{(r)} = \hat{A}_{N-r+1}^{N-r+1}$ and the remaining coefficients satisfy the linear system

$$\sum_{l=2}^{M-N+1} B_l^{(r)} E_s^{(r-1)[l]}(\xi) = \sum_{j=1}^{r-1} \hat{A}_s^{N-r+j+1} \epsilon_j^{(r)} + \hat{A}_s^{N-r+1}, \quad s \in [N - r + 2, M],$$

where the coefficients $E_s^{(r-1)[l]}(\xi)$ are rational functions in ξ . Due to the compatibility of the above linear system for almost all $\xi > 1$ and the regularity properties of the coefficients, there immediately follow both the uniqueness for almost all $\xi > 1$ and the regularity properties in ξ for $B_l^{(r)}$ and $\epsilon_j^{(r)}$. Again, using Lemmas 2, 8 and 10, we immediately get the required asymptotic estimates for the coefficients $B_l^{(r)}$ ($l \in [2, M - N + 1]$) and $\epsilon_j^{(r)}$, ($j \in [r - 1]$) as in (62) and (63), respectively, when $\xi \gg 1$. In particular, substituting $\vec{t} = \vec{0}$ in (67), we have $\sum_{l=1}^{M-N+1} \hat{B}_l^{(r)} = 1$.

b) The set of conditions

$$\Psi^{(r)}(\lambda_1^{(r)}, \vec{t}) = e^{\theta_{N-r+1}(\vec{t})}, \quad \Psi^{(r)}(\lambda_n^{(r)}, \vec{t}) = \Psi^{(r-1)}(\alpha_n^{(r-1)}, \vec{t}), \quad n \in [2, M - N + 1],$$

is equivalent to

$$\hat{B}_n^{(r)} \prod_{j \neq n}^{M-N+1} (\lambda_n^{(r)} - \lambda_j^{(r)}) = \prod_{k=1}^{M-N} (\lambda_n^{(r)} - b_k^{(r)}), \quad n \in [1, M - N + 1].$$

Again, using Lemma 7 in the case $c_j = \mathring{B}_j^{(r)}$, and Lemma 3, we immediately obtain the required estimates for the regularity and for position of the poles $b_k^{(r)}$ ($k \in [M - N]$) as well as the leading order estimates for ($\xi \gg 1$) as in (64).

c) Finally, from Lemmas 9, 10 and Theorem 5, we get the required estimates for the regularity, the sign and the leading order term expansion of $\Psi^{(r)}(\alpha_n, \vec{t})$, for $n \in [2, M - N + 1]$. \square

In Theorem 6 we provide the explicit construction of the normalized vacuum wave function, with the properties required in Theorem 3. This remark completes the proof of Theorem 3.

6. ANALYTIC PROPERTIES OF THE EFFECTIVE DIVISOR

6.1. Zero divisor for the vacuum wave function. In Theorem 6 we provide an explicit estimate of the position of the vacuum pole divisor (64). Let us now present some estimates for the position of vacuum zero divisor for sufficiently small times. We do not use directly these formulas in our paper, but they may be useful in future investigations.

Corollary 2. *Let $\xi \gg 1$ be fixed, $\Gamma = \Gamma(\xi)$ be as in Definition 1 and $\Psi(\zeta, \vec{t})$ be the vacuum wavefunction of Theorem 3. Then in each finite oval $\Omega_{r,n}$, ($r \in [N]$, $n \in [M - N]$), $\Psi(\zeta, \vec{t})$ possesses exactly one simple pole $b_n^{(r)}(\xi)$, whose position is independent of \vec{t} , and exactly one simple zero $\chi_n^{(r)}(\xi; \vec{t})$. In particular*

- (1) $b_n^{(r)}(\xi) \in]\lambda_{n+1}^{(r)}, \lambda_n^{(r)}[\subset \Gamma_r \cap \Omega_{r,n}$;
- (2) $\chi_n^{(r)}(\xi; \vec{0}) = b_n^{(r)}(\xi)$;
- (3) $\chi_n^{(r)}(\xi; \vec{t}) \in]\lambda_{n+1}^{(r)}, \lambda_n^{(r)}[\subset \Gamma_r \cap \Omega_{r,n}$, for all \vec{t} ;
- (4) Assume that only a finite number of times is different from zero: $t_j = 0$ for $j > j_0$, and all times t_1, t_2, \dots, t_{j_0} lie in a compact domain K_0 . Then we have the following asymptotic expansion in $\xi \gg 1$:

$$(68) \quad \chi_n^{(r)}(\xi; \vec{t}) = - \frac{\sum_{l=1}^n \hat{B}_l^{(r)} V_l^{(r)}(\vec{t})}{\sum_{l=1}^{n+1} \hat{B}_l^{(r)} V_l^{(r)}(\vec{t})} \xi^{2(j-1)} (1 + O(\xi^{-1})) = - \frac{\sum_{j=N-r+1}^{N-r+n} \frac{\Delta_{[j; N-r+n+1, \dots, N+n-1]} e^{\theta_j}}{\Delta_{[N-r+n+1, \dots, N+n-1]}} \xi^{2(j-1)} (1 + O(\xi^{-1})).$$

Proof. The number of poles $b_k^{(r)}$ is equal to the number of ovals and their position is computed in Proposition 3 and in Theorem 6. Therefore the number of zeroes of $\Psi(\zeta, \vec{t})$ is equal to the number of ovals. For $\vec{t} = (0, 0, \dots)$, by definition, the zeroes coincide with the divisor points, and their positions continuously depend on \vec{t} . A zero could leave a real oval only if it collides with another zero coming from another oval. That is impossible since $\Psi^{(r)}(\lambda_j^{(r)}, \vec{t}) > 0$, for all \vec{t} , with $r \in [N]$, $j \in [M - N + 1]$ (see Theorem 6). It means that for all times \vec{t} each zero remains in the same open interval $] \lambda_n^{(r)}, \lambda_{n+1}^{(r)}[$.

All terms $V_l^{(r)}(\vec{t})$ are of order 1 in ξ for $(t_1, \dots, t_{j_0}) \in K_0$. Let us write the function $\Psi^{(r)}(\zeta, \vec{t})$ as a sum of simple fractions:

$$\Psi^{(r)}(\zeta, \vec{t}) = f_{r,\xi}(\vec{t}) + \sum_{k=1}^{M-N} \frac{\psi_k^{(r)}(\vec{t})}{\zeta - b_k^{(r)}},$$

where $f_{r,\xi}(\vec{t})$ is as in (43) and the positions of the poles are given by (64). Therefore

$$\psi_k^{(r)}(\vec{t}) = \xi^{2k-2} \cdot \left(\frac{\sum_{j=1}^k \hat{B}_j^{(r)} V_j^{(r)}(\vec{t}) - \left(\sum_{j=1}^k \hat{B}_j^{(r)} \right) V_{k+1}^{(r)}(\vec{t})}{\left(\sum_{j=1}^{k+1} \hat{B}_j^{(r)} \right)^2} \hat{B}_{k+1}^{(r)} \right) (1 + O(\xi^{-1})),$$

and inside the interval $[\lambda_{k+1}^{(r)}, \lambda_k^{(r)}]$ we have

$$(69) \quad \Psi^{(r)}(\zeta, \vec{t}) = \xi^{2k-2} \cdot \left(\frac{\sum_{j=1}^{k+1} \hat{B}_j^{(r)} V_j^{(r)}(\vec{t})}{\sum_{j=1}^{k+1} \hat{B}_j^{(r)}} - \frac{\psi_k^{(r)}(\vec{t})}{\zeta - b_k^{(r)}} \right) (1 + O(\xi^{-1})).$$

We complete the proof solving the equation $\Psi^{(r)}(\zeta, \vec{t}) = 0$ and using the approximation (69). \square

We remark that the condition that each zero of $\Psi(P, \vec{t})$ lies in a well-defined open interval $[\lambda_{j+1}^{(r)}, \lambda_j^{(r)}]$ for all \vec{t} , is natural since $\Psi(P, \vec{t})$ represents a vacuum wave function: no collision is possible for the zero divisor in this case!

6.2. Zero divisor for the normalized Darboux transformed wave function. We now provide some estimates on the zero and pole divisors of the normalized wavefunction $\tilde{\Psi}(\zeta, \vec{t})$. The pole (effective) divisor has been characterized in Theorem 4. For the zero divisor $\mathcal{D}_\xi(\vec{t})$ we adopt the following notation. By construction, for any fixed \vec{t} , its restriction to Γ_0 is $\mathcal{D}^{(0)}(\vec{t}) = \{\gamma_k^{(0)}(\vec{t}), k \in [N]\}$ and it coincides with the set of solutions to (12)

$$(\gamma_l^{(0)})^N(\vec{t}) - w_1(\vec{t})(\gamma_l^{(0)})^{N-1}(\vec{t}) - \dots - w_{N-1}(\vec{t})\gamma_l^{(0)} - w_N(\vec{t}) = 0, \quad l \in [N].$$

The restriction of the zero divisor to Γ_r is $\mathcal{D}_\xi^{(r)}(\vec{t}) = \{\gamma_l^{(r)}(\vec{t}), l \in [M - N - 1]\}$ and it coincides with the set of solutions to

$$D^{(k)}\Psi^{(r)}(\zeta, \vec{t}) = 0, \quad r \in [N].$$

In the following theorem we estimate the position of the zero divisor in the case in which only a finite number of times in \vec{t} may be different from zero, and moreover they vary in a neighborhood of $\vec{0} = (0, \dots, 0)$. In particular we give the explicit estimate for the position of the effective divisor \mathcal{D}_ξ .

Theorem 7. (Estimate of the position of divisor \mathcal{D}_ξ) Let $\xi \gg 1$ and let $D\Psi^{(r)}(\zeta, \vec{t})$, $r \in [N]$, as above. Assume that only a finite number of times may be different from zero: $t_j = 0$ for $j > j_0$, and all times t_1, t_2, \dots, t_{j_0} lie in a compact domain K_0 containing the point $(t_1, \dots, t_{j_0}) = (0, \dots, 0)$. Then for $\xi \gg 1$, generically the following asymptotic expansion holds for the zeroes of $D\Psi^{(r)}(\zeta, \vec{t})$, $\vec{t} \in K_0$:

$$(70) \quad \gamma_n^{(r)}(\vec{t}) = - \frac{\sum_{l=1}^n \hat{B}_l^{(r)} DV_l^{(r)}(\vec{t})}{\sum_{l=1}^{n+1} \hat{B}_l^{(r)} DV_l^{(r)}(\vec{t})} \xi^{2(n-1)} (1 + O(\xi^{-1})) = - \frac{\sum_{j=N-r+1}^{N-r+n} \frac{\Delta_{[j; N-r+n+1, \dots, N+n-1]} P^{(0)}(\kappa_j) e^{\theta_j}}{\Delta_{[N-r+n+1, \dots, N+n-1]}}}{\sum_{j=N+r-1}^{N-r+n+1} \frac{\Delta_{[j; N-r+n+2, \dots, N+n]} P^{(0)}(\kappa_j) e^{\theta_j}}{\Delta_{[N-r+n+2, \dots, N+n]}}} \xi^{2(n-1)} (1 + O(\xi^{-1})),$$

where $P^{(0)}(\kappa_j) = \prod_{l=1}^N (\kappa_j - \gamma_l^{(0)}(\vec{t}))$, $j \in [M]$. In particular, for $(t_1, \dots, t_{j_0}) = (0, \dots, 0) \in K_0$ we have the estimate of the effective divisor \mathcal{D}_ξ .

In Corollaries 3 and 4 we control the position of the both the pole and the zero divisor on each sheet. Indeed, during the time evolution the divisor points can pass through the double points $X \in \Gamma_{r_1} \cap \Gamma_{r_2}$ only in pairs and coming from different sheets ($r_1 \neq r_2$), because of the properties of $\tilde{\Psi}(P, \vec{t})$ settled in Theorem 2 (see Figure 9).




FIGURE 9. A pair of divisor points passes through a double point.

In Theorem 4 we have shown that any finite oval contains exactly one point of the zero divisor $\mathcal{D}_\xi(\vec{t})$. Let us provide additional information about the positions of the zero divisor points for any fixed real \vec{t} .

Corollary 3. Characterization of the zero divisor $\mathcal{D}_\xi(\vec{t})$ For any fixed $\xi \gg 1$ and for any real \vec{t} , we have

- (1) $\mathcal{D}^{(0)}(\vec{t}) \subset]\kappa_1, \kappa_M[$ and $\#(\mathcal{D}^{(0)}(\vec{t}) \cap]\kappa_1, \kappa_M[) = N$;
- (2) There is at most one divisor point in each interval $[\kappa_j, \kappa_{j+1}]$, $j \in [M-1]$;
- (3) For any $r \in [N]$, $\mathcal{D}_\xi^{(r)}(\vec{t}) \subset]\lambda_{M-N+1}^{(r)}, \alpha_{M-N+1}^{(r)}[$ and $\#(\mathcal{D}_\xi^{(r)}(\vec{t}) \cap]\lambda_{M-N+1}^{(r)}, \alpha_{M-N+1}^{(r)}[) = M - N - 1$;
- (4) For any $r \in [N]$, there is at most one divisor point in each interval $[\lambda_{j+1}^{(r)}, \lambda_j^{(r)}]$, $j \in [M-N]$;
- (5) For any $r \in [N]$, there is at most one divisor point in each interval $[\alpha_j^{(r)}, \alpha_{j+1}^{(r)}]$, $j \in [M-N]$, where $\alpha_1^{(r)} = \lambda_1^{(r)}$;
- (6) For any $r \in [N]$,

$$s^{(r)}(\vec{t}) \equiv \#(\mathcal{D}_\xi^{(r)}(\vec{t}) \cap]\lambda_1^{(r)}, \alpha_{M-N+1}^{(r)}[) \leq \min\{N - r, M - N - r\}.$$

For any fixed real \vec{t} , we have the complete control of the position of the divisor $\mathcal{D}_\xi^{(r)}(\vec{t})$ and in particular it is possible to estimate the position of the effective divisor.

Corollary 4. (Counting the number of positive poles and zeros on each sheet) For any fixed \vec{t} , the number of negative and positive divisor points in $\mathcal{D}_\xi^{(r)}(\vec{t})$ is uniquely determined for all $r \in [N]$ from $\mathcal{D}^{(0)}(\vec{t})$. Indeed let \vec{t} be fixed and define

$$\begin{aligned} s^{(0)} &\equiv s^{(0)}(\vec{t}) = \#(\mathcal{D}^{(0)}(\vec{t}) \cap]\kappa_N, \kappa_M[); \\ s^{(r)} &\equiv s^{(r)}(\vec{t}) = \#(\mathcal{D}_\xi^{(r)}(\vec{t}) \cap]\lambda_1^{(r)}, \alpha_{M-N+1}^{(r)}[); \quad r \in [N]. \end{aligned}$$

- (1) $s^{(r)}$ is a decreasing function of r , $r \in [0, N]$ and $s^{(N)} = 0$;
- (2) $s^{(r)} \leq \min\{N - r, M - N - r\}$, for all $r \in [0, N]$;
- (3) If $s^{(0)} = 1$, then $s^{(r)} = 0$ for any $r \in [N]$.
- (4) If $s^{(0)} > 1$ and $\#(\mathcal{D}^{(0)} \cap]\kappa_{N-1}, \kappa_N[) = 1$, then $s^{(1)} = s^{(0)}$; otherwise $s^{(1)} = s^{(0)} - 1$;
- (5) Let $r \in [2, N]$ be fixed and suppose that $s^{(r-1)} \geq 1$. Then:
if $\#(\mathcal{D}^{(0)} \cap]\kappa_{N-r+1}, \kappa_{N-r+2}[) = 1$, then $s^{(r)} = s^{(r-1)}$;
otherwise $s^{(r)} = s^{(r-1)} - 1$.

Corollary 5. Under the hypotheses of the Theorem 4 and for any fixed \vec{t} , the following holds true:

- (1) $D\Psi^{(0)}(\kappa_M, \vec{t}) > 0$, $(-1)^N D\Psi^{(0)}(\kappa_1, \vec{t}) > 0$;
- (2) $(-1)^r D\Psi^{(r)}(\alpha_{M-N+1}^{(r)}, \vec{t}) > 0$, for all $r \in [N-1]$;
- (3) $\tilde{\Psi}^{(0)}(\kappa_M, \vec{t}) > 0$, $\tilde{\Psi}^{(0)}(\kappa_1, \vec{t}) > 0$;




FIGURE 10. The topological scheme of spectral curve for soliton data $Gr^{TP}(2,4)$, $\Gamma(\xi)$ (left) is the partial normalization the plane algebraic curve (right), which is the rational degeneration of the genus 4 M-curve in [76]. The ovals in the nodal plane curve are labeled as in the real part of its partial normalization.

$$(4) \quad \tilde{\Psi}^{(r)}(\alpha_{M-N+1}^{(r)}, \vec{t}) > 0, \text{ for all } r \in [N].$$

Remark 14. In [39], Malanyuk states that if A is an element of $Gr^{TNN}(N, M)$ then, for $j \in [N]$, $\gamma_j^{(0)}(\vec{0})$ are real, distinct and lie in $[\kappa_1, \kappa_M]$. Our estimates improve such result in the case $Gr^{TP}(N, M)$ and are optimal.

7. $\Gamma(\xi)$ AND THE VACUUM DIVISOR FOR SOLITON DATA IN $Gr^{TP}(2,4)$

In this section we construct the rational curve and the vacuum pole divisor associated to generic soliton data in $Gr^{TP}(2,4)$. The degenerate curve $\Gamma(\xi)$ in Definition [4] is the partial normalization of the nodal plane curve in [75], which is the rational degeneration of the genus 4 M-curve in [76]. Let us recall that generic curves of sufficiently high genus can not be represented as plane curves without self-intersections [22], [3], therefore the partial normalization is generically unavoidable. We plot both the topological model and the partial normalization for this example in Figure [10].

7.1. $\Gamma(\xi)$ and its desingularization for generic soliton data in $Gr^{TP}(2,4)$. According to Definition [4], the degenerate curve $\Gamma(\xi)$ is obtained gluing three copies of \mathbb{CP}^1 : Γ_0 , Γ_1 and Γ_2 .

On Γ_0 we have 5 real marked points $\kappa_1 < \kappa_2 < \kappa_3 < \kappa_4$ and P_0 . Let $\tilde{\zeta} \equiv \zeta$ the local coordinate on Γ_0 such that $\tilde{\zeta}^{-1}(P_0) = 0$ and let us denote with the same symbol the phases and their local coordinates, $\tilde{\zeta}(\kappa_j) = \kappa_j$, $j \in [4]$. To simplify the expressions below and without loss of generality, we take $\tilde{\zeta}(\kappa_1) = 0$.

On Γ_1 , we have 6 real ordered marked points, with ζ -coordinates: $\zeta(\alpha_3^{(1)}) = \xi > \zeta(\alpha_2^{(1)}) = \xi^{-1} > \zeta(\lambda_1^{(1)}) = 0 > \zeta(\lambda_2^{(1)}) = -1 > \zeta(\lambda_3^{(1)}) = -\xi^2$ and $\zeta^{-1}(Q_1) = 0$. The fractional linear change of coordinates $\zeta \mapsto \tilde{\zeta}$ is uniquely defined by the conditions that the $\tilde{\zeta}$ coordinates of the marked points $\lambda_j^{(1)} \in \Gamma_1$, $j \in [3]$, coincide with the $\tilde{\zeta}$ coordinates of $\kappa_{j+1} \in \Gamma_0$: $\tilde{\zeta}(\lambda_1^{(1)}) = \kappa_2$, $\tilde{\zeta}(\lambda_2^{(1)}) = \kappa_3$ and $\tilde{\zeta}(\lambda_3^{(1)}) = \kappa_4$:

$$(71) \quad \tilde{\zeta} = \frac{\xi^2 \kappa_2 (\kappa_4 - \kappa_3) - \zeta [\xi^2 \kappa_4 (\kappa_3 - \kappa_2) - \kappa_3 (\kappa_4 - \kappa_2)]}{\xi^2 (\kappa_4 - \kappa_3) - \zeta [\xi^2 (\kappa_3 - \kappa_2) - (\kappa_4 - \kappa_2)]}$$

Then, it is easy to verify that

$$(72) \quad \begin{aligned} \tilde{\zeta}(\alpha_2^{(1)}) &= \frac{\xi^2 \kappa_2 (\kappa_4 - \kappa_3) - (\xi - 1) \kappa_3 (\kappa_4 - \kappa_2)}{\xi^2 (\kappa_4 - \kappa_3) - (\xi - 1) (\kappa_4 - \kappa_2)} = \kappa_2 - \frac{(\kappa_3 - \kappa_2)(\kappa_4 - \kappa_2)}{\xi(\kappa_4 - \kappa_3)} + O(\xi^{-2}) \\ \tilde{\zeta}(\alpha_3^{(1)}) &= \frac{\xi \kappa_4 (\kappa_3 - \kappa_2) - \kappa_3 (\kappa_4 - \kappa_2)}{\xi (\kappa_3 - \kappa_2) - (\kappa_4 - \kappa_2)} = \kappa_4 + \frac{(\kappa_4 - \kappa_3)(\kappa_4 - \kappa_2)}{\xi(\kappa_3 - \kappa_2)} + O(\xi^{-2}) \\ \tilde{\zeta}(Q_1) &= \frac{\xi^2 \kappa_4 (\kappa_3 - \kappa_2) - \kappa_3 (\kappa_4 - \kappa_2)}{\xi^2 (\kappa_3 - \kappa_2) - (\kappa_4 - \kappa_2)} = \kappa_4 + \frac{(\kappa_4 - \kappa_3)(\kappa_4 - \kappa_2)}{\xi^2 (\kappa_3 - \kappa_2)} + O(\xi^{-4}). \end{aligned}$$

On Γ_2 , we have 4 real ordered marked points, with ζ -coordinates: $\zeta(\lambda_1^{(2)}) = 0 > \zeta(\lambda_2^{(2)}) = -1 > \zeta(\lambda_3^{(2)}) = -\xi^2$ and $\zeta^{-1}(Q_2) = 0$. The fractional linear change of coordinates $\zeta \mapsto \tilde{\zeta}$ is uniquely defined by the condition that the $\tilde{\zeta}$ coordinates at the double points $\lambda_j^{(2)} \in \Gamma_2$, $j \in [3]$ respectively coincide with the $\tilde{\zeta}$ coordinates of $\kappa_1 \in \Gamma_0$, $\alpha_j^{(1)} \in \Gamma_1$: $\tilde{\zeta}(\lambda_1^{(2)}) = 0$, $\tilde{\zeta}(\lambda_2^{(2)}) = \tilde{\zeta}(\alpha_2^{(1)})$ and $\tilde{\zeta}(\lambda_3^{(2)}) = \alpha_3^{(1)}$:

$$(73) \quad \tilde{\zeta} = \frac{(\xi^2 - 1)\tilde{\zeta}(\alpha_2^{(1)})\tilde{\zeta}(\alpha_3^{(1)})\zeta}{\xi^2(\tilde{\zeta}(\alpha_2^{(1)}) - \tilde{\zeta}(\alpha_3^{(1)})) - \zeta(\tilde{\zeta}(\alpha_3^{(1)}) - \xi^2\tilde{\zeta}(\alpha_2^{(1)}))},$$

with $\tilde{\zeta}(\alpha_2^{(1)})$, $\tilde{\zeta}(\alpha_3^{(1)})$ as in (72). Then, it is easy to verify that

$$(74) \quad \tilde{\zeta}(Q_2) = \frac{(\xi^2 - 1)\tilde{\zeta}(\alpha_3^{(1)})\tilde{\zeta}(\alpha_2^{(1)})}{\xi^2\tilde{\zeta}(\alpha_2^{(1)}) - \tilde{\zeta}(\alpha_3^{(1)})} = \tilde{\zeta}(\alpha_3^{(1)}) + \frac{\tilde{\zeta}(\alpha_3^{(1)})(\tilde{\zeta}(\alpha_3^{(1)}) - \tilde{\zeta}(\alpha_2^{(1)}))}{\xi^2\tilde{\zeta}(\alpha_2^{(1)})} + O(\xi^{-4})$$

and

$$\tilde{\zeta}(Q_2) - \tilde{\zeta}(\alpha_3^{(1)}) = \frac{(\kappa_4 - \kappa_2)\kappa_4}{\xi^2\kappa_2} + O(\xi^{-3}).$$

Then, in the local coordinates $(\tilde{\mu}, \tilde{\zeta})$, the reducible real algebraic curve $\Gamma(\xi)$ corresponding to the gluing rules of Definition 4 is the following plane nodal curve

$$(75) \quad \tilde{\mu} \cdot (\tilde{\mu} - p_0(\tilde{\zeta})) \cdot (\tilde{\mu} + p_2(\tilde{\zeta})) = 0,$$

where $p_0(\tilde{\zeta}) = \prod_{j=1}^3 (\tilde{\zeta} - \kappa_j)$, $p_2(\tilde{\zeta}) = \frac{\kappa_2 \kappa_3 \kappa_4}{\tilde{\zeta}(\alpha_2^{(1)})\tilde{\zeta}(\alpha_3^{(1)})} (\tilde{\zeta} - \tilde{\zeta}(\alpha_2^{(1)}))(\tilde{\zeta} - \tilde{\zeta}(\alpha_3^{(1)}))$. (75) is the rational degeneration of the M-curve, which is the normalization of the following nodal curve:

$$(76) \quad \Gamma(\xi)_\delta : \quad \tilde{\mu} \cdot (\tilde{\mu} - p_0(\tilde{\zeta})) \cdot (\tilde{\mu} + p_2(\tilde{\zeta})) - \delta^2 \left(\frac{p_0(\tilde{\zeta}) + p_2(\tilde{\zeta})}{\tilde{\zeta}} \right)^2 \left(\kappa_4 + \frac{1}{2\xi} - \tilde{\zeta} \right) = 0, \quad |\delta| \ll 1.$$

It is easy to verify that the curve (76) has genus 4 for generic values of κ_j s and $\xi \gg 1$. By construction it also possesses the maximum number of ovals.

7.2. The leading order coefficients and vectors of the vacuum wavefunction. For completeness we compute the coefficients and the vectors of Theorem 5 for points in $Gr^{\text{TP}}(2, 4)$. The totally positive matrix in banded form defined in (80) expressed in function of the FZ-basis $x_{i,s}$ is

$$A = \begin{pmatrix} 1 & \frac{x_{2,2} + x_{1,2}x_{2,1}}{x_{1,1}x_{1,2}} & \frac{x_{2,2}}{x_{1,2}} & 0 \\ 0 & 1 & x_{1,1} & x_{1,2} \end{pmatrix},$$

and \hat{A} is the corresponding normalized totally positive matrix, obtained dividing each A_j^i by the sum of the elements on the i -th row (see Remark 9). Then the bases $E^{(0)}$, $E^{(1)}$ and the

coefficients $B_j^{(1)}, B_j^{(2)}, j \in [3]$ from the Principal Algebraic Lemma and Theorem 5 can be easily calculated:

$$\begin{aligned} \hat{E}^{(0)} &= \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \hat{E}^{(1)} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & \frac{1}{1+x_{1,1}} & \frac{x_{1,1}}{1+x_{1,1}} & 0 \end{pmatrix}, \\ \hat{B}_1^{(1)} &= \frac{1}{1+x_{1,1}+x_{1,2}}, \quad \hat{B}_2^{(1)} = \frac{x_{1,1}}{1+x_{1,1}+x_{1,2}}, \quad \hat{B}_3^{(1)} = \frac{x_{1,2}}{1+x_{1,1}+x_{1,2}} \\ (77) \quad \hat{B}_1^{(2)} &= \hat{A}_1^1 = \frac{x_{1,2}x_{1,1}}{x_D^{(2)}}, \quad \hat{B}_2^{(2)} = \frac{\Delta_{[2,3]}}{\hat{A}_3^2} = \frac{x_{2,1}x_{1,2}}{x_D^{(2)}}, \quad \hat{B}_3^{(2)} = \frac{\Delta_{[3,4]}(\hat{A}_2^2 + \hat{A}_3^2)}{\hat{A}_3^2 \hat{A}_4^2} = \frac{x_{1,1}x_{2,2} + x_{2,2}}{x_D^{(2)}}, \end{aligned}$$

where $\Delta_{[2,3]} = \hat{A}_2^1 \hat{A}_3^2 - \hat{A}_3^1 \hat{A}_2^2$, $\Delta_{[3,4]} = \hat{A}_3^1 \hat{A}_4^2$ and $x_D^{(2)} = x_{1,1}(x_{1,2} + x_{2,2}) + x_{1,2}x_{2,1} + x_{2,2}$.

7.3. The vacuum divisor for soliton data in $Gr^{\text{TP}}(2, 4)$. Let us calculate now the vacuum wave function from Theorem 6 for this example. In the local coordinate ζ , the vacuum wavefunction takes the form

$$\begin{aligned} \Psi^{(0)}(\zeta, \vec{t}) &= \exp(\theta(\zeta, \vec{t})), & \zeta \in \Gamma_0, \\ \Psi^{(1)}(\zeta, \vec{t}) &= \frac{(\xi^2 + \zeta)(\zeta + 1)e^{\theta_2(\vec{t})} + (\xi^2 + \zeta)\zeta x_{1,1}e^{\theta_3(\vec{t})} + \zeta(\zeta + 1)x_{1,2}e^{\theta_4(\vec{t})}}{\zeta^2(1+x_{1,1}+x_{1,2}) + \zeta(\xi^2[1+x_{1,1}] + x_{1,2} + 1) + \xi^2}, & \zeta \in \Gamma_1, \\ \Psi^{(2)}(\zeta, \vec{t}) &= \frac{(\zeta + 1)(\zeta + \xi^2)e^{\theta_1(\vec{t})} + B_2^{(2)}\zeta(\zeta + \xi^2)V_2^{(2)}(\xi, \vec{t}) + B_3^{(2)}\zeta(\zeta + 1)V_3^{(2)}(\xi, \vec{t})}{C^{(2)}(\zeta^2 + \nu_1^{(2)}\zeta + \nu_2^{(2)})}, & \zeta \in \Gamma_2 \end{aligned}$$

with

$$\begin{aligned} V_2^{(2)}(\xi, \vec{t}) &= \Psi^{(1)}(\xi^{-1}, \vec{t}) = \frac{(1+\xi^3)e^{\theta_2(\vec{t})} + x_{1,1}(\xi^2 - \xi + 1)e^{\theta_3(\vec{t})} + x_{1,2}e^{\theta_4(\vec{t})}}{\xi^3 + \xi^2 x_{1,1} - \xi x_{1,1} + x_{1,1} + x_{1,2} + 1} \\ &= \left(e^{\theta_2(\vec{t})} + \frac{x_{1,1}}{\xi} e^{\theta_3(\vec{t})} + \frac{x_{1,2}}{\xi^3} e^{\theta_4(\vec{t})} \right) (1 + O(\xi^{-1})), \\ V_3^{(2)}(\xi, \vec{t}) &= \Psi^{(1)}(\xi, \vec{t}) = \frac{(\xi + 1)e^{\theta_2(\vec{t})} + \xi x_{1,1}e^{\theta_3(\vec{t})} + x_{1,2}e^{\theta_4(\vec{t})}}{\xi(x_{1,1} + 1) + x_{1,2} + 1} = \left(\frac{e^{\theta_2(\vec{t})} + x_{1,1}e^{\theta_3(\vec{t})}}{x_{1,1} + 1} + \frac{x_{1,2}e^{\theta_4(\vec{t})}}{(1+x_{1,1})\xi} \right) (1 + O(\xi^{-1})), \\ B_2^{(2)} &= \frac{(\xi x_{1,2}x_{2,1} - x_{1,2}x_{2,1} - x_{2,2})(\xi^3 + \xi^2 x_{1,1} - \xi x_{1,1} + x_{1,1} + x_{1,2} + 1)}{(\xi - 1)^2 \xi^2 x_{1,1} x_{1,2}} = \frac{x_{2,1}}{x_{1,1}} + O(\xi^{-1}), \\ B_3^{(2)} &= \frac{(\xi^2 x_{2,2} - \xi(x_{1,2}x_{2,1} + x_{2,2}) + x_{1,2}x_{2,1} + x_{2,2})(\xi(x_{1,1} + 1) + x_{1,2} + 1)}{\xi x_{1,1} x_{1,2} (\xi - 1)^2} = \frac{x_{2,2}(1+x_{1,1})}{x_{1,1} x_{1,2}} + O(\xi^{-1}), \\ \epsilon_1^{(2)} &= \frac{\xi^2 x_{2,2} - \xi x_{1,2}x_{2,1} + x_{1,2}x_{2,1} + x_{2,2}}{\xi^2 x_{1,1} x_{1,2} (\xi - 1)} = \frac{x_{2,2}}{x_{1,1} x_{1,2} \xi} + O(\xi^{-2}), \\ C^{(2)} &= 1 + B_2^{(2)} + B_3^{(2)}, \quad \nu_1^{(2)} = \frac{\xi^2}{C^{(2)}}, \quad \nu_2^{(2)} = \frac{\xi^2(1+B_2^{(2)}) + 1 + B_3^{(2)}}{C^{(2)}}, \end{aligned}$$

so that, in agreement with the estimates in Theorem 6, we have

$$\frac{1}{C^{(2)}} = \hat{B}_1^{(2)} + O(\xi^{-1}), \quad \frac{B_2^{(2)}}{C^{(2)}} = \hat{B}_2^{(2)} + O(\xi^{-1}), \quad \frac{B_3^{(2)}}{C^{(2)}} = \hat{B}_3^{(2)} + O(\xi^{-1}),$$

with $\hat{B}_j^{(2)}, j \in [3]$, as in (77).

The vacuum poles in Γ_1 in the local coordinate ζ take the form

$$\begin{aligned} \zeta(b_1^{(1)}) &= -\frac{1}{1+x_{1,1}} + \frac{x_{1,1}x_{1,2}}{\xi^2(1+x_{1,1})^3} + O(\xi^{-4}), \in]-1, 0[, \\ \zeta(b_2^{(1)}) &= -\frac{1+x_{1,1}}{1+x_{1,1}+x_{1,2}}\xi^2 - \frac{x_{1,1}x_{1,2}}{(1+x_{1,1})(1+x_{1,1}+x_{1,2})} - \frac{x_{1,1}x_{1,2}}{\xi^2(1+x_{1,1})^3} + O(\xi^{-4}) \in]-\xi^2, -1[, \end{aligned}$$

and, using (71), in the local coordinates $\tilde{\zeta}$ we have

$$\begin{aligned} \tilde{\zeta}(b_1^{(1)}) &= -\frac{\kappa_2 x_{1,1}(\kappa_4 - \kappa_3) + \kappa_3(\kappa_4 - \kappa_2)}{x_{1,1}(\kappa_4 - \kappa_3) + \kappa_4 - \kappa_2} + \frac{x_{1,1}(1+x_{1,1}+x_{1,2})(\kappa_4 - \kappa_3)(\kappa_4 - \kappa_2)(\kappa_3 - \kappa_2)}{\xi^2(1+x_{1,1})(x_{1,1}(\kappa_4 - \kappa_3) + \kappa_4 - \kappa_2)^2} + O(\xi^{-4}), \in]\kappa_2, \kappa_3[\\ \tilde{\zeta}(b_2^{(1)}) &= \kappa_4 - \frac{(\kappa_4 - \kappa_2)(\kappa_4 - \kappa_3)x_{1,2}}{\xi^2(1+x_{1,1})(\kappa_3 - \kappa_2)} + O(\xi^{-4}) \in]\kappa_3, \kappa_4[. \end{aligned}$$

Similarly, the vacuum poles in Γ_2 in the local coordinate ζ take the form

$$\begin{aligned}\zeta(b_1^{(2)}) &= -\frac{x_{1,1}}{x_{1,1}+x_{2,1}} + \frac{x_{1,1}(x_{1,1}x_{1,2}x_{2,1}+x_{1,2}x_{2,1}-x_{2,2})}{\xi x_{1,2}(x_{1,1}+x_{2,1})^2} + O(\xi^{-2}) \in]-1, 0[, \\ \zeta(b_2^{(2)}) &= -\frac{x_{1,2}(x_{1,1}+x_{2,1})}{x_{1,2}(x_{1,1}+x_{2,1})+x_{2,2}(1+x_{1,1})}\xi^2 + c_1^{(2)}\xi + O(1) \in]-\xi^2, -1[, \end{aligned}$$

with

$$\begin{aligned}c_1^{(2)} &= \frac{x_{1,1}^2(x_{1,2}^2x_{2,1}+x_{1,2}x_{2,1}x_{2,2}-x_{1,2}x_{2,2})+x_{1,2}^2(x_{1,1}x_{2,1}^2+x_{1,1}x_{2,1}-x_{1,1}x_{2,2}+x_{2,1}^2-x_{2,1}x_{2,2})}{(x_{1,1}(x_{1,2}+x_{2,2})+x_{1,2}x_{2,1}+x_{2,2})^2} + \\ &+ \frac{x_{1,1}x_{1,2}x_{2,1}x_{2,2}-2x_{1,1}x_{1,2}x_{2,2}-x_{1,1}x_{2,2}^2-x_{1,2}x_{2,1}x_{2,2}-x_{2,2}^2}{(x_{1,1}(x_{1,2}+x_{2,2})+x_{1,2}x_{2,1}+x_{2,2})^2}, \end{aligned}$$

so that, using (73), in the local coordinates $\tilde{\zeta}$ we have

$$\begin{aligned}\tilde{\zeta}(b_1^{(2)}) &= \frac{\kappa_2\kappa_4x_{1,1}}{\kappa_4x_{11}+x_{21}(\kappa_4-\kappa_2)} - \frac{(\kappa_4-\kappa_3)(\kappa_4-\kappa_2)}{(\kappa_3-\kappa_2)\xi}, \in]0, \tilde{\zeta}(\alpha_2^{(1)})[, \\ \tilde{\zeta}(b_2^{(2)}) &= \tilde{\zeta}(\alpha_3^{(1)}) - \frac{x_{2,2}(1+x_{11})\kappa_4(\kappa_4-\kappa_2)}{\xi^2x_{1,2}\kappa_2(x_{1,1}+x_{2,1})} + O(\xi^{-3}) \in]\tilde{\zeta}(\alpha_2^{(1)}), \tilde{\zeta}(\alpha_3^{(1)})]. \end{aligned}$$

The effective divisor may be easily computed applying the Darboux transformation generated by $f^{(1)}(\vec{t}) = e^{\theta_2} + x_{1,1}e^{\theta_3} + x_{1,2}e^{\theta_4}$ and $f^{(2)}(\vec{t}) = e^{\theta_1} + \frac{x_{2,2}+x_{1,2}x_{2,1}}{x_{1,1}x_{1,2}}e^{\theta_2} + \frac{x_{2,2}}{x_{1,2}}e^{\theta_3}$.

APPENDIX A. POINTS IN $Gr^{\text{TP}}(N, M)$ AND TOTALLY POSITIVE MATRICES IN CLASSICAL SENSE

In this Section we provide the characterization of the positivity properties of the banded matrices introduced in Definition 5. We show that such matrices are totally positive in classical sense (see Proposition 4) and we parametrize $Gr^{\text{TP}}(N, M)$ using the local coordinates $x_{r,s}$, $r \in [N]$, $s \in [M - N]$, which are naturally associated to our construction (see Proposition 5). We believe that all results presented in this Appendix are known to experts, but we failed to find an appropriate reference for some of them.

Points in the totally positive Grassmannian $Gr^{\text{TP}}(N, M)$ may be represented by real $N \times M$ matrices with all maximal $(N \times N)$ minors strictly positive. $Gr^{\text{TP}}(N, M)$ is the top cell in the sense of Postnikov's decomposition [48] of the totally non-negative Grassmannian $Gr^{\text{TP}}(N, M)$.

Let us discuss the relations between total positivity of matrices in the classical sense and the property of total positivity in the Grassmannian.

Definition 6. *We recall that a matrix B is called totally positive (respectively strictly totally positive) if all minors of all orders of B are non-negative (respectively positive) [47].*

It is easy to establish the following natural connection between points of $Gr^{\text{TP}}(N, M)$ and $N \times (M - N)$ strictly totally positive matrices (see [48]): let the $N \times M$ matrix A , represent a point in $Gr^{\text{TP}}(N, M)$. Then, using the standard elementary operations on rows it can be uniquely transformed to reduced row echelon form:

$$(78) \quad A^{\text{RRE}} = \left[\begin{array}{cccc|cccc} 1 & \cdots & 0 & 0 & 0 & \pm b_{N1} & \pm b_{N2} & \cdots & \pm b_{NM-N} \\ & \ddots & & & & & & \cdots & \\ 0 & \cdots & 1 & 0 & 0 & b_{31} & b_{32} & \cdots & b_{3M-N} \\ 0 & \cdots & 0 & 1 & 0 & -b_{21} & -b_{22} & \cdots & -b_{2M-N} \\ 0 & \cdots & 0 & 0 & 1 & b_{11} & b_{12} & \cdots & b_{1M-N} \end{array} \right]$$

where the matrix B

$$(79) \quad B = \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1M-N} \\ b_{21} & b_{22} & \cdots & b_{2M-N} \\ b_{31} & b_{32} & \cdots & b_{3M-N} \\ \vdots & \vdots & \ddots & \vdots \\ b_{N1} & b_{N2} & \cdots & b_{NM-N} \end{bmatrix}$$

is strictly totally positive.

A convenient characterization of strictly totally positive matrices is the following.

Theorem 8. (Theorem 2.3 page 39, [47]) *B is strictly totally positive if and only if all k -th order minors of B composed by the first k rows and k consecutive columns, and also all k -th order minors of B composed by the first k columns and k consecutive rows are strictly positive for $k = 1, \dots, \min\{N, M-N\}$.*

The number of such minors is $N \times (M - N)$ and they form a basis of coordinates for strictly totally positive $N \times (M - N)$ matrices, since all of the other minors of B may be expressed in terms of subtraction-free rational functions of such coordinates. Since any maximal minor of A^{RRE} is expressed as a minor of B , also all of the maximal minors of A^{RRE} are expressed as subtraction free rational functions of such coordinates, that is they form a totally positive basis in Fomin–Zelevinsky sense [17].

In [51], Talaska studies the problem of reconstructing an element $A \in Gr^{\text{TNN}}(N, M)$ from a subset of its Plücker coordinates $\Delta_I(A)$. For each cell in the Gelfand–Serganova decomposition of $Gr^{\text{TNN}}(N, M)$ (see for instance [48] for necessary definitions), she characterizes a minimal set of Plücker coordinates $T(L)$ sufficient to reconstruct the corresponding element using Postnikov boundary measurement map and Le–diagrams [48]. In this way, she constructs a totally positive basis in Fomin–Zelevinsky sense $T(L)$ associated to the Le–diagram [48] of any point in $Gr^{\text{TNN}}(N, M)$.

It is straightforward to check that for points in $Gr^{\text{TP}}(N, M)$, Talaska’s basis of minors coincides with the basis in Theorem 8.

For our purposes it is convenient to transform the matrix $A^{(\text{RRE})}$ to the **banded form**:

$$(80) \quad A = \begin{bmatrix} 1 & A_2^1 & A_3^1 & A_4^1 & \cdots & A_{M-N+1}^1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 1 & A_3^2 & A_4^2 & \cdots & A_{M-N+1}^2 & A_{M-N+2}^2 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & 1 & A_4^3 & \cdots & A_{M-N+1}^3 & A_{M-N+2}^3 & A_{M-N+3}^3 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 1 & \cdots & \cdots & \cdots & A_{M-2}^{N-1} & A_{M-1}^{N-1} & 0 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 1 & \cdots & \cdots & A_{M-2}^N & A_{M-1}^N & A_M^N \end{bmatrix}$$

Here all elements A_j^i with $j < i$ or $j > M - N + i$ are 0.

This transformation can be achieved by applying the Gauss elimination process starting from the last column.¶

In Proposition 4 we show that A as in (80) is a totally positive matrix in classical sense. For the proof we need the following result

¶ We observe that this transformation from the reduced row echelon form to the banded form corresponds to left multiplication by a $N \times N$ upper triangular matrix with unit determinant, therefore it preserves the point of the Grassmannian.

Lemma 5. Let A be a $N \times M$ matrix in the banded form (80) representing a point of $Gr^{TP}(N, M)$. Consider all $n \times n$ submatrices consisting of consecutive rows and arbitrary columns in increasing order: $A_{[j_1, j_2, \dots, j_n]}^{[i, i+1, \dots, i+n-1]}$ for all $n \in [N]$, $i \in [N - n + 1]$. Then their determinants are non-negative:

$$\Delta_{[j_1, j_2, \dots, j_n]}^{[i, i+1, \dots, i+n-1]} \geq 0.$$

Moreover, this determinant is positive if and only if the corresponding submatrix has no zero columns or rows. In particular, all elements A_j^i with $i \leq j \leq M - N + i$ are positive.

Proof. If the submatrix has a zero column, then the associated minor is zero. The condition that the matrix has no zero columns means exactly that $j_1 \geq i$, $j_n \leq M - N + i + n - 1$. Then

$$\Delta_{[j_1, j_2, \dots, j_n]}^{[i, i+1, \dots, i+n-1]} = \Delta_{[1, 2, \dots, i-1, j_1, j_2, \dots, j_n, M-N+i+n, \dots, M]} > 0.$$

In particular, for $i \leq j \leq M - N + i$,

$$A_j^i = \Delta_{[1, 2, \dots, i-1, j, M-N+i+1, \dots, M]} > 0.$$

(Condition $i \leq j \leq M - N + i$ guarantees that this minor has no repeating columns and the columns are in increasing order). \square

Proposition 4. Let A be a $N \times M$ matrix in the banded form (80) representing a point of $Gr^{TP}(N, M)$. Then the matrix A is totally positive.

Proof. We know already that all maximal $(N \times N)$ minors are strictly positive. By Theorem 2.13 in [47], page 56, Lemma 5 implies that the matrix A is totally positive. \square

The following proposition shows that the representation of a point in $Gr^{TP}(N, M)$ through a totally positive matrix in banded form as in (80) is naturally linked to the strictly totally positive $N \times (M - N)$ matrix B defined in (79) and gives another criterion to check the total positivity property.

Proposition 5. Let A be a matrix in banded form with $A_i^i = 1$, $i \in [N]$, and $A_j^i = 0$ if and only if $j < i$ or $j > M - N + i$, with $i \in [N]$, $j \in [M]$. Let

$$(81) \quad x_{r,s} = \Delta_{[N-r+1+s, \dots, N+s]}^{[N-r+1, \dots, N]}(A), \quad r \in [N], \quad s \in [M - N].$$

Then A represents a point of $Gr^{TP}(N, M)$ if and only if $x_{r,s} > 0$, $\forall r \in [N]$, $\forall s \in [M - N]$. Moreover in such a case

$$(82) \quad x_{r,s} = \begin{cases} \Delta_{[s, \dots, s+r-1]}^{[1, \dots, r]}(B), & r \leq s \leq M - N - k + 1, \\ \Delta_{[r, \dots, r+s-1]}^{[1, \dots, s]}(B), & s < r \leq N - s, \end{cases}$$

with B as in (79).

Proof. The minors $x_{r,s}$ may be transformed to maximal $N \times N$ minors of A , so they have to be all positive. Let now A be in banded form with all $x_{r,s} > 0$ as defined in (81) and put it in RRE form. By definition it takes the form as in (78) with pivot set $\{1, \dots, N\}$. Let B be the associated matrix as in (79). Then the minors of B formed by the first r rows and consecutive columns and the minors formed by the first r columns and r consecutive rows, by construction, are just the $x_{r,s}$ minors of the matrix A ($r \in [N]$, $s \in [M - N]$) as in (82).

Then by Theorem 8, B is strictly totally positive if and only if $x_{r,s}$ are all positive and, in such case, A represents a point $Gr^{TP}(N, M)$. \square

Remark 15. The coordinates $x_{r,s}$ coincide with the basis of minors in Lemma 5 and with a totally positive basis in Fomin–Zelevinsky sense, and we shall refer to them simply as the **FZ**-basis.

The following Corollary is the key observation which allows to express the recursive construction of the M-curve and of the wavefunction in invariant form.

Corollary 6. *Let A be the banded totally positive matrix defined above and representing a given point in $Gr^{TP}(N, M)$. Then all of its minors of any order are either zero because they contain a zero row or a zero column, or they are subtraction-free rational expressions in the FZ-basis $x_{r,s}$. In particular the minors of A formed by the last r rows and r columns are subtraction free rational expressions of the elements $x_{l,s}$, $l \in [r]$, $s \in [M - N]$ of the FZ-basis.*

We also require the following version of Fekete's Lemma (see [47], page 37), adapted to our setting:

Lemma 6. *Let $N \leq M$ and assume A to be a $N \times M$ banded matrix in the form (80) with the following properties:*

- (1) *Consider the submatrix \hat{A} obtained from A by removing the first row and the first column. All $N - 1$ -order minors of \hat{A} are positive.*
- (2) *All N -order minors of A composed from consecutive columns are also positive.*

Then all N -order minors of A are positive.

APPENDIX B. LEMMAS FOR THE PROOF OF THEOREM 6

The proof of Theorem 6 requires a series of analytic estimates which we provide in this Appendix. We assume here that $\xi \gg 1$.

In the next Lemma for a fixed $r \in [N]$ we assume that the vacuum wave function at infinity is a linear combination of its values at the double points λ_j with positive coefficients, and we show that there exists an unique collection of corresponding divisor points b_k located in proper intervals, we provide the estimates on the positions of the points b_k , and for all marked points α_n we compute the vacuum wave function at leading order in ξ for all phases. The points α_n are the double points attached to the next component of the spectral curve.

Lemma 7. *Let $c_n > 0$, $n \in [M - N + 1]$ and such that $\sum_{n=1}^{M-N+1} c_n = 1$. Let $\lambda_1 = 0$, $\lambda_k = -\xi^{2(k-2)}$, $k = 2, \dots, M - N + 1$ and define*

$$C_n(\zeta) = c_n \frac{\prod_{j \neq n} (\zeta - \lambda_j)}{\prod_{k=1}^{M-N} (\zeta - b_k)}, \quad n \in [M - N + 1].$$

Then

$$C_n(\lambda_j) = \delta_j^n \quad \forall j, n \in [M - N + 1],$$

for uniquely defined poles $b_k = b_k(\xi) \in]\lambda_{k+1}, \lambda_k[$, $k \in [M - N]$, such that for $\xi \gg 1$,

$$b_k(\xi) = -\frac{\sum_{j=1}^k c_j}{\sum_{j=1}^{k+1} c_j} \xi^{2(k-1)} (1 + O(\xi^{-1})).$$

Moreover, in such case $\forall \zeta \in \mathbb{C}$, $\sum_{n=1}^{M-N+1} C_n(\zeta) = 1$ and

$$(83) \quad C_j(\pm \xi^{2s-5}) = \begin{cases} \frac{c_j}{\sum_{l=1}^{s-1} c_l} \cdot (1 + O(\xi^{-1})) & j \in [2, s-1], \\ \pm \frac{c_j}{(\sum_{l=1}^{s-1} c_l)} \cdot \frac{(1 + O(\xi^{-1}))}{\xi^{2(j-s)+1}} & j \in [s, M - N + 1]. \end{cases}$$

Proof. Let $P(\zeta) = \prod_{k=1}^{M-N} (\zeta - b_k)$. Then $C_j(\lambda_j) = 1$ if and only if $P(\lambda_j) = c_j \prod_{k \neq j} (\lambda_j - \lambda_k)$, $j \in [M-N+1]$. Thanks to the positivity of the coefficients c_j , $P(\lambda_j)$ and $P(\lambda_{j+1})$ have opposite signs $j \in [M-N]$ so that poles $b_k \in]\lambda_{k+1}, \lambda_k[$, $k \in [M-N]$.

By construction $Q(\zeta) = \sum_{j=1}^{M-N+1} C_j(\zeta)$ is a rational function of degree less than or equal to $M-N$ and takes the value 1 in $M-N+1$ points, from which we conclude that it is constant to 1 everywhere.

The estimate for the leading order expansion of b_k , $k \in [M-N]$, for $\xi \gg 1$, follows from the fact that, for any $l \in [M-N]$, the l -th symmetric product in b_k s is a linear combination of the l -th symmetric products in λ_l for $l \neq j$, $j \in [M-N+1]$, that is

$$\hat{\pi}_l(b_1, \dots, b_{M-N}) \equiv \sum_{1 \leq j_1 < j_2 < \dots < j_l \leq M-N} \left(\prod_{s=1}^l b_{j_s} \right) = \sum_{j=1}^{M-N+1} c_j \hat{\pi}_l(\lambda_1, \dots, \hat{\lambda}_j, \dots, \lambda_{M-N+1}) \\ \sum_{j=1}^{M-N+1} c_j \left(\sum'_{1 \leq j_1 < j_2 < \dots < j_l \leq M-N+1} \left(\prod_{s=1}^l \lambda_{j_s} \right) \right) = \left(\sum_{j=1}^{M-N+1-l} c_j \right) \xi^{p(l)} + l.o.t.,$$

where

$$p(l) = 2 \sum_{j=M-N-l}^{M-N-1} j = l(2M - 2N - 1 - l),$$

from which we easily get the assertion on the leading order behavior of the poles.

Finally the estimate on the asymptotic behavior of $C_j(\alpha_s)$, $s \in [2, M-N]$ easily follows taking into account of the leading orders of λ_j s and b_k s. \square

In the next Lemma we use the fact that the vacuum wave function at infinity $\Psi_\infty^{(r)}(\vec{t})$ is a linear combination of its known values at the double points $\alpha_n^{(r-1)}$ with unknown coefficients $B_n^{(r)}$. We impose that $\Psi_\infty^{(r)}(\vec{t})$ is equal to the heat hierarchy solution $f^r(\vec{t})$ associated to the $N-r+1$ -th row of the banded matrix \hat{A} plus small correction, which is a linear combination of heat hierarchy solutions associated to rows below. Then the resulting linear system uniquely defines both the coefficients $B_n^{(r)}$ and the small correcting constants $\epsilon_k^{(r)}$. Moreover, if ξ is sufficiently big, the coefficients $B_n^{(r)}$, $\epsilon_k^{(r)}$ are positive and may be explicitly estimated (see Formulas (87)-(88)). Let us point out that the proof of positivity of $B_n^{(r)}$ is essentially based on the Principal Algebraic Lemma.

Lemma 8. *Let $r \in [2, N]$ be fixed and $\xi > 1$. Let $\alpha_n^{(r-1)}$ ($n \in [2, M-N+1]$) as in (36), and*

$$\Psi_\infty^{(r)}(\vec{t}) = \hat{A}_{N-r+1}^{N-r+1} e^{\theta_{N-r+1}} + \sum_{n=2}^{M-N+1} B_n^{(r)} \Psi^{(r-1)}(\alpha_n^{(r-1)}, \vec{t}),$$

for some $B_n^{(r)} \in \mathbb{R}$ and

(84)

$$\Psi^{(r-1)}(\alpha_n^{(r-1)}, \vec{t}) = \sum_{j=1}^M E_j^{(r-1)[n]}(\xi) e^{\theta_j(\vec{t})} = \left\{ \sum_{j=N-r+2}^{N-r+n} \sigma_{n,j}^{(r-1)} e^{\theta_j} + \sum_{j=N-r+n+1}^{N+n-2} \frac{\sigma_{n,j}^{(r-1)} e^{\theta_j}}{\xi^{j-N+r-n-1}} + \right. \\ \left. + \sum_{j=N+n-1}^M \frac{\sigma_{n,j}^{(r-1)} e^{\theta_j}}{\xi^{r-1+2(j-N-n+1)}} \right\} \times \frac{(1+O(\xi^{-1}))}{\sum_{s=N-r+2}^{N-r+n} \sigma_{n,s}^{(r-1)}},$$

where for all $n \in [2, M-N+1]$, $j \in [N-r+2, M]$, $\sigma_{n,j}^{(r-1)} > 0$ are constants independent of ξ , and, moreover:

$$(85) \quad \sigma_{n,j}^{(r-1)} = \begin{cases} \Delta_{[j; N-r+n+1, N-r+n+2, \dots, N+n-2]} & \text{if } j \in [N-r+2, N-r+n] \\ \Delta_{[N-r+n+1, N-r+n+2, \dots, N+n-2; j]} & \text{if } j \in [N+n-1, M]. \end{cases}$$

Then the requirement

$$(86) \quad \Psi_\infty^{(r)}(\vec{t}) = \sum_{j=N-r+1}^M \left(\hat{A}_j^{N-r+1} + \sum_{k=1}^{r-1} \hat{A}_j^{N-r+k+1} \epsilon_k^{(r)} \right) e^{\theta_j},$$

uniquely defines $B_n^{(r)} = B_n^{(r)}(\xi)$, $n \in [2, M - N + 1]$, and $\epsilon_k^{(r)} = \epsilon_k^{(r)}(\xi)$, $k \in [r - 1]$, which are rational in ξ and strictly positive for all $\xi \gg 1$. Moreover the following estimates hold true

$$(87) \quad B_n^{(r)} = \frac{\Delta_{[N-r+n, \dots, N+n]} \left(\sum_{s=N-r+2}^{N-r+n} \Delta_{[s; N-r+n+1, \dots, N+n-1]} \right)}{\Delta_{[N-r+n, \dots, N+n-1]} \Delta_{[N-r+n+1, \dots, N+n]}} (1 + O(\xi^{-1}));$$

$$(88) \quad \epsilon_k^{(r)} = \frac{\sigma_{M-N+1, M-r+k+1}^{(r-1)} \cdot \hat{A}_{M-r+1}^{N-r+1}}{\sigma_{M-N+1, M-r+1}^{(r-1)} \cdot \hat{A}_{M-r+k+1}^{N-r+k+1}} \cdot \frac{1}{\xi^k} (1 + O(\xi^{-1})).$$

Proof. The proof is straightforward since the linear system associated to (86) in $B_j^{(r)}$, $\epsilon_k^{(r)}$ is clearly compatible for $\xi \gg 1$, the coefficients are rational functions in ξ . Let us define

$$(89) \quad \hat{\sigma}_{n,j}^{(r-1)} = \frac{\sigma_{n,j}^{(r-1)}}{\sum_{s=N-r+2}^{N-r+n} \sigma_{n,s}^{(r-1)}}, \quad \forall n \in [2, M - N + 1], j \in [N - r + 2, M],$$

then, for $\xi \gg 1$, the linear system may be expressed as

$$\sum_{\hat{j}=\hat{n}}^{M-N} \hat{\sigma}_{\hat{j}+1, N-r+\hat{n}+1}^{(r-1)} B_{\hat{j}+1}^{(r)} - \sum_{k=1}^{r-1} \epsilon_k^{(r)} \hat{A}_{N-r+\hat{n}+1}^{N-r+k+1} = \hat{A}_{N-r+\hat{n}+1}^{N-r+1} + O(\xi^{-1}), \quad \hat{n} \in [M - N],$$

$$\sum_{j=s}^{r-1} \frac{\hat{\sigma}_{M-N+1+s-j, M-N+s}^{(r-1)}}{\xi^j} B_{M-N+1+s-j}^{(r)} (1 + O(\xi^{-1})) - \sum_{l=s}^{r-1} \epsilon_l^{(r)} \hat{A}_{M-r+1+s}^{N-r+1+l} = 0, \quad s \in [r - 1].$$

Using the Principal Algebraic Lemma and Theorem 5, we easily conclude that, at leading order in ξ the above system is equivalent to the linear system

$$\hat{\Omega} \hat{c} = \hat{p},$$

in the unknowns $\hat{c} = [B_2^{(r)}, \dots, B_{M-N+1}^{(r)}, \epsilon_1^{(r)}, \dots, \epsilon_{r-1}^{(r)}]^T$, where $\hat{p} = [\hat{A}_{N-r+2}^{(N-r+1)}, \dots, \hat{A}_{M-r+1}^{(N-r+1)}, 0, \dots, 0]^T$ and $\hat{\Omega}$ is the $(M - N + r - 1) \times (M - N + r - 1)$ matrix, such that for $\hat{n} \in [M - N]$:

$$\hat{\Omega}_{\hat{j}}^{\hat{n}} = \begin{cases} \hat{\sigma}_{\hat{j}+1, N-r+\hat{n}+1}^{(r-1)}, & \hat{j} \in [\hat{n}, M - N] \\ 0 & \hat{j} \in [\hat{n} - 1], \\ \hat{A}_{N-r+\hat{n}+1}^{M+r-1-\hat{j}}, & \hat{j} \in [M - N + 1, M - N + r - 1], \end{cases}$$

and for $\hat{n} \in [M - N + 1, M - N + r - 1]$

$$\hat{\Omega}_{\hat{j}}^{\hat{n}} = \begin{cases} 0 & \hat{j} \in M - N - 1] \\ \frac{\hat{\sigma}_{M-N+1, N-r+\hat{n}+1}^{(r-1)}}{\xi^{M-N-\hat{n}}}, & \hat{j} = M - N \\ \hat{A}_{N-r+\hat{n}+1}^{M+r-1-\hat{j}}, & \hat{j} \in [M - N + 1, M - N + r - 1], \end{cases}$$

that is

$$\hat{\Omega} = \begin{bmatrix} \frac{\sigma_{2,N-r+2}^{(r-1)}}{\sigma_{2,N-r+2}^{(r-1)}} & \frac{\sigma_{3,N-r+2}^{(r-1)}}{\sigma_{3,N-r+2}^{(r-1)}} & \dots & \dots & \frac{\sigma_{M-N+1,N-r+2}^{(r-1)}}{\sum_{j=N-r+2}^{M-r+1} \sigma_{M-N+1,j}^{(r-1)}} & \hat{A}_{N-r+2}^{N-r+2} & 0 & \dots & 0 \\ 0 & \frac{\sigma_{3,N-r+2}^{(r-1)}}{\sigma_{3,N-r+2}^{(r-1)}} & \dots & \dots & \frac{\sigma_{M-N+1,N-r+3}^{(r-1)}}{\sum_{j=N-r+2}^{M-r+1} \sigma_{M-N+1,j}^{(r-1)}} & \hat{A}_{N-r+3}^{N-r+2} & \hat{A}_{N-r+3}^{N-r+3} & 0 \dots & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \dots & 0 & \frac{\sigma_{s,N-r+s}^{(r-1)}}{\sum_{j=N-r+2}^{M-r+1} \sigma_{s,j}^{(r-1)}} & \dots & \frac{\sigma_{M-N+1,N-r+s}^{(r-1)}}{\sum_{j=N-r+2}^{M-r+1} \sigma_{s,j}^{(r-1)}} & \hat{A}_{N-r+s}^{N-r+2} & \hat{A}_{N-r+s}^{N-r+3} & \dots & \hat{A}_{N-r+s}^N \\ \vdots & \ddots & \ddots & \ddots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & \dots & \dots & \dots & 0 & \frac{\sigma_{M-N+1,M-r+1}^{(r-1)}}{\sum_{j=N-r+2}^{M-r+1} \sigma_{M-N+1,j}^{(r-1)}} & \hat{A}_{M-r+1}^{N-r+2} & \hat{A}_{M-r+1}^{N-r+3} & \dots & \hat{A}_{M-r+1}^N \\ 0 & 0 & \dots & \dots & 0 & \frac{\sigma_{M-N+1,M-r+2}^{(r-1)}}{\xi \left(\sum_{j=N-r+2}^{M-r+1} \sigma_{M-N+1,j}^{(r-1)} \right)} & \hat{A}_{M-r+2}^{N-r+2} & \hat{A}_{M-r+2}^{N-r+3} & \dots & \hat{A}_{M-r+2}^N \\ 0 & 0 & \dots & \dots & 0 & \frac{\sigma_{M-N+1,M-r+2}^{(r-1)}}{\xi^2 \left(\sum_{j=N-r+2}^{M-r+1} \sigma_{M-N+1,j}^{(r-1)} \right)} & 0 & \hat{A}_{M-r+1}^{N-r+3} & \dots & \hat{A}_{M-r+1}^N \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \dots & \dots & 0 & \frac{\sigma_{M-N+1,M}^{(r-1)}}{\xi^{r-1} \left(\sum_{j=N-r+2}^{M-r+1} \sigma_{M-N+1,j}^{(r-1)} \right)} & 0 & \dots & 0 & \hat{A}_M^N \end{bmatrix}.$$

Then the coefficients

$$B_n^{(r)}(\xi) = \hat{B}_n^{(r)} (1 + O(\xi^{-1})), \quad n \in [2, M - N + 1]$$

where $\hat{B}_n^{(r)}$ are as in Theorem 5, while $\epsilon_k^{(r)} = O(\xi^{-k})$, $r \in [N - i]$ and at leading order are as in (87) and (88). In particular, if $\sigma_{n,j}^{(r-1)}$ are all positive then also $\hat{B}_l^{(r)}(\xi) > 0$, $l \in [2, M - N + 1]$ and $\epsilon_k^{(r)}(\xi) > 0$, $k \in [r - 1]$ for all $\xi \gg 1$. \square

In the next Lemma we refine the results of Theorem 5, namely at double points $\alpha_n^{(r)}$ we estimate the vacuum wave function at the leading order in ξ for all phases. We recall that in Theorem 5 only the coefficients in front of the dominant phases were computed.

Lemma 9. *Let $r \in [2, N]$ be fixed. Let λ_j ($j \in [M - N + 1]$) as in (35), $\alpha_n^{(r-1)}$ ($n \in [2, M - N + 1]$) as in (36),*

$$\Psi^{(r)}(\zeta, \vec{t}) = C_1(\zeta) e^{\theta_{N-r+1}} + \sum_{n=2}^{M-N+1} C_n(\zeta) \Psi^{(r-1)}(\alpha_n^{(r-1)}, \vec{t}),$$

with $\Psi^{(r-1)}(\alpha_n^{(r-1)}, \vec{t})$ as in (84),

$$C_n(\zeta) = \hat{B}_n^{(r)} \frac{\prod_{j \neq n}^{M-N+1} (\zeta - \lambda_j)}{\prod_{k=1}^{M-N} (\zeta - b_k^{(r)})}, \quad n \in [M - N + 1],$$

with

$$\hat{B}_n^{(r)}(\xi) = \begin{cases} \hat{A}_{N-r+1}^{N-r+1} & n = 1 \\ \frac{B_n^{(r)}(\xi)}{1 + \sum_{k=1}^{r-1} \epsilon_k^{(r)}(\xi)}, & n \in [2, M - N + 1] \end{cases},$$

with $B_n^{(r)}(\xi)$, $\epsilon_k^{(r)}(\xi)$ as in Lemma 8, and $b_k^{(r)}(\xi)$, ($k \in [M - N]$) as in Lemma 7 with $c_n = \hat{B}_n^{(r)}(\xi)$. Let $\hat{\sigma}_{n,j}^{(r-1)}$ as in (89), with $\sigma_{n,j}^{(r-1)} > 0$ as in Lemma 8.

Then, for $\alpha_n^{(r)}$ ($n \in [2, M - N + 1]$) as in (36), we have

$$(90) \quad \Psi^{(r)}(\alpha_n^{(r)}, t) = \left(\sum_{j=N-r+1}^{N-r+n-1} \hat{\sigma}_{n,r}^{(r)} e^{\theta_j} + \sum_{j=N-r+n}^{N+n-2} \frac{\hat{\sigma}_{n,j}^{(r)}}{\xi^{j-N+r-n+1}} e^{\theta_j} + \sum_{j=n+N-2}^M \frac{\hat{\sigma}_{n,j}^{(r)}}{\xi^{2(j-N-n+2)+r}} e^{\theta_j} \right) (1 + O(\xi^{-1})),$$

for uniquely defined positive constants $\sigma_{n,s}^{(r)}$ such that, for any $n \in [2, M - N + 1]$,

$$(91) \quad \hat{\sigma}_{n,s}^{(r)} = \begin{cases} \frac{\hat{B}_1^{(r)}}{\sum_{j=1}^{n-1} \hat{B}_j^{(r)}}, & s = N - r + 1, \\ \frac{\sum_{j=s+r-N}^{n-1} \hat{B}_j^{(r)} \cdot \hat{\sigma}_{j,s}^{(r-1)}}{\sum_{i=1}^{n-1} \hat{B}_i^{(r)}}, & s \in [N - r + 2, N - r + n - 1], \\ \frac{\hat{B}_{n-1}^{(r)} \cdot \hat{\sigma}_{n-1,s}^{(r-1)} + \hat{B}_n^{(r)} \cdot \hat{\sigma}_{n,s}^{(r-1)}}{\sum_{i=1}^{n-1} \hat{B}_i^{(r)}}, & s \in [N - r + n, N + n - 2], \\ \frac{\sum_{j=n}^{s-N+1} \hat{B}_j^{(r)} \cdot \hat{\sigma}_{j,s}^{(r-1)}}{\sum_{i=1}^{n-1} \hat{B}_i^{(r)}}, & s \in [N + n - 1, M]. \end{cases}$$

Finally, by construction,

$$\sum_{k=N-r+1}^{N-r+n-1} \hat{\sigma}_{n,k}^{(r)} = 1, \quad \forall n \in [2, M - N + 1].$$

The proof of the above Lemma is straightforward and follows by direct inspection of the leading order in ξ for each phase θ_s , $s \geq N - r + 1$, using the definition of $\Psi^{(r)}(\zeta, \vec{t})$, and the asymptotic expansions of $C^{(r)}(\alpha_s^{(r)})$, as in (83), with $c_j = \hat{B}_j^{(r)} = \hat{B}_j^{(r)} (1 + O(\xi^{-1}))$ and of $\Psi^{(r-1)}(\alpha_s^{(r-1)}, \vec{t})$ as in (84).

Remark 16. Lemmas 8 and 9 allow to compute the coefficients $B_n^{(r)}$, $\epsilon_k^{(r)}$ and $\sigma_{n,s}^{(r)}$ recursively in $r \in [N]$, starting from the case $r = 1$ computed directly in Proposition 3.

The coefficients $\hat{B}_n^{(1)}$, $\sigma_{n,k}^{(1)}$ are all positive for $n \in [2, M - N + 1]$, $s \in [N - r + 2, M]$ by the same Proposition 3. Moreover $\epsilon_k^{(r)}$ and $\sigma_{n,k}^{(r)}$ respectively in (88) and in (91) are subtraction free rational expressions in $\hat{B}_n^{(r)}$, $\sigma_{n,k}^{(r-1)}$ and the matrix entries of \hat{A} . The total positivity property of the matrix \hat{A} ensures that $\hat{B}_n^{(r)} > 0$, thanks to Theorem 5. As a consequence we get that also all $\epsilon_k^{(r)} > 0$ and $\sigma_{n,s}^{(r)} > 0$.

In Theorem 5, we have computed $\sigma_{n,s}^{(r)}$ for $s \in [N - r + 1, N - r + n]$, $n \in [2, M - N + 1]$ (see (51)). In the next Lemma we compute explicitly these coefficients also for $s \in [N + n - 1, M]$, $n \in [2, M - N + 1]$.

Lemma 10. Let $r \in [2, N]$ and suppose that $\sigma_{n,s}^{(r-1)}$, $\hat{B}_n^{(r)}$ are as in (85) and (52), respectively. Then, for any $n \in [2, M_N + 1]$, we have

$$(92) \quad \sigma_{n,j}^{(r)} = \begin{cases} \Delta_{[j; N-r+n, N-r+n+1, \dots, N+n-2]}, & \text{if } j \in [N - r + 1, N - r + n - 1] \\ \Delta_{[N-r+n, N-r+n+1, \dots, N+n-2; j]}, & \text{if } j \in [N + n - 1, M]. \end{cases}$$

Proof. The case $j \in [N - r + 1, N - r + n - 1]$, $n \in [2, M - N + 1]$ is just (51) which is proven using Lemmas 3 and 4. The case $j \in [N + n - 1, M]$, $n \in [2, M - N + 1]$ follows in a similar way using the identity

$$(93) \quad \sum_{n=k}^{M-N+1} \frac{\Delta_{[N-r+n, \dots, N+n-1]} \cdot \Delta_{[N-r+n+1, \dots, N+n-2; j]}}{\Delta_{[N-r+n+1, \dots, N+n-1]} \cdot \Delta_{[N-r+n, \dots, N+n-2]}} = \frac{\Delta_{[N-r+k, \dots, N+k-2; j]}}{\Delta_{[N-r+k, \dots, N+k-2]}},$$

for $r \in [N - 1]$, $k \in [2, M - N + 1]$, $j \in [N + k - 1, M]$, which may be proven recursively along the same lines as for (56). \square

REFERENCES

- [1] Abenda S., “On a family of KP multi-line solitons associated to rational degenerations of real hyperelliptic curves and to the finite non-periodic Toda hierarchy”, *J.Geom.Phys.* **119** (2017), 112-138.
- [2] Abenda S.; Grinevich P.G., “KP theory, plane-bipartite networks in the disk and rational degenerations of M-curves”, [arXiv:1801.00208](#).
- [3] Arbarello E.; Cornalba M.; Griffiths P.A., “Geometry of algebraic curves. Volume II. With a contribution by Joseph Daniel Harris”, *Grundlehren der Mathematischen Wissenschaften* **268**, Springer, Heidelberg, (2011), xxx+963 pp.
- [4] Arkani-Hamed N.; Bourjaily J.L.; Cachazo F.; Goncharov A.B.; Postnikov A.; Trnka J., “Scattering Amplitudes and the Positive Grassmannian”, [arXiv:1212.5605](#).
- [5] Arkani-Hamed N.; Bourjaily J.L.; Cachazo F.; Goncharov A.B.; Postnikov A.; Trnka J., “Grassmannian geometry of scattering amplitudes”, Cambridge University Press, Cambridge, (2016), ix+194 pp.
- [6] Biondini, G.; Kodama, Y., “On a family of solutions of the Kadomtsev-Petviashvili equation which also satisfy the Toda lattice hierarchy”, *J. Phys. A* **36** (2003), no. 42, 10519-10536.
- [7] Boiti, M.; Pempinelli, F.; Pogrebkov, A.K.; Prinari, B., “Towards an inverse scattering theory for non-decaying potentials of the heat equation”, *Inverse Problems* **17** (2001), 937-957.
- [8] Boiti, M.; Pempinelli, F.; Pogrebkov, A. K., “Properties of the solitonic potentials of the heat operator”, *Theoret. and Math. Phys.*, **168**, (2011), no. 1, 865-874.
- [9] Cavaretta, A. S. Jr.; Dahmen, W. A.; Micchelli, C. A.; Smith, P. W., “A factorization theorem for banded matrices”, *Linear Algebra Appl.* **39** (1981), 229-245.
- [10] Chakravarty, S.; Kodama, Y., “Classification of the line-soliton solutions of KPII”, *J. Phys. A* **41** (2008), no. 27, 275209, 33 pp.
- [11] Chakravarty, S.; Kodama, Y., “Soliton solutions of the KP equation and application to shallow water waves”, *Stud. Appl. Math.* **123** (2009), no. 1, 83-151.
- [12] de Boor, C.; Pinkus, A., “The approximation of a totally positive band matrix by a strictly banded totally positive one”, *Linear Algebra Appl.* **42** (1982), 81-98.
- [13] Dickey, L. A., “Soliton equations and Hamiltonian systems”, Second edition. *Advanced Series in Mathematical Physics*, **26**, World Scientific Publishing Co., Inc., River Edge, NJ, (2003), xii+408 pp.
- [14] Dimakis, A.; Müller-Hoissen, F., “KP line solitons and Tamari lattices”, *J. Phys. A* **44** (2011), no. 2, 025203, 49 pp.
- [15] Dubrovin, B. A.; Krichever, I. M.; Novikov, S. P., “Integrable systems. Dynamical systems, IV”, *Encyclopaedia Math. Sci.*, **4**, Springer, Berlin, (2001), 177-332.
- [16] Dubrovin, B. A.; Natanzon S. M., “Real theta-function solutions of the Kadomtsev-Petviashvili equation”, *Izv. Akad. Nauk SSSR Ser. Mat.*, **52** (1988), no. 2, 267-286.
- [17] Fomin, S.; Zelevinsky, A., “Double Bruhat cells and total positivity”, *J. Amer. Math. Soc.* **12** (1999), no. 2, 335-380.
- [18] Fomin, S.; Zelevinsky, A., “Cluster algebras I: foundations.”, *J. Am. Math. Soc.* **15** (2002), 497-529.
- [19] Gantmacher F.R.; Krein M.G., “Sur les matrices oscillatoires”, *C.R.Acad.Sci.Paris* **201** (1935), 577-579.
- [20] Gekhtman, M.; Shapiro, M.; Vainshtein, A., “Cluster algebras and Poisson geometry”, *Mathematical Surveys and Monographs*, **167**, American Mathematical Society, Providence, RI, (2010), xvi+246 pp.
- [21] Goncharov, A.B.; Kenyon, R., “Dimers and cluster integrable systems”, *Ann. Sci. c. Norm. Supr. (4)* **46** (2013), no. 5, 747-813.
- [22] Griffiths P.; Harris J., “Principles of Algebraic Geometry”, John Wiley & Sons, (1978).
- [23] Gross, B.H.; Harris, J., “Real algebraic curves”, *Ann. Sci. cole Norm. Sup. série 4*, **14** (1981), no. 2, 157-182.
- [24] Harnack, A., “Über die Vieltheiligkeit der ebenen algebraischen Curven”, *Math. Ann.* **10** (1876), 189-199.
- [25] Hirota, R., “The direct method in soliton theory”, *Cambridge Tracts in Mathematics*, **155**, Cambridge University Press, Cambridge, (2004), xii+200 pp.
- [26] Kadomtsev, B.B.; Petviashvili, V.I., “On the stability of solitary waves in weakly dispersive media”, *Sov. Phys. Dokl.* **15** (1970), 539-541.
- [27] Kenyon, R.; Okounkov, A.; Sheffield, S., “Dimers and amoebae”, *Ann. of Math. (2)* **163** (2006), no. 3, 1019-1056.
- [28] Kodama, Y.; Williams, L.K., “KP solitons, total positivity, and cluster algebras”, *Proc. Natl. Acad. Sci. USA* **108** (2011), no. 22, 8984-8989.

- [29] Kodama, Y.; Williams, L., “The Deodhar decomposition of the Grassmannian and the regularity of KP solitons”, *Adv. Math.* **244** (2013), 979-1032.
- [30] Kodama, Y.; Williams, L., “KP solitons and total positivity for the Grassmannian”, *Invent. Math.* **198** (2014), no. 3, 637-699.
- [31] Krichever, I. M., “An algebraic-geometric construction of the Zakharov-Shabat equations and their periodic solutions”. (Russian) *Dokl. Akad. Nauk SSSR* **227** (1976), no. 2, 291-294.
- [32] Krichever, I. M., “Integration of nonlinear equations by the methods of algebraic geometry”, (Russian) *Funkcional. Anal. i Priloen.* **11** (1977), no. 1, 15-31, 96.
- [33] Krichever, I.M., “Spectral theory of nonstationary Schroedinger operators and non stationary Peierls model.”, *Funk.anal. i pril.* **20** (1986), no. 3, 42–54.
- [34] Krichever, I.M.; Vaninsky, K.L., “The periodic and open Toda lattice.” *AMS/IP Stud. Adv. Math.*, **33**, Amer. Math. Soc., Providence, RI, (2002), 139–158.
- [35] Lam, T., “Dimers, webs, and positroids”, *J. Lond. Math. Soc. (2)* **92** (2015), no. 3, 633–656.
- [36] Lusztig, G., “Total positivity in reductive groups.” *Lie theory and geometry*, 531-568, *Progr. Math.*, **123**, Birkhäuser Boston, Boston, MA, (1994).
- [37] Lusztig, G., “Total positivity in partial flag manifolds”, *Represent. Theory* **2** (1998), 70-78.
- [38] Lusztig, G., “A survey of total positivity”, *Milan J. Math.* **76** (2008), 125-134.
- [39] Malanyuk, T.M., “A class of exact solutions of the Kadomtsev-Petviashvili equation”, *Russian Math. Surveys* **46** (1991), no. 3, 225-227.
- [40] Marsh, R.J.; Rietsch, K., “Parametrizations of flag varieties”, *Represent. Theory* **8** (2004), 212-242.
- [41] Matveev, V.B., “Some comments on the rational solutions of the Zakharov-Schabat equations”, *Letters in Mathematical Physics*, **3** (1979), 503–512.
- [42] Matveev, V. B.; Salle, M.A., “Darboux transformations and solitons.”, *Springer Series in Nonlinear Dynamics*. Springer-Verlag, Berlin, (1991).
- [43] Miwa, T.; Jimbo, M.; Date, E., “Solitons. Differential equations, symmetries and infinite-dimensional algebras”, *Cambridge Tracts in Mathematics*, **135**, Cambridge University Press, Cambridge, (2000), x+108 pp.
- [44] Natanzon S.M., “Moduli of real algebraic surfaces, and their superanalogues. Differentials, spinors, and Jacobians of real curves”, *Russian Mathematical Surveys*, **54** (1999), no 6, 1091-1147.
- [45] Novikov, S.P.; Manakov, S. V.; Pitaevskii, L. P.; Zakharov, V. E., “Theory of solitons. The inverse scattering method”, *Contemporary Soviet Mathematics. Consultants Bureau [Plenum]*, New York, (1984), xi+276 pp.
- [46] Petrowsky, I., “ On the Topology of Real Plane Algebraic Curves”, *Annals of Mathematics Second Series*, **39** (1938), no 1, 189-209.
- [47] Pinkus, A., “Totally positive matrices”, *Cambridge Tracts in Mathematics*, **181**, Cambridge University Press, Cambridge, (2010) xii+182 pp. ISBN: 978-0-521-19408-2.
- [48] Postnikov A., “Total positivity, Grassmannians, and networks.”, [arXiv:math/0609764](https://arxiv.org/abs/math/0609764) [math.CO].
- [49] Sato, M., “Soliton equations as dynamical systems on infinite-dimensional Grassmann manifold”, in: *Non-linear PDEs in Applied Sciences (US-Japan Seminar, Tokyo)*, P. Lax and H. Fujita eds., North-Holland, Amsterdam , (1982), 259-271.
- [50] Schoenberg J., “Über variationsvermindende lineare Transformationen”, *Mathematische Zeitschrift* **32**, (1930), 321-328.
- [51] Talaska, K., “Combinatorial formulas for q -coordinates in a totally nonnegative Grassmannian”, *J. Combin. Theory Ser. A* **118** (2011), no. 1, 58-66.
- [52] Veselov, A.P.; Novikov, S.P., “Finite-zone, two-dimensional, potential Schrödinger operators. Explicit formulas and evolution equations”, *Soviet Math. Dokl.* **30** (1984), 588-591.
- [53] Veselov, A.P.; Novikov, S.P., “Finite-zone, two-dimensional Schrödinger operators. Potential operators”, *Soviet Math. Dokl.* **30** (1984), 705-708.
- [54] Viro O. Ya., “Real plane algebraic curves: constructions with controlled topology”, *Leningrad Math. J.* **1** (1990), no. 5, 1059-1134.
- [55] Whitney, A. M., “A reduction theorem for totally positive matrices”, *J. Analyse Math.* **2**, (1952), 88–92.
- [56] Zakharov V.E.; Shabat A.B., “A scheme for integrating the nonlinear equations of mathematical physics by the method of the inverse scattering problem. I”, *Functional Analysis and Its Applications* **8** (1974), no 3, 226-235.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI BOLOGNA, ITALY, SIMONETTA.ABENDA@UNIBO.IT

L.D.LANDAU INSTITUTE FOR THEORETICAL PHYSICS, PR. AK SEMENOVA 1A, CHERNOGOLOVKA, 142432, RUSSIA, PGG@LANDAU.AC.RU, LOMONOSOV MOSCOW STATE UNIVERSITY, FACULTY OF MECHANICS AND MATHEMATICS, RUSSIA, 119991, MOSCOW, GSP-1, 1 LENINSKIYE GORY, MAIN BUILDING,, MOSCOW INSTITUTE OF PHYSICS AND TECHNOLOGY, 9 INSTITUTSKIY PER., DOLGOPRUDNY, MOSCOW REGION, 141700, RUSSIA.