

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Use of tandem mass spectrometry (LC-MS-MS) for the measurement of thyroid hormones in dogs with spontaneous hypothyroidism

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Carotenuto G, L.F. (2019). Use of tandem mass spectrometry (LC-MS-MS) for the measurement of thyroid hormones in dogs with spontaneous hypothyroidism. JOURNAL OF VETERINARY INTERNAL MEDICINE, 33(2), 68-68.

Availability:

This version is available at: <https://hdl.handle.net/11585/640860> since: 2018-08-23

Published:

DOI: <http://doi.org/>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

1. [Use of Tandem Mass Spectrometry \(LC-MS/MS\) for the Measurement of Thyroid Hormones in Dogs with Spontaneous Hypothyroidism](#)
G. Carotenuto; F. Lunetta; S. De Arcangeli; E. Malerba; F. Del Baldo; S. Corradini; F. Fracassi

Search Result #1: **Use of Tandem Mass Spectrometry (LC-MS/MS) for the Measurement of Thyroid Hormones in Dogs with Spontaneous Hypothyroidism** [Click to go to the TOP](#)

Author(s): G. Carotenuto; F. Lunetta; S. De Arcangeli; E. Malerba; F. Del Baldo; S. Corradini; F. Fracassi
Address (URL):

Use of Tandem Mass Spectrometry (LC-MS/MS) for the Measurement of Thyroid Hormones in Dogs with Spontaneous Hypothyroidism
28th ECVIM-CA Congress, 2018
G. Carotenuto; F. Lunetta; S. De Arcangeli; E. Malerba; F. Del Baldo; S. Corradini; F. Fracassi
University of Bologna, Ozzano dell'Emilia, Italy

In human medicine, liquid chromatography tandem mass spectrometry (LC-MS/MS) is actually considered the "gold standard" for measurement of many hormone concentrations, and it is widely used in clinical practice; its diagnostic performance has never been investigated in dogs with hypothyroidism (DWH).

The aim of this study was to determine whether serum concentrations of fT_4 , fT_3 , rT_3 , $3.3-T_2$, $3.5-T_2$, measured with LC-MS/MS, were able to differentiate DWH (n=13) from dogs with non-thyroidal illness (DNTI) (n=12), septic dogs (SD) (n=12), and healthy dogs (HD) (n=12).

Hypothyroidism was diagnosed based on consistent clinical signs, laboratory findings, total T_4 (TT_4) and cTSH concentrations below and above the reference interval (RI), respectively; in dogs with normal cTSH, a rhTSH stimulation test was performed to confirm the diagnosis. In DNTI, hypothyroidism was excluded upon a negative result of a rhTSH stimulation test. SD were diagnosed based on alteration of temperature, cardiac and respiratory frequency, differential leukocyte count and C-reactive protein concentration above RI. HD were considered healthy upon history and physical examination. Hormone evaluations were performed with LC-MS/MS on surplus serum stored at $-80^{\circ}C$. TT_4 and cTSH were measured using a validated immunoassay (Immulite®).

Non-significant differences considering signalment, age and body weight were found between groups. Median TT_4 and fT_4 serum concentrations were significantly higher ($p<0.001$) in HD compared to DNTI, DWH and SD. Median fT_3 serum concentration was significantly lower in DWH and DNTI compared to SD ($p<0.001$ and $p=0.0091$, respectively) and HD ($p<0.001$ and $p=0.0024$, respectively). Median rT_3 serum concentration was significantly lower in DWH compared to SD ($p=0.0141$) and HD ($p=0.0128$). Median $3.3-T_2$ serum concentration was significantly higher in DWH compared to DNTI ($p=0.0038$) and HD ($p=0.0447$). There were non-significant differences regarding median $3.5-T_2$ serum concentrations among the dogs of the four groups.

Using the ROC curve analysis to differentiate DWH from DNTI+SD, an AUC of 0.86 ($p=0.003$), 0.76 ($p=0.009$) and 0.75 ($p=0.012$) was obtained for fT_3 , fT_4 and TT_4 , respectively. Values of $fT_3 < 0.61$ pmol/L better discriminated hypothyroidism with 69% sensitivity (95%CI: 39–91%), 83% specificity (95%CI: 63–95%) and accuracy of 0.86 (95%CI: 0.74–0.98).

Although serum fT_3 and fT_4 (LC-MS/MS) have shown better performances than the serum TT_4 (Immulite®) in identifying DWH, the overlap between DWH and DNTI+SD was unfortunately relevant also for the thyroid hormone measurements with LC-MS/MS. Despite the introduction of new analytical methods, the use of dynamic tests (e.g., rhTSH stimulation test) remains the better method to discriminate DWH from DNTI.

DISCLOSURES

No disclosures to report