



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Computational challenges in Astrochemistry

This is the submitted version (pre peer-review, preprint) of the following publication:

Published Version:

Biczysko, M., Bloino, J., Pizzarini, C. (2018). Computational challenges in Astrochemistry. WILEY INTERDISCIPLINARY REVIEWS. COMPUTATIONAL MOLECULAR SCIENCE, 8(3), e1349/1-e1349/38 [10.1002/wcms.1349].

Availability:

This version is available at: <https://hdl.handle.net/11585/655711.2> since: 2019-01-22

Published:

DOI: <http://doi.org/10.1002/wcms.1349>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final accepted manuscript of:


M. Biczysko, J. Bloino, C. Puzzarini. Computational Challenges in Astrochemistry. *WIREs Comput Mol Sci*, 8, e1349 (2018). DOI:

Available at: <https://doi.org/10.1002/wcms.1349>

© 2017 Wiley Periodicals, Inc.

Computational Challenges in Astrochemistry

Malgorzata Biczysko, Julien Bloino, Cristina Puzzarini*



TOC - Cosmic evolution from simplicity to complexity: the crucial challenges in the computational spectroscopy of astroCOMs can be successfully overcome by combining state-of-the-art quantum-mechanical approaches with ad hoc treatments of the nuclear motion

Computational Challenges in Astrochemistry

Malgorzata Biczysko^a, Julien Bloino^b, Cristina Puzzarini^{c*}

^a *International Center for Quantum and Molecular Structures, Shanghai University, China*

^b *Institute of Chemistry of Organometallic Compounds, CNR, Pisa, Italy*

^c *Department of Chemistry "Giacomo Ciamician", University of Bologna, Italy*

Article Type:

Advanced Review

Abstract

Cosmic evolution is the tale of progressive transition from simplicity to complexity. The newborn universe starts with the simplest atoms formed after the Big Bang and ends with "astronomical complex organic molecules" (astroCOMs). Understanding the chemical evolution of the universe is one of the main aims of astrochemistry, with the starting point being the knowledge whether a molecule is present in the astronomical environment under consideration and, if so, its abundance. However, the interpretation of astronomical detections and the identification of molecules are not all straightforward. In particular, molecular species characterized by large amplitude motions represent a major challenge for molecular spectroscopy and, in particular, for computational spectroscopy. More in general, for flexible systems the conformational equilibrium needs to be taken into account and accurately investigated. It is shown that crucial challenges in the computational spectroscopy of astroCOMs can be successfully overcome by combining state-of-the-art quantum-mechanical approaches with *ad hoc* treatments of the nuclear motion, thus demonstrating that the rotational and vibrational features can be predicted with the proper accuracy. The second key step in astrochemistry is understanding how astroCOMs are formed and how they chemically evolve toward more complex species. The challenges in the computational chemistry of astroCOMs are related to the derivation of feasible formation routes in the typically harsh conditions (extremely low temperature and density) of the interstellar medium, as well as the understanding of the chemical evolution of small species toward macromolecules. Once all intermediate species and the transition states connecting them in a plausible formation route are identified, state-of-the-art computational approaches allow obtaining structure, energies, spectroscopic properties and thermochemical data for all of them with the required accuracy. The thermochemical

*Correspondence to: cristina.puzzarini@unibo.it

information is then used to compute kinetic rate constants to obtain reliable input parameters for astrochemical models.

Introduction: The interstellar molecular complexity

For many years, the interstellar medium (ISM) was considered too hostile for organic species to be formed. However, roughly forty years ago, the first molecules containing carbon chains and rings were discovered. With time, a rapidly increasing number of molecules showing some significant complexity has been observed in space. Indeed, the detection of almost 200 molecules in interstellar or circumstellar shells (see, for example, the Cologne Database for Molecular Spectroscopy (CDMS)^{1,2}) suggests that the ISM is characterized by a rich chemistry. Understanding the chemical evolution of the universe is one of the main aims of astrochemistry,^{3,4} with the starting point being the knowledge whether a molecule is present in the astronomical environment under consideration and, if so, its abundance. The second step is understanding how these molecular species are formed and how they evolve chemical-ly toward more complex species. In this scenario, molecular spectroscopy and computational chemistry play a crucial role.⁵⁻¹³ On one side, astronomical observations of spectroscopic signatures provide the unequivocal proof of the presence of chemical species; on the other side, quantum chemistry allows shedding light on their formation/transformation mechanisms, which are nowadays still object of debate and often not experimentally investigable.

Similarly to astronomical observations that started with the detection of the smallest di-/triatomic molecules, the computational support in the field of astrochemistry began with highly accurate theoretical predictions of ro-vibrational transitions of small (di- and triatomic) molecules, thus leading to extended line lists, which are important for the study of hot atmospheres, particularly those of exoplanets, brown dwarfs, and cool stars.¹⁴ However, new challenges are posed by the large amount of data that nowadays need to be analyzed. In fact, the advent of large instantaneous bandwidth receivers characterized by high spectral resolution has opened up the possibility of large spectral surveys with a uniform sensitivity. If on the one hand, they are clearly the ideal mean to obtain a complete census of the species emitting in the recorded spectral survey, on the other hand, their identification is particularly challenging, thus requiring the increasing support from laboratory (both experimental and theoretical) studies. In fact, these new opportunities start to explore areas where the reference experimental laboratory data are lacking.

Among others, the ultimate goal of the detection of prebiotic and biological molecules in the astrochemical environments is one of the most appealing challenges of astrochemistry.¹⁵ In recent years, the Prebiotic Interstellar Molecular Survey (PRIMOS) project at the Green Bank Telescope (GBT)^{16,17} contributed to the detection of over a dozen of new interstellar molecules (containing 6 to 11 atoms), also including the first example of chiral molecule, methyloxirane.¹⁸ In view of the opportunities opened by the Atacama Large Millimeter/submillimeter Array of radiotelescopes (ALMA), working in the 31–950 GHz frequency range, and by the Mid-Infrared Instrument (MIRI), part of the James Webb Space Telescope (JWST, to be launched in 2018), it is necessary to extend reference data to reach higher frequencies to support astronomical observations. To give an idea, the rotational signatures of molecules showing a certain degree of complexity are often obtained from low-frequency laboratory measurements, well below 100 GHz. Moreover, even for small and widely studied molecules, experimental studies have not yet completely and unequivocally determined the whole far-to-mid range IR spectrum, with some bands (even including fundamental transitions) still missing or not measured with sufficient accuracy or in isolated, “space-like” conditions.¹⁹ Finally, emission spectra of relevance to astrochemical studies for samples composed of mixtures of unknown components are still quantitatively resolved only for a limited number of bands.

The focus of the present review is on the quantum-chemical support to the investigation of the so-called astronomical complex organic molecules, hereinafter referred to as astroCOMs, which are molecules containing at least six atoms, including carbon, hydrogen and, in most cases, at least another heavy element. Examples of astroCOMs, discussed in more details in the following, are gathered in Figure 1. Even if aminoacids and nucleobases have been found in meteorites (see, for example, Refs.^{20,21}), they have not yet been unequivocally observed in the ISM, clearly showing the importance of improved reference spectroscopic data for their detection in different space environments. In this respect, it is encouraging that nowadays it is possible to evaluate structural, thermodynamic and spectroscopic properties of medium-sized molecules with an accuracy that, until recently, could be reached only for very small rigid systems.^{22–25} Highly accurate results can be nowadays obtained by means of composite and/or hybrid models, thus paving the way toward the complete spectro-











































					
	Formamide	Oxirane	<i>trans</i> -Acrolein	<i>cis</i> -Acrolein	Acetyl cyanide
Formula	HC(O)NH ₂	C ₂ H ₄ O	C ₃ H ₄ O	C ₃ H ₄ O	CH ₃ COCN
N atoms	6	7	8	8	8
Mass	45	44	56	56	69
μ (Deb)	4.0	1.9	3.2	2.6	3.6
Detected in ISM or CSE	 Sagittarius B2	 Sagittarius B2	 Sagittarius B2		
Year	 1971	 1997	 2004		
Other Astro implications	fairly typical hot core molecule, also galactic center, Solar-type protostar; Isotopologues: ¹³ C and D	hot core regions, galactic center (10–20 K), low-mass protostar	galactic center (10–20 K), low-mass protostar	less stable by 712 cm ⁻¹	expected to be present in the ISM
					
	<i>gauche</i> -Ethyl mercaptan	<i>trans</i> -Ethyl mercaptan	<i>Tc</i> -Pyruvic acid	Glycine Ip/ttt	Glycine IIn/cc
Formula	CH ₃ CH ₂ SH	CH ₃ CH ₂ SH	C ₂ CH ₃ O ₂ OH	H ₂ NCH ₂ COOH	H ₂ NCH ₂ COOH
N atoms	9	9	10	10	10
Mass	62	62	88	75	75
μ (Deb)	1.8	1.8	2.4	1.2	5.7
Detected in ISM or CSE	 Orion KL			 Sgr B2(N), Orion KL, ..	
Year	 2014			 2003	
Other Astro implications	2016, Sgr B2(N)	Less stable by 162 cm ⁻¹	molecule with 3 O atoms not yet discovered	?? Sgr B2(N), ?? Orion KL;  meteorites  stardust	Less stable by 312 cm ⁻¹
					
	Methyl oxirane	Pyrimidine	Uracil	Cytosine	Phenalenyl ^{+/-}
Formula	CH ₃ CHCH ₂ O	C ₄ H ₄ N	C ₂ CH ₃ O ₂ OH	C ₄ H ₅ N ₃ O	C ₁₃ H ₉
N atoms	10	10	12	13	22
Mass	58	80	88	75	165
μ (Deb)	2.3	2.4	4.4	6.6	0.0
Detected in ISM or CSE	 Sagittarius B				
Year	 2016				
Other Astro implications	First chiral molecule	 N-heterocycles in Titan's atmosphere;  meteorites	 meteorites		 Aromatic ions in Titan's atmosphere

Figure 1: The astroCOMs (or potential astroCOMs) addressed in this review: molecular formula and structure together with the relevant information.

scopic characterization of astroCOMs to be then detected in the interstellar medium through their rotational and vibrational features.^{15,25} We will discuss the successful strategies that allow the determination of bond distances, rotational constants, conformational enthalpies and vibrational frequencies with accuracies better than 0.001 Å, 20 MHz, 1 kJ mol⁻¹ and 10 cm⁻¹, respectively. Particular emphasis will be put on the cases illustrating that theoretical data allow to reassess astronomical observations and provide new insights on possible discoveries of yet-undetected astroCOMs.^{15,19} Possible pathways to overcome some of the remaining challenges related to the molecular complexity and flexibility, which are due to the conformational equilibria and large amplitude motions, will be also discussed. The same theoretical schemes can be also applied to the investigation of the chemical evolution,²⁶ thus allowing for reliable thermochemical and kinetic information on the formation processes of astroCOMs in various environments, including cold interstellar clouds.

Computational spectroscopy of astroCOMs

Almost 200 molecules have been detected in space, with at least 175 molecules observed unambiguously in the interstellar medium or circumstellar shells and about 60 molecules detected in extragalactic sources (see, for example, CDMS^{1,2}). Our knowledge on the universe chemical inventory has been obtained and continuously updated by means of astronomical observations, which have revealed a plethora of molecular species,²⁷ thus demonstrating the rich chemistry of the universe. Molecules can exist in a wide range of astrophysical environments, from the extremely cold regions between stars to the atmospheres of stars themselves. Focusing on their detection, the spectroscopic techniques of interest are rotational and infrared spectroscopies. In fact, gas-phase species have been mostly discovered via their rotational signatures, with the corresponding frequencies going from the millimeter-wave region up to far-infrared. However, spectroscopies in the infrared domain play a crucial role, for example, in the investigation of planetary atmospheres.

As a preliminary remark, we should emphasize the fact that many different theoretical models have been proposed in the literature for the computation of spectroscopic properties, ranging from highly accurate levels of theory²⁸⁻³³ to more practical methods. Regarding

the methodologies applicable to astroCOMs for their spectroscopic characterization, two popular approaches, either based on variational^{34–37} or perturbative^{38–56} schemes should be mentioned, with several programs offering the possibility of performing anharmonic rovibrational calculations, at different levels of sophistication.^{30,34,45,57–65} The approach chosen in the present review is based on vibrational perturbation theory (VPT). To have a better understanding of the challenges to face in accurate calculations on molecular systems of interest in astrochemistry, let us first recall some key aspects of the underlying theory, with special reference to spectroscopic parameters and nuclear motion effects.

Theory

First, we will assume that the Born-Oppenheimer approximation stands, so the electronic and nuclear degrees of freedom can be separated. The rovibrational Hamiltonian $\mathcal{H}_{\text{rovib}}$ is then defined as

$$\mathcal{H}_{\text{rovib}} = \frac{\hbar^2}{2hc} \sum_{\tau,\eta} \mu_{\tau\eta} (J_\tau - \pi_\tau)(J_\eta - \pi_\eta) + \frac{1}{2} \sum_{i=1}^N \omega_i p_i^2 + V + U \quad (1)$$

where τ and η are Cartesian axes of the molecule-fixed frame, $\boldsymbol{\mu}$ is the effective inverse molecular inertia tensor, and \mathbf{J} and $\boldsymbol{\pi}$ are, respectively, the rotational and vibrational angular momentum operators. V is the potential energy operator and U is a mass-dependent contribution

$$U = -\frac{\hbar^2}{8hc} \sum_{\tau} \mu_{\tau\tau}, \quad (2)$$

which vanishes for linear systems. The form of the vibrational angular momentum operator $\boldsymbol{\pi}$ is

$$\boldsymbol{\pi} = \sum_{i,j=1}^N \boldsymbol{\zeta}_{ij} \sqrt{\frac{\omega_i}{\omega_j}} q_i p_j. \quad (3)$$

In the equations above, \mathbf{q} and \mathbf{p} refer to the vectors of the dimensionless vibrational coordinates and their conjugate momenta, respectively. N is the number of vibrational normal modes ($N = 3N_a - 5$ for linear systems, $N = 3N_a - 6$ otherwise, with N_a the number of atoms). Finally, $\boldsymbol{\zeta}$ is the matrix of Coriolis couplings.

Both $\boldsymbol{\mu}$ and V can be expanded as Taylor series of the normal coordinates \mathbf{q} about the

equilibrium geometry:

$$\boldsymbol{\mu} = \boldsymbol{\mu}_{\tau\eta}^{\text{eq}} - \sum_{i=1}^N \boldsymbol{\mu}_{\tau\tau}^{\text{eq}} \mathbf{a}_{i,\tau\eta} \mu_{\eta\eta}^{\text{eq}} q_i + \frac{3}{4} \sum_{\varsigma} \sum_{i,j=1}^N \boldsymbol{\mu}_{\tau\tau}^{\text{eq}} \mathbf{a}_{i,\tau\varsigma} \mu_{\varsigma\varsigma}^{\text{eq}} \mathbf{a}_{j,\varsigma\eta} \mu_{\eta\eta}^{\text{eq}} q_i q_j + \dots \quad (4)$$

$$V = \frac{1}{2} \sum_{i=1}^N \omega_i q_i^2 + \frac{1}{6} \sum_{i,j,k=1}^N k_{ijk} q_i q_j q_k + \frac{1}{24} \sum_{i,j,k,l=1}^N k_{ijkl} q_i q_j q_k q_l + \dots \quad (5)$$

with

$$\begin{aligned} \mathbf{a}_{i,\tau\eta} &= \frac{\partial \mathbf{I}_{\tau\eta}^{\text{eq}}}{\partial q_i} \\ k_{ijk} &= \frac{\partial^3 V}{\partial q_i \partial q_j \partial q_k} \\ k_{ijkl} &= \frac{\partial^4 V}{\partial q_i \partial q_j \partial q_k \partial q_l} \end{aligned}$$

k_{ijk} and k_{ijkl} are commonly referred to as the cubic and quartic force constants, respectively, and ω_i is the wavenumber of mode i .

Once $\boldsymbol{\mu}$ and V in Eq. (1) are replaced by their Taylor expansions, it is possible to recast the rovibrational Hamiltonian as a series of terms \mathcal{H}_{fg} based on the powers f of the vibrational operators (\mathbf{q} and \mathbf{p}) and g of the rotational operators (\mathbf{J}):

$$\begin{aligned} \mathcal{H}_{\text{vibrot}} &= \mathcal{H}_{20} + \mathcal{H}_{30} + \mathcal{H}_{40} + \dots \\ &+ \mathcal{H}_{21} + \mathcal{H}_{31} + \mathcal{H}_{41} + \dots \\ &+ \mathcal{H}_{02} + \mathcal{H}_{12} + \mathcal{H}_{22} + \dots \end{aligned} \quad (6)$$

For the sake of clarity, the subscript ‘‘vibrot’’ has been dropped in the right-hand side of this definition. In the following, we will also assume that the molecule is oriented in such a way that its principal moments of inertia are collinear with the Cartesian axes of the reference frame, the so-called Eckart orientation, in order to minimize the vibrational-rotational couplings. Under this condition, the equilibrium effective inverse molecular inertia tensor $\boldsymbol{\mu}^{\text{eq}}$ can be replaced by the equilibrium molecular rotational constants vector \mathbf{B}^{eq} defined as

$$\mathbf{B}_{\tau}^{\text{eq}} = \frac{\hbar^2}{2hc\mathbf{I}_{\tau\tau}^{\text{eq}}} . \quad (7)$$

In order to derive analytic formulas for the simulation of rotational and vibrational spectra, two major formalisms can be followed, namely the Rayleigh-Schrödinger perturbation

theory and the van Vleck contact transformation (CT).^{42,66} The latter is based on the transformation of the Schrödinger equation, with the Hamiltonian redefined as:

$$\tilde{\mathcal{H}} = e^{iS}\mathcal{H}e^{-iS} , \quad (8)$$

where S is a Hermitian operator so that e^{iS} is unitary.

$\tilde{\mathcal{H}}$ results to be diagonal in a given basis ϕ , with the diagonal elements E being the eigenvalues of \mathcal{H} . What is of interest in this context is that the contact transformation method gives a separation of the vibration-rotation Hamiltonian, which makes it possible to discuss the effective rotational Hamiltonian for an individual vibrational level, or a block of near-degenerate vibrational levels.

Rotational Hamiltonian

The terms $\tilde{\mathcal{H}}_{0g}$, with $g = 2, 4, 6, \dots$, in the effective Hamiltonian are the pure rotational and centrifugal contributions, which describe the rotational energy levels for the vibrational ground state. The complete treatment can be found in the literature⁴²; here, we recall only the key aspects of interest.

As stated above, the vibrational contact transformation leads to the rotational Hamiltonian

$$\tilde{\mathcal{H}}_{rot} = \mathcal{H}_{02} + \tilde{\mathcal{H}}_{04} + \tilde{\mathcal{H}}_{06} , \quad (9)$$

where $\tilde{\mathcal{H}}_{04}$ and $\tilde{\mathcal{H}}_{06}$ are the quartic and sextic centrifugal terms. $\mathcal{H}_{02} = \tilde{\mathcal{H}}_{02}$ is the rigid-rotor Hamiltonian:

$$\mathcal{H}_{02} = \sum_{\tau} B_{\tau}^{\text{eq}} J_{\tau}^2 , \quad (10)$$

where B_{τ}^{eq} has been defined in Eq. (7). From a computational point of view, this is straightforwardly obtained from the geometry optimization. The quartic centrifugal Hamiltonian is defined as

$$\tilde{\mathcal{H}}_{04} = \frac{1}{4} \sum_{\tau\eta\varsigma\varrho} \tau_{\tau\eta\varsigma\varrho} J_{\tau} J_{\eta} J_{\varsigma} J_{\varrho} , \quad (11)$$

where the tensor $\tau_{\tau\eta\varsigma\varrho}$ depends only on the harmonic part of the potential energy surface (PES). To obtain the quartic centrifugal distortion parameters actually employed, further contact transformations with purely rotational operators (thus diagonal in the vibrational

quantum numbers) are then required. The sextic centrifugal distortion constants are derived from the term $\tilde{\mathcal{H}}_{06}$, which involves harmonic, anharmonic, and Coriolis perturbation terms for its calculation. Therefore, from a computational point of view, anharmonic force field computations are needed for their determination. To relate the experimental parameters to combinations of $\tau_{\tau\eta\zeta\rho}$ ($\tau_{\tau\eta\zeta\rho\epsilon\iota}$ in the case of sextics), it is necessary to further, completely reduce the Hamiltonian. Different results are then obtained depending on the reduction chosen.^{42,67,68} To give an example, the rotational Hamiltonian up to the sextics for a rigid asymmetric-top molecule in the Watson A-reduction form^{67,68} is:

$$\begin{aligned}
\tilde{\mathcal{H}}_{rot}^{(A)} = & \sum_{\tau} B_{\tau}^{(A)} J_{\tau}^2 \\
& - \Delta_J (J^2)^2 - \Delta_{JK} J^2 J_z^2 - \Delta_K J_z^4 \\
& - \frac{1}{2} [(\delta_J J^2 + \delta_K J_z^2), (J_+^2 + J_-^2)]_+ \\
& + \Phi_J (J^2)^3 + \Phi_{JK} (J^2)^2 J_z^2 + \Phi_{KJ} J^2 J_z^4 + \Phi_K J_z^6 \\
& + \frac{1}{2} [(\phi_J (J^2)^2 + \phi_{JK} J^2 J_z^2 + \phi_K J_z^4), (J_+^2 + J_-^2)]
\end{aligned} \tag{12}$$

where J^2 and J_{\pm} are the total angular momentum and the ladder operators, respectively, with $[X, Y]_+$ representing the anticommutator. The Δ 's and δ 's are the quartic centrifugal distortion constants, while the Φ 's and ϕ 's are the sextic ones.⁶⁸

The Hamiltonian given for example in Eq. (12) applies to the rigid-rotor approximation case. However, another important issue is the effect of molecular vibrations. The operators $\tilde{\mathcal{H}}_{22}$, $\tilde{\mathcal{H}}_{42}$, $\tilde{\mathcal{H}}_{24}$, ... in Eq. (6) contain the terms describing the dependence of the rotational and centrifugal constants on the vibrational quantum numbers. The quantities multiplying J_{τ}^2 in Eq. (6) are actually the effective rotational constants and contain the contributions beyond the rigid-rotor harmonic-oscillator (RRHO) approximation. Their effects on rotational motion can be conveniently described by means of vibrational perturbation theory, and we refer the reader to, for example, Refs.^{42,67} for a detailed treatment. While there are no corrections at the first order in VPT, at the second order (VPT2) three contributions appear that, expressed by means of the usual contact-transformation method,⁴² give:⁴⁰

$$B_{\tau}^v = B_{\tau}^{\text{eq}} - \sum_{i=1}^N \alpha_{i,\tau} \left(v_i + \frac{d_i}{2} \right), \tag{13}$$

where the superscript v denotes a specific vibrational state and the sum is taken over all fundamental vibrational modes i , with v_i its quantum number and d_i its degeneracy order. The $\alpha_{i,\tau}$'s are the so-called vibration-rotation interaction constants and contain three contributions: the first one is a corrective term related to the moment of inertia, the second one is due to the Coriolis interactions, and the last is an anharmonic correction. Therefore, from a computational point of view, anharmonic force field calculations are required to correct the equilibrium rotational constants for vibrational effects.

Vibrational Hamiltonian

For the simulation of vibrational spectra, a purely vibrational hamiltonian (\mathcal{H}_{f0}) is commonly used, which also includes the zeroth-order contribution from the effective inverse molecular inertia tensor:

$$\frac{\hbar^2}{2hc} \sum_{\tau,\eta} \mu_{\tau\eta}^{\text{eq}} \pi_\tau \pi_\eta = \sum_{\tau} B_{\tau}^{\text{eq}} \sum_{i,j,k,l=1}^N \zeta_{ij,\tau} \zeta_{kl,\tau} \sqrt{\frac{\omega_i \omega_k}{\omega_j \omega_l}} q_i p_j q_k p_l \quad (14)$$

As the name implies, the second-order vibrational perturbation theory is based on Taylor expansions of the harmonic potential (V) and vibrational (ε) energies, as well as the vibrational wavefunction, up to the second order.³⁸ Considering the terms that are only functions of the normal coordinates or their conjugate momenta in Eq. (6), the vibrational Hamiltonian up to the fourth power of the operators of \mathbf{q} , \mathbf{p} is then defined as

$$\mathcal{H}_{\text{vib}} = \mathcal{H}_{20} + \mathcal{H}_{30} + \mathcal{H}_{40} \quad (15)$$

$$\begin{aligned} &= \frac{1}{2} \sum_{i=1}^N \omega_i (p_i^2 + q_i^2) + \frac{1}{6} \sum_{i,j,k=1}^N k_{ijk} q_i q_j q_k + \frac{1}{24} \sum_{i,j,k,l=1}^N k_{ijkl} q_i q_j q_k q_l \\ &+ \sum_{\tau} B_{\tau}^{\text{eq}} \sum_{i,j,k,l=1}^N \zeta_{ij,\tau} \zeta_{kl,\tau} \sqrt{\frac{\omega_i \omega_k}{\omega_j \omega_l}} q_i p_j q_k p_l + U \end{aligned} \quad (16)$$

Proper support of the normal modes degeneracy in VPT2 requires some specific treatments, thus leading to more complex formulas than for non-degenerate modes (see Ref.⁵⁴ for a recent overview on the extension to symmetric and linear tops). For this reason, and to not clutter the discussions, the normal mode degeneracy will be ignored in the following.

At the VPT2 level, a single formula is needed to compute the energy (ε_m , in cm^{-1}) of any vibrational state $|m\rangle$,

$$\varepsilon_m = \varepsilon_0 + \sum_{i=1}^N v_i^m \omega_i + \sum_{i,j=1}^N \chi_{ij} \left[v_i^m v_j^m + \frac{1}{2}(v_i^m + v_j^m) \right] \quad (17)$$

where v_i^m is the number of quanta associated to mode i in state $|m\rangle$ and ω_i its harmonic wavenumber. ε_0 is the zero-point vibrational energy, defined as

$$\begin{aligned} \varepsilon_0 = & \sum_{i=1}^N \frac{\omega_i}{2} + \sum_{i,j=1}^N \frac{k_{iijj}}{32} - \sum_{i,j,k=1}^N \left[\frac{k_{iik}k_{jjk}}{32\omega_k} + \frac{k_{ijk}^2}{48(\omega_i + \omega_j + \omega_k)} \right] \\ & - \sum_{\tau} \frac{B_{\tau}^{\text{eq}}}{4} \left[1 - \sum_{i=1}^{N-1} \sum_{j=i+1}^N \{ \zeta_{ij,\tau} \}^2 \frac{(\omega_i - \omega_j)^2}{\omega_i \omega_j} \right] \end{aligned} \quad (18)$$

and χ is the matrix containing the anharmonic contributions, with elements given by

$$16\chi_{ii} = k_{iiii} - \frac{5k_{iii}^2}{3\omega_i} - \sum_{\substack{j=1 \\ j \neq i}}^N \frac{(8\omega_i^2 - 3\omega_j^2)k_{iij}^2}{\omega_j(4\omega_i^2 - \omega_j^2)} \quad (19)$$

$$\begin{aligned} 4\chi_{ij} = & k_{iijj} - \frac{2\omega_i k_{iij}^2}{(4\omega_i^2 - \omega_j^2)} - \frac{2\omega_j k_{ijj}^2}{(4\omega_j^2 - \omega_i^2)} - \frac{k_{iii}k_{ijj}}{\omega_i} - \frac{k_{jjj}k_{iij}}{\omega_j} \\ & + \sum_{\substack{k=1 \\ k \neq i,j}}^N \left[\frac{2\omega_k(\omega_i^2 + \omega_j^2 - \omega_k^2)k_{ijk}^2}{\Delta_{ijk}} - \frac{k_{iik}k_{jjk}}{\omega_k} \right] + \frac{4(\omega_i^2 + \omega_j^2)}{\omega_i \omega_j} \sum_{\tau} B_{\tau}^{\text{eq}} \{ \zeta_{ij,\tau} \}^2 \end{aligned} \quad (20)$$

with

$$\Delta_{ijk} = \omega_i^4 + \omega_j^4 + \omega_k^4 - 2(\omega_i^2 \omega_j^2 + \omega_i^2 \omega_k^2 + \omega_j^2 \omega_k^2)$$

Transition energies from the ground state are straightforwardly obtained from Eq. (17) as “ $\nu_m = \varepsilon_m - \varepsilon_0$ ”.

Conversely, defining the anharmonic transition moments for the properties of interest (here the electric dipole) is more cumbersome for two major reasons. First, the property must also be expanded as a Taylor series, beyond the first-order derivative. This is referred to as the electrical anharmonicity, by contrast with the mechanical anharmonicity from the definition of the wavefunction. Next, at variance with the energy, it is not possible to derive a single, compact analytic formula, like Eq. (17), applicable to any transition. Multiple formulas are needed, depending on the number of quanta and excited modes separating the initial (generally the ground state) and final states (fundamental, first overtones, 2-quanta

combinations and so on). Explicit equations for the transitions, including the properties function of the momenta conjugate of the normal coordinates, can be found in Refs. ^{46,52,55,69,70}. Instead of building compact formulas, an alternative, more general strategy is to build the integrals at runtime, during the calculations themselves, using either Rayleigh-Schrödinger⁷¹ or van Vleck⁴⁸ formalisms. However, the computational cost and memory requirements to store the data necessary for the integrals are significantly higher than those required by the analytic forms. A presentation of their characteristic features is beyond the scope of the present contribution and will not be done here.

A quick inspection of Eqs. (19) and (20) is sufficient to identify simple conditions for which the VPT2 energies will be excessively high or even infinite, that is, when a denominator becomes exceedingly small. Those are known as Fermi resonances (FRs), divided in type I ($\omega_i \approx 2\omega_j$) and type II ($\omega_i \approx \omega_j + \omega_k$). Besides the limit case of an actual equality, which means that at least one element of the matrix χ is infinite, a near resonance can be sufficient to obtain unphysical results due to an excessive contribution from anharmonicity. This is a well-known issue of VPT2, which has been extensively studied in the literature^{42,47,51,72–83} and needs to be correctly addressed for a successful application of this method, *a fortiori* in the study of astroCOMs. A major difficulty lies in the definition of the resonance conditions. A very simple strategy to adopt is to define manually the resonances, based on the final vibrational energies. However, it does not scale well with the system size and is not sufficiently robust for a systematic use. A more efficient way is to automatize the identification process, which requires formalizing the concept of Fermi resonances. This can be achieved through different schemes.^{47,51,54,72,74–76,81–83} A preliminary remark, which is also a reason for the variety of approaches proposed in the literature, is that most of them depend ultimately on thresholds defined empirically, thus introducing a degree of arbitrariness. However, this can be balanced to some extent by combining multiple criteria. A simple definition, and the starting point of many selection schemes, is the magnitude of the energy difference between the states (fundamental and 2-quanta overtones or combinations) in the denominator. Such a basic test, however, neglects the magnitude of the overall term, which also depends on the cubic force field in the numerator. As a result, determining a single threshold applicable to every case is unpractical. A more robust solution is to complement this initial evaluation

with a control of the magnitude of the potentially resonant term and its contribution to the anharmonic correction. We will mention here only the test proposed by Martin and coworkers, which is based on the deviation of this term from the variational energies of a model variational treatment of a system including only the two states involved in the resonance (for instance the states with harmonic energies “ ω_i ” and “ $\omega_j + \omega_k$ ” for a type II resonance).⁷⁴ More details can be found in the original paper. Terms identified as resonant are removed from the perturbative treatment for the calculation of the energy. This approach is named deperturbed VPT2 (DVPT2). However, this also means that the VPT2 treatment is truncated since terms are missing. This can be corrected through an *ad hoc* variational step, which reintroduces the previously discarded terms, using the DVPT2 vibrational energies as references. Different acronyms have been used to designate the overall procedure. Here, it is referred to as the generalized VPT2 (GVPT2). While GVPT2 is able to provide accurate results, it still relies directly on one or more empirical criteria for the identification of Fermi resonances. To overcome this limitation, a two-step procedure has been introduced to select resonances, the absolute frequency difference must be lower than a threshold Δ_ω^{1-2} and the deviation from the variational model must be higher than a chosen value K^{1-2} . In some cases, small variations in the force constants (harmonic and anharmonic), like during a benchmark of electronic structure calculation methods or over a series of geometries along a reaction path, can be sufficient to modify the set of resonances, with a potentially notable impact on the final vibrational energies. An alternative way, proposed by Kuhler, Truhlar and Isaacson, is to replace every potentially resonant term with a non-resonant form derived from a model system comprising only the two states involved in each case.⁷⁵ A strong advantage of this method, denoted degeneracy-corrected PT2 (DCPT2), is that it does not use any threshold. However, the transformation is based on a number of approximations that introduce some inaccuracy for each replaced terms, which can be a problem far from resonance, where VPT2 behaves correctly. This shortcoming can be partially corrected by introducing a transition function which will mix the DCPT2 and VPT2 results for each potentially resonant terms, thus favoring VPT2 far from resonance. More details on this approach, called hybrid DCPT2-VPT2 (HDCPT2), can be found in Ref.⁵¹.

Finally, we should mention another type of resonances, collectively called Darling-Dennison

resonances (DDRs).^{48,49,54,72,76,78,83–86} This term describes couplings of states through the second-order vibrational Hamiltonian, missing in standard VPT2 calculations. They originally refer to interactions between overtones, in particular for the CH and OH stretching modes, following the original study of Darling and Dennison on the water molecule.⁸⁴ The term has now a broader meaning and is also used to designate the so-called second-order interactions between fundamentals (“1-1”), 2-quanta states (“2-2”, “2-11”, “11-11”, where “2” represents an overtone and “11” a combination), and fundamentals with 3-quanta states (“1-3”, “1-21”, “1-111”), the last case being needed if 3-quanta transitions are also included in the anharmonic calculations.^{49,76} These resonances are commonly treated through a variational procedure, analogous to that used for FRs. In order to limit this step to relevant couplings, a two-step analysis similar to that introduced for FRs can be used, based on the absolute energy difference between the states (below a given threshold Δ_{ω}^{A-B}) and the magnitude of the off-diagonal term to be used in the variational step (higher than a chosen parameter, K^{A-B}). An hurdle specific to DDRs is the potential presence of Fermi resonances in the variational terms themselves. Depending on the method used for the identification of FRs in energy calculations, it may be possible or impractical to extend it straightforwardly to the case of DDRs. Further discussions on this problem can be found in Refs.^{76,83,87} In the following, GVPT2 will refer to the full variational treatment, including corrections to Fermi and Darling-Dennison resonances.

At variance, the problem of resonances in intensity calculations has been more scarcely addressed in the literature, to some extent due to the limited number of programs supporting them.^{48,56,70,73,83} Since they are related to the mechanical anharmonicity (wavefunction), it is possible to use the analysis made for the energy on the transition moments as well. An important difference, however, is the impact of Darling-Dennison resonances, which can lead to incorrect intensities if not identified correctly. It is especially important for modes or states (for 3-quanta transitions) with near-equal harmonic energies, like for instance, the CH symmetric and antisymmetric stretching modes of a methyl group. Depending on the protocol followed for the definition of DDRs, it may be necessary to complement it with an *ad hoc* test targeted to handle the most critical cases (for instance of near-equal energies).^{56,87} States coupled through a Darling-Dennison interaction need to be removed from the VPT2

formulas of the transition moments, thus resulting in fully deperturbed results where both FRs and DDRs are removed. These states are then projected on the GVPT2 states using the eigenvectors obtained from the energy calculation.⁷⁰

From a computational perspective, for most of the examples chosen in this review, the suites of quantum chemical programs CFOUR⁵⁷ (high-level theoretical models, rotational spectroscopy) and GAUSSIAN⁶¹ (vibrational spectroscopy, thermodynamics, anharmonic contributions) have been used. Regarding VPT2 methods, vibrational energies and IR intensities have been calculated with GVPT2 and the thermodynamic properties with HDCPT2. Moreover, simulations of fully anharmonic vibrational spectra have required the preceding computation of a semi-diagonal quartic PES and a semi-diagonal cubic property surface (PS), which could be effectively derived through numerical differentiation along the normal coordinates of analytic harmonic force constants (for the PES) and first derivatives of the properties (for the PS). To be more specific, the property of interest here, in the context of astroCOMs and the simulation of infrared spectra, is the electric dipole moment.

Computational methodology

The basic requirements for the theoretical support in the field of astrochemistry are highly accurate predictions of the equilibrium molecular structures of astroCOMs, their rotational and vibrational spectroscopic parameters, and the abundance of different possible isomers or conformers for specific space-like conditions. To reach the required accuracy, the best possible quantum-mechanical (QM) methodologies must be employed in the description of the electronic structure, and then combined with solutions of the nuclear motion problem beyond the rigid-rotor harmonic-oscillator approximation. If, on the one hand, electronic structure calculations should be pushed to the limit in order to determine very accurate equilibrium structures, on the other hand, it is also true that the portion of the PES around the equilibrium, i.e., the force constants of Eq. (5) up to the fourth order, need to be described at the highest possible level for a proper inclusion of the anharmonic effects in spectroscopic and thermodynamic properties.

The key point to obtain accurate results in predicting molecular structures and harmonic force fields is to reduce as much as possible the errors due to the truncation of both basis set

and wavefunction, the so-called one- and N-electron errors, respectively. To this purpose, composite schemes that evaluate the contributions important to reach high accuracy at the best possible level and combine them through the additivity approximation, have been set up.^{88–93}

The so-called “gradient” scheme,^{91,92} implemented in the quantum-chemical program package CFour,⁵⁷ mainly involves calculations at the coupled cluster (CC) singles and doubles with perturbative triples corrections, CCSD(T), level.⁹⁴ However, the full CC singles, doubles and triples (CCSDT)^{95–97} and the CC singles, doubles, triples, quadruples (CCSDTQ)⁹⁸ models can also be employed. The correlation-consistent polarized cc-p(C)VnZ basis sets^{99–102} are usually used in conjunction with this scheme. Within the “gradient” scheme, the energy gradient to be minimized in the geometry optimization procedure is set up according to the specific needs, the target accuracy, and the dimension of the system under consideration. To recover the error due to the basis-set truncation, an extrapolation to the complete basis set (CBS) limit can be performed. Since extrapolation to the CBS limit is carried out within the frozen core approximation, core-valence correlation (CV) effects are accounted for by adding the energy gradient corresponding to the difference of all-electron and frozen-core CCSD(T) calculations in the same basis set. In a similar manner, corrections from a full treatment of triples (fT) and quadruples (fQ) can be included using small basis sets for their evaluation (usually of double-, triple- ζ quality). The overall energy gradient is given by the following expression:

$$\begin{aligned} \frac{dE_{\text{CBS+core+fT+fQ}}}{dx} = & \frac{dE^\infty(\text{HF-SCF})}{dx} + \frac{d\Delta E^\infty(\text{CCSD(T)})}{dx} \\ & + \frac{d\Delta E(\text{CV})}{dx} + \frac{d\Delta E(\text{full-T})}{dx} + \frac{d\Delta E(\text{full-Q})}{dx}. \end{aligned} \quad (21)$$

From literature studies based on comparisons with experiment (see, for example, Refs.^{23,24,91,92,103–105} and references therein), the accuracy obtainable with the scheme presented above in terms of structural parameters is better than 0.001 Å for bonds and 0.1 degrees for angles. It is particularly noteworthy that the loss of accuracy when the contributions from the full treatment of triple and quadruple excitations in the cluster expansion are ignored is very limited. The importance of this observation lies in the fact that, while the CCSD(T)/CBS+CV+fT+fQ level is computationally very expensive, the CCSD(T)/CBS+CV one is affordable also for

Table 1: Statistical analysis of the relative errors (in %) in the computed rotational constants with respect to experimentally determined equilibrium \mathbf{B}^{eq} and vibrational ground-state \mathbf{B}^0 values^a.

computational approach	$\bar{\Delta}^b$	$\bar{\Delta}_{\text{abs}}^c$	Δ_{std}^d
With respect to \mathbf{B}^{eq}			
fc-CCSD(T)/cc-pV6Z	-0.277	0.296	0.158
fc-CCSD(T)/CBS	-0.253	0.274	0.141
fc-CCSD(T)/cc-pV6Z + CV	0.041	0.057	0.100
fc-CCSD(T)/cc-pV6Z + CV + ΔT	0.071	0.087	0.110
fc-CCSD(T)/cc-pV6Z + CV + ΔT + ΔQ	-0.027	0.063	0.072
fc-CCSD(T)/CBS + CV + ΔT + ΔQ	-0.003	0.041	0.071
With respect to \mathbf{B}^0			
fc-CCSD(T)/CBS	0.380	0.499	0.643
fc-CCSD(T)/CBS + CV	0.690	0.762	0.590
fc-CCSD(T)/CBS + CV + ΔT + ΔQ	0.632	0.710	0.594
fc-CCSD(T)/CBS + CV + ΔT + ΔQ + ΔB_{vib}^e	0.007	0.057	0.124

^a Data from Ref. ⁹³.

^b Mean error.

^c Mean absolute error.

^d Standard deviation.

^e Vibrational corrections evaluated using Eq. (13) at the CCSD(T)/cc-pCVQZ level of theory with all electrons correlated.

medium-sized molecules, like glycine²² and pyruvic acid.²³ In terms of geometrical parameters, it is well recognized to perform within 0.001–0.002 Å for distances and 0.05–0.1 degrees for angles (see, for example, Refs. ^{22,23,91,92,106}). In Ref. ⁹³, the “gradient” scheme was tested in terms of the accuracy reachable in predicting rotational constants, with the relevant results being summarized in Table 1. The overall conclusion is that the so-called CCSD(T)/CBS+CV level of theory leads to equilibrium rotational constants that agree well within 0.1% with experiment. However, Table 1 also points out that for keeping the same accuracy when the comparison is performed with the experimental ground state rotational constants (\mathbf{B}^0), it is mandatory to include the vibrational corrections (see Eq. (13)).

To extend the applicability of composite approaches to larger molecules, an effective

solution is provided by the so-called “geometry” scheme, which is based on the assumption that the additivity approximation can be directly applied to geometrical parameters and that they show the same behavior as the energy (see, for example, Refs.^{107–110}). Analogously to the “gradient” scheme, the various contributions are evaluated separately at the highest possible level and then combined in order to obtain best theoretical estimates. Unlike the “gradient” scheme, which requires some specific software implementations to be used, the “geometry” scheme is directly applicable and is definitely less computationally expensive. To further extend the size of systems amenable to accurate structural determinations, it is necessary to further reduce the computational cost. In recent years, a very effective approach, which mainly involves geometry optimizations using Møller-Plesset second-order theory (MP2)¹¹¹ and basis sets of triple- and quadruple- ζ quality, has been set up.^{22,25,110,112} The starting point is the geometry optimization at the CCSD(T)/cc-pVTZ level within the frozen-core approximation, while all contributions required to improve its accuracy are computed at the MP2 level. As discussed in Ref.¹⁰⁹, to take into account basis-set truncation effects, the CBS limit is evaluated by making use of the consolidated n^{-3} extrapolation form¹¹³ applied to MP2/cc-pVTZ and MP2/cc-pVQZ optimized geometries. The effect due to core-valence electron correlation is included by means of the corresponding correction, $\Delta r(\text{CV})$, derived as the difference between the geometrical parameters optimized at the MP2/cc-pCVTZ level correlating all and only valence electrons. On the whole, within the “geometry” scheme, the best-estimated equilibrium structure is determined as:

$$r(\text{best}) = r(\text{CCSD(T)/VTZ}) + \Delta r(\text{CBS}) + \Delta r(\text{CV}) + \Delta r(\text{aug}) . \quad (22)$$

This approach is denoted “cheap” geometry scheme²⁴ to stress its reduced computational cost. While there is no theoretical justification for the inclusion of diffuse function effects ($\Delta r(\text{aug})$) once the extrapolation to the CBS limit is performed, the latter correction is often introduced to recover on an empirical basis the limitations affecting extrapolation procedures carried out with small- to medium-sized basis sets. Based on the comparison with best-estimated equilibrium structures obtained using the “gradient” scheme as well as with experiment,^{19,22–24,110} the overall conclusion is that the so-called “cheap” geometry scheme is robust, reliable and accurate, even when dealing with rather flexible molecules like

glycine²² and pyruvic acid.²³ For bond distances, for instance, it is noted that the maximum absolute deviation with respect to CCSD(T)/CBS+CV is smaller than 0.001 Å.

According to Eq. (13) and to the results summarized in Table 1, the evaluation of the rotational constants to be used in the actual prediction of the rotational spectrum involves the determination of the vibration-rotation interaction constants, with the latter requiring in turn the computation of the cubic force field. On the basis of the literature on this topic (see, for example, Refs.^{25,93,106,110,112} and references therein), the level of theory required ranges from density functional theory (DFT) in conjunction with double- ζ basis set to CCSD(T) in combination with quadruple- ζ set.

Moving to harmonic force fields, a composite approach similar to the “cheap” geometry scheme for equilibrium structures is employed to obtain best estimates:

$$\omega(\text{best}) = \omega(\text{CCSD(T)/VTZ}) + \Delta\omega(\text{CBS}) + \Delta\omega(\text{CV}) + \Delta\omega(\text{aug}) , \quad (23)$$

where $\omega(\text{CCSD(T)/VTZ})$ denotes the harmonic frequencies at the CCSD(T)/cc-pVTZ level in a normal coordinates representation. The CBS, CV and aug corrections are defined in the same way as for geometries, with the additional necessary condition being related to the normal modes correspondence for all employed levels of theory. On the basis of the available literature,^{22-24,114} this scheme is able to provide harmonic frequencies with an accuracy of 1–2 cm^{-1} with respect to CCSD(T)/CBS+CV,²⁴ with improvements of 5–10 cm^{-1} with respect to CCSD(T)/cc-pVTZ.^{22-24,114} An analogous composite scheme has also been derived to determine best estimates for the IR intensities,^{22,24,114} $I(\text{best})$, within the double-harmonic approximation. Since extrapolation schemes have not been formulated yet for such a property, Eq. (23) has been reformulated as follows:

$$I(\text{best}) = I(\text{CCSD(T)/VTZ}) + \Delta I(\text{CV}) + \Delta I(\text{QZ-TZ}) + \Delta I(\text{aug}) , \quad (24)$$

where $\Delta I(\text{QZ-TZ})$ is the correction due to the “MP2/cc-pVQZ - MP2/cc-pVTZ” difference, and the other contributions are defined similarly to frequencies.

As a byproduct of the harmonic force fields evaluated at different levels of theory in the frame of the so-called “cheap” scheme, best estimates for quartic centrifugal-distortion constants can be derived using the same composite scheme defined for IR intensities in Eq. (24).

While there is remarkable literature concerning the accuracy and the convergence behavior of computed rotational constants, little has been done by contrast on the centrifugal-distortion constants. In Ref. ¹¹⁵, the quartic and sextic centrifugal-distortion constants were evaluated at the HF-SCF, CCSD and CCSD(T) levels in conjunction with a hierarchical series of basis sets, ranging from triple- to quintuple- ζ quality. With respect to the CCSD(T)/aug-cc-pCVnZ calculations, it is noted that both quartic and sextic centrifugal-distortion constants show a monotonic trend while enlarging the set, thus allowing conclusions to be drawn regarding the convergence to the basis-set limit and the error due to the basis-set truncation. On these grounds, the CCSD(T)/aug-cc-pCV5Z values have a residual basis-set error no larger than 0.2%. By comparison of the constants evaluated at the HF-SCF, CCSD and CCSD(T) levels of theory, reliable centrifugal-distortion constants can already be obtained at the HF-SCF level, with discrepancies on average of 10–15% with respect to CCSD. Another observation is that the effects of triples are small, the differences between CCSD(T) and CCSD being on average of 1–2%. Therefore, the effect of higher excitations are expected to be almost negligible. Since the CCSD(T) method provides results converged with respect to the truncation of the wavefunction and the composite scheme mentioned above is sufficient to recover the error due to the basis-set truncation, the application of the latter scheme leads to a quantitative estimate of the quartic centrifugal-distortion constants. Due to the very small magnitude of the sextic centrifugal-distortion terms, any level of theory is sufficient to get good predictions.

The evaluation of anharmonic corrections requires an extensive sampling of the potential energy and property surfaces, thus leading to strong limitations in the level of theory affordable. To achieve high accuracy, the so-called hybrid QM/QM' approaches need to be employed. These models combine an evaluation of the equilibrium structure, as well as of the harmonic PES and PS, at the highest possible QM level (including composite schemes) with anharmonic corrections computed by means of less expensive QM' techniques. In particular, the combination of CCSD(T) with DFT functionals, thus resulting in a plethora of hybrid CC/DFT approaches, has been well-tested for small- to medium-sized closed- and open-shell molecular systems, also including several potential astroCOMs. ^{15,23,24,53,116–118}

The definition of hybrid models requires to evaluate, as a first step, the performance of

computationally less-demanding QMs and then to select those suitable for obtaining anharmonic corrections at the desired accuracy. In this respect, it is crucial to assure that these methodologies predict reliable equilibrium structures, harmonic vibrational wavenumbers (ω), normal modes (\mathbf{q}), and IR intensities (I_{IR}). An appealing factor of the methodologies rooted in the density functional theory¹¹⁹ is that they are able to provide results with a reasonably good accuracy, in particular in comparison to the computationally more demanding MP2¹¹¹ method. Among the many methods available, hybrid or double hybrid functionals, like B3LYP¹²⁰ and B2PLYP,¹²¹ combined with basis sets of double- and triple- ζ quality, also containing polarization and diffuse functions, are demonstrated to be particularly effective. Nevertheless, it is of crucial importance to verify the accuracy of the DFT equilibrium structures first. In addition to the CCSD(T)/CBS+CV geometries, whose performance has been discussed above, an effective and reliable testing ground is offered by the so-called semi-experimental (SE) equilibrium structures ($r_{\text{SE}}^{\text{eq}}$).¹²² The latter are based on a combination of experimental ground state rotational constants and computed vibrational corrections for different isotopic species and permit the determination of structural parameters accurate within 0.001 Å and 0.1 degrees for bond lengths and angles, respectively (see, for example, Refs.^{106,123,124}). As an illustration of the accuracy reachable by DFT approaches, Table 2 reports the mean (MAE) and maximum ($|\text{MAX}|$) absolute errors for the bond lengths of a set of small- and medium-sized, semi-rigid and flexible astroCOMs, also including the uracil nucleobase and the glycine aminoacid. With regard to the accessibility of accurate molecular structures, it is worthwhile noting that a database collecting accurate $r_{\text{SE}}^{\text{eq}}$ geometries for a large number of small- and medium-sized species has been recently set up.^{125,126}

Table 2 confirms the high accuracy of the structural parameters obtained from the CCSD(T)/CBS+CV composite approach and the “cheap” geometry scheme, with average and maximum discrepancies of about 0.001 and 0.003–0.004 Å, respectively. In particular, the latter model represents a significant improvement over CCSD(T)/cc-pVTZ, with MAE and $|\text{MAX}|$ smaller by about 70%, at a limited additional cost due to MP2 computations. The high accuracy of the so-defined “cheap” equilibrium structures is particularly promising for the astrochemical detection of larger astroCOMs, also including small biomolecules. Concerning less expensive QM models, which can be employed for the generation of the

Table 2: Accuracy of equilibrium bond lengths computed by different QM methods^a.

astroCOM	B3LYP ^b	B2PLYP ^c	MP2 ^c	CCSD(T) ^d	“cheap” ^e	bestCC ^f
			MAE wrt r_{SE}^{eq}			
oxirane	0.0059	0.0024	0.0035	0.0048	0.0014	0.0008
methyloxirane	0.0068	0.0020	0.0026	0.0035	0.0011	0.0009
acrolein	0.0053	0.0013	0.0022	0.0040	–	0.0006
pyruvic acid	0.0074	0.0038	0.0048	0.0053	0.0015	0.0011
pyrimidine	0.0054	0.0008	0.0024	0.0043	0.0006	0.0006
glycine	0.0075	0.0033	0.0035	0.0045	0.0009	0.0010
uracile	0.0063	0.0026	0.0041	0.0049	0.0012	–
ALL	0.0064	0.0023	0.0033	0.0045	0.0011	0.0008
			MAX wrt r_{SE}^{eq}			
oxirane	0.007	0.006	0.009	0.007	0.003	0.001
methyloxirane	0.008	0.007	0.009	0.005	0.003	0.002
acrolein	0.009	0.004	0.009	0.007	–	0.002
pyruvic acid	0.013	0.007	0.010	0.007	0.003	0.003
pyrimidine	0.007	0.001	0.004	0.007	0.001	0.001
glycine	0.013	0.008	0.008	0.010	0.003	0.004
uracile	0.013	0.008	0.008	0.008	0.003	–
ALL	0.013	0.008	0.010	0.010	0.003	0.004

^a Mean (MAE) and maximum (|MAX|) absolute deviations (in Å) with respect to the semi-experimental equilibrium structure for oxirane,^{19,24} methyloxirane,²⁴ *trans*- and *cis*-acrolein,^{118,125} *Tc*-pyruvic acid,²³ pyrimidine,^{125,127} Ip/ttt and IIn/cc glycine conformers^{22,105,128} and uracile,^{110,117,129} complemented by data from this work.

^b Computed with the SNSD⁵³ basis set.

^c Computed with the aug-cc-pVTZ basis set.

^d Computed with the cc-pVTZ basis set.

^e Obtained by means of the “cheap” geometry scheme, see text.

^f CCSD(T)/CBS+CV equilibrium structure (see text) obtained using the cc-pVTZ and cc-pVQZ basis sets for the CBS extrapolation and the cc-pCVTZ set for the core-correlation correction.

anharmonic PES and PS for molecular systems composed of dozens of atoms, it is evident that the double-hybrid B2PLYP structures are on average slightly better than their MP2 counterparts, at a comparable computational cost.

As a matter of fact, benchmark studies suggest that for larger systems, whenever the evaluation of the harmonic PES by means of composite schemes becomes prohibitively expensive, the B2PLYP functional in conjunction with basis sets of at least triple- ζ quality provides an effective route for both rotational and vibrational spectroscopy investigations.^{23,24,53,118,127,130} In particular, for several astroCOMs, it has been shown that vibrational wavenumbers in agreement with experiment within 6–12 cm^{-1} can be obtained from anharmonic GVPT2 calculations based on B2PLYP/aug-cc-pVTZ harmonic PESs,^{24,118,129–131} to be compared with an accuracy of 4–10 cm^{-1} obtained when using harmonic PESs of CCSD(T)/CBS+CV quality.^{24,117,118,130,131} Concerning IR intensities, the B2PLYP/aug-cc-pVTZ results agree with the best-estimated values within 1–2 km mol^{-1} , thus showing the same accuracy as MP2 computations.²⁴

The overall good accuracy demonstrated by B2PLYP in the description of the harmonic PESs and PSs fully justifies its use in the evaluation of anharmonic contributions. For fundamental bands, the anharmonic corrections on the wavenumbers agree within 2–7 cm^{-1} with those obtained at the CCSD(T) level, and within 0.5 km mol^{-1} for the IR intensities.^{24,130} Therefore, hybrid CC/DFT models with anharmonic corrections at the B2PLYP/aug-cc-pVTZ level give highly accurate results for both vibrational wavenumbers and IR intensities. In particular, this leads for the latter to an agreement within 1 km mol^{-1} with experimental integrated cross sections, which means an error smaller than 5% and 10% for the strongest and weakest signals, respectively.^{114,131} Concerning line positions, it should be noted that the only large discrepancies noted, of order of 20 cm^{-1} , involve intra-molecular hydrogen bonds, as for instance the $\nu(\text{O-H})_{\text{HB}}$ band of the *Tc*-pyruvic acid.²³ However, even in these cases, the discrepancies of B2PLYP anharmonic wavenumbers with respect to experiment can be entirely related to the harmonic part, at least for the fundamental transitions.²³

Finally, we can conclude that the least expensive B3LYP model clearly does not provide the accuracy required for the prediction of equilibrium structures and properties in the field of astrochemistry, with MAEs an order of magnitude larger than those obtainable with

the best theoretical approaches. This has also an impact on the harmonic frequencies that therefore show too large errors.^{53,130} However, it should be noted that, among the various DFT models, the B3LYP functional still represents a reliable and accurate approach for the evaluation of anharmonic contributions to be coupled to higher-level methods, provided that the normal modes are sufficiently similar. Extensive benchmark studies have shown that B3LYP/SNSD computations can be applied to evaluate accurate vibrational corrections to the rotational constants^{25,125} and anharmonic corrections to vibrational wavenumbers and IR intensities.^{22,24,53,131} By contrast, many more recent functionals show significantly larger discrepancies with respect to experiment for both harmonic frequencies and anharmonic corrections.^{53,130} Thus, the use of newly-developed DFT models in the field of astrochemistry needs an extensive validation first.⁸⁷

While B3LYP molecular structures are not sufficiently accurate to be used in spectroscopic application, it has been shown that combining B3LYP/SNSD structures and thermodynamic properties with CCSD(T)/CBS+CV electronic energies is a reliable approach to assess the population of conformers.²² In more details, it is apparent that relative electronic energies computed at the best theoretical and B3LYP/SNSD equilibrium structures usually differ by less than 0.1 kJ mol⁻¹. Thus, the B3LYP/SNSD model represents an effective route for extensive conformational searches that, once followed by accurate estimates of the electronic energies, allows identifying the most stable conformers of astroCOMs on which highly accurate theoretical investigations should be focused.

Spectroscopy of astroCOMs: Oxirane as a test case

Starting from its detection in the Interstellar Medium¹³² as one of the first cyclic molecules, the potential role in astrochemistry of oxirane has been addressed in several publications (see, for example, Refs.^{15,19,133–135}), with special reference to Titan’s (the largest moon of Saturn) atmosphere. In Ref.¹³³, Raulin and coworkers demonstrated that experiments designed to simulate the upper atmosphere of Titan, i.e., a high voltage applied to a mixture containing methane (CH₄, ~2%), nitrogen (98% N₂), and traces of oxygen (0.01% CO) at low temperature (100–150 K), can lead to the production of oxirane. Furthermore, Bernstein and Lynch

demonstrated a close correlation between the IR spectra of oxirane, among others, and the major unidentified infrared bands (UIRs).¹³⁶ The relevance of the oxiranes’ family is also pointed out by the recent, first detection of a chiral molecule, methyloxirane, in a cold, extended molecular shell around the embedded, massive protostellar clusters in the Sagittarius B2 star-forming region.¹⁸ In view of its broad astrochemical relevance and the fact that its spectroscopy has been very well characterized from a computational and experimental point of view, we use oxirane as a test case to introduce and discuss the computational approach described in the previous section.

Definition of the protocol

The equilibrium rotational constants have been directly derived from the equilibrium geometry obtained by means of the CCSD(T)/CBS+CV “gradient” scheme (see Eq. (21)) in conjunction with large basis sets (cc-pVQZ and cc-pV5Z for the extrapolation to the CBS limit and cc-pCVQZ for the CV contribution). To predict the ground-state rotational constants, these best-estimated equilibrium rotational constants have been then subsequently augmented by vibrational and electronic corrections at the CCSD(T)/cc-pVQZ and CCSD(T)/aug-cc-pVQZ levels, respectively. The electronic contributions are due to the electronic-distribution contribution to the moments of inertia and are thus connected to the rotational g-factor (see, for example, Ref.¹⁰⁶). Quartic and sextic centrifugal-distortion constants have been obtained by means of composite schemes, with the one employed for the quartic terms adopted to derive the best-estimated harmonic wavenumbers (in a normal coordinate representation) as well. In detail, the CCSD(T)/CBS+CV+aug scheme has been used, with the extrapolation to the CBS limit based on results at the CCSD(T)/cc-pVTZ and CCSD(T)/cc-pVTQZ levels, the CV contribution evaluated at the CCSD(T)/cc-pCVTZ level, and the ‘aug’ correction determined from the comparison of CCSD(T)/aug-cc-pVTZ and CCSD(T)/cc-pVTZ values. For sextics, a composite approach analogous to the previous one has been employed, however neglecting the ‘aug’ contribution.

The anharmonic vibrational wavenumbers and IR intensities have been computed using the GVPT2 model applied to hybrid “CC(Best)/CC” anharmonic force fields: the harmonic force field obtained at the CCSD(T)/CBS+CV+aug has been combined with anharmonic

Table 3: Ground-state spectroscopic parameters of oxirane.

Parameter		Best estimate ^a	Calculated ^b	Scaled ^c	Experiment ^d
A	(MHz)	25528.216	25540.56	25491.12	25483.86060(96)
B	(MHz)	22121.162	22135.67	22119.82	22120.87122(88)
C	(MHz)	14110.762	14118.44	14101.74	14097.82440(55)
Δ_J	(kHz)	20.537	20.33	20.11	20.68572(77)
Δ_{JK}	(kHz)	21.630	19.62	21.49	20.91000(216)
Δ_K	(kHz)	26.725	27.09	27.22	27.59143(275)
δ_J	(kHz)	6.104	6.12	6.19	6.210757(211)
δ_K	(kHz)	17.654	17.14	17.38	18.10778(120)

^a Equilibrium rotational constants corresponding to a best-estimated CCSD(T)/CBS+CV equilibrium structure obtained using the cc-pVQZ and cc-pV5Z basis sets for the CBS extrapolation and the cc-pCVQZ set for the core-correlation correction. Corrections to rotational constants: Vibrational at the CCSD(T)/cc-pVQZ level and electronic at the CCSD(T)/aug-cc-pVQZ level. Best-estimated quartic centrifugal-distortion constants obtained with the CCSD(T)/CBS+CV+aug scheme. For details, see Ref.¹⁹.

^b Equilibrium rotational constants corresponding to a CCSD(T)/CBS+CV equilibrium structure obtained using the cc-pVTZ and cc-pVQZ basis sets for the CBS extrapolation and the cc-pCVTZ set for the core-correlation correction. Vibrational corrections to rotational constants at the B3LYP/SNSD level. Quartic centrifugal-distortion constants at the CCSD(T)/cc-pCVTZ level (all electron).

^c Spectroscopic parameters evaluated according to footnote *b* scaled using those of methyloxirane computed at the same level and the corresponding experimental data.¹³⁸

^d Experiment: Ref.¹³⁹.

terms at either the CCSD(T)/cc-pVTZ or CCSD(T)/cc-pVQZ level. For an exhaustive account, the reader is referred to Ref.¹⁹. For comparison purposes, the harmonic force field from the “cheap” scheme has also been considered.¹³⁷ The anharmonic PESs and PSs have been computed employing different QM methods: B3LYP/SNSD, B2PLYP/aug-cc-pVTZ, CCSD(T)/cc-pVTZ and the CCSD(T)/cc-pVQZ.

Rotational Spectroscopy

The rotational spectroscopy parameters of oxirane computed using the protocol described above are summarized in Table 3, where they are compared to the available experimental data. We note a very good agreement between our best estimates and experiment, with discrepancies on the order of 0.1%, 2%–3%, and 3%–4% for rotational, quartic, and sextic centrifugal-distortion constants, respectively.

As mentioned in the Introduction, the focus is on the spectroscopic signatures. Our best-

estimated parameters (first column of Table 3) lead to the prediction of rotational transitions with a relative accuracy better than 0.1% in the centimeter/millimeter-wave region, with this upper limit increasing up to 0.15% in the far infrared (FIR) region. This means that our computed parameters can predict rotational frequencies of 100 GHz and 2 THz with an uncertainty, in the worst case, of 100 MHz and 3 GHz, respectively. Figure 2 shows two portions, at ~ 85 and ~ 104 GHz, of the rotational spectrum of oxirane. Comparison with experiment confirms the expected accuracy: the discrepancies in relative terms range from 0.002% ($J = 6_{2,4} - 6_{1,5}$) to 0.09% ($J = 2_{2,0} - 1_{1,1}$). To further improve the predictive capability of our computational protocol, a scaling procedure can be applied:

$$X_{\text{scal}}^{\text{oxi}} = X_{\text{calc}}^{\text{oxi}} \times (X_{\text{exp}}^{\text{ref}}/X_{\text{calc}}^{\text{ref}}), \quad (25)$$

where X denotes a generic spectroscopic parameter. The superscripts “oxi” and “ref” refer to oxirane and the reference compound, respectively; “scal”, “exp”, and “calc” denote the scaled, experimental, and quantum-chemically calculated values for X . The idea at the basis of this empirical refinement is that the discrepancies affecting the computed spectroscopic constants are in some way systematic and thus, chosen a reference species that shares some similarity in the electronic structure with oxirane, a sort of scaling factor can be derived. We avoid the definition of a unique scaling factor, but instead we obtain one for each parameter, even if we can point out that the resulting scaling factors are generally very similar to one another. The empirical scaling has demonstrated to be extremely effective if the reference molecule is an isotopic species, as shown in Ref.¹⁹ for mono-deuterated oxirane (with the main isotopic species used as reference compound). However, in the present case, for demonstrative purposes, methyloxirane has been chosen as the reference molecule. For both oxirane and methyloxirane, the equilibrium rotational constants have been determined at the CCSD(T)/CBS+CV level using the cc-pVTZ and cc-pVQZ basis sets for the CBS extrapolation and the cc-pCVTZ set for the core-correlation correction, successively augmented by vibrational corrections at the B3LYP/SNSD level.²⁴ The parameters of oxirane have been scaled according to Eq. (25), using the experimental data recently reported for methyloxirane.¹³⁸ The calculated and scaled spectroscopic constants (second and third column of Table 3, respectively) are compared to experiment (fourth column). A significant

improvement is noted; for rotational constants, for example, the discrepancy is reduced to 0.03% or even lower. This improvement in turn leads to better predictions of the rotational transitions with respect to the best-estimated level, as seen in panel *b*) of Figure 2, with deviations below 0.01–0.02%. We finally note that the empirical scaling permits the determination of rotational-parameter values with spectroscopic accuracy when the quantum-chemical protocol described above cannot be applied.




Figure 2: Comparison of the best-estimated (see footnote *a* of Table 3, in red), scaled (in black) and experimental (Ref.¹³⁹, in blue) rotational spectrum of oxirane in the *a*) ~85 GHz and *b*) ~104 GHz frequency regions.

Vibrational Spectroscopy

The GVPT2 fundamental wavenumbers and IR intensities of oxirane computed using the protocol described above are compared in Table 4 with the most accurate gas-phase experimental results. Notably, a very good agreement with experiment is achieved, with mean absolute errors and maximum deviations of about 5.5–7.5 cm^{-1} and 9–13 cm^{-1} , respectively, for all hybrid models when the harmonic part is evaluated by means of a composite scheme accounting for extrapolation to the CBS limit and core correlation effects (either “bestCC”, i.e., CCSD(T)/CBS+CV+aug, or the “cheap” scheme). We note that when the “bestCC” harmonic part is employed, the MAE and |MAX| values are within 5.3–5.6 cm^{-1} and 9–11 cm^{-1} , respectively, for all hybrid models, regardless of the way the anharmonic correction has been evaluated, from expensive CCSD(T)/cc-pVQZ computations or from B3LYP/SNSD. Based

on the results presented in Table 4, but also on several studies available on this topic (see, for example, Refs.^{23,114,131}), CC/QM’ hybrid force fields, where CC stands for a CCSD(T)-based composite approach, are able to predict transition wavenumbers with an accuracy of about 5–10 cm⁻¹ and 10–20 cm⁻¹ for fundamental and non-fundamental bands, respectively. The latter error estimate for overtones and combination bands is in line with the discrepancy between the theoretical “bestCC”/CCSD(T)/cc-pVQZ (2922 cm⁻¹) and experimental (2931 cm⁻¹) wavenumbers for the intense band corresponding to the combination $\nu_2+\nu_{10}$.¹⁹ Concerning IR intensities, we note in all cases a good qualitative agreement, while a more quantitative analysis is precluded due to large error bars on most data caused by the limited resolution of most experiments.¹⁴⁰ Nevertheless, where high-resolution data are available, namely for ν_2 , ν_5 , and ν_{10} ,^{141,142} theoretical results agree well within 10%. Furthermore, we note that the reliability of our computed CC/QM’ fully anharmonic IR intensities has been recently demonstrated not only for fundamentals, but also for overtones and combination bands.^{114,131} The overall conclusion is that the high accuracy of GVPT2 CC/QM’ results allow an accurate and complete characterization of the IR spectrum of oxirane in terms of band positions and intensities, also including overtones and combination bands. The theoretical predictions of all intense transitions, from mid-IR (MIR) to near-IR (NIR), represent a valuable support for the detection of oxirane and other astroCOMs in new space environments through their vibrational signatures. In particular, the “cheap” composite scheme can be effectively applied to the investigation of astroCOMs up to about a dozen of non-hydrogen atoms. We also note an overall good quality of the B2PLYP/aug-cc-pVTZ computations, with mean absolute errors below 10 cm⁻¹ and maximum discrepancies below 20 cm⁻¹. This finding is of interest for astrochemical studies of significantly larger systems, for which harmonic PES computations at the CCSD(T)/cc-pVTZ level are not feasible.

As an example, we present a comparison between the simulated emission spectra of oxirane in the 3–13 μm range (800–6000 cm⁻¹)¹⁹ and the astronomical observations of the planetary nebula NGC 7027, as reported in a uniform database (2.4–45.4 micron spectra) from the Infrared Space Observatory Short Wavelength Spectrometer (ISO-SWS).¹⁴⁶ Figure 3 shows selected parts of the spectra, in the 3.2–3.8 μm and 10.5–13.0 μm ranges, along with all identified line fluxes¹⁴⁷ marked by vertical grey lines, thus clearly showing that sev-

Table 4: Theoretical predictions of GVPT2 wavenumbers (ν / cm^{-1}) and infrared intensities ($I_{\text{IR}} / \text{km mol}^{-1}$) for oxirane fundamental transitions, compared to experiment.

Mode	Symm.	B2PLYP ^a		"cheap" ^b		bestCC ^c		Experiment		Assignment						
		ν	I_{IR}	ν	I_{IR}	ν	I_{IR}	ν	I_{IR}		ν	I_{IR}^g	Ref.			
ν_1	A1	3031	9.5	3025	11.3	3018	13.8	3027	16.4	3024	15.9	3021	15.9	3018	143	CH ₂ sym stretch
ν_2	A1	1500	1.2	1489	1.1	1487	1.2	1492	0.9	1489	1.1	1490	1.2	1498	141,144	CH ₂ scissor
ν_3	A1	1272	14.2	1265	14.0	1264	13.8	1270	14.2	1268	13.8	1266	13.6	1270	140,144,145	ring breathing
ν_4	A1	1124	0.4	1118	0.6	1115	0.4	1120	0.2	1119	0.3	1115	0.2			CH ₂ wagging
ν_5	A1	866	68.7	869	72.7	869	72.7	876	72.0	875	68.7	873	68.9	877	142,144 ^a	Ring deformation
ν_6	A2	3057	0.0	3061	0.0	3058	0.0	3064	0.0	3062	0.0	3065	0.0			CH ₂ asym stretch
ν_7	A2	1163	0.0	1155	0.0	1156	0.0	1157	0.0	1156	0.0	1160	0.0			CH ₂ rocking
ν_8	A2	1028	0.0	1025	0.0	1019	0.0	1027	0.0	1026	0.0	1027	0.0			CH ₂ twisting
ν_9	B1	3026	23.2	3016	19.1	3009	29.5	3017	30.2	3015	31.4	3009	31.4	3007	140,143	CH ₂ asym stretch
ν_{10}	B1	1478	0.1	1464	0.4	1462	0.3	1465	0.1	1463	0.1	1461	0.1	1471	141,144	CH ₂ scissor
ν_{11}	B1	1147	0.2	1145	0.1	1150	0.2	1145	0.2	1144	0.3	1154	0.2			CH ₂ wagging
ν_{12}	B1	809	10.4	809	9.1	809	8.6	818	9.3	817	8.8	816	8.3	822	140,144	Ring deformation
ν_{13}	B2	3071	44.3	3074	32.3	3071	31.9	3077	32.0	3075	35.0	3076	34.2	3066	140,143	CH ₂ asym stretch
ν_{14}	B2	1149	3.6	1143	4.0	1141	4.2	1146	4.3	1145	4.1	1145	4.1			CH ₂ twisting
ν_{15}	B2	812	0.1	807	0.4	806	0.5	807	0.3	807	0.2	810	0.3	808	140,144	CH ₂ rocking
	MAE ⁱ	8.4		7.3		6.2		5.3		5.6		5.4				
	MAX ⁱ	18.9		12.8		12.9		11.0		9.0		10.0				

^a Computed with the aug-cc-pVTZ basis set.

^b Harmonic wavenumbers and IR intensities obtained by means of the "cheap" composite scheme. For details, see Ref. 137.

^c CCSD(T)/CBS+CV+aug harmonic wavenumbers and IR intensities (see text). For details, see Ref. 19.

^d Computed with the cc-pVTZ basis set.

^e Computed with the SNSD basis set⁵³.

^f Computed with the cc-pVQZ basis set.

^g Numbers given in parenthesis are the experimental errors.

^h Total intensity in the spectral region 750–950 cm^{-1} corresponding to the ν_5 , ν_{12} , and ν_{15} fundamental transitions.¹⁴²

ⁱ Mean absolute error (MAE) and maximum absolute deviations (|MAX|) with respect to experimental wavenumbers.

eral, still unidentified IR transitions (UIR) could be assigned. From Figure 3, we note that the group of oxirane transitions related to the CH₂ stretching fundamental vibrations match well the observed UIR feature at 3.3 μm ; similarly the one at 3.4 μm is well represented by the strong combination band $\nu_2 + \nu_{10}$. In particular, a reliable prediction of the latter requires fully anharmonic computations in order to assess the band position and absolute intensity. Nevertheless, this good match cannot be conclusive for the detection of oxirane because the broad features in the ISO-SWS database are most likely resulting from CH stretching vibrations of several astroCOMs. In this respect, a good agreement between our best-estimated spectra and less computationally demanding models (in particular based on B2PLYP) in reproducing the main spectral features opens a path to reliable simulations of more complex, aromatic and aliphatic molecules suggested to be present in circumstellar and interstellar nebulae.¹⁴⁸ More characteristic of oxirane is its most intense band at 11.45 μm (873 cm^{-1}), which is related to the ring deformation, and for which the good match with the 11.2–11.3 μm UIR feature provides a more plausible support to oxirane detection. This band can also be chosen to distinguish between oxirane and methyloxirane, which shows weaker features at 10.53 and 12.05 μm (950 cm^{-1} and 830 cm^{-1}) that are related to the ring deformation and the ring C-C stretching, respectively.²⁴ Unfortunately, this band cannot be used for the possible identification of oxirane in Titan’s atmosphere, as the infrared spectra recorded by Cassini’s (CIRS) instrument over the 10–1500 cm^{-1} range show a strong absorption at 11.5–12.5 μm (800–860 cm^{-1}) due to the ethane ν_9 band, thus hiding any close-lying transitions.

Challenges in computational spectroscopy of astroCOMs

Spectral surveys consisting in a series of spectra covering, at millimeter/submillimeter wavelengths, several dozens of GHz with a uniform sensitivity are the ideal mean to obtain a complete census of the species emitting in the investigated astronomical object. The first step of their analysis thus consists in identifying the various lines, with the main difficulty being related to those molecules responsible for hundreds of them. These are species that are characterized by large amplitude motions (LAM) that very frequently occur at low fre-




Figure 3: Comparison between the simulated (colour) spectra of oxirane and the observed (black) emission spectra of planetary nebula NGC 7027 (Sloan et al.¹⁴⁶). Selected wavelength ranges: (a) the 3.2–3.8 μm and (b) 10.5–13.0 μm . The observed Identified line fluxes reported by Salas et al.¹⁴⁷ are marked by the grey lines. Computations with hybrid models are denoted as QM/QM', with CCvxz (x=t,q) standing for CCSD(T)/cc-pVXZ. Theoretical spectra have been convoluted with a Lorentzian function with half-width at half-maximum of 1 cm^{-1} .

quencies. Typical examples are methanol, methyl formate or dimethyl ether, and the large number of lines is often due to rotational transitions in low-frequency vibrational modes (like those associated to the methyl rotor). In addition to the difficulty of astronomical detection and identification, these molecules represent a major challenge for molecular spectroscopy and, in particular, for computational spectroscopy.

Another important issue is the identification of conformers and their relative abundances, which might become crucial when astronomical observations involve the hotter regions of the ISM, like star-forming sources known as “hot cores”. Among the astroCOMs, in relation to the issue of the origin of life, particular interest is devoted to those showing a prebiotic character. These species are usually flexible systems and therefore their conformational equilibrium needs to be taken into account and accurately investigated. Inaccurate or wrong evaluation of the relative conformational energies may give misleading results and conclusions.

Large Amplitude Motion

A large number of unidentified lines, especially in spectra of the hot cores associated with star formation, are due to rotational transitions in the low-lying excited vibrational states of a relatively small number of molecules, denoted as “astrophysical weeds” (see, for example, Ref.¹⁴⁹ and references therein). From a computational point of view, it is necessary to characterize the rotational spectra in the vibrational ground state and in the low-lying vibrational states, and understand how they are affected by LAM(s).

The recent computational spectroscopic characterization of acetyl cyanide (CH_3COCN) and its enolic species¹⁵⁰ provides the opportunity to point out the accuracy obtainable in the computation of the required spectroscopic parameters. Acetyl cyanide belongs to those medium-sized molecules (6–10 atoms), detected or expected to exist in the ISM, that show a potential prebiotic character because present in plausible chemical routes to simple biomolecules, like amino acids or simple peptides. Therefore, a lot of effort has been put in the identification of these compounds, their precursors, or their fragments. Although acetyl cyanide has not been detected yet, it is expected to be present in the ISM. Its electronic ground state has been already well characterized experimentally, thus giving us the opportunity to demonstrate how highly accurate QM approaches are able to deal with the challenges




Figure 4: The potential energy profile of acetyl cyanide along the torsional angle (see, the inset) evaluated at the CCSD(T)/CBS+CV level using CCSD(T)/cc-pVTZ optimized geometries.

offered by hindered methyl-rotor in rotational spectroscopy.

Acetyl cyanide shows a methyl torsional motion. The internal rotation of the $-\text{CH}_3$ group relative to the $-\text{COCN}$ frame can be described by the torsional angle α (see Figure 4), which leads to three equivalent configurations. Therefore, the potential energy function with respect to α is periodic and can be modelled by the expression:

$$V(\alpha) = \frac{V_3}{2} [1 - \cos(3\alpha)] . \quad (26)$$

In the limit of an infinite barrier, each torsional state v is three-fold degenerate. For a finite barrier, the tunneling effect leads to a splitting of this threefold degeneracy (because of the probability of tunneling through the barrier) into two levels, a non-degenerate A level and a doubly-degenerate E level (see Figure 4). As a consequence, in the rotational spectrum we observe rotational transitions in the A and E torsional sublevels for each vibrational

state; in other words, the rotational spectrum of acetyl cyanide in its vibrational ground state results from the overlap of the rotational spectrum of the A state and that of the E state. Therefore, to predict the overall spectrum the rotational constants for both A and E sublevels are needed.

In Ref.¹⁵⁰, the CCSD(T)/CBS+CV “gradient” scheme has been employed to accurately predict the equilibrium rotational constants, while the vibrational corrections have been evaluated using the VPT2 approach with a cubic force field computed at the MP2/aug-cc-pVTZ level, which also provided the sextic centrifugal-distortion constants. The “cheap” geometry scheme has been used instead to generate an harmonic force field, which allowed in turn the evaluation of the quartic centrifugal-distortion terms. To predict the rotational constants of the E state, it is necessary to calculate the V_3 barrier. For this purpose, the potential energy function along the torsional angle has been determined at the CCSD(T)/cc-pVTZ level by carrying out geometry optimizations with the α value kept fixed (also known as a relaxed scan), as shown in Figure 4. In a subsequent step, the energy evaluation has been improved by single-point calculations at the CCSD(T)/CBS+CV level on the CCSD(T)/cc-pVTZ optimized structures, thus leading to a V_3 value of 420.1 cm^{-1} , in good agreement with the experimental value ($V_3 = 404.867(90) \text{ cm}^{-1}$).¹⁵¹ The computed spectroscopic parameters are collected in Table 5, where they are compared to their experimental counterparts. Despite a slight overestimation of the splitting between A and E rotational constants, the overall accuracy is very satisfactory. In Table 5, the prediction of the rotational constants for the lowest vibrational states ($v_{18} = 1$ and $v_{12} = 1$) together with the available experimental data is also reported. Making use of Eq. (25), the computed data have also been scaled based on the experimental values for the vibrational ground state. For all rotational constants, the discrepancies between experiment and theory, in relative terms, are well below 0.1%, a value which can be further improved (to $\sim 0.03\%$) by applying an effective empirical strategy such as the scaling procedure. In Ref.¹⁵⁰, a comparison between the computed and experimental centrifugal-distortion constants is also provided. For the best-estimated quartic centrifugal-distortion constants, the average discrepancy with respect to experiment is about 2.7%, which can be viewed as a very good result because of the presence of a large amplitude motion and the smallness of these constants. The sextic centrifugal-distortion constants

are extremely small, the largest ones being only a few tenths of Hz. Therefore, an average discrepancy in relative terms with respect to experiment of 16% can be considered a good outcome.

The large amplitude vibrations cannot be accurately described by perturbative approaches, and require an expansion of the PES beyond the quartic order. A possible solution, albeit limited to small molecules (i.e., less than ten vibrational degrees of freedom), is to carry out fully variational computations of the ro-vibrational energy levels.^{28–30,152} However, more effective models are required for larger systems, like the astroCOMs. In this respect, it is convenient to retain a perturbative treatment and separate the degrees of freedom related to small-amplitude vibrations and LAMs. These can then be treated separately or with hybrid schemes,¹⁵³ like those based on the reaction path/surface Hamiltonian.^{154–156} It should be noted that, in order to reduce the couplings between LAMs and other vibrational degrees of freedom, internal coordinate-based approaches are more suitable. Nevertheless, for molecules showing both LAM and small-amplitude vibrations, standard VPT2 computations based on Cartesian-based normal coordinates for the whole system allow mapping all molecular vibrations and represents a feasible route, for instance, to check whether there are any Fermi interactions between large- and small- amplitude vibrations.¹⁵⁷

Another interesting example in the field of astrochemistry¹⁵⁸ is represented by ethyl mercaptan¹⁵⁷ (EtSH), with the *gauche* conformer only slightly more stable than the *trans* one with an energy difference of 162 cm⁻¹. The conformational flexibility of this molecule is related to the H-S-C-C torsional angle (see Figure 1), which leads to a potential energy profile with barriers within 350–500 cm⁻¹. In EtSH, there is a second LAM, which corresponds to the internal rotation of the methyl group and is characterized by a significantly larger barrier of about 1200 cm⁻¹. Both conformers show dipole moments of the same order of magnitude (about 1.8 Debye), which render possible their detection in the ISM by means of radioastronomical observations either in millimeter-/submillimeter-wave or IR ranges. In Ref.¹⁵⁷, in order to provide the required spectroscopic characterization, the H-S-C-C torsional motion and the internal rotation of the methyl group have been excluded from the perturbative description and treated variationally. For this purpose, a two-dimensional PES has been computed at the CCSD(T)/aug-cc-pVTZ level of theory, and then the couplings

Table 5: Computed and experimental rotational constants (MHz) of acetyl cyanide for the vibrational ground state and for the $v_{18} = 1$ and $v_{12} = 1$ vibrational states.

Parameter	<i>A</i> state		<i>E</i> state		Scaled ^c
	Best theo. ^a	Experiment ^b	Best theo. ^a	Experiment ^b	
Ground state					
A_0 / MHz	10193.6	10185.37774(26)	10193.0	10185.13561(34)	
B_0 / MHz	4157.3	4157.48341(11)	4157.0	4157.37176(10)	
C_0 / MHz	3003.2	3002.76275(11)	3003.2	3002.76390(12)	
D_J / kHz	0.790	0.816733(84)			
D_{JK} / kHz	17.916	17.80532(26)			
D_K / kHz	-7.029	-6.6543(10)			
d_J / kHz	0.256	0.266340(26)			
d_K / kHz	10.382	10.42700(48)			
H_J / Hz	3.410×10^{-4}	$4.51(21) \times 10^{-4}$			
H_{JK} / Hz	0.202	0.17658(16)			
H_{KJ} / Hz	-0.437	-0.37141(51)			
H_K / Hz	0.285	0.2519(10)			
h_J / Hz	1.714×10^{-4}	$2.183(54) \times 10^{-4}$			
h_{JK} / Hz	0.104	0.09265(15)			
h_K / Hz	0.479	0.4374(10)			
$v_{18} = 1$ (126.06 cm ⁻¹)					
A_0 / MHz	10195.3	10191.8472(22)			10187.1
B_0 / MHz	4149.2	4149.5333(13)			4149.4
C_0 / MHz	3001.1	3000.69275(94)			3000.6
$v_{12} = 1$ (176 cm ⁻¹)					
A_0 / MHz	10176.9	10157.64443(76)			10168.7
B_0 / MHz	4176.8	4175.90448(37)			4176.9
C_0 / MHz	3006.3	3005.52426(35)			3005.9

^a Equilibrium rotational constants (\mathbf{B}^{eq}) from CCSD(T)/CBS+CV equilibrium structure augmented by vibrational corrections ($\Delta\mathbf{B}^v$) at the MP2/aug-cc-pVTZ level. Best-estimated quartic centrifugal-distortion constants from the “cheap” scheme. Sextic centrifugal-distortion constants at the MP2/aug-cc-pVTZ level. For details, see Ref.¹⁵⁰.

^b Ref.¹⁵¹

^c Scaled parameters obtained by employing Eq. (25).

Table 6: Theoretical predictions of the large-amplitude fundamentals (ω and ν in cm^{-1}), as obtained from the 2-D variational and GVPT2 treatments, compared to experiment.

astroCOMs	Harmonic ^a	2D-variational ^b	GVPT2 ^a	Experiment ^c	Assignment
	ω	ν	ν	ν	
<i>gauche</i> -EtSH	201	189.61	191	193.0	(-)SH torsion
		188.13		191.8	(+)SH torsion
	259	254.07	251	247.5	(-)CH torsion
		254.05			(+)CH torsion
<i>trans</i> -EtSH	179	155.74	171	158.0	SH torsion
	255	245.20	248		CH torsion

^a Computations at the CCSD/cc-pVTZ level, see Ref. ¹⁵⁷.

^b Computations at the CCSD(T)/aug-cc-pVTZ level, see Ref. ¹⁵⁷.

^c Experimental wavenumbers as reported in Ref. ¹⁵⁷.

with other vibrational degrees of freedom have been taken into account through an effective correction of their contribution to the zero-point vibrational energy (ZPVE). The reader is referred to Ref. ¹⁵⁷ for all details. These computations allowed the accurate determination of the torsional energy levels, including the tunneling splitting observed for the *gauche* conformer. The results are summarized in Table 6, which highlights the large differences between the *gauche* and *trans* ethyl mercaptan spectral features in the FIR region. In particular, the fundamental transitions for the SH and CH torsions of *gauche*-EtSH are localized at 188.87 cm^{-1} with a splitting of 1.47 cm^{-1} and at 254.06 cm^{-1} with a splitting of 0.02 cm^{-1} , respectively. They clearly do not superimpose with those of *trans*-EtSH, which are shifted by 32 cm^{-1} and 9 cm^{-1} , respectively. In Table 6, the variational computations can be compared with the GVPT2 results, thus showing that the perturbative treatment provides reasonable frequencies for the CH torsion of both conformers and for the SH torsion of *gauche*-EtSH. This is not the case for the SH torsion of *trans*-EtSH, for which the first excited energy level lies above the torsional barrier. ¹⁵⁷

The problem of a reliable treatment of large-amplitude modes is not only related to their direct detection in the FIR region, but also to the coupling with other vibrations that can

be observed in the MIR range. In this respect, the LAM can contaminate significantly the VPT2 treatment, thus leading to large errors even for the intense, high frequency vibrations. This issue can be overcome by employing reduced-dimensionality perturbative models, where LAMs are excluded from the VPT2 treatment. The efficiency of such an approach is evinced in Table 7, which collects some examples for the *gauche* conformer of EtSH, discussed above, and the nucleobase cytosine. For the latter, the large amplitude motion is the out-of-plane wagging vibration of the NH₂ group (see Figure 1). The torsional and wagging LAMs coupled with the H-X (X=C, S, N) stretching vibrations lead to the incorrect prediction of the latter by means of GVPT2. At variance, in the reduced-dimensionality GVPT2 models (i.e., the so-called “SkipPT2” approach), the LAMs and all related cubic and quartic force constants are excluded from the perturbative treatment. As evident from Table 7, this treatment leads to the accurate prediction of the intense IR bands related to the C-H stretching vibrations of *gauche*-EtSH and to the N-H stretchings of cytosine. For *gauche*-EtSH, with computations at the CCSD(T)/cc-pVTZ level, the so-called “SkipPT2” approach leads to MAE and |MAX| of 4.5 cm⁻¹ and 7 cm⁻¹, respectively. A slightly lower accuracy is observed for cytosine (MAE=15 cm⁻¹ and |MAX|=21 cm⁻¹), as expected from QM computations at the B3LYP level.¹²⁹ As noted before, the accuracy can indeed be improved with GVPT2 computations based on CC/DFT hybrid force fields. However, it might be unclear *a priori* whether the LAMs contaminate the GVPT2 treatment of high-frequency vibrations or not. To address this point, we can consider the case of methyloxirane that shows a barrier for the internal rotation of the methyl group of 893 cm⁻¹.¹³⁸ It has been shown that in this case, the full GVPT2 treatment using the “cheap”/B3LYP hybrid force field leads to highly accurate wavenumbers, with a mean absolute deviation of 4.8 cm⁻¹ and a maximum error below 18 cm⁻¹ for all fundamental vibrations, even for the C-H stretchings and the torsion of the methyl group.²⁴ Interestingly, the reduced model with the methyl torsion excluded from VPT2 shows results very similar to the full GVPT2 approach for all C-H stretching vibrations, but with improved mean and maximum errors with respect to experiment, reduced to 4 cm⁻¹ and 8 cm⁻¹, respectively. This indicates that the reduced-dimensionality, LAM-free GVPT2 models represent a reliable approach for accurate vibrational computations and should be preferred in presence of LAMs, especially if the actual impact of the latter on the

Table 7: Theoretical predictions of high-frequency vibrations (ω and ν in cm^{-1}) coupled with LAM, compared to experiment.

astroCOMs	Harmonic	GVPT2 Full	GVPT2 Skip ^a	Experiment ^b	Assignment
	ω	ν	ν	ν	
<i>gauche</i> -EtSH ^c	3110	3033	2969	2967	CH stretch
	3130	2958	2987	2980	CH ₂ stretch
cytosine ^d	3597	3473	3438	3441	NH ₂ sym stretch
	3617	3452	3450	3472	N ₁ H stretch
	3733	3604	3540	3564	NH ₃ asym stretch
methyloxirane ^e	3039	2961	2959	2942	CH ₃ sym stretch
	3110	3015	3015		CH ₂ sym stretch
	3117	2978	2973	2974	CH ₃ asym stretch
	3120	2993	2991	2995	CH ₃ asym stretch
	3139	3001	2998	3001	CH stretch
	3204	3058	3058	3051	CH ₂ asym stretch
MAE ^f	142	21	9		
MAX ^f	169	66	24		

^a Low-frequency large-amplitude modes excluded from the GVPT2 treatment.

^b Experimental wavenumbers as compiled in: Ref. ¹⁵⁷ for *gauche*-ethyl mercaptan, Ref. ¹²⁹ for cytosine, and Ref. ²⁴ for methyloxirane.

^c Computations at the CCSD(T)/cc-pVTZ level, see Ref. ¹⁵⁷.

^d Computations at the B3LYP/SNSD level, see Ref. ¹²⁹.

^e Computations at the “cheap”/B3LYP level see Refs. ^{24,56}

^f Mean absolute error (MAE) and maximum absolute deviations (|MAX|) (in cm^{-1}) with respect to the experimental wavenumbers.

rest of the system cannot be assessed univocally.

Conformers

As mentioned above, the rotational spectrum of molecular species that exhibit large amplitude motions can be a challenge for both computational and experimental molecular spectroscopies. AstroCOMs of increasing size and complexity are often characterized by high flexibility due to rotations around single bonds. On general grounds, the presence of LAMs implies molecular flexibility, which in turn leads to the existence of several conformers. The analysis of the rotational and vibrational spectra might therefore be hampered by the concomitant presence of different conformers, thus requiring the accurate determination of their

relative energies. The situation can be further complicated by the fact that the relative energies of the lowest-excited vibrational states of the most stable conformer might be similar to those of the vibrational ground states of less stable species.

A noteworthy example is the rotational spectrum of glycine ($\text{NH}_2\text{CH}_2\text{COOH}$), the smallest amino-acid, which has been found in several carbonaceous meteorites, and is one of the two most abundant amino acids in Stardust-returned foil samples,²⁰ but its presence in the ISM has not yet been confirmed. As pointed out in Ref.¹⁵, regarding its search in hot-core sources (i.e., in environment at $T \sim 200$ K) the second most stable conformer, IIn, is more important than the first one, glycine Ip. In fact, at a temperature of about 200 K, despite the relative population of the conformer IIn being only $\sim 10\%$, its rotational spectrum is more intense than that of the most abundant Ip conformer. This is due to the larger dipole moment components of glycine IIn with respect to those of glycine Ip (see Figure 1). However, the correct evaluation of their relative energy is of fundamental importance. Figure 5 shows how the relative intensity of the rotational spectrum changes depending on the energy difference between Ip and IIn. The CDMS database reports a value of about 700 cm^{-1} ,¹⁵⁹ which implies at $T = 200$ K a population for the IIn conformer lower than 1% (using the relative energy instead of the free energy to estimate the population ratio). The consequence is that, as seen in the left side of Figure 5, the rotational spectra of the two conformers have approximately the same intensity. On the other hand, if a value of 312 cm^{-1} , obtained at the CCSD(T)/CBS+CV level (also including ZPVE correction),²² is considered, the rotational spectrum of the IIn conformer results to be about four times more intense than that of glycine Ip (see right side of Figure 5). While the accurate evaluation of the relative energy of glycine conformers is addressed in the following, we point out that the results presented here have an important impact on the attempt of identifying glycine in space. The significant population of glycine IIn already at 200 K is important for the detection of glycine in the ISM; in particular, on the basis of the recent literature that asserts that glycine could reach detectable levels in cold solar-type system precursors¹⁶⁰ or hot-core sources.¹⁶¹ The rotational spectrum of the most stable conformer of glycine is expected to have peak intensities greater than 10 mK in the ALMA band 2 (68–90 GHz) for detections from a solar-type system precursor, like the L1544 pre-stellar core.¹⁶⁰ For pre-stellar core systems, due to their characteristic

very low temperatures, it is fully justified to consider that only the most stable conformer, Ip, is present. However, only glycine Ip has also been considered for the simulation of the emission spectra of the hot-core source NGC 6334 IRS1,¹⁶¹ with a characteristic temperature of 200 K.




Figure 5: Simulation of the rotational spectra of glycine Ip and IIn at $T = 200$ K in the 80–250 GHz frequency range. Spectroscopic parameters are taken from the CDMS database,¹⁵⁹ while the population ratio is based on an energy difference of either 700 cm^{-1} (left, Ref.¹⁵⁹) or 312 cm^{-1} (right, Ref.²²).

More exhaustively, with respect to the notation used above, the glycine conformers are denoted using, in addition to the number for the stability order and the “p,n” labels for the planarity or non-planarity of the backbone, respectively, the “c,t” labels for indicating the *cis* or *trans* orientation with respect to the three torsional angles. According to this notation, the three experimentally observed conformers are Ip/ttt, IIn/ccc and IIIp/tct (see Ref.²² for extended discussion). Among the torsional degrees of freedom, those around the C–N and C–C bonds are also large amplitude motions. In addition to the problem of getting correct spectroscopic properties as described above, they represent also a challenging aspect for the accurate calculation of thermochemical properties.^{51,162–165} In this case as well, a feasible route is to partition the molecular Hamiltonian between large- and small-amplitude motions, which can be treated at different levels of accuracy.¹⁶⁵ In this respect, within the hindered-rotor anharmonic-oscillator model (HRAO), the modes identified as large-amplitude, hindered-rotor motions¹⁶² are treated separately and considered inactive

in the vibrational perturbative computations. The HRAO approach, in conjunction with HDCPT2 computations and simple perturbation theory (SPT),¹⁶⁶ represents a viable route for reliable computations of thermodynamic properties.⁵¹

Table 8 collects the theoretical prediction of the gas-phase thermodynamic properties of the glycine conformers at 15 K, 200 K and 410 K. These three temperatures have been selected as they correspond to different types of astrochemical environments: from the cold interstellar regions to the hot cores, and finally toward the temperature limits for the detection of glycine before its decomposition at 506 K. All predictions are based on the best estimates of relative electronic energies at the CCSD(T)/CBS+CV level in conjunction with thermodynamic properties derived from either the standard RRHO model or the more accurate HRAO scheme, with the anharmonic PES computed at the “cheap”/B3LYP level.²² Boltzmann populations of the Ip/ttt and IIn/ccc conformers at different temperatures, derived from the data collected in CDMS,¹⁵⁹ are also reported for comparison purposes. All data agree that only the most stable conformer can be present at 15 K, despite differences in the prediction of relative energies up to 5 kJ mol⁻¹. The situation is much different at higher temperatures. In particular, we can note that the RRHO model strongly overestimates the population of the IIIp/tct rotamer, which is predicted to have a population of about 10% at 200 K and over 50% at 410 K, thus meaning a population larger than that of the Ip/ttt conformer. At variance, the full HRAO approach is able to provide a reasonable relative free energy for this conformer at all temperatures, in good agreement with experimental laboratory observations.¹⁶⁷ The differences between the results obtained with RRHO and HRAO are significant also for the IIn/ccc conformer, for which our best theoretical estimates predict non-negligible abundances of about 10% at 200 K and 20% at 410 K.

As addressed for the detection of rotational features, also in the case of IR observations it is necessary to identify the most stable conformers and to assess the probability of their detection in the astronomical environments, which might vary considerably depending on space-like conditions and spectral ranges. Considering the detection of glycine by means of its signatures in the MIR region, and according to the discussion above, we should conclude that the Ip/ttt conformer gives the strongest signals due to its largest population at all temperatures and to the fact that the most characteristic bands show similar intensity.

Of interest to the present discussion is that extremely accurate results have been obtained by applying the GVPT2 approach to the “cheap”/B3LYP hybrid force field, thus providing accurate predictions for the most intense C=O stretching vibration (the so-called Amide I), i.e., with an accuracy of $\pm 5 \text{ cm}^{-1}$ for all glycine conformers.²² This finding is of relevance to the detection of glycine, and other prebiotic molecules, in complex astrophysical environments, thus allowing the distinction of the different conformers in the fingerprint spectral range.¹⁵

Challenges in computational chemistry of astroCOMs

The evidence that a fairly rich chemistry is taking place across the Universe and the discovery of biomolecular precursors in the interstellar space also arise the issue of the terrestrial origins of life, whose elucidation is one of the goals of astrochemistry. Two alternative theories have been suggested so far: (1) exogenous and (2) endogenous synthesis. In the first theory, prebiotic molecules came from space, the carriers being comets, asteroids and meteorites.¹⁶⁸ The rationale behind this suggestion is that plenty of astroCOMs have been observed in interstellar clouds. In the frame of this theory, it is of fundamental importance not only to discover prebiotic species in space, but also to understand how they could be produced in the typically harsh conditions (extremely low temperature and density) of the ISM. In the endogenous theory, the synthesis of simple organic molecules having a potential relation to the origin of life occurred directly on our planet starting from simple parent molecules in the atmosphere, liquid water and various energy sources. The Urey-Miller experiment was a milestone in this theory.^{169,170} In this context, Titan has been postulated to represent a model of primitive Earth.¹⁷¹ Therefore, the organic chemistry in Titan’s atmosphere could be directly linked to prebiotic organic synthesis in the atmosphere of our primitive planet.

Formation routes

As briefly addressed above, one of the challenges of astrochemistry is to understand how the molecular complexity gradually evolves from simple atomic or diatomic species to astroCOMs, and toward biological macromolecules, the building blocks of life. Focusing on

Table 8: Theoretical^a thermodynamic properties (kJ mol⁻¹) of the three experimentally observed glycine conformers.

	CDMS ^b						$\Delta E+ZPVE^c$						ΔG								
	Ip/ttt		IIn/ccc		%		Ip/ttt		IIn/ccc		%		IIIp/tct		Ip/ttt		IIn/ccc		IIIp/tct		
	kJ mol ⁻¹	%	kJ mol ⁻¹	%	kJ mol ⁻¹	%	kJ mol ⁻¹	%	kJ mol ⁻¹	%	kJ mol ⁻¹	%	kJ mol ⁻¹	%	kJ mol ⁻¹	%	kJ mol ⁻¹	%	kJ mol ⁻¹	%	
15K																					
RRHO	0	100	8.37	<0.01	0	100	3.82	<0.01	7.48	<0.01	0	100	3.82	<0.01	7.34	<0.01	7.34	<0.01	7.34	<0.01	
HRAO ^{d,e}					0	100	3.73	<0.01	7.94	<0.01	0	100	3.72	<0.01	7.87	<0.01	7.87	<0.01	7.87	<0.01	
200K																					
RRHO		99.4		0.6		90.0		9.0		1.0		82.5		4.24		6.4		3.34		11.1	
HRAO ^{d,e}						89.7		9.5		0.8		88.7		3.53		10.6		8.08		0.7	
410K																					
RRHO		92.1		7.9		66.9		22.7		7.7		38.3		5.41		7.9		-1.17		53.9	
HRAO ^{d,e}						66.8		23.4		6.8		75.1		4.41		20.6		9.72		4.3	

^a Conformational energies with respect to the Ip/ttt conformer. All thermodynamic properties have been computed at 1 atm.

^b Ref.¹⁵⁹.

^c CCSD(T)/CBS+CV energy computed at the corresponding optimized geometry and corrected for ZPVE at the “cheap”/B3LYP level.

^d The two lowest vibrations have been described by hindered-rotor contributions computed by means of an automatic procedure.¹⁶²

^e Contributions computed by means of the HDCPT⁵¹ model using the hybrid “cheap”/B3LYP force field in conjunction with simple perturbation theory (SPT)^{51,166} (see text for details).

astroCOMs, a good understanding of the possible formation/destruction pathways and of the associated thermochemistry and kinetics is necessary to fully exploit the currently available and future observational data as well as to prepare new observations. Since laboratory experiments are often not able to reproduce adequately the extreme conditions that characterize the ISM, accurate state-of-the-art computational approaches play a fundamental role to derive possible reaction mechanisms. Therefore, quantum-chemical calculations are increasingly exploited. Indeed, quantum chemistry has demonstrated to be a powerful tool to obtain reliable input parameters for astrochemical models, which in turn lead to simulated abundances (See for instance Refs.^{172,173} for further details). Therefore, reaction schemes with all relevant pathways need to be formulated, thus requiring the identification of all intermediate species, from reactants to products, as well as of the transition states connecting them. In practical terms, accurate electronic structure calculations should be carried out for every chemical species involved in the reaction scheme in order to obtain structure, energies, spectroscopic properties and from them thermochemical data. The latter are then fed into a suitable tool to compute kinetic rate constants.

A survey of the available literature reveals that much information is still lacking and only a small fraction of the elementary reactions has been characterized. The mechanisms leading to astroCOMs in star forming regions are still object of debate, although it is clear that the synthesis can proceed via chemical reactions on the surface of (icy) dust grains or in the gas phase.¹⁷⁴ Both synthetic routes represent great challenges for quantum chemistry, with the key point being the very accurate evaluation of the energetics: the reaction to occur in the harsh conditions typical of the ISM requires that all transition states lie below the reactant asymptotes. The vicinity of the energy levels of the transition states and the reactant asymptotes is a real computational challenge because the status of the transition states as ‘emerged’ or ‘submerged’ can fall within the uncertainty of the calculations. In this respect, the recent investigation of the gas-phase formation route of formamide provides a significative example.²⁶ Formamide (HCONH_2) has a peculiar role in prebiotic chemistry because it can generate both genetic and metabolic molecules (see for instance Ref.¹⁷⁵). According to Ref.²⁶, the $\text{NH}_2 + \text{H}_2\text{CO}$ reaction proceeds with the formation of a bound intermediate. Along the reaction path, a van der Waals complex is formed and a transition




Figure 6: Proposed path for the approach of $\text{NH}_2 + \text{formaldehyde (H}_2\text{CO)}$. Relative electronic CCSD(T)/CBS+CV energies (in kJ mol^{-1}) in black. In the inset: the relative energy of the TS5 transition state at different levels of theory is given. See text.

state has to be surmounted to reach the addition intermediate (see Figure 6). Then, a second transition state connects the bound intermediate to the final products. It should be noted that, for this particular case, the relative energy of the transition state is above the reactant asymptotes. However, the issue here is whether this transition state is significantly above the reactants or not. The results at different levels of theory are collected in Figure 6. It is noteworthy that at the CCSD(T)/CBS+CV level, the energy of the transition state is slightly above the energy of the reactants' asymptote. However, when a full treatment of triple (fT) and quadruple excitations (fQ) is also included, thus leading to the CCSD(T)/CBS+CV+fT and CCSD(T)/CBS+CV+fT+fQ approaches, the barrier related to this transition state decreases significantly, dropping from 3.77 kJ mol⁻¹ to 2.05 kJ mol⁻¹ with full-T and to 1.67 kJ mol⁻¹ with full-T and full-Q. From Figure 6, the inadequacy of the CCSD(T)/maug-cc-pVTZ level of theory is evident, while the good performance of the B2PLYP functional in conjunction with D3 dispersion correction¹⁷⁶, even if clearly due to a compensation of errors, is encouraging for the investigation of larger systems.

Chemical evolution

Recent measurements have revealed that Titan's atmosphere contains significant quantities of carbon ($\sim 2\%$ CH₄) and nitrogen (98% N₂), and traces of oxygen (~ 50 ppm CO).¹⁷¹ Therefore, the photochemical production of complex molecules containing C, N, O, and H is possible. Due to its reducing atmosphere and its rich and complex organic chemistry (see, Ref.¹⁷⁷ and references therein), Titan is thought to represent a model of primitive Earth. The investigation of Titan's atmosphere might therefore provide a unique opportunity to shed light on the organic evolution undertaken in the atmosphere of early Earth. Furthermore, Cassini measurements have revealed the presence of numerous and large carbocations and carbanions in Titan's upper atmosphere (see, Ref.¹⁷⁷ and references therein). No other planets or satellites in our solar system have been found to bear the same level of chemical complexity as observed on Titan. There is an increasing complexity moving from the upper atmosphere to the ground: prebiotic molecules in Titan's upper atmosphere evolve to lead to the formation of macromolecules, the subsequent step is the assembly in a haze layer followed by the precipitation and sedimentation of complex organics on the surface, where




Figure 8: Simulated IR spectra of $C_{13}H_9^+$ and $C_{13}H_9^-$. Anharmonic wavenumbers and intensities were computed at the B3LYP/6-31+G(d,p) level of theory and the stick spectra convoluted by means of Lorentzian broadening functions with half-widths at half maximum of 2 cm^{-1} .

the final organic catalysis in the hydrocarbon lakes occurs.

In Ref.¹⁷⁷, reaction mechanisms have been proposed to describe the growth process of the complex macromolecules observed by the mass spectrometers onboard Cassini. Figure 7 shows the initial steps of the formation of macromolecules, focusing on plausible reaction routes from simple aromatic molecules to complex polyaromatic compounds. Once confirmed, the proposed reaction mechanisms will provide the first step toward understanding how aromaticity (also including the formation of heterocycles) was achieved in the atmosphere of primitive Earth. In particular, in Ref.¹⁷⁷, the phenalenyl cation $C_{13}H_9^+$ and its anion $C_{13}H_9^-$ were identified as important intermediates in the process of molecular growth. An important confirmation of the plausibility of the proposed mechanisms would be the astronomical observation of the spectroscopic features of $C_{13}H_9^+$ and $C_{13}H_9^-$, for which however there is a great paucity of spectroscopic information. Being highly symmetric, these molecular species lack a permanent electric dipole moment and their detection relies only on IR spectroscopy. In this respect, the great opportunity offered by JWST should be pointed out. In Figure 8, the fully anharmonic IR spectra of the phenalenyl cation and anion, computed at the B3LYP/6-31+G(d,p) level, are shown. For this level of theory, from the literature on

this topic based on the comparison with experiment and state-of-the-art calculations (see, for example, Refs. ^{53,117,178}), a mean absolute error of 10–12 cm^{-1} and maximum discrepancies of 30 cm^{-1} are expected, which provide a reasonable guidance for astronomical detections, with possible improvements through hybrid schemes as discussed above. However, the phenalenyl cation and anion represent a complex case for VPT2 calculations. First, as symmetric tops, a proper account of degeneracies, ignored in the present discussions, is necessary. Second, the out-of-plane ring deformation mode acts as a LAM, thus contaminating the anharmonic results, and must be adequately decoupled from the rest of the system, in a way similar to the “SkipPT2” scheme seen before. Finally, being large-sized molecules, they also bear a significant number of resonances, in particular the Darling-Dennison ones, which are rather sensitive to the quality of the electronic structure calculation and also need to be properly assessed and corrected.

Concluding remarks

Some of the most challenging aspects in the field of computational chemistry applied to astrochemistry have been reviewed. The molecular complexity in space was first addressed to explain the importance of the support of computational spectroscopy to astrochemical investigations. To fully exploit the possibilities offered by computational spectroscopy in the study of gaseous species, from the micro- to the infrared- wavelengths, a comprehensive summary of the underlying ro-vibrational theory was first given, followed by a description of state-of-the-art protocols for the simulation of small-to-medium sized astroCOMs with the required accuracy. The methodology and its different technical aspects were then illustrated through a test case, oxirane - an example of semi-rigid astroCOM, demonstrating the accuracy of theoretical predictions for rotational and vibrational spectroscopic features. Subsequently, the challenges in the computational spectroscopy of astroCOMs have been faced: by means of selected examples, the most relevant aspects have been introduced, discussed and solved. Finally, we moved to deal with the challenges in the computational chemistry of astroCOMs. While spectroscopy offers the opportunity to study, discover and quantify molecular species in space, computational chemistry allows understanding how astroCOMs could be produced

in the typically harsh conditions of the interstellar medium. The challenges in evaluating feasible formation routes of relevant prebiotic species as well as in understanding the chemical evolution of small species toward macromolecules have been addressed. To summarize, we demonstrated that, even for cases that are challenging for both experiment and theory, the use of the proper combination of state-of-the-art QM approaches and effective anharmonic treatments of rotational and vibrational problems leads to the quantitative prediction of the spectroscopic and thermochemical properties. The chosen examples and their discussion have shown that rotational constants, which are the leading terms in rotational spectroscopy, can be predicted with an accuracy, in relative terms, better than 0.1% and that this accuracy can be further improved (by about one order of magnitude) by means of an empirical scaling procedure. Moving to vibrational signatures, the typical MAE and |MAX| for fundamentals are better than 10 cm^{-1} and 15 cm^{-1} , respectively. Concerning energetics, the importance of accurately determining energy differences of conformers has been well evinced, with the state-of-the-art computational approaches presented here able to provide gas-phase thermodynamic properties with an accuracy better than 1 kcal mol^{-1} .

In the last years, a close collaboration between astronomers and spectroscopic laboratories have allowed the identification of thousands of unknown lines (due, in most cases, to “weeds”). However, in addition to neutral, closed-shell molecules, ionic and radical species are also present in space. They provide an additional challenge in astrochemistry. Their experimental investigation is strongly hampered by the difficulties in their on-the-fly production and their instability, and thus is not always feasible. On the other hand, state-of-the-art QM approaches are able to compute spectroscopic parameters with the same accuracy as demonstrated for closed-shell molecules. Computational spectroscopic characterizations can therefore be used to guide laboratory experiments, and when the latter are not viable, calculations can be further improved by the empirical scaling procedure described above and directly used to support astronomical searches. However, from a theoretical point of view, new developments are required in order to improve the predictive capabilities because of the lack of some implementations. To give an example, to the best of our knowledge, a CCSD(T) implementation for the computation of electron spin-nuclear spin coupling constants, important for the prediction of the rotational spectrum of radicals, is still missing.

Acknowledgements

This work has been supported in Bologna by MIUR ‘PRIN 2015’ funds (project “STARS in the CAOS (Simulation Tools for Astrochemical Reactivity and Spectroscopy in the Cyberinfrastructure for Astrochemical Organic Species)” - Grant Number 2015F59J3R) and by the University of Bologna (RFO funds), and in Shanghai by National Natural Science Foundation of China (Grant No. 91641128). The SMART@SNS Laboratory (<http://dreams.sns.it>) is gratefully acknowledged for the utilization of computer resources. The support of the COST CMTS-Actions CM1405 (MOLIM: MOLEcules In Motion) and CM1401 (Our Astro-Chemical History) is acknowledged. The authors also acknowledge Dr. Lorenzo Spada for fruitful discussions.

References

1. H. S. P. Müller, F. Schlöder, J. Stutzki, and G. Winnewisser, *J. Mol. Struct.* **742**, 215 (2005).
2. C. P. Endres, S. Schlemmer, P. Schilke, J. Stutzki, and H. S. Müller, *J. Mol. Spectrosc.* **327**, 95 (2016), ISSN 0022-2852, new Visions of Spectroscopic Databases, Volume II, URL <http://www.sciencedirect.com/science/article/pii/S0022285216300340>.
3. E. Herbst and J. T. Yates, *Chem. Rev.* **113**, 8707 (2013).
4. W. A. Brown, *Phys. Chem. Chem. Phys.* **16**, 3343 (2014), URL <http://dx.doi.org/10.1039/C4CP90004A>.
5. S. Schlemmer, H. Mutschke, T. Giesen, and C. Jäger, eds., *Laboratory Astrochemistry : From Molecules through Nanoparticles to Grains* (Wiley-VCH Verlag GmbH & Co., 2015), ISBN 9783527408894.
6. D. E. Woon and E. Herbst, *Astrophys. J. Suppl. Ser.* **185**, 273 (2009), URL <http://stacks.iop.org/0067-0049/185/i=2/a=273>.
7. Q. Yu, J. M. Bowman, R. C. Fortenberry, J. S. Mancini, T. J. Lee, T. D. Crawford, W. Klemperer, and J. S. Francisco, *J. Phys. Chem. A* **119**, 11623 (2015).
8. C. W. B. Jr, E. Peeters, and L. J. Allamandola, *Astrophys. J.* **697**, 311 (2009), URL <http://stacks.iop.org/0004-637X/697/i=1/a=311>.
9. R. Peverati, P. P. Bera, T. J. Lee, and M. Head-Gordon, *Astrophys. J.* **830**, 128 (2016), URL <http://stacks.iop.org/0004-637X/830/i=2/a=128>.
10. B. Finney, R. C. Fortenberry, J. S. Francisco, and K. A. Peterson, *J. Chem. Phys.* **145**, 124311 (2016), URL <http://dx.doi.org/10.1063/1.4963337>.
11. R. C. Fortenberry, *Int. J. Quantum Chem.* **117**, 81 (2017), ISSN 1097-461X, URL <http://dx.doi.org/10.1002/qua.25180>.

12. C. Puzzarini, *Int. J. Quantum Chem.* **117**, 129 (2017), ISSN 1097-461X, URL <http://dx.doi.org/10.1002/qua.25284>.
13. A. Candian and C. J. Mackie, *Int. J. Quantum Chem.* **117**, 146 (2017), ISSN 1097-461X, URL <http://dx.doi.org/10.1002/qua.25292>.
14. J. Tennyson and S. N. Yurchenko, *Int. J. Quantum Chem.* **117**, 92 (2017), ISSN 1097-461X, URL <http://dx.doi.org/10.1002/qua.25190>.
15. V. Barone, M. Biczysko, and C. Puzzarini, *Acc. Chem. Res.* **48**, 1413 (2015), URL <http://dx.doi.org/10.1021/ar5003285>.
16. *Access to the entire primos data set, specifics on the observing strategy, and overall frequency coverage information is available.*, (last visited: 5 May 2017), URL <http://www.cv.nrao.edu/~aremijan/PRIMOS/>.
17. J. L. Neill, M. T. Muckle, D. P. Zaleski, A. L. Steber, B. H. Pate, V. Lattanzi, S. Spezzano, M. C. McCarthy, and A. J. Remijan, *Astrophys. J.* **755**, 153 (2012), URL <http://stacks.iop.org/0004-637X/755/i=2/a=153>.
18. B. A. McGuire, P. B. Carroll, R. A. Loomis, I. A. Finneran, P. R. Jewell, A. J. Remijan, and G. A. Blake, *Science* **352**, 1449 (2016), ISSN 0036-8075, URL <http://science.sciencemag.org/content/352/6292/1449>.
19. C. Puzzarini, M. Biczysko, J. Bloino, and V. Barone, *Astrophys. J.* **785**, 107 (2014), URL <http://stacks.iop.org/0004-637X/785/i=2/a=107>.
20. J. E. Elsila, D. P. Glavin, and J. P. Dworkin, *Meteorit. Planet. Sci.* **44**, 1323 (2009), ISSN 1945-5100.
21. M. P. Callahan, K. E. Smith, H. J. Cleaves, J. Ruzicka, J. C. Stern, D. P. Glavin, C. H. House, and J. P. Dworkin, *Proc. Nat. Acad. Sci. USA* **108**, 13995 (2011), URL <http://www.pnas.org/content/108/34/13995.abstract>.
22. V. Barone, M. Biczysko, J. Bloino, and C. Puzzarini, *Phys. Chem. Chem. Phys.* **15**, 10094 (2013), URL <http://dx.doi.org/10.1039/C3CP50439E>.

23. V. Barone, M. Biczysko, J. Bloino, P. Cimino, E. Penocchio, and C. Puzzarini, *J. Chem. Theory Comput.* **11**, 4342 (2015), URL <http://dx.doi.org/10.1021/acs.jctc.5b00580>.
24. V. Barone, M. Biczysko, J. Bloino, and C. Puzzarini, *J. Chem. Phys.* **141**, 034107 (2014), URL <http://scitation.aip.org/content/aip/journal/jcp/141/3/10.1063/1.4887357>.
25. C. Puzzarini, M. Biczysko, V. Barone, L. Largo, I. Peña, C. Cabezas, and J. L. Alonso, *J. Phys. Chem. Lett.* **5**, 534 (2014).
26. F. Vazart, D. Calderini, C. Puzzarini, D. Skouteris, and V. Barone, *J. Chem. Theory Comput.* **12**, 5385 (2016).
27. A. G. G. M. Tielens, *Rev. Mod. Phys.* **85**, 1021 (2013), URL <https://link.aps.org/doi/10.1103/RevModPhys.85.1021>.
28. J. Pesonen and L. Halonen, *Adv. Chem. Phys.* (John Wiley & Sons, Inc., 2003), vol. 125, chap. Recent Advances in the Theory of Vibration-Rotation Hamiltonians, pp. 269–349, ISBN 9780471428022, URL <http://dx.doi.org/10.1002/0471428027.ch4>.
29. A. G. Császár, C. Fábri, T. Szidarovszky, E. Mátyus, T. Furtenbacher, and G. Czakó, *Phys. Chem. Chem. Phys.* **14**, 1085 (2012), URL <http://dx.doi.org/10.1039/C1CP21830A>.
30. E. Mátyus, G. Czakó, and A. G. Császár, *J. Chem. Phys.* **130**, 134112 (2009), URL <http://scitation.aip.org/content/aip/journal/jcp/130/13/10.1063/1.3076742>.
31. J. Tennyson, *Wiley Interdiscip. Rev. Comput. Mol. Sci.* **2**, 698 (2011), URL <http://dx.doi.org/10.1002/wcms.94>.
32. T. Carrington Jr., *J. Chem. Phys.* **146**, 120902 (2017), URL <http://dx.doi.org/10.1063/1.4979117>.
33. V. Szalay, *J. Chem. Phys.* **146**, 124107 (2017), URL <http://dx.doi.org/10.1063/1.4978686>.

34. S. Carter, A. R. Sharma, J. M. Bowman, P. Rosmus, and R. Tarroni, *J. Chem. Phys.* **131**, 224106 (2009), URL <http://link.aip.org/link/?JCP/131/224106/1>.
35. G. Rauhut and T. Hrenar, *Chem. Phys.* **346**, 160 (2008), URL <http://www.sciencedirect.com/science/article/pii/S0301010408000748>.
36. O. Christiansen, *Phys. Chem. Chem. Phys.* **14**, 6672 (2012), URL <http://dx.doi.org/10.1039/C2CP40090A>.
37. T. K. Roy and R. B. Gerber, *Phys. Chem. Chem. Phys.* **15**, 9468 (2013), URL <http://dx.doi.org/10.1039/C3CP50739D>.
38. H. H. Nielsen, *Rev. Mod. Phys.* **23**, 90 (1951), URL <http://link.aps.org/doi/10.1103/RevModPhys.23.90>.
39. G. Amat and H. H. Nielsen, *J. Mol. Spectrosc.* **2**, 152 (1958), ISSN 0022-2852, URL <http://www.sciencedirect.com/science/article/pii/0022285258900705>.
40. I. M. Mills, in *Molecular Spectroscopy: Modern Research*, edited by K. N. Rao and C. W. Mathews (Academic Press, New York, 1972), chap. 3.2, pp. 115–140, URL <http://www.sciencedirect.com/science/article/pii/B9780125806404500133>.
41. I. M. Mills, in *Theoretical Chemistry: Volume 1*, edited by R. N. Dixon (The Royal Society of Chemistry, 1974), vol. 1, pp. 110–159, ISBN 978-0-85186-754-0, URL <http://dx.doi.org/10.1039/9781847557124-00110>.
42. M. R. Aliev and J. K. G. Watson, in *Molecular Spectroscopy: Modern Research*, edited by K. N. Rao (Academic Press, 1985), vol. 3, pp. 1–67, ISBN 978-0-12-580643-5, URL <http://www.sciencedirect.com/science/article/pii/B9780125806435500063>.
43. D. A. Clabo Jr., W. D. Allen, R. B. Remington, Y. Yamaguchi, and H. F. Schaefer III, *Chem. Phys.* **123**, 187 (1988), URL <http://www.sciencedirect.com/science/article/pii/0301010488872719>.
44. W. D. Allen, Y. Yamaguchi, A. G. Császár, D. A. Clabo Jr., R. B. Remington, and H. F. Schaefer III, *Chem. Phys.* **145**, 427 (1990).

45. F. Gaw, A. Willetts, N. Handy, and W. Green, *Advances in Molecular Vibrations and Collision Dynamics* (JAI Press, 1992), vol. 1, chap. SPECTRO - a program for derivation of spectroscopic constants from provided quartic force fields and cubic dipole fields.
46. J. Vázquez and J. F. Stanton, *Mol. Phys.* **104**, 377 (2006).
47. V. Barone, *J. Chem. Phys.* **122**, 014108 (2005), URL <http://link.aip.org/link/?JCP/122/014108/1>.
48. S. V. Krasnoshchekov, E. V. Isayeva, and N. F. Stepanov, *J. Phys. Chem. A* **116**, 3691 (2012), <http://pubs.acs.org/doi/pdf/10.1021/jp211400w>, URL <http://pubs.acs.org/doi/abs/10.1021/jp211400w>.
49. A. M. Rosnik and W. F. Polik, *Mol. Phys.* **112**, 261 (2014), <http://dx.doi.org/10.1080/00268976.2013.808386>, URL <http://dx.doi.org/10.1080/00268976.2013.808386>.
50. M. Ringholm, D. Jonsson, R. Bast, B. Gao, A. J. Thorvaldsen, U. Ekström, T. Helgaker, and K. Ruud, *J. Chem. Phys.* **140**, 034103 (2014), URL <http://scitation.aip.org/content/aip/journal/jcp/140/3/10.1063/1.4861003>.
51. J. Bloino, M. Biczysko, and V. Barone, *J. Chem. Theory Comput.* **8**, 1015 (2012), <http://pubs.acs.org/doi/pdf/10.1021/ct200814m>, URL <http://pubs.acs.org/doi/abs/10.1021/ct200814m>.
52. J. Bloino and V. Barone, *J. Chem. Phys.* **136**, 124108 (2012), URL <http://link.aip.org/link/?JCP/136/124108/1>.
53. V. Barone, M. Biczysko, and J. Bloino, *Phys. Chem. Chem. Phys.* **16**, 1759 (2014), URL <http://dx.doi.org/10.1039/C3CP53413H>.
54. M. Piccardo, J. Bloino, and V. Barone, *Int. J. Quantum Chem.* **115**, 948 (2015), ISSN 1097-461X, URL <http://dx.doi.org/10.1002/qua.24931>.

55. J. Bloino, J. Phys. Chem. A **119**, 5269 (2015), <http://dx.doi.org/10.1021/jp509985u>, URL <http://dx.doi.org/10.1021/jp509985u>.
56. J. Bloino, M. Biczysko, and V. Barone, J. Phys. Chem. A **119**, 11862 (2015), <http://dx.doi.org/10.1021/acs.jpca.5b10067>, URL <http://dx.doi.org/10.1021/acs.jpca.5b10067>.
57. J. F. Stanton, J. Gauss, M. E. Harding, and P. G. Szalay, CFour a quantum chemical program package (2011), with contributions from A. A. Auer, R. J. Bartlett, U. Benedikt, C. Berger, D. E. Bernholdt, Y. J. Bomble, O. Christiansen, M. Heckert, O. Heun, C. Huber, T.-C. Jagau, D. Jonsson, J. Jusélius, K. Klein, W. J. Lauderdale, D. Matthews, T. Metzroth, L. A. Mueck, D. P. O'Neill, D. R. Price, E. Prochnow, C. Puzzarini, K. Ruud, F. Schiffmann, W. Schwalbach, S. Stopkiewicz, A. Tajti, J. Vázquez, F. Wang, J. D. Watts and the integral packages MOLECULE (J. Almloef and P. R. Taylor), PROPS (P. R. Taylor), ABACUS (T. Helgaker, H. J. Aa. Jensen, P. Jørgensen, and J. Olsen), and ECP routines by A. V. Mitin and C. van Wüllen. For the current version, see <http://www.cfour.de>.
58. DALTON, a molecular electronic structure program, Release DALTON2013 (2014), see <http://daltonprogram.org>.
59. M. W. Schmidt, K. K. Baldrige, J. A. Boatz, S. T. Elbert, M. S. Gordon, J. H. Jensen, S. Koseki, N. Matsunaga, K. A. Nguyen, S. Su, et al., J. Comput. Chem. **14**, 1347 (1993), ISSN 1096-987X, URL <http://dx.doi.org/10.1002/jcc.540141112>.
60. M. S. Gordon and M. W. Schmidt, *Theory and Applications of Computational Chemistry: the first forty years* (Elsevier, 2005), chap. Advances in electronic structure theory: GAMESS a decade later, pp. 1167–1189.
61. M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, et al., *Gaussian 16 Revision A.03* (2016), gaussian Inc. Wallingford CT.

62. H.-J. Werner, P. J. Knowles, G. Knizia, F. R. Manby, M. Schütz, P. Celani, T. Korona, R. Lindh, A. Mitrushenkov, G. Rauhut, et al., *Molpro, version 2012.1, a package of ab initio programs* (2012), see <http://www.molpro.net>.
63. M. Valiev, E. J. Bylaska, N. Govind, K. Kowalski, T. P. Straatsma, H. J. J. Van Dam, D. Wang, J. Nieplocha, E. Apra, T. L. Windus, et al., *Comput. Phys. Commun.* **181**, 1477 (2010), ISSN 0010-4655, URL <http://www.sciencedirect.com/science/article/pii/S0010465510001438>.
64. Y. Shao, L. F. Molnar, Y. Jung, J. Kussmann, C. Ochsenfeld, S. T. Brown, A. T. Gilbert, L. V. Slipchenko, S. V. Levchenko, D. P. O'Neill, et al., *Phys. Chem. Chem. Phys.* **8**, 3172 (2006), URL <http://dx.doi.org/10.1039/B517914A>.
65. P. Carbonnière, A. Dargelos, and C. Pouchan, *Theor. Chem. Acc.* **125**, 543 (2010), ISSN 1432-881X, URL <http://dx.doi.org/10.1007/s00214-009-0689-7>.
66. J. H. Van Vleck, *Phys. Rev.* **33**, 467 (1929), URL <http://link.aps.org/doi/10.1103/PhysRev.33.467>.
67. J. K. G. Watson, *J. Chem. Phys.* **48**, 4517 (1968), URL <http://scitation.aip.org/content/aip/journal/jcp/48/10/10.1063/1.1668020>.
68. J. K. G. Watson, in *Vibrational spectra and structure, vol.6. A series of advances*, edited by J. Durig (Elsevier, Amsterdam, Netherlands, 1977), pp. 1–89, ISBN 0-444-41588-2.
69. A. Willets, N. C. Handy, W. H. Green Jr., and D. Jayatilaka, *J. Phys. Chem.* **94**, 5608 (1990), URL <http://pubs.acs.org/doi/abs/10.1021/j100377a038>.
70. J. Vázquez and J. F. Stanton, *Mol. Phys.* **105**, 101 (2007).
71. I. M. Konen, I. B. Pollack, E. X. J. Li, M. I. Lester, M. E. Varner, and J. F. Stanton, *J. Chem. Phys.* **122**, 094320 (2005), URL <http://scitation.aip.org/content/aip/journal/jcp/122/9/10.1063/1.1854094>.
72. R. D. Amos, N. C. Handy, W. H. Green, D. Jayatilaka, A. Willets, and P. Palmieri, *J. Chem. Phys.* **95**, 8323 (1991), URL <http://link.aip.org/link/?JCP/95/8323/1>.

73. P. Bouř, J. Phys. Chem. **98**, 8862 (1994), URL <http://pubs.acs.org/doi/abs/10.1021/j100087a008>.
74. J. M. L. Martin, T. J. Lee, P. M. Taylor, and J.-P. François, J. Chem. Phys. **103**, 2589 (1995).
75. K. M. Kuhler, D. G. Truhlar, and A. D. Isaacson, J. Chem. Phys. **104**, 4664 (1996), URL <http://dx.doi.org/10.1063/1.471161>.
76. J. M. L. Martin and P. M. Taylor, Spectrochimica Acta Part A: Molecular Spectroscopy **53**, 1039 (1997).
77. A. Miani, E. Cané, P. Palmieri, A. Trombetti, and N. C. Handy, J. Chem. Phys. **112**, 248 (2000), URL <http://scitation.aip.org/content/aip/journal/jcp/112/1/10.1063/1.480577>.
78. V. Hänninen and L. Halonen, Mol. Phys. **101**, 2907 (2003), <http://dx.doi.org/10.1080/00268970310001598740>, URL <http://dx.doi.org/10.1080/00268970310001598740>.
79. D. Begue, P. Carbonniere, and C. Pouchan, J. Phys. Chem. A **109**, 4611 (2005), <http://pubs.acs.org/doi/pdf/10.1021/jp0406114>, URL <http://pubs.acs.org/doi/abs/10.1021/jp0406114>.
80. E. Cané, A. Miani, and A. Trombetti, J. Phys. Chem. A **111**, 8218 (2007), <http://dx.doi.org/10.1021/jp071610p>, URL <http://dx.doi.org/10.1021/jp071610p>.
81. D. A. Matthews and J. F. Stanton, Mol. Phys. **107**, 213 (2009), <http://www.tandfonline.com/doi/pdf/10.1080/00268970902769463>, URL <http://www.tandfonline.com/doi/abs/10.1080/00268970902769463>.
82. M. R. Hermes and S. Hirata, J. Chem. Phys. **139**, 034111 (2013), URL <http://scitation.aip.org/content/aip/journal/jcp/139/3/10.1063/1.4813123>.

83. S. V. Krasnoshchekov, E. V. Isayeva, and N. F. Stepanov, *J. Chem. Phys.* **141**, 234114 (2014), URL <http://scitation.aip.org/content/aip/journal/jcp/141/23/10.1063/1.4903927>.
84. B. T. Darling and D. M. Dennison, *Phys. Rev.* **57**, 128 (1940), URL <http://link.aps.org/doi/10.1103/PhysRev.57.128>.
85. K. K. Lehmann, *Mol. Phys.* **66**, 1129 (1989), URL <http://dx.doi.org/10.1080/00268978900100751>,
<http://dx.doi.org/10.1080/00268978900100751>.
86. D. A. Matthews, J. Vázquez, and J. F. Stanton, *Mol. Phys.* **105**, 2659 (2007), <http://dx.doi.org/10.1080/00268970701618424>, URL <http://dx.doi.org/10.1080/00268970701618424>.
87. J. Bloino, A. Baiardi, and M. Biczysko, *Int. J. Quantum Chem.* **116**, 1543 (2016), ISSN 1097-461X, URL <http://dx.doi.org/10.1002/qua.25188>.
88. A. G. Császár, W. D. Allen, and H. S. III, *J. Chem. Phys.* **108**, 9751 (1998).
89. J. A. Montgomery Jr., M. J. Frisch, J. W. Ochterski, and G. A. Petersson, *J. Chem. Phys.* **110**, 2822 (1999).
90. A. Tajti, P. G. Szalay, A. G. Császár, M. Kállay, J. Gauss, E. F. Valeev, B. A. Flowers, J. Vázquez, and J. F. Stanton, *J. Chem. Phys.* **121**, 11599 (2004).
91. M. Heckert, M. Kállay, and J. Gauss, *Mol. Phys.* **103**, 2109 (2005).
92. M. Heckert, M. Kállay, D. P. Tew, W. Klopper, and J. Gauss, *J. Chem. Phys.* **125**, 044108 (pages 10) (2006), URL <http://link.aip.org/link/?JCP/125/044108/1>.
93. C. Puzzarini, J. Heckert, and J. Gauss, *J. Chem. Phys.* **128**, 194108 (2008).
94. K. Raghavachari, G. W. Trucks, J. A. Pople, and M. Head-Gordon, *Chem. Phys. Lett.* **157**, 479 (1989), ISSN 0009-2614, URL <http://www.sciencedirect.com/science/article/pii/S0009261489873956>.

95. J. Noga and R. J. Bartlett, *J. Chem. Phys.* **86**, 7041 (1987).
96. G. E. Scuseria and H. F. Schaefer III, *Chem. Phys. Lett.* **152**, 382 (1988).
97. J. D. Watts and R. J. Bartlett, *J. Chem. Phys.* **93**, 6104 (1993).
98. K. M. and P. R. Surján, *J. Chem. Phys.* **115**, 2945 (2001).
99. T. H. Dunning Jr., *J. Chem. Phys.* **90**, 1007 (1989), URL <http://scitation.aip.org/content/aip/journal/jcp/90/2/10.1063/1.456153>.
100. A. K. Wilson, T. van Mourik, and J. Dunning, T. H., *J. Mol. Struct. Theochem* **388**, 339 (1996).
101. D. E. Woon and T. H. Dunning Jr., *J. Chem. Phys.* **103**, 4572 (1995), URL <http://link.aip.org/link/?JCP/103/4572/1>.
102. K. A. Peterson and T. H. Dunning Jr., *J. Chem. Phys.* **117**, 10548 (2002).
103. C. Puzzarini, *Phys. Chem. Chem. Phys.* **15**, 6595 (2013), URL <http://dx.doi.org/10.1039/C3CP44301A>.
104. S. Thorwirth, L. A. Mück, J. Gauss, F. Tamassia, V. Lattanzi, and M. C. McCarthy, *J. Phys. Chem. Lett.* **2**, 1228 (2011).
105. V. Barone, M. Biczysko, J. Bloino, and C. Puzzarini, *Phys. Chem. Chem. Phys.* **15**, 1358 (2013), URL <http://dx.doi.org/10.1039/C2CP43884D>.
106. C. Puzzarini, J. F. Stanton, and J. Gauss, *Int. Rev. Phys. Chem.* **29**, 273 (2010), <http://dx.doi.org/10.1080/01442351003643401>, URL <http://dx.doi.org/10.1080/01442351003643401>.
107. J. M. Martin and P. R. Taylor, *Chem. Phys. Lett.* **248**, 336 (1996), ISSN 0009-2614, URL <http://www.sciencedirect.com/science/article/pii/0009261495013539>.
108. J. Demaison, L. Margulès, and J. E. Boggs, *Struct. Chem.* **14**, 159 (2003), ISSN 1572-9001, URL <http://dx.doi.org/10.1023/A:1022138431967>.

109. C. Puzzarini, J. Phys. Chem. A **113**, 14530 (2009), URL <http://pubs.acs.org/doi/abs/10.1021/jp9040327>.
110. C. Puzzarini and V. Barone, Phys. Chem. Chem. Phys. **13**, 7189 (2011), URL <http://dx.doi.org/10.1039/C0CP02636K>.
111. C. Møller and M. S. Plesset, Phys. Rev. **46**, 618 (1934), URL <http://link.aps.org/doi/10.1103/PhysRev.46.618>.
112. C. Puzzarini, M. Biczysko, V. Barone, I. Pena, C. Cabezas, and J. L. Alonso, Phys. Chem. Chem. Phys. **15**, 16965 (2013), URL <http://dx.doi.org/10.1039/C3CP52347K>.
113. T. Helgaker, W. Klopper, H. Koch, and J. Noga, J. Chem. Phys. **106**, 9639 (1997), URL <http://link.aip.org/link/?JCP/106/9639/1>.
114. A. Pietropolli Charmet, P. Stoppa, S. Giorgianni, J. Bloino, N. Tasinato, I. Carnimeo, M. Biczysko, and C. Puzzarini, J. Phys. Chem. A **121**, 3305 (2017), pMID: 28398055, URL <http://dx.doi.org/10.1021/acs.jpca.7b02060>.
115. C. Puzzarini, G. Cazzoli, and J. Gauss, J. Chem. Phys. **137**, 154311 (2012), URL <http://scitation.aip.org/content/aip/journal/jcp/137/15/10.1063/1.4758316>.
116. C. Puzzarini, M. Biczysko, and V. Barone, J. Chem. Theory Comput. **6**, 828 (2010), URL <http://pubs.acs.org/doi/abs/10.1021/ct900594h>.
117. C. Puzzarini, M. Biczysko, and V. Barone, J. Chem. Theory Comput. **7**, 3702 (2011), <http://pubs.acs.org/doi/pdf/10.1021/ct200552m>, URL <http://pubs.acs.org/doi/abs/10.1021/ct200552m>.
118. C. Puzzarini, E. Penocchio, M. Biczysko, and V. Barone, J. Phys. Chem. A **118**, 6648 (2014), URL <http://pubs.acs.org/doi/abs/10.1021/jp503672g>.
119. K. Burke, J. Chem. Phys. **136**, 150901 (2012), URL <http://scitation.aip.org/content/aip/journal/jcp/136/15/10.1063/1.4704546>.

120. A. D. Becke, J. Chem. Phys. **98**, 5648 (1993), URL <http://link.aip.org/link/?JCP/98/5648/1>.
121. S. Grimme, J. Chem. Phys. **124**, 034108 (2006).
122. P. Pulay, W. Meyer, and J. E. Boggs, J. Chem. Phys. **68**, 5077 (1978), URL <http://scitation.aip.org/content/aip/journal/jcp/68/11/10.1063/1.435626>.
123. F. Pawłowski, P. Jørgensen, J. Olsen, F. Hegelund, T. Helgaker, J. Gauss, K. L. Bak, and J. F. Stanton, J. Chem. Phys. **116**, 6482 (2002), URL <http://link.aip.org/link/?JCP/116/6482/1>.
124. J. Demaison, J. E. Boggs, and A. G. Császár, *Equilibrium Molecular Structures: From Spectroscopy to Quantum Chemistry* (CRC Press, 2010), ISBN 9781439811351, URL <https://books.google.it/books?id=7DZB0ITw0BsC>.
125. M. Piccardo, E. Penocchio, C. Puzzarini, M. Biczysko, and V. Barone, J. Phys. Chem. A **119**, 2058 (2015).
126. E. Penocchio, M. Piccardo, and V. Barone, J. Chem. Theory Comput. **11**, 4689 (2015).
127. M. Biczysko, J. Bloino, G. Brancato, I. Cacelli, C. Cappelli, A. Ferretti, A. Lami, S. Monti, A. Pedone, G. Prampolini, et al., Theor. Chem. Acc. **131**, 1 (2012), ISSN 1432-881X, URL <http://dx.doi.org/10.1007/s00214-012-1201-3>.
128. V. Barone, M. Biczysko, J. Bloino, and C. Puzzarini, J. Chem. Theory Comput. **9**, 1533 (2013), <http://pubs.acs.org/doi/pdf/10.1021/ct3010672>, URL <http://pubs.acs.org/doi/abs/10.1021/ct3010672>.
129. T. Fornaro, M. Biczysko, S. Monti, and V. Barone, Phys. Chem. Chem. Phys. **16**, 10112 (2014).
130. M. Biczysko, P. Panek, G. Scalmani, J. Bloino, and V. Barone, J. Chem. Theory Comput. **6**, 2115 (2010), <http://pubs.acs.org/doi/pdf/10.1021/ct100212p>, URL <http://pubs.acs.org/doi/abs/10.1021/ct100212p>.

131. I. Carnimeo, C. Puzzarini, N. Tasinato, P. Stoppa, A. P. Charmet, M. Biczysko, C. Cappelli, and V. Barone, *J. Chem. Phys.* **139**, 074310 (2013), URL <http://link.aip.org/link/?JCP/139/074310/1>.
132. J. E. Dickens, W. M. Irvine, M. Ohishi, M. Ikeda, S. Ishikawa, A. Nummelin, and . Hjalmarson, *Astrophys. J.* **489**, 753 (1997), URL <http://stacks.iop.org/0004-637X/489/i=2/a=753>.
133. J.-M. Bernard, P. Coll, A. Coustenis, and F. Raulin, *Planet. Space Sci.* **51**, 1003 (2003), ISSN 0032-0633, surfaces and Atmospheres of the Outer Planets their Satellites and Ring Systems, URL <http://www.sciencedirect.com/science/article/pii/S0032063303001314>.
134. P. Coll, J.-M. Bernard, R. Navarro-Gonzalez, and F. Raulin, *Astrophys. J.* **598**, 700 (2003), URL <http://stacks.iop.org/0004-637X/598/i=1/a=700>.
135. C. J. Bennet, Y. Osamura, M. D. Lebar, and R. I. Kaiser, *Astrophys. J.* **634**, 711 (2005).
136. L. S. Bernstein and D. K. Lynch, *Astrophys. J.* **704**, 226 (2009), URL <http://stacks.iop.org/0004-637X/704/i=1/a=226>.
137. C. Puzzarini, A. Ali, M. Biczysko, and V. Barone, *Astrophys. J.* **792**, 118 (2014), URL <http://stacks.iop.org/0004-637X/792/i=2/a=118>.
138. A. Mesko, L. Zou, P. B. Carroll, and S. L. W. Weaver, *J. Mol. Spectrosc.* **335**, 49 (2017), ISSN 0022-2852, spectroscopy and Inter/Intramolecular Dynamics in Honor of Walther Caminati - Part 1, URL <http://www.sciencedirect.com/science/article/pii/S0022285216303563>.
139. C. Medcraft, C. D. Thompson, E. G. Robertson, D. R. T. Ap-padoo, and D. McNaughton, *Astrophys. J.* **753**, 18 (2012), URL <http://stacks.iop.org/0004-637X/753/i=1/a=18>.

140. T. Nakanaga, J. Chem. Phys. **74**, 5384 (1981), URL <http://scitation.aip.org/content/aip/journal/jcp/74/10/10.1063/1.440967>.
141. F. K. Tchana, M. Ngom, A. Perrin, J.-M. Flaud, W. Lafferty, S. Ndiaye, and E. Ngom, J. Mol. Spectrosc. **292**, 1 (2013), ISSN 0022-2852, URL <http://www.sciencedirect.com/science/article/pii/S0022285213001136>.
142. F. K. Tchana, J.-M. Flaud, W. Lafferty, and M. Ngom, Mol. Phys. **112**, 1633 (2014).
143. W. J. Lafferty, J. M. Flaud, F. Kwabia Tchana, and J. M. Fernandez, Mol. Phys. **111**, 1983 (2013), URL <http://www.tandfonline.com/doi/abs/10.1080/00268976.2013.775516>.
144. J.-M. Flaud, W. Lafferty, F. K. Tchana, A. Perrin, and X. Landsheere, J. Mol. Spectrosc. **271**, 38 (2012), ISSN 0022-2852, URL <http://www.sciencedirect.com/science/article/pii/S0022285211002840>.
145. D. K. Russell and R. Wesendrup, J. Mol. Spectrosc. **217**, 59 (2003), ISSN 0022-2852, URL <http://www.sciencedirect.com/science/article/pii/S0022285202000103>.
146. G. C. Sloan, K. E. Kraemer, S. D. Price, and R. F. Shipman, Astrophys. J. Suppl. Ser. **147**, 379 (2003), URL <http://stacks.iop.org/0067-0049/147/i=2/a=379>.
147. J. Bernard Salas, S. R. Pottasch, D. A. Beintema, and P. R. Wesselius, Astron. Astrophys. **367**, 949 (2001), URL <https://doi.org/10.1051/0004-6361:20000435>.
148. S. Kwok and Y. Zhang, Nature **479**, 80 (2011), ISSN 0028-0836, URL <http://dx.doi.org/10.1038/nature10542>.
149. B. Tercero, I. Kleiner, J. Cernicharo, H. V. L. Nguyen, A. Lopez, and G. M. M. Caro, Astrophys. J. Lett. **770**, L13 (2013), URL <http://stacks.iop.org/2041-8205/770/i=1/a=L13>.
150. A. Bellili, R. Linguerrì, M. Hochlaf, and C. Puzzarini, J. Chem. Phys. **143**, 184314 (2015), URL <http://dx.doi.org/10.1063/1.4935493>.

151. A. Kraśnicki, L. Pszczółkowski, and Z. Kisiel, J. Mol. Spectrosc. **260**, 57 (2010), ISSN 0022-2852, URL <http://www.sciencedirect.com/science/article/pii/S0022285209002628>.
152. A. O. Mitrushchenkov, J. Chem. Phys. **136**, 024108 (2012), URL <http://scitation.aip.org/content/aip/journal/jcp/136/2/10.1063/1.3672162>.
153. A. Baiardi, J. Bloino, and V. Barone, J. Chem. Theory Comput. **13**, DOI:10.1021/acs.jctc.7b00236 (2017).
154. W. H. Miller, N. C. Handy, and J. E. Adams, J. Chem. Phys. **72**, 99 (1980), URL <http://scitation.aip.org/content/aip/journal/jcp/72/1/10.1063/1.438959>.
155. T. Carrington and W. H. Miller, J. Chem. Phys. **81**, 3942 (1984), URL <http://scitation.aip.org/content/aip/journal/jcp/81/9/10.1063/1.448187>.
156. D. P. Tew, N. C. Handy, and S. Carter, J. Chem. Phys. **125**, 084313 (2006), URL <http://scitation.aip.org/content/aip/journal/jcp/125/8/10.1063/1.2338891>.
157. M. L. Senent, C. Puzzarini, R. Dominguez-Gomez, M. Carvajal, and M. Hochlaf, J. Chem. Phys. **140**, 124302 (2014).
158. L. Kolesniková, B. Tercero, J. Cernicharo, J. L. Alonso, A. M. Daly, B. P. Gordon, and S. T. Shipman, Astrophys. J. Lett. **784**, L7 (2014), URL <http://stacks.iop.org/2041-8205/784/i=1/a=L7>.
159. *Glycine, aminoacetic acid, conformer ii*, (last visited: 5 May 2017), URL <https://www.astro.uni-koeln.de/cgi-bin/cdmsinfo?file=e075512.cat>.
160. I. Jimenez-Serra, L. Testi, P. Caselli, and S. Viti, Astrophys. J. Lett. p. arXiv:1404.5849 (2014).
161. R. T. Garrod, Astrophys. J. **765**, 60 (2013), URL <http://stacks.iop.org/0004-637X/765/i=1/a=60>.
162. P. Y. Ayala and H. B. Schlegel, J. Chem. Phys. **108**, 2314 (1998), URL <http://scitation.aip.org/content/aip/journal/jcp/108/6/10.1063/1.475616>.

163. J. Zheng, T. Yu, E. Papajak, I. M. Alecu, S. L. Mielke, and D. G. Truhlar, *Phys. Chem. Chem. Phys.* **13**, 10885 (2011), URL <http://dx.doi.org/10.1039/C0CP02644A>.
164. J. Zheng and D. G. Truhlar, *J. Chem. Theory Comput.* **9**, 2875 (2013), URL <http://pubs.acs.org/doi/abs/10.1021/ct400231q>.
165. D. Skouteris, D. Calderini, and V. Barone, *J. Chem. Theory Comput.* **12**, 1011 (2016), PMID: 26765363.
166. D. G. Truhlar and A. D. Isaacson, *J. Chem. Phys.* **94**, 357 (1991), URL <http://link.aip.org/link/doi/10.1063/1.460350>.
167. S. G. Stepanian, I. D. Reva, E. D. Radchenko, M. T. S. Rosado, M. L. T. S. Duarte, R. Fausto, and L. Adamowicz, *J. Phys. Chem. A* **102**, 1041 (1998), URL <http://pubs.acs.org/doi/abs/10.1021/jp973397a>.
168. C. Chyba and C. Sagan, *Nature* **355**, 125 (1992).
169. S. L. Miller, *Science* **117**, 528 (1953).
170. S. L. Miller and H. C. Urey, *Science* **130**, 245 (1959).
171. F. Raulin, C. McKay, J. Lunine, and T. Owen, *Titan's Astrobiology* (Springer Netherlands, Dordrecht, 2010), pp. 215–233, ISBN 978-1-4020-9215-2.
172. V. Barone, C. Latouche, D. Skouteris, F. Vazart, N. Balucani, C. Ceccarelli, and B. Lefloch, *Mon. Not. R. Astron. Soc. Lett.* **453**, L31 (2015), URL <http://dx.doi.org/10.1093/mnrasl/slv094>.
173. D. Skouteris, F. Vazart, C. Ceccarelli, N. Balucani, C. Puzzarini, and V. Barone, *Mon. Not. R. Astron. Soc. Lett.* **468**, L1 (2017), URL <http://dx.doi.org/10.1093/mnrasl/slx012>.
174. A. Bacmann, V. Taquet, A. Faure, C. Kahane, and C. Ceccarelli, *Astron. Astrophys.* **541**, L12 (2012), URL <https://doi.org/10.1051/0004-6361/201219207>.

175. R. Saladino, G. Botta, S. Pino, G. Costanzo, and E. Di Mauro, *Chem. Soc. Rev.* **41**, 5526 (2012), URL <http://dx.doi.org/10.1039/C2CS35066A>.
176. S. Grimme, J. Antony, S. Ehrlich, and H. Krieg, *J. Chem. Phys.* **132**, 154104 (2010), URL <http://scitation.aip.org/content/aip/journal/jcp/132/15/10.1063/1.3382344>.
177. A. Ali, E. S. Jr., D. Chornay, B. Rowe, and C. Puzzarini, *Planet. Space Sci.* **109–110**, 46 (2015), ISSN 0032-0633, URL <http://www.sciencedirect.com/science/article/pii/S0032063315000161>.
178. G. N. Ten, V. V. Nechaev, and S. V. Krasnoshchekov, *Opt. Spectrosc.* **108**, 37 (2010), ISSN 1562-6911, URL <http://dx.doi.org/10.1134/S0030400X10010078>.

FURTHER READING

Astrochemistry

1. S. Yamamoto. “Introduction to Astrochemistry Chemical Evolution from Interstellar Clouds to Star and Planet Formation”, Springer (2017).
2. D. A. Williams, S. Viti. “Observational Molecular Astronomy Exploring the Universe Using Molecular Line Emissions”, Cambridge University Press (2013).
3. S. Kwok. “Organic Matter in the Universe”, Wiley (2012).
4. S. Schlemmer, T. Giesen, H. Mutschke (Eds). “Laboratory Astrochemistry: From Molecules through Nanoparticles to Grains”, Wiley (2011).
5. J. Tennyson. “Astronomical Spectroscopy. An Introduction to the Atomic and Molecular Physics of Astronomical Spectra”, 2nd Edition, World Scientific (2011).
6. “The Astrochymist” website: <http://www.astrochymist.org/>

Quantum Mechanics

1. T. Helgaker, P. Jørgensen, J. Olsen. “Electronic-Structure Theory”, Wiley (2000).
2. A. Szabo, N. S. Ostlund, “Modern Quantum Chemistry”, MacMillan Publishing Co., Inc. (1982).
3. I. Shavitt, R. J. Bartlett, “Many-Body Methods in Chemistry and Physics”, Cambridge University Press (2009)

Spectroscopy

1. M. Quack, F. Merkt. “Handbook of High-resolution Spectroscopy”, Wiley (2011).
2. V. Barone (Ed). “Computational Strategies for Spectroscopy, from Small Molecules to Nano Systems”, John Wiley & Sons (2011).
3. S. Califano, “Vibrational States”, John Wiley & Sons (1976)
4. G. Herzberg, “Molecular Spectra and Molecular Structure: I. Spectra of Diatomic Molecules”, D. Van Nostrand Company (1950)
5. E. B. Wilson Jr., J. C. Decius, P. C. Cross, “Molecular Vibrations”, McGraw-Hill Book Company, Inc. (1955)