



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Identification of QTLs linked to fruit quality traits in apricot (*Prunus armeniaca* L.) and biological validation through gene expression analysis using qPCR

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

García-Gómez, B.E., Salazar, J.A., Dondini, L., Martínez-Gómez, P., Ruiz, D. (2019). Identification of QTLs linked to fruit quality traits in apricot (*Prunus armeniaca* L.) and biological validation through gene expression analysis using qPCR. *MOLECULAR BREEDING*, 39(2), 1-19 [10.1007/s11032-018-0926-7].

Availability:

This version is available at: <https://hdl.handle.net/11585/669312> since: 2019-02-25

Published:

DOI: <http://doi.org/10.1007/s11032-018-0926-7>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

Identification of QTLs linked to fruit quality traits in apricot (*Prunus armeniaca* L.) and biological validation through gene expression analysis using qPCR

Beatriz Ester García-Gómez · Juan Alfonso Salazar · Luca Dondini · Pedro Martínez-Gómez · David Ruiz

Abstract Nine important fruit quality traits—including fruit weight, stone weight, fruit diameter, skin ground colour, flesh colour, blush colour, firmness, soluble solids content and acidity content—were studied for two consecutive years in two F₁ apricot progeny derived from the crosses ‘Bergeron’ × ‘Currot’ (B×C) and ‘Goldrich’ × ‘Currot’ (G×C). Results showed great segregation variability between populations, which was expected because of the polygenic nature and quantitative inheritance of all the studied traits. In addition, some correlations were observed among the fruit quality traits studied. QTL (quantitative trait loci) analysis was carried out using the phenotypic data and genetic linkages maps of ‘B×C’ and ‘G×C’ obtained with SSR and SNP markers. The most significant QTLs were localised in LG4 for soluble solids content and in LG3 for skin and flesh colour. In LG4, we can highlight the presence of candidate genes involved in D-glucose and D-mannose

binding, while in LG3, we identified *MYB* genes previously linked to skin colour by other authors. In order to clearly identify the candidate genes responsible for the analysed traits, we converted the QTLs into expression QTLs and analysed the abundance of transcripts in the segregating genotypes ‘GC 2–11’ and ‘GC 3–7’ from the G×C population. Using qPCR, we analysed the gene expression of nine candidate genes associated with the QTLs identified, including transcription factors (*MYB 10*), carotenoid biosynthesis genes (*LOX 2*, *CCD1* and *CCD4*), anthocyanin biosynthesis genes (*ANS*, *UFGT* and *F3’5’H*), organic acid biosynthesis genes (*NAD ME*) and ripening date genes (*NAC*). Results showed variable expression patterns throughout fruit development and between contrasted genotypes, with a correlation between validated genes and linked QTLs. The *MYB10* gene was the best candidate gene for skin colour. In addition, we found that monitoring *NAC* expression is a good RNA marker for evaluating ripening progression.

Beatriz Ester García-Gómez and Juan Alfonso Salazar contributed equally to this work.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s11032-018-0926-7>) contains supplementary material, which is available to authorized users.

B. E. García-Gómez · J. A. Salazar · P. Martínez-Gómez (✉) · D. Ruiz
Departamento de Mejora Vegetal, CEBAS-CSIC, PO Box 164,
E-30100 Espinardo, Murcia, Spain
e-mail: pmartinez@cebas.csic.es

L. Dondini (✉)
Dipartimento di Scienze Agrarie, Università degli Studi di
Bologna, Via Fanin 46, 40127 Bologna, Italy
e-mail: luca.dondini@unibo.it

Keywords Apricot · *Prunus armeniaca* · Fruit quality · Breeding QTL · Candidate gene · qPCR

Introduction

Apricot (*Prunus armeniaca* L.) is one of the most important and desirable of the temperate tree fruits, with a total world production of around 3.88 million tonnes in 2016 (<http://www.fao.org/faostat/en/#data>). It is the third stone fruit in terms of worldwide economic importance

after peach and plum. Fruit quality is a key factor for apricot consumption, and the priorities of apricot breeding programmes are therefore good fruit taste, improved attractiveness and good postharvest behaviour among new varieties (Infante et al. 2011). These features are required for competitiveness in the globalised markets. In addition, demand is increasing for high levels of carotenoids and polyphenols in new varieties, since these compounds have a positive effect on health due to their antioxidant and defensive activity against a variety of degenerative diseases (Hertog et al. 1993; Machlin 1995; Van den Berg et al. 2000; Bazzano et al. 2002).

Fruit quality is a complex human concept including sensory properties (appearance, texture, taste and aroma), nutritional values, chemical compounds, mechanical properties and functional properties (Infante et al. 2008). A high number of pomological traits influence the fruit quality in apricot. In addition, most apricot fruit quality traits—such as soluble solids content, acidity, skin colour, blush colour and firmness—are polygenic and quantitatively inherited (Ruiz and Egea 2008; Ruiz et al. 2008, 2010; Salazar et al. 2013, 2014, 2016). Several QTLs linked to these fruit quality traits have been described in this species (Ruiz et al. 2010; Salazar et al. 2013, 2014).

The final goal of identifying these QTL regions is to convert conventional QTLs into expression QTLs (eQTLs) in order to clearly identify candidate genes responsible for the analysed traits. The eQTLs are genetic regions identified by applying QTL localisation methods to data on the abundance of transcripts in segregating genotypes (Druka et al. 2010). eQTLs are then derived from polymorphisms in the genome that result in differential measurable transcript levels (Salazar et al. 2014), where a correlation between genotype and expression levels is detected (Conesa et al. 2016). These associations can help us unravel the genetic bases of complex traits with a composite of interacting genes, such as colour (Sugiyama et al. 2014; Huang et al. 2014) or seed size (Ye et al. 2014). The biological validation of QTLs through gene expression analysis using qPCR has recently been assayed in *Vitis vinifera* (Huang et al. 2014), *Citrus* sp. (Sugiyama et al. 2014), *Brassica napus* (Lang et al. 2017) and *Rubus idaeus* (Simpson et al. 2017). To date, however, eQTL studies have not been performed in *Prunus* species to the best of our knowledge.

This work consists of a 2-year inheritance study of the most important fruit quality traits (fruit weight, stone weight, fruit diameter, fruit skin colour, flesh colour, blush colour, firmness, soluble solids and acidity) in two F₁ apricot progeny with a common male parent. In addition, the identification of stable QTLs enabled us to assay candidate genes located inside these QTL regions through gene expression analysis using qPCR.

Material and methods

Germplasm

The plant material assayed included two F₁ apricot progeny from the crosses between ‘Bergeron’ × ‘Currot’ (B×C) and ‘Goldrich’ × ‘Currot’ (G×C) with 187 and 200 seedlings, respectively. The ‘B×C’ and ‘G×C’ seedlings were planted in field conditions in 2009. ‘Bergeron’ is a French cultivar characterised by a late ripening time, high fruit weight, an attractive orange fruit colour, medium firmness, medium sugar content and acidity, high carotenoid levels and a good canning and cold storage capacity. ‘Goldrich’ is a North American cultivar from the cross ‘Sunglo’ × ‘Perfection’ characterised by a medium ripening time, an orange fruit colour, high fruit weight, high firmness, medium sugar content, high acidity and high carotenoid levels. Finally, the common male parent ‘Currot’ is a traditional Spanish cultivar characterised by a very early ripening time, low fruit weight, a light yellow colour, medium to low firmness, high sugar content, low acidity and low carotenoid levels. In addition, within the ‘G×C’ progeny, two genotypes—‘GC 2–11’ and ‘GC 3–7’—were selected for qPCR analysis. These genotypes were selected because they showed similar phenology and fruit characteristics with the exception of the fruit colour. ‘GC 2–11’ is self-compatible and early blooming and has an intermediate-sized oblong fruit with yellow skin, intense red blush and a yellow flesh colour. ‘GC 3–7’ is also self-compatible and early blooming with an intermediate-sized oblong fruit but with orange skin, intense red blush and a light orange flesh colour.

Phenotyping protocols

Several fruit quality traits were analysed for a period of two consecutive years (2012 and 2013). These traits

included physical traits (fruit weight, stone weight, fruit diameter, fruit colour and firmness) and biochemical traits (soluble solids and acidity). Fifty fruits per genotype were collected at the optimal ripening state, based on their skin ground colour (fully-coloured) and firmness (Sims and Comin 1963), in order to then select 12 fruits per genotype for fruit quality evaluations. Fruit and stone weight were measured using a Blauscal digital scale (model AH-600), with an accuracy of 0.01 g. Fruit skin colour, flesh colour, and blush colour were determined with a Minolta Chroma Meter (CR-300; Minolta, Ramsey, NJ, USA) tri-stimulus colour analyser calibrated to a white porcelain reference plate using a CIELAB scale with colour space coordinates L^* , a^* and b^* . To assess colour, we used the Hue angle [$H^\circ = \arctangent(b^*/a^*)$] parameter, which was determined around the equatorial region (Brown and Walker 1990). Firmness was quantified by a compression test in Newtons (N) using a Lloyd press (model LR10K; Fareham, Hants, UK). Soluble solid content (SS) was measured in °Brix, calibrated as the percentage of sucrose at 20 °C using a digital Optic Iyymen System (model DR-101). Finally, acidity was determined using 2 g of homogenised sample diluted in 30 ml of distilled water, and the values were obtained as grams of malic acid/100 ml, since this is the dominant organic acid in apricots (Souty et al. 1990).

All of the evaluated traits were analysed in 12 fruits per genotype and year. As for acidity and soluble solids, we evaluated these traits in three replicates resulting from the pool of 12 fruits. We used SPSS 12.0 software for Windows (Chicago, USA) to perform the statistical analyses. Frequency histograms were constructed for each fruit quality trait using the average value of each seedling in two different years. ANOVA analysis was performed to highlight differences among genotypes and years for those traits that met the criteria of normality according to Shapiro–Wilk, while the traits that did not meet the criteria of normality were analysed by the Kruskal–Wallis non-parametric test. Bivariate correlations between different traits were calculated with row data from the 2 years using the Pearson correlation coefficient.

In addition, samples were collected for qPCR analysis during the fruit development and ripening periods at three different times: A, at the beginning of veraison; B, during colour change and advanced veraison; and C, when the fruit was mature and showed its full colour state.

Marker identification and analysis

The new SSR and SNP markers assayed were located in scaffolds 3 and 4 of the *Prunus* reference genome v1.0 (<http://www.rosaceae.org/peach/genome>; Verde et al. 2013), where different QTLs linked to fruit quality traits have been described in peach (Eduardo et al. 2011; Fresnedo-Ramírez et al. 2015), apricot (Salazar et al. 2014) and Japanese plum (Salazar et al. 2017). These markers were identified by mapping the transcriptome reads obtained by NGS reads in BAM format, available after an RNA-Seq approach on five different cDNA libraries from ‘Rojo Pasión’, two from ‘Z506-7’ (Salazar et al. 2015) and two different cDNA libraries from ‘Bergeron’ × ‘Currot’. The reads were processed by SAMtools (Li et al. 2009) and displayed in alignment with the peach genome by Tablet (Milne et al. 2013) with the *Prunus* reference genome. The sequences were uploaded to the Phytozome website, and mask options were used to identify and eliminate repetitive domains. Total genomic DNA was extracted from young expanded leaves using the CTAB procedure described by Doyle and Doyle (1987).

For mapping, 25 new SNPs were selected and organised using 1 SNPLex, and 16 new SSRs were analysed by multiplex. SSR amplifications were performed according to multiplex PCR protocol as described by Campoy et al. (2010), using tag F primer labelled with FAM, TAM or HEX fluorescent dyes and unlabelled tag R primer. A volume of 3 µl of genomic DNA (concentration 10 ng/µl) was used in a 10-µl reaction mix. For fragment analysis, 3 µl of each PCR was mixed with 7 ml of formamide and 0.2 µl of GeneScan 500 LIZ® standard and sequenced using an ABI Prism 3730 DNA Analyser (Applied Biosystems, MA, USA). SSR peaks were visualised using Peak Scanner 1.0 software. The SNP design and analysis was performed according to Salazar et al. (2015). One independent SNPLex array was designed using MassARRAY Assay Design 3.1 software (available for the Sequenom platforms) in which the 25 selected SNPs were included.

Linkage analysis and QTL identification

Genetic linkage maps were constructed by JoinMap.3 (Van Ooijen 2006), and QTL identification was carried out using MapQTL ver. 4. Linkage groups 3 and 4 were reconstructed using the Kosambi function with a

frequency of recombination of 0.4 and a minimum LOD value over 3. As for QTL identification, phenotypic and genotypic data were analysed by first performing a test of 1000 permutations to designate the LOD threshold reference of $\alpha < 0.05$, $\alpha < 0.01$ and $\alpha < 0.001$ (genomewide) for each fruit quality trait using QTL detection interval mapping (parametric test) and the Kruskal–Wallis test (non-parametric test). In addition, the strongest marker cofactors of each QTL were identified by ‘Automatic Cofactor Selection’, and Multiple QTL analysis was applied in order to identify the most significant markers of the QTL interval. Linkage maps and QTL intervals were drawn using MapChart 2.3 software (Voorrips 2002) and the LOD colour gradient by Harry Plotter (Java software).

Biological validation of eQTLs through gene expression analysis using qPCR

Total RNA was extracted from the fruit mesocarps using a modified PowerPlant® RNA Isolation Kit (www.mobio.com), including a treatment with DNase I during extraction with an On-Spin Column DNase I Kit® (www.mobio.es). The cDNA was synthesised using SuperScript III Reverse Transcriptase (Thermo Fisher Scientific). To investigate the expression pattern, we evaluated nine candidate genes associated with the identified QTLs using qPCR, including transcription factors (*MYB 10*), carotenoid biosynthesis genes (*LOX 2*, *CCD1* and *CCD4*), anthocyanin biosynthesis genes (*ANS*, *UFGT* and *F3'5'H*), organic acid biosynthesis genes (*NAD ME*) and ripening date genes (*NAC*). Real-time qPCR experiments were performed with a StepOnePlus Real-Time PCR System (Applied Biosystems). Specific primers were designed based on previously obtained sequences in apricot libraries using Primer3 software (Supplementary Table 1) or validated using primers designed and assayed in peach.

We evaluated qPCR efficiency using the standard curve method. For all real-time qPCR reactions, we made a 10- μ l mix including 5 μ l Power SYBR® Green PCR Master Mix (Applied Biosystems), 0.5 μ l of each primer (5 μ M) and 2 μ l of cDNA (2.5 ng/ μ l). Each sample was analysed in triplicate from a pool of fruits collected in 2017. *SI8*, *RPL12* and *CYP1* were analysed as candidate housekeeping genes using the RefFinder web tool (Xie et al. 2012) (<http://leonxie.esy.es/RefFinder/?type=reference>). R version 3.4.2

with default setting was used to perform the statistical analyses of differential gene expression (RStudio 2015). Frequency histograms were constructed for each gene using the average value of each genotype and ripening state. Before checking the normality, homocestadicity and sample independence. We used the Kruskal–Wallis rank sum test (Hollander et al. 2013) for non-parametric distributions to determine the differences between the medians.

Results

Descriptive phenotypic analysis

The phenotypical characterisation was focused on phenotyping the plant material and studying the inheritance of the most important fruit quality traits in the apricot progeny ‘Bergeron’ \times ‘Currot’ (‘B \times C’) and ‘Goldrich’ \times ‘Currot’ (‘G \times C’). We did this by assaying 187 and 200 seedlings, respectively (Fig. 1 and Supplementary Table 2). In both progeny, there were significant differences by genotype and year for all evaluated traits (Supplementary Tables 3 and 4). Different frequency histograms were also constructed using the average data for each seedling, trait and year, which revealed normal distribution among the majority of traits studied in both progeny (Fig. 1). The Shapiro–Wilk normality test corroborated this normality (Supplementary Table 5). We can therefore confirm the polygenic nature and quantitative inheritance of all the traits studied.

A low fruit diameter influence was observed in both ‘B \times C’ and ‘G \times C’ populations, which may be due to the small size of ‘Currot’. Furthermore, the fruit skin colour of the seedlings was closer to that of ‘Currot’ (the parent in both populations). Intermediate inheritance with respect to parents was especially observed in seedlings in terms of acidity level (g malic acid/100 ml) in the ‘G \times C’ population, which may be due to high acidity from ‘Goldrich’ (2.6–3 g/100 ml) and low acidity from ‘Currot’ (1–1.4 g/100 ml). By contrast, for soluble solid content, we found transgressive values both above and below those observed in the parents (7–18 °Brix) in both progeny (Fig. 1).

Finally, according to the Pearson correlation analysis, we were able to confirm a high correlation between years for all evaluated traits, especially for fruit skin colour (0.670**), flesh colour (0.738**), acidity (0.764**) and soluble solids (0.720**). Results show




Fig. 1 Distribution among 187 seedlings of 'Bergeron' × 'Currot' (orange bars) and 200 seedlings of 'Goldrich' × 'Currot' (red bars) F_1 apricot progeny of the following fruit quality traits: ripening

date, fruit weight, stone weight, fruit diameter, skin colour, blush colour, flesh colour, firmness, soluble solids and acidity for the years 2012 and 2013

that these traits are quite balanced between years. We also found correlations among different traits that

occurred every year, such as correlations between skin and flesh colour (0.808*) and a low correlation

between acidity and fruit skin colour (-0.238^{**}) (Table 1).

Marker-trait analyses

New genetic markers were added to the genetic maps of two F₁ apricot progeny derived from the cross ‘Bergeron’ × ‘Currot’ (‘B×C’) and ‘Goldrich’ × ‘Curr’ (‘G×C’) previously reported by Salazar et al. (2016). All of these markers were designed with the aim to saturate the LG3 and LG4 of both progeny, where important phenology and fruit quality QTLs were found after preliminary analyses. Thus, a total of new 53 molecular markers (SSRs and SNPs) were mapped in LG3 and LG4 of the ‘B×C’ population, while 67 markers (SSRs and SNPs) were mapped in the same LGs for the ‘G×C’ population. Finally, increased mapping coverage was obtained for the ‘B×C’ (841.4 cM) and ‘G×C’ (789.5 cM) populations with respect to the 809.2 cM and 775.8 cM previously reported (Supplementary Table 6). These new molecular markers allow us to saturate the relevant genomic regions in LG3 and LG4, and some of them are potential markers for marker-assisted selection (MAS). In addition, integrated linkage groups for both parents and populations were generated for LG3 and LG4 to obtain more saturated genomic regions for potential phenology and fruit quality QTLs.

Moreover, in the integrated analysis of genotypic and phenotypic data, we identified several QTLs for each quality trait in both progeny (Supplementary Fig. 1 and Fig. 2; Table 2). These QTL results corroborate the great variability and polygenic nature of each pomological trait including different genome regions associated with morphological quantitative traits (fruit weight, stone weight and fruit diameter), morphological qualitative traits (skin colour, blush colour and flesh colour) and organoleptic traits (flesh firmness, soluble solids and acidity).

Morphological quantitative traits (stone and fruit weight and diameter)

We observed different QTLs linked to fruit (FW) and stone (SW) weight traits in LGs 1, 2, 3, 5, 6 and 7 in both the ‘B×C’ and ‘G×C’ populations (Supplementary Fig. 1 and Fig. 2). In ‘G×C’ progeny, however, the most significant QTLs for FW, SW and fruit diameter (CAL) were located in LG1,

which indicates that all of these traits are highly related, while in ‘B×C’ progeny, the stone weight QTL was linked to LG6 (Table 2). The LG6 of ‘Bergeron’ thus seems to be highly related to SW for both years, especially in the middle region chromosome close to the S6_12174633 and _22539524 markers, reaching a percentage of variance explanation (PEV) value of around 20%. Concerning the ‘G×C’ progeny, the most important QTLs for SW were on LG1 and LG2, for FW on LG1 and for CAL on LG1 and LG3 with a PEV value above 20%. The most significant marker in ‘G×C’ progeny for these traits was S1_38382339 in LG1, which reached LOD values of 7.41 and 9.48 for FW and CAL, respectively, with a PEV value close to 30%. As for the SW QTL, a PEV value close to 40% was reached for the S1_38382339 marker in ‘Goldrich’, and another SW QTL was located in LG2 around S2_18992724 with a PEV value close to 20%, coinciding with ‘Currot’ in the ‘B×C’ population. In ‘Currot’, almost no QTLs were detected for these traits, particularly in ‘B×C’ progeny.

Morphological qualitative traits (skin, flesh and blush colour)

Different skin (SKC), flesh (FLSC) and blush colour (BLSC) QTLs were located along linkage groups 1, 2, 3, 5 and 6 in both ‘B×C’ and ‘G×C’ populations (Supplementary Fig. 1 and Fig. 2). The most significant QTLs for the ‘B×C’ population were located in LG1 for BLSC and in LG6 for FLSC (Table 2). For the ‘G×C’ population, however, the most significant QTLs linked to fruit colour were identified in LG3 for SKC and FLSC (Table 2). In ‘Bergeron’, a QTL for BLSC was located in LG1 around the S1_14071595 and S1_20067081 markers, reaching a LOD value of around 7 and a PEV of around 30%.

As for the ‘G×C’ population, as mentioned above, the most important SKC QTLs were located in LG3 of ‘Goldrich’, and S3_18629805 and S3_18842927 were the most significant markers with LOD values over 20 and a PEV value close to 50%. The same SNPs were also associated with red blush colour and FLSC but with lower LOD values (Supplementary Fig. 2 and Table 2). In Fig. 2, we can see the great influence SKC has at the end of LG3, especially in ‘G×C’ progeny.

Table 1 Pearson correlation coefficients for different fruit quality traits for the years 2012 and 2013 in 187 apricot seedlings of ‘Bergeron’ × ‘Currot’ and 200 apricot seedlings ‘Goldrich’ × ‘Currot’ populations

	‘Bergeron’ × ‘Currot’								
	FW	SW	CAL	SKC	BLSC	FLSC	FIRM	SS	MALIC
Fruit weight (FW)	<i>0.498**</i>	0.644**	0.915**	−0.044	−0.004	0.008	0.280**	0.000	−0.101
Stone weight (SW)	0.573**	<i>0.772**</i>	0.515**	−0.260**	−0.066	−0.316**	0.387**	−0.101	−0.073
Fruit diameter (CAL)	0.905**	0.437**	<i>0.491**</i>	0.297**	0.216**	0.391**	−0.172*	−0.168*	−0.254**
Skin colour (SKC)	0.123	−0.061	0.060	<i>0.660**</i>	0.505**	0.645**	−0.343**	−0.051	0.052
Blush colour (BLSC)	0.183*	0.136	0.083	0.390**	<i>0.424**</i>	0.308**	−0.227**	−0.253**	−0.179*
Flesh colour (FLSC)	0.208**	−0.070	0.143	0.722**	0.270**	<i>0.738**</i>	−0.259**	−0.014	−0.038
Firmness (FIRM)	0.023	0.363**	0.176*	−0.201**	−0.002	−0.179*	<i>0.638**</i>	−0.114	0.156*
Soluble solids (SS)	−0.220**	−0.268**	−0.009	−0.098	−0.241**	−0.164*	0.056	<i>0.720**</i>	0.197**
Acidity (MALIC)	−0.235**	0.023	−0.143	−0.238**	−0.065	−0.165*	−0.076	0.434**	<i>0.687**</i>
	‘Goldrich’ × ‘Currot’								
	FW	SW	CAL	SKC	BLSC	FLSC	FIRM	SS	MALIC
Fruit weight (FW)	<i>0.515**</i>	0.757**	0.952**	−0.002	0.093	0.040	0.328**	0.082	−0.002
Stone weight (SW)	0.735**	<i>0.452**</i>	0.663**	−0.060	0.085	−0.062	0.435**	0.082	0.206**
Fruit diameter (CAL)	0.932**	0.628**	<i>0.542**</i>	0.208**	0.175*	0.246**	0.087	−0.200**	−0.145*
Skin colour (SKC)	0.118	−0.134	0.046	<i>0.670**</i>	0.550**	0.786**	−0.262**	0.012	−0.007
Blush colour (BLSC)	0.088	−0.112	0.128	0.542**	<i>0.371**</i>	0.378**	−0.052	−0.097	−0.032
Flesh colour (FLSC)	0.162*	−0.102	0.121	0.808**	0.374**	<i>0.733**</i>	−0.139	0.074	−0.044
Firmness (FIRM)	0.142*	0.285**	0.278**	−0.232**	−0.225**	−0.179*	<i>0.583**</i>	−0.061	0.213**
Soluble solids (SS)	−0.205**	−0.181*	0.069	−0.051	−0.176*	−0.041	0.287**	<i>0.603**</i>	0.182*
Acidity (MALIC)	−0.119	0.055	−0.066	−0.101	−0.022	−0.063	0.218**	0.276**	<i>0.764**</i>

The correlation is significant at the 0.05 (*) and 0.01 levels (**). The diagonal italicized line shows the correlation between years. Below the diagonal line, the results correspond to the 2012 correlation while those above the line show the correlation for 2013

Organoleptic traits

Regarding fruit firmness in the ‘B×C’ population, we only found linked QTLs in LG4, reaching a LOD value of 11.27 and a PEV value of 35% (Supplementary Fig. 1 and Table 2). This QTL is located in the middle region of chromosome 4 close to S4_10035210, S4_13001709 and SSR4_13182815 (Table 2). In the ‘G×C’ population, however, fruit firmness QTLs were detected in LGs 1, 2, 3, 4, and 6 (Supplementary Fig. 2). In ‘Goldrich’, the most important QTL was identified in LG 6, showing a PEV above 30% for the marker S6_5217065. As for the ‘Currot’ parent, LGs 2 and 3 seem to be strongly related to firmness, and S2_12550486, SSR3_15343739 and S3_15588939 are the most significant markers with a PEV value of around 20% (Table 2). In ‘G×C’ progeny, especially for LG1 and LG3, firmness and reddish skin colour may be inherited

together from ‘Goldrich’, which is characterised by a high level of firmness and an intense fruit skin colour.

Sugar content is a relevant trait for fruit quality and together with acidity contributes to the fruit taste, influencing consumer perceptions. QTLs for soluble solid (SS) content were identified along LGs 2, 3 and 4 (Supplementary Fig. 1 and Fig. 2), although the most significant QTLs were identified in LG4 for both populations. In ‘B×C’ and ‘G×C’ progeny, SS QTLs were consistent for both years, and S4_10035210, UDAP439, SSR4_13182815 and S4_13642265 (Fig. 2) were the most important markers.

QTL intervals in the other LGs reached lesser significance values than in LG4 for both populations. In the integrated LG4, in the QTL region for soluble solids, markers of both parents are included. The results show a greater influence downstream of SSR UDAP439, highlighting the QTL interval between S4_9061773




Fig. 2 LOD gradient scores by interval mapping analysis for skin colour in ‘G×C’ in LG3 and soluble solids in both the ‘B×C’ and ‘G×C’ populations in LG4. On the left side of each chromosome,

the centimorgan (cM) scale is shown, and below each chromosome, the LOD range is shown by colour gradient. The assayed candidate genes are indicated with arrows in bold and italics

and S4_11947345 (Fig. 2). LG4 undoubtedly has the greatest influence on this trait.

Acidity (the malic acid concentration) also plays an important role in fruit quality and flavour. In ‘B×C’ progeny, different acidity QTLs were identified in LGs 1 and 8 (Supplementary Fig. 1). The most important QTLs, however, were located at the beginning of LG8 in ‘Bergeron’ (in a region close to the marker S8_13630387), reaching a PEV value above 30% for both years. In the LG8 of ‘Currot’, on the other hand, the nearest marker linked to acidity was S8_4773161, reaching a PEV value close to 60% (Table 2). In the ‘G×C’ population, different QTLs were localised in LGs 2, 5, 6 and 8 (Supplementary Fig. 2), and the most significant QTLs were those obtained in LG2 and LG8 of ‘Goldrich’ and ‘Currot’, respectively. In ‘Goldrich’, in LG2, the markers S2_18992724 and UDAp456 reached significant LOD values of around

7 and a PEV value of 20% (Etienne et al. 2002a; Xi et al. 2016).

As for LG8, QTLs for both years were located at the end of the LG in ‘Currot’, involving the markers S8_17732612 and UDP98409 (Supplementary Fig. 1 and Fig. 2). This QTL interval reached a LOD value of around 7 and a PEV value above 20% (Table 2). In the ‘Goldrich’ parent, the nearest marker linked to acidity was S8_15339816, reaching a PEV value of over 20%. The QTLs linked to acidity were thus consistent for both populations and years, especially in LG8, so it would be interesting to explore this chromosome more in depth in conjunction with this trait.

Candidate genes and biological validation

The results of qPCR analysis show that *S18*, *RPL12* and *CYP1* are the best housekeeping genes. *S18* was

Table 2 Summary of the most significant markers linked to QTLs of fruit quality traits [ripening time (RT), stone weight (SW), firmness (FIRM), blush colour (BLSC), soluble solids (SS) and acidity (MALIC)] by interval mapping (IM) and multiple QTL

mapping (MQM) in a F1 apricot progeny of ‘Bergeron’ × ‘Currot’ and ‘Goldrich’ × ‘Currot’ during the years 2012 (_12) and 2013 (_13). LOD threshold for QTL intervals: ** $\alpha < 0.05$, *** $\alpha < 0.01$

‘Bergeron’						
Trait-Year	LG	QTL interval (IM)	Location	Nearest marker (MQM)	lod	% expl
SW_12	6	[36.7;52.8]**	41.1	S6_22539524	5.47	21.2
FLSC_12	6	[41.1;52.8]**	41.1	S6_22539524	4.77	20.7
FIRM_12	4	[33.6;56.3]***	46.0	S4_13001709	6.56	21.2
SS_12	4	[32.3;42.7]***	38.7	S4_10035210	7.75	26.2
MALIC_12	8	[0.0;10.1]**	0.0	S8_13630387	3.47	33.2
SW_13	6	[10.0;21.7]**	21.7	S6_12174633	5.14	22.9
SW_13	6	[36.7;41.1]**	41.1	S6_22539524	5.61	20.8
BLSC_13	1	[39.9;46.5]***	43.2	S1_14071595	7.09	29.0
BLSC_13	1	[39.0;46.5]***	44.0	S1_20067081	7.34	30.9
FIRM_13	4	[33.6;56.3]***	38.7	S4_10035210	11.27	35.0
FIRM_13	4	[33.6;56.3]***	44.7	SSR4_13182815	7.92	24.5
SS_13	4	[33.6;56.3]***	38.7	S4_10035210	9.4	29.6
SS_13	4	[33.6;56.3]***	44.7	SSR4_13182815	7.17	22.6
MALIC_13	8	[0.0;10.1]**	0.0	S8_13630387	4.71	45.8
‘Goldrich’						
SKC_12	3	[15.4;43.7]***	37.8	S3_17969374	19.23	42.5
SKC_12	3	[15.4;43.7]***	40.4	S3_18842927	22.98	50.0
FLSC_12	3	[15.4;43.7]***	37.8	S3_17969374	17.3	38.9
FLSC_12	3	[15.4;43.7]***	40.4	S3_18842927	18.79	43.2
FIRM_12	6	[0.9;9.6]**	4.6	S6_5217065	5.23	33.5
FW_13	1	[50.4;64.9]**	64.9	S1_38382339	7.41	28.8
SW_13	1	[50.4;64.9]*	64.9	S1_38382339	5.38	38.7
CAL_13	1	[35.4;64.9]**	64.9	S1_38382339	9.48	29.4
SW_13	2	[25.9;35.3]**	35.3	S2_18992724	8.7	21.9
SKC_13	3	[27.3;43.7]***	40.4	S3_18842927	21.95	47.0
SKC_13	3	[27.3;43.7]***	40.6	S3_18629805	21.97	47.0
FLSC_13	1	[50.4;64.9]**	64.9	S1_38382339	5.97	49.0
FLSC_13	3	[27.3;43.7]***	40.4	S3_18842927	14.56	34.5
FLSC_13	3	[27.3;43.7]***	40.6	S3_18629805	14.57	34.5
SS_13	4	[26.6;49.5]***	38.6	S4_13642265	10.56	27.8
SS_13	4	[26.6;49.5]***	40.2	SSR4_13182815	10.18	25.1
MALIC_13	2	[25.9;35.3]**	33.7	UDAp456	7.15	21.5
MALIC_13	2	[25.9;35.3]**	35.3	S2_18992724	7.5	20.6
MALIC_13	8	[17.0;26.0]**	21.0	S8_15339816	5.34	22.4
‘Currot’ (from the BxC Map)						
FIRM_12	4	[25.8;64.8]**	47.5	S4_13001709	6.55	21.2
FIRM_12	4	[25.8;64.8]**	47.9	SSR4_13182815	6.49	21.0
SS_12	4	[25.8;45.2]***	38.8	S4_10035210	6.91	23.4
SS_12	4	[25.8;45.2]***	40.2	UDAp439	7.57	24.6
FIRM_13	4	[30.8;64.8]***	47.5	S4_13001709	8.21	25.5

Table 2 (continued)

‘Bergeron’						
Trait-Year	LG	QTL interval (IM)	Location	Nearest marker (MQM)	lod	% expl
FIRM_13	4	[30.8;64.8]***	47.9	SSR4_13182815	7.92	24.5
SS_13	4	[37.9;52.9]***	38.8	S4_10035210	8.45	26.7
SS_13	4	[37.9;52.9]***	40.2	UDAp439	8.38	26.7
MALIC_13	8	[0.0;3.1]**	0.0	S8_4773161	4.1	59.4
‘Currot’ (from the GxC Map)						
CAL_12	3	[0.0;2.5]**	0.0	S3_899478	4.56	29.6
SKC_12	3	[42.5;84.4]***	67.6	S3_19398598	22.84	52.0
FLSC_12	3	[42.5;84.4]***	67.6	S3_19398598	18.93	44.5
FIRM_12	3	[47.1;65.2]***	52.8	SSR3_15343739	7.86	20.9
FIRM_12	3	[47.1;65.2]***	53.0	S3_15588939	8.17	21.2
SS_12	4	[26.1;64.1]***	46.2	UDAp439	8.1	21.7
SW_13	2	[21.1;53.6]***	38.6	S2_18992724	8.19	20.7
SKC_13	3	[53.5;84.4]***	67.6	S3_19398598	21.25	47.3
FLSC_13	3	[60.2;84.4]***	67.6	S3_19398598	15.52	37.3
FIRM_13	2	[16.6;38.6]***	21.1	S2_12550486	7.64	21.8
SS_13	4	[31.1;64.1]***	44.7	S4_10035210	9.35	23.4
SS_13	4	[31.1;64.1]***	46.2	UDAp439	10.85	27.3
MALIC_13	2	[16.6;48.6]**	21.1	S2_12550486	7.17	20.7
MALIC_13	8	[38.7;44.8]**	44.8	S8_17732612	7.86	20.8

discarded because of the high number of repetitions found in the apricot genome. *RPL12* and *CYP1* were used as reference genes for data normalisation (Niu et al. 2014), and the levels of relative expression were calculated by the $2^{-\Delta\Delta C_t}$ method (Pfaffl 2001), taking the Ct value from the ‘GC 2-11-A’ sample as the reference expression level and using both housekeeping genes in the normalisation of gene expression (Vandesompele et al. 2002).

The spatial and temporal expression of structural genes in anthocyanin biosynthesis is determined by the combination and interaction between R2R3-MYB, basic helix–loop–helix (*bHLH*) and WD40-type transcription factors (TFs), forming the MBW complex (Dare et al. 2008). Six MYB10-like and three *bHLH*-like TFs have been identified in peach. Of the six peach sequences potentially regulating anthocyanin biosynthesis, only three have been found to be expressed in apricot fruits: *MYB10.1* (ppa026640m), *MYB10.2* (ppa016711m), known *PpMYB10* (Ravaglia et al. 2013) and *MYB10.3* (ppa020385m). In the available peach genome

sequence, the three *MYB10s* (*MYB10.1*, *MYB10.2* and *MYB10.3*) are all located within 80 kb in LG3. In the current study, the maximum expression of these genes was found during flowering for all three genes. In the mesocarp, on the other hand, expression during ripening was only relevant for *MYB10.1* and *MYB10.3*, which were detectable in the mesocarp and peel and were highest in the mesocarp around the stone. *MYB10* showed higher expression in ‘GC 2-11’ than in ‘GC 3-7’, however (Fig. 3). The Kruskal–Wallis rank sum test confirmed statistical differences between genotypes that decreased during the fruit ripening process in both of them although not significantly (p value_{state} = 0.1979 and p value_{genotype} = 0.0004868),

At the beginning of LG5, a QTL for SKC was described close to S5_8566413 and AMPA105 in ‘G’ and ‘C’. The *ANS* gene is located near this QTL (Supplementary Fig. 1 and Fig. 2; Supplementary Table 1). The analysis of differential expression by qPCR showed statistically significant differences between states and genotypes (p value_{state} = 0.03631 and p value_{genotype} =




Fig. 3 Gene expression analysis using qPCR of candidate genes related to carotenoid biosynthesis (*LOX2*, *CCD1* and *CCD4*); anthocyanin biosynthesis (*MYB10*, *bHLH*, *DFR*, *F3'*

5'H, *UFGT* and *ANS*), organic acid biosynthesis (*NAD ME*) and ripening date (*NAC*) linked to QTLs

0.01185) being upregulated in 'GC 3-7' and reaching the maximum expression at the beginning of the ripening process (Fig. 3).

Flavonoid 3'-hydroxylase (*F3'H*) and flavonoid 3',5'-hydroxylase (*F3'5'H*), which are P450 enzymes, catalyse the hydroxylation of dihydrokaempferol (DHK) to form (2*R*,3*R*)-dihydroquercetin and dihydromyricetin, respectively.

As we mentioned above, a QTL for SKC was described close to S5_8566413 and AMPA105 in 'G' and 'C' the beginning of LG5. The gene *F3'5'H* is located near this QTL, where we also found the *ANS* gene (Supplementary Fig. 1 and Fig. 2; Supplementary Table 1). The analysis of differential expression by qPCR showed statistically significant differences

between genotypes but not between ripening states (p value_{state} = 0.1678 and p value_{genotype} = 0.003892). There was no minimal expression of *F3'5'H* in the 'GC 2-11' genotype. In 'GC 3-7', on the other hand, the expression of *F3'5'H* increased during the ripening process and reached its maximum expression in ripe fruit (Fig. 3).

The *UFGT* gene was located in the middle of LG6 in 'B', very close to the FLSC QTL (Supplementary Fig. 1 and Fig. 2; Supplementary Table 1). The expression of *UFGT* showed a sharp increase with a steep slope between veraison and maturity. The Kruskal–Wallis rank sum test confirmed statistically significant differences between genotypes and ripening states (p value_{state} = 0.03631 and p value_{genotype} = 0.005411). Furthermore,

the differences between genotypes were most significant in the expression of *UFGT*, which was upregulated in the ‘GC 3-7’ genotype during the entire fruit ripening process (Fig. 3).

The acidity trait is defined by the presence of a NAD-dependent malic enzyme (*NAD-ME*) close to LG2 QTL related to the malic acid in ‘G’ and ‘C’ (Supplementary Figs. 1 and 2; Supplementary Table 1). During the ripening process, the expression of *NAD-ME* decreased in both apricot genotypes (Fig. 3). No statistically significant differences were found between states, but there were statistically significant differences between genotypes (p value_{state} = 0.1284 and p value_{genotype} = 0.0234). Furthermore, the decrease in *NAD-ME* correlated with the increase in malic acid during fruit ripening in both genotypes.

LOX 2 genes were found in the BLSC, MALIC, SS and FIRM QTL regions in LG2. The expression of *LOX 2* was highest at the beginning of the ripening process (Fig. 3) and decreased until the fruit was completely mature in both genotypes (Supplementary Figs. 1 and 2; Supplementary Table 1). Statistical analysis confirmed significant differences between ripening states but not between genotypes (p value_{state} = 0.0005111 and p value_{genotype} = 0.6911).

In the regions linked to BSKC, FLSC and SKC in LG1 and LG2, two genes related to carotenoid synthesis, including carotenoid cleavage dioxygenase 1 (*CCD1*) and carotenoid cleavage dioxygenase 4 (*CCD4*), were identified (Supplementary Figs. 1 and 2; Supplementary Table 1). In our linkage map, *CCD1* was located at the beginning of LG1, in the middle of a FLSC QTL in ‘C’. *CCD1* was located in LG2 inside the FLSC QTL in ‘C’, but the expression did not change during the ripening process or between genotypes (p value_{state} = 0.2963 and p value_{genotype} = 0.05763). *CCD4* was located in LG1, inside BLSC and FLSC in ‘B’ in the same position described in peach (Adami et al. 2013). The expression of *CCD4* was upregulated (p value_{state} = 0.002117 and p value_{genotype} = 0.04694) to a significantly greater extent in the ‘GC 2-11’ genotype than in ‘GC 3-7’, and the highest levels occurred during veraison (Fig. 3). These results correlated with the flesh colour of each genotype—either light yellow in ‘GC 2-11’ or orange in ‘GC 3-7’—at the end of the ripening process.

Finally, *NAC* was located inside the SS QTL in LG4 for all of the tested genotypes. Furthermore, it was upregulated more in ‘GC 3-7’ than in ‘GC 2-11’ during

the ripening process (Fig. 3), which correlated with the MD of each genotype. The analysis of differential expression by qPCR showed statistically significant differences between states and genotypes (p value_{state} = 0.002754 and p value_{genotype} = 0.04694), which correlated with the increase in SS during the ripening process in both genotypes and also with the early maturation date of ‘GC 3-7’ compared to that of ‘GC 2-11’.

Discussion

Descriptive phenotypic analysis

The presence of transgressive trait values in the phenotype evaluation is probably due to the influence of the genetic background of the parents, which largely determines the segregation patterns obtained in the offspring (Salazar et al. 2013). In addition, the traits evaluated show the influence of genetic background and intermediate inheritance due to the transmission of co-dominant genes. We can thus confirm the polygenic nature and quantitative inheritance of the studied traits as well as the large variability of the seedlings in each population and the good correlation between years, as found by Salazar et al. (2013). Moreover, some correlations, such as acidity and fruit skin colour, could indicate that low skin colour (H°) or a more reddish skin colour could be related to high acidity levels. Ruiz et al. (2008) obtained similar results. In addition, Ruiz et al. (2005b) linked the carotenoid content to titratable acidity, obtaining a Pearson correlation of around 0.40. However, despite the fact that significant correlations have been identified between skin colour, acidity and carotenoid content, we can assert that these correlations were quite low.

Marker-trait analyses

Morphological quantitative traits (stone and fruit weight and diameter)

Some of our results agree with previous results from Quilot et al. (2004) and Eduardo et al. (2011), who located different QTLs for fruit weight in LGs 5 and 6, respectively. Eduardo et al. (2011) also linked the SSR marker UDP412 located in LG6 to fruit weight in the ‘Contender’ × ‘Ambra’ peach population. Campoy et al. (2015) placed a major QTL for fruit weight in LG5 of

two cherry F_1 progeny and identified some candidate genes in the *Prunus persica* genome v1.0.

There are few studies to date about fruit diameter. In a different peach progeny, for instance, Abbott et al. (1998) detected QTLs in LG1 and LG8, and Cantín et al. (2010) in LG4. Wang et al. (2000) described QTLs related to fruit size in LG2 and LG6 in cherry. De Franceschi et al. (2013) identified 23 FW2.2/CNR (cell number regulator) genes involved in fruit size along the peach genome in all linkage groups. These authors located two of the CNR genes for cherry (PavCNR12 and PavCNR20) in the QTL interval of LGs 2 and 6. In our populations, we also observed important QTLs for these traits in the same LGs and in similar positions to the CNR genes described by De Franceschi et al. (2013).

Recently, Fresnedo-Ramírez et al. (2016) studied fruit diameter (FD) and fruit weight (FW) as the most important yield traits using a pedigree analysis under a Bayesian framework. These authors used a pedigree of 464 individuals from different founders, breeding lines, commercial varieties, phantom parents and progeny. They identified five QTLs explaining 29 and 17% of phenotypic variance for FD and FW, respectively. However, the most relevant QTLs were identified on LGs 6 and 7 for two consecutive years linked to FD and FW in agreement with our results in ‘B×C’ progeny. These approaches could be interesting for studying the most relevant genetic components of fruit quality traits in any breeding programme (Fresnedo-Ramírez et al. 2016).

Morphological qualitative traits (skin, flesh and blush colour)

As we have seen, the QTLs for these traits are located in several linkage groups, which indicates the polygenic nature of the traits. This trend has also been observed in other *Prunus* species. In peach, for example, QTLs for skin colour have been located in LG2 (Quarta et al. 2000); in LGs 2 and 6 (Verde et al. 2002); in LG5 (Quilot et al. 2004); in LGs 3, 4, 6 and 7 (Eduardo et al. 2011); and in LG5 (Illa et al. 2011). In cherry, two QTLs were located in LGs 3 and 6 (Sooriyapathirana et al. 2010). Similarly, several authors have localised flesh colour QTLs in LG1 (Bliss et al. 2002; Martínez-García et al. 2013), LGs 1 and 3 (Quilot et al. 2004) and LG3 (Abbott et al. 1998; Illa et al. 2011) in peach as well as in LGs 3 and 8 (Sooriyapathirana et al. 2010) in cherry. In apricot progeny, Socquest-Juglard et al. (2012) located QTLs for fruit skin colour in LG3, and Ruiz et al. (2010)

located skin colour QTLs at the end of the same LG in the ‘Goldrich’ × ‘Moniquí’ population, coinciding with ‘Goldrich’ in our population. Other authors such as Quilot et al. (2004) and Martínez-García et al. (2013) have located QTLs associated with red and yellow flesh on LG1.

Other studies in apricot have shown QTLs for flesh colour in LGs 1 and 6 (Salazar et al. 2013). Concerning candidate gene identification, a *PavMYB10* gene [homologous to MdMYB10 in apple and to AtPAP1 in *Arabidopsis* (Espley et al. 2007)] was found at the end of LG 3, where the major QTL region for cherry skin and flesh colour is located. This suggests that this gene may be one of the main determinants of skin and flesh colour in cherry (Sooriyapathirana et al. 2010). For apple, the MdMYB10 transcription factor is involved in anthocyanin accumulation and is responsible for red skin and flesh colour (Espley et al. 2007; Chagné et al. 2007; Takos et al. 2006; Ban et al. 2007). These transcription factors in apple have been located in LG9, which is collinear with LG3 in peach (Illa et al. 2011). In our study, we identified the presence of an MYB gene located in the same QTL region as in cherry, close to S3_18629805 (Fig. 3). These data confirm synteny among the species belonging to the *Rosaceae* family.

Organoleptic traits

To date, few authors have referenced firmness QTLs. We would like to highlight those published in peach by Cantín et al. (2010) and Ogundiwin et al. (2009) in LGs 1, 4, 5, 7 and 8, as well as the major QTLs detected by Campoy et al. (2015) in LG6 in cherry progeny. The presence of several QTLs along different LGs undoubtedly indicates the polygenic nature of this trait and of the majority of fruit quality traits in general.

Regarding soluble solid content, many authors have described QTLs for single sugars like sucrose, fructose and glucose distributed throughout the peach LGs (Abbott et al. 1998; Dirlewanger et al. 1999; Quarta et al. 2000; Verde et al. 2002; Etienne et al. 2002a; Quilot et al. 2004; Cantín et al. 2010; Illa et al. 2011; Eduardo et al. 2011). In other species such as cherry, Wang et al. (2000) detected different QTLs in LGs 6 and 7, while in apricot, Ruiz et al. (2010) placed different QTLs in LGs 3 and 4 in agreement with Salazar et al. (2013), who identified QTLs in LGs 3, 4 and 5. In addition, in our progeny, the soluble solid content seems to be linked to ripening time and the fruit development

period, because all of these traits are mainly influenced by linkage group 4 in the same genomic region (Dirlewanger et al. 2012; Salazar et al. 2013; Pirona et al. 2013).

Similarly, several authors have detected acidity QTLs in almost all peach LGs (Dirlewanger et al. 1999; Etienne et al. 2002a; Quilot et al. 2004; Cantín et al. 2010; Eduardo et al. 2011). In apricot, Ruiz et al. (2010) and Salazar et al. (2013) located acidity QTLs in LGs 6, 7, 8 and LGs 1, 2 and 4, respectively. As we have seen, it is difficult to find a specific region related to fruit acidity, and it would be necessary to saturate these QTL regions to define more specific loci linked to this trait. In apricot, LG2 in ‘Goldrich’ and LG8 in ‘Currot’ seem to be the most important LGs involved in acidity levels.

Finally, in our QTL analysis, the most relevant results are linked to fruit skin colour in LG3, soluble solid content in LG4 and acidity content in LGs 2 and 8. This allows us to establish a frame of reference in the search for major genes linked to fruit quality traits in apricot using molecular markers. We have to consider, however, that the most consistent and significant QTLs for skin colour and soluble solid content were identified in LGs 3 and 4, respectively (Fig. 3). In addition, new markers mapped in LG4 could be linked to the ripening time QTL previously reported by Salazar et al. (2016), where SSR4_13182815 is related to soluble solid content, reaching a PEV value of over 10, which could be used as a potential molecular marker for MAS (Table 2).

These molecular tools could be used for aligning specific regions of different species and for developing markers to be used for breeding purposes. Fruit skin colour offers a good example: this trait seems to be controlled by a specific region (and probably by the same gene) in the species belonging to the *Rosaceae* family. This has been reported in peach (Frett et al. 2014), apricot (Ruiz et al. 2010; Socquest-Juglard et al. 2012), cherry (Sooriyapathirana et al. 2010), plum (Salazar et al. 2017), apple (Espley et al. 2007; Chagné et al. 2007) and pear (Pierantoni et al. 2010).

Candidate genes and biological validation

Gene expression using qPCR showed variable expression patterns throughout the fruit development period with a relationship between validated candidate genes and fruit quality traits linked to QTLs (Supplementary Fig. 1 and Fig. 2; Supplementary Table 1). The qPCR results also showed the clear implication of the assayed

genes in the ripening process, and there was a clear correlation between DNA and RNA results. We observed differential expression of *ANS*, *CCD4*, *UFGT*, *F3'5'H*, *NAD-ME* and *NAC* genes between the two assayed genotypes. In addition, we observed clear expression differences during the ripening process in *NAC*, *ANS*, *UFGT*, *CCD4* and *LOX 2*. *ANS*, *UFGT* and *F3'5'H* and *LOX 2* were upregulated in the ‘GC 3-7’ genotype and were responsible for the flesh, skin and blush colour; *NAC* was also upregulated in the ‘GC 3-7’ genotype and was responsible for maturity date. In ‘GC 2-11’, *CCD4* and *MYB10* were upregulated and were responsible for flesh, skin and blush colour. *NAD-ME*, on the other hand, was upregulated in ‘GC 3-7’, decreased during the ripening process and was responsible for the acidity content.

The transcription factor *MYB10* was upregulated in ‘GC 2-11’ in fruits with yellow flesh and skin and decreased during the fruit ripening period. We speculate that the anthocyanin biosynthetic pathway may be controlled by transcriptional complexes in apricot, and the transcriptional complexes are probably the orthologs of MYB regulators that can induce the transcription of anthocyanin pathway genes (Zhou et al. 2012, 2015). In Japanese plum (*P. salicina* L.), TFs of the *R2R3MYB* subfamily of the MYB family were linked to the regulation of anthocyanin biosynthesis. Analysis shows that a sustained increase in the expression of *PsMYB10* began in S2 in the skin of all the red cultivars and continued until S4. Furthermore, *PsMYB10* showed the highest positive correlation with anthocyanin accumulation and *ANS* and *UFGT* gene expression, suggesting a putative function of *PsMYB10* in the regulation of transcriptional control during anthocyanin biosynthesis. On the other hand, there is a significant negative correlation between anthocyanin accumulation, *ANS* and *UFGT* gene expression and the highest expression of *PsMYB1* in all yellow tissues (Gonzalez et al. 2016). In addition, epigenetic mechanisms such as promoter methylation of *R2R3 MYB* genes have been shown to play an important role in the regulation of anthocyanin accumulation in of apple (Telias et al. 2011) and pear fruits (Wang et al. 2013). Therefore, orthologs *R2R3-MYB* and epigenetic modification must be thoroughly researched in order to deeply understand the anthocyanin biosynthetic pathway in apricot.

In apricot fruit, the un-blushed skin colour, green at the beginning, became red in conjunction with chlorophyll degradation and anthocyanin accumulation, which

seems to occur early in fruit development and maturation. As the apricots ripened, the anthocyanin concentration increased in the skin, reaching a maximum and then decreasing toward the end of the maturation phase (Bureau et al. 2009).

The apricot species presents large variability in fruit colour, ranging from white ('Moniqui') and orange ('Goldrich') to fruits largely covered with a strong red blush ('Orange Red'). Anthocyanins are the most important pigments responsible for red coloration in apricots. The anthocyanin biosynthetic pathway is a major branch of flavonoid metabolism. Flavonoid biosynthesis genes like *ANS*, *F3'5'H* and *UFGT* are highly conserved among species and are organised in several branches, leading to the production of different flavonoids. Some branches are species-specific, whereas others are almost ubiquitous. The major anthocyanin compounds found in apricot fruit skin are cyanidin-3-O-rutinoside followed by cyanidin-3-O-glucoside and peonidin-3-O-rutinoside (Bureau et al. 2009; Gouble et al. 2005; Jo et al. 2015; Ruiz et al. 2005a; Ruiz et al. 2005b; Ruiz et al. 2008; Ayour et al. 2016; Campbell et al. 2013; Ruiz and Egea 2008; Gomez and Ledbetter 1997; Marty et al. 2005).

F3'H and *F3'5'H* determine the hydroxylation pattern of the B-ring of flavonoids and anthocyanins, which are necessary for leucocyanidin and leucodelphinidin production, respectively. These are the key enzymes that determine the anthocyanin structures and therefore colour (Gonzalez et al. 2016; Tanaka et al. 2008).

During fruit development, apricot undergoes a continuous accumulation of organic acids, which are used as respiratory substrates (Etienne et al. 2002b). All organic acids, including oxalate, tartrate, quinate, malate, citrate, fumarate and total organic acids, mostly increased during the early state of fruit development and decreased until fruits were fully ripe. Quinate, malate and citrate are the predominant organic acids throughout the whole fruit development and ripening period. As for the ratio of whole organic acids, malate is the first major organic acid in apricot, and the ratio of malate in the fruit decreased during development. Malate synthase (*MS*) is the key enzyme for the synthesis of malic acid. Malic enzymes (malate oxidoreductases) catalyse the oxidative decarboxylation of malate to produce pyruvate. In apricot, malate could be metabolised by the NADP-malic enzyme (*NADP-ME*) and the NAD-malic enzyme (*NAD-ME*). The decrease in *MS* and increase in *NADP-ME* and *NAD-ME* were together responsible for the malate decrease throughout the

development and ripening process, which further confirms the correlation between the activity of these enzymes and malate accumulation (Xi et al. 2016).

In plants, lipoxygenase (*LOX*) is closely related to fruit ripening and senescence. Additionally, fruit quality traits like fruit firmness, ethylene production and soluble solid content have been found to be affected by the differential expression of *LOX* under different storage conditions (Guo et al. 2017). In the fatty acid pathway, unsaturated fatty acids linoleic acid (18:2) and linolenic acid (18:3) can be converted into hydroperoxides by *LOX*, which is consistent with the corresponding aroma products of the metabolic flux (Xi et al. 2016). *PpaLOX2.2* has been identified in peach and may be required for fruit ripening during storage (Guo et al. 2017). Furthermore, *LOX 2* was determined as a putative candidate gene for transforming aldehyde from fatty acids by β -oxidation, yielding the precursors for the terpene pathway, which is related to carotenoid content (Eduardo et al. 2012).

In the edible portion of apricot, β -carotene is the main pigment followed by β -cryptoxanthin and γ -carotene (Ruiz et al. 2005b). The mechanism that controls carotenoid accumulation is largely unknown. Recently, two different regulatory mechanisms have been postulated. One is focused on carotenoid degradation, and the other is focused on sink capacity. Carotenoids can be cleaved into volatile apocarotenoids in fruit by *CCD*, generating flavour compounds (Tanaka et al. 2008). The differential expression of *CCD4* has thus been proposed as being the major determinant in the accumulation of carotenoids in peach fruit flesh (Brandi et al. 2011). In apricot, a rapid, significant increase in *CCD* activity has been found during the fruit development process in both the peel and pulp (Zhang et al. 2010; Xi et al. 2016). *CCD4* is the gene controlling flesh colour in peach, and its expression results in the degradation of carotenoids in white-fleshed genotypes, while the yellow colour arises as a consequence of its inactivation when the yellow phenotype originating from at least three independent mutations disrupts *CCD4* function, thus preventing carotenoid degradation (Adami et al. 2013). *NAC* is one of the largest plant transcription factor families and is a key regulator of developmental programs and stress response. *NAC* has been linked to maturity date (MD), where a few QTLs were detected with a higher explained variation. Fine mapping of the MD locus in peach identified the candidate gene (ppa008301m) encoding a transcription factor of the

NAC family as a possible causal gene (Pirona et al. 2013).

In our linkage map, there is a major QTL for SS where *NAC* is located. As we have mentioned, this is related to the soluble solid content and seems to be linked to ripening time and the fruit development period, because all of these traits are mainly influenced by the same region in LG4 (Supplementary Fig. 1 and Fig. 2). This is likely due to a major pleiotropic effect of MD masking the identification of other QTLs for different traits (Dirlewanger et al. 2012; Salazar et al. 2013; Pirona et al. 2013; Balogh et al. 2018). In this case, with a longer ripening process, fruit quality traits have more time to develop and fruits have more time to accumulate sugars and acids (Eduardo et al. 2010). A sequence variant in a *NAC* candidate gene with a 9 bp insertion (compared with the *Prunus persica* v1.0. reference genome) was shown to co-segregate with the MD trait and can be used in selecting early-maturity genotypes. The marker development on this sequence polymorphism provides a convenient molecular tool for discriminating between early- and late-ripening individuals in breeding populations (Pirona et al. 2013). The MD trait could thus be tentatively regarded as a co-dominant marker. These data indicate that MD could be controlled by the same locus within the *Prunus* species and suggest that selection for this trait should be efficient and rapidly integrated into breeding programmes (Dirlewanger et al. 2012).

Conclusions

Several QTLs related to fruit quality traits— such as fruit weight and stone weight, skin and flesh colour, firmness, soluble solid content and acidity—were identified in both ‘B×C’ and G×C’ populations. The most significant QTLs for soluble solid content were localised in LG4 and for skin and flesh colour in LG3. We can highlight the presence of candidate genes for the soluble solid QTL in LG4 that are involved in di-glucose and D-mannose binding (ppa001122m, ppa000854m and ppb001660m). Concerning skin colour, a gene encoding for an MYB transcription factor (previously linked to skin colour by other authors) was identified in the QTL region in LG3. The markers located in these QTL regions could be used for marker-assisted selection for all of the abovementioned traits. All of the genes linked to colour, acidity or

ripening date located in these QTLs had an expression level in agreement with the observed phenotype in the apricot fruits. The potentially easiest traits for marker assisted selection are those related to Mendelian inheritance, such as skin colour, which has been reported in LG3 of peach, apricot and plum, and ripening time, which has been reported in LG4 in the same species. Results also show the clear implication of the assayed genes in the ripening process: the qPCR data confirm the QTL analyses, and the correlation is clear between the results collected on DNA and those observed with the RNA approach. The *MYB10* gene is the best candidate gene for skin colour and was validated using qPCR expression. In addition, monitoring *NAC* expression is a good RNA marker for evaluating ripening progression. These results are of great interest for gene function validation and molecular breeding in apricot together with the development of PCR markers linked to skin colour for breeding programmes and the development of RNA markers for monitoring apricot fruit ripening.

Author contributions LD, PM-G and D.R. participated in the design and coordination of the study. D.R. and J.A.S. performed the phenotypic evaluation. B.E.G-G and J.A.S. Salazar carried out the SSR and SNP analyses, and B.E.G-G carried out the qPCR analysis. B.E.G-G, J.A.S., LD, PM-G and D.R. participated in the manuscript elaboration and discussion.

Funding information This study was financially assisted by the Seneca Foundation of the Region of Murcia (Saavedra Fajardo Postdoctoral fellow 20397/SF/17) during the stay of Juan A. Salazar in Murcia. This study was also supported by the “Apricot breeding” project of the Spanish Ministry of Economy and Competitiveness (AGL2013-41452-R) and the project “Breeding stone fruit species assisted by molecular tools” of the Seneca Foundation of the Region of Murcia (19879/GERM/15).

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- Abbott AG, Rajapakse S, Sosinski B, Lu ZX, Sossey-Alaoui K, Gannavarapu M, Reighard G, Ballard RE, Baird WV, Callahan A (1998) Construction of saturated linkage maps of peach crosses segregating for characters controlling fruit quality, tree architecture and pest resistance. *Acta Hort* 465: 41–49
- Adami M, De Franceschi P, Brandi F, Liverani A, Giovannini A, Rosati C, Dondini L, Tartarini S (2013) Identifying a carotenoid cleavage dioxygenase (*ccd4*) gene controlling yellow/

- white fruit flesh color of peach. *Plant Mol Biol Reporter* 31 (5):1166–1175
- Ayour J, Sagar M, Alfeddy MN, Taourirte M, Benichou M (2016) Evolution of pigments and their relationship with skin color based on ripening in fruits of different Moroccan genotypes of apricots (*Prunus armeniaca* L.). *Scientia Hort* 207:168–175
- Balogh E, Halász J, Szani Z, Hegedus A (2018) Correspondence between maturity date and molecular variations in a NAC transcription factor of diploid and polyploid *Prunus* species. *Turkish J Agric Forest* 42:136–144
- Ban Y, Honda C, Hatsuyama Y, Igarashi M, Bessho H, Moriguchi T (2007) Isolation and functional analysis of a MYB transcription factor gene that is a key regulator for the development of red coloration in apple skin. *Plant Cell Physiol* 48: 958–970
- Bazzano LA, He J, Ogden LG, Loria CM, Vupputuri S, Myers L, Whelton PK (2002) Fruit and vegetable intake and risk of cardiovascular disease in US adults the first national health and nutrition examination survey epidemiologic follow-up study. *Am J Clin Nutr* 76:93–99
- Bliss FA, Arulsekhar S, Foolad MR, Becerra V, Gillen AM, Warburton ML, Dandekar AM, Kocsisne GM, Mydin KK (2002) An expanded genetic linkage map of *Prunus* based on an interspecific cross between almond and peach. *Genome* 45:520–529
- Brandi F, Bar E, Mourgues F, Horváth G, Turcsi E, Giuliano G, Rosati C (2011) Study of “Redhaven” peach and its white-fleshed mutant suggests a key role of CCD4 carotenoid dioxygenase in carotenoid and norisoprenoid volatile metabolism. *BMC Plant Biol* 11(1):24
- Brown GS, Walker TD (1990) Indicators of maturity in apricots using biplot multivariate analysis. *J Sci Food Agri* 53:321–331
- Bureau S, Catherine M, Renard GC, Reich M, Ginies C, Audergon JM (2009) Change in anthocyanin concentrations in red apricot fruits during ripening. *Food Sci Technol* 42(1):372–377
- Campbell OE, Merwin IA, Padilla-Zakour OI (2013) Characterization and the effect of maturity at harvest on the phenolic and carotenoid content of Northeast USA apricot (*Prunus armeniaca*) varieties. *J Agric Food Chem* 61(51): 12700–12710
- Campoy JA, Martínez-Gómez P, Ruiz D, Rees J, Celton JM (2010) Developing microsatellite multiplex and megaplex PCR systems for high throughput characterization of breeding progenies and linkage maps spanning the apricot genome. *Plant Mol Biol Report* 28:560–568
- Campoy JA, Le Dantec L, Barreneche T, Dirlewanger E, Quero-García J (2015) New insights into fruit firmness and weight control in sweet cherry. *Plant Mol Biol Report* 33:783–796
- Cantín CM, Crisosto CH, Ogundwin EA, Gradziel T, Torrents J, Moreno MA, Gorgocena Y (2010) Chilling injury susceptibility in an intra-specific peach [*Prunus persica* (L.) Batsch]. *Postharvest Biol Technol* 58:79–87
- Chagné D, Carlisle CM, Blond C, Volz RK, Whitworth CJ, Oraguzie NC, Crowhurst RN, Allan AC, Espley RV, Hellen RP, Gardiner SE (2007) Mapping a candidate gene (MdMYB10) for red flesh and foliage colour in apple. *BMC Gen* 8:212
- Conesa A, Madrigal P, Tarazona S, Gomez-Cabrero D, Cervera A, McPherson A, Szcześniak MW, Gaffney DJ, Elo LL, Zhang X, Mortazavi A (2016) A survey of best practices for RNA-seq data analysis. *Genome Biol* 17:13
- Dare AP, Schaffer RJ, Lin-Wang K, Allan AC, Hellens RP (2008) Identification of a cis-regulatory element by transient analysis of co-ordinately regulated genes. *Plant Meth* 4:17
- De Franceschi P, Stegmeir T, Cabrera A, van der Knaap E, Rosyara UR, Sebolt AM, Dondini L, Dirlewanger E, Quero-García J, Campoy JA, Iezzoni AF (2013) Cell number regulator genes in *Prunus* provide candidate genes for the control of fruit size in sweet and sour cherry. *Mol Breed* 32: 311–326
- Dirlewanger E, Quero-García J, Le Dantec L, Lambert P, Ruiz D, Dondini L, Illa E, Quilot-Turion B, Audergon JM, Tartarini S, Letourmy P, Arús P (2012) Comparison of the genetic determinism of two key phenological traits, flowering and maturity dates, in three *Prunus* species: peach, apricot and sweet cherry. *Heredity* 109(5):280–292
- Dirlewanger E, Moing A, Rothan C, Svanella L, Pronier V, Guye A, Plomion C, Monet R (1999) Mapping QTLs controlling fruit quality in peach (*Prunus persica* (L.) Batsch). *Theor Appl Genet* 98:18–31
- Doyle JJ, Doyle JL (1987) A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem Bull* 19:11–15
- Druka A, Potokina E, Luo Z, Jiang N, Chen X, Kearsy M, Waugh R (2010) Expression quantitative trait loci analysis in plants. *Plant Biotech J* 8:10–27
- Eduardo I, Pacheco I, Chietera G, Bassi D, Pozzi C, Vecchietti A, Rossini L (2010) QTL analysis of fruit quality traits in two peach intraspecific populations and importance of maturity date pleiotropic effect. *Tree Genet Gen* 7(2):323–335
- Eduardo I, Pacheco I, Chietera G, Bassi D, Pozzi C, Vecchietti A, Rossini L (2011) QTL analysis of fruit quality traits in two peach intraspecific populations and importance of maturity date pleiotropic effect. *Tree Genet Gen* 7:323–335
- Eduardo I, Chietera G, Pirona R, Pacheco I, Troglio M, Banchi E, Bassi D, Rossini L, Vecchietti A, Pozzi C (2012) Genetic dissection of aroma volatile compounds from the essential oil of peach fruit: QTL analysis and identification of candidate genes using dense SNP maps. *Tree Genet Gen* 9(1):189–204
- Espley RV, Hellens RP, Puterill J, Kutty-Amma S, Allan AC (2007) Red colouration in apple fruit is due to the activity of a MYB transcription factor, MdMYB10. *Plant J* 49:414–427
- Etienne C, Rothan C, Moing A, Plomion C, Bodnes C, Svanella-Dumas L, Cosson P, Pronier V, Monet R, Dirlewanger E (2002a) Candidate gene and QTLs for sugar and organic acid content in peach. *Theor Appl Genet* 105:145–159
- Etienne C, Moing A, Dirlewanger E, Raymond P, Monet R, Rothan C (2002b) Isolation and characterization of six peach cDNAs encoding key proteins in organic acid metabolism and solute accumulation: involvement in regulating peach fruit acidity. *Physiol Plantarum* 114(2):259–270
- Fresnedo-Ramírez J, Bink MC, van de Weg E, Famula TR, Crisosto CH, Frett TJ, Gradziel TM (2015) QTL mapping of pomological traits in peach and related species breeding germplasm. *Mol Breed* 35:1–19
- Fresnedo-Ramírez J, Frett TJ, Sandefur PJ, Salgado-Rojas A, Clark JR, Gasic K, Peace CP, Anderson N, Hartmann TP,

- Byrne DH, Bink MCAM, van de Weg E, Crisosto CH, Gradziel TM (2016) QTL mapping and breeding value estimation through pedigree-based analysis of fruit size and weight in four diverse peach breeding programs. *Tree Genet Gen* 12:25
- Frett TJ, Reighard GL, Okie WR, Gasic K (2014) Mapping quantitative trait loci associated with blush in peach [*Prunus persica* (L.) Batsch]. *Tree Genet Gen* 10:367–381
- Gomez E, Ledbetter CA (1997) Development of volatile compounds during fruit maturation: characterization of apricot and plum x apricot hybrids. *J Sci Food Agri* 74(4):541–546
- Gonzalez M, Salazar E, Cabrera S, Olea P, Carrasco B (2016) Analysis of anthocyanin biosynthesis genes expression profiles in contrasting cultivars of Japanese plum (*Prunus salicina* L.) during fruit development. *Gene Expr Patterns* 21(1):54–62
- Guo S, Song Z, Ma R, Yang Y, Yu M (2017) Genome-wide identification and expression analysis of the lipoxygenase gene family during peach fruit ripening under different post-harvest treatments. *Acta Physiol Plantarum* 39(5):111
- Gouble B, Bureau S, Grotte M, Reich M, Reling P, Audergon JM (2005) Apricot postharvest ability in relation to ethylene production: influence of picking time and cultivar. In *Proceedings of the 5th International Postharvest Symposium, Vols 1-3*, edited by F. Mencarelli and P. Tonutti, 127–133. Leuven 1: International Society Horticultural Science
- Hertog MGL, Hollman PCH, Katan MB, Kromhout D (1993) Estimation of daily intake of potentially anticarcinogenic flavonoids and their determinants in adults in the Netherlands. *Nutr Cancer* 20:21–29
- Hollander, M., Wolfe, D. A., & Chicken, E. (2013). *Nonparametric statistical methods* (Vol. 751). John Wiley & Sons
- Huang YF, Violet S, Guiraud JL, Torregrosa L, Bertrand Y, Cheynier V, This P, Terrier N (2014) A negative MYB regulator of proanthocyanidin accumulation, identified through expression quantitative locus mapping in the grape berry. *New Phytol* 201(3):795–809
- Illa I, Eduardo I, Audergon JM, Barale F, Dirlwanger E, Li X, Moing A, Lambert P, Le Dantec L, Gao Z, Poëssel JL, Pozzi C, Rossini L, Vecchiotti A, Arús P, Howad W (2011) Saturating the *Prunus* (stone fruits) genome with candidate genes for fruit quality. *Mol Breed* 28:667–682
- Infante R, Martínez-Gómez P, Predieri S (2008) Quality oriented fruit breeding: peach [*Prunus persica* (L.) Batsch]. *J Food Agri Environ* 6:342–356
- Infante R, Martínez-Gómez P, Predieri S (2011) Breeding for fruit quality in *Prunus*. In: Jenks MA, Bebeli PJ (eds) *Breeding for fruit quality*. Ed. Wiley and Blackwell, New York (USA), pp 201–229
- Jo Y, Lian S, Cho JK, Choi H, Chu H, Cho WK (2015) De novo transcriptome assembly of two different apricot cultivars. *Gen Data* 6:275–276
- Lang L, Xu A, Ding J, Zhang Y, Zhao N, Tian ZS, Liu YP, Wang Y, Liu X, Liang FH (2017) Quantitative trait locus mapping of salt tolerance and identification of salt-tolerant genes in *Brassica napus* L. *Front Plant Sci* 8:100
- Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Durbin R (2009) The sequence alignment/map format and SAMtools. *Bioinformatics* 25(16):2078–2079
- Machlin LJ (1995) Critical assessment of the epidemiological data concerning the impact of antioxidant nutrients on cancer and cardiovascular disease. *Crit Rev Food Sci Nutr* 35:41–50
- Martínez-García PJ, Fresnedo-Ramírez J, Parfitt DE, Gradziel TM, Crisosto CH (2013) Effect prediction of identified SNPs linked to fruit quality and chilling injury in peach [*Prunus persica* (L.) Batsch]. *Plant Mol Biol* 81:175–188
- Marty I, Bureau S, Sarkissian G, Gouble B, Audergon JM, Albagnac G (2005) Ethylene regulation of carotenoid accumulation and carotenogenic gene expression in colour-contrasted apricot varieties (*Prunus armeniaca*). *J Exp Botany* 56(417):1877–1886
- Milne I, Stephen G, Bayer M, Cock PJ, Pritchard L et al (2013) Using tablet for visual exploration of second-generation sequencing data. *Brief Bioinform* 14:193–202
- Niu J, Zhu B, Cai J, Li P, Wang L, Dai H, Qiu L, Yu H, Ha D, Zhao H, Zhang H, Lin S (2014) Selection of reference genes for gene expression studies in Siberian apricot (*Prunus sibirica* L.) germplasm using quantitative real-time PCR. *PLoS One* 9(8):e103900
- Ogundiwin EA, Peace CP, Gradziel TM, Parfitt DE, Bliss FA, Crisosto CH (2009) A fruit quality gene map of *Prunus*. *BMC Gen* 10:587
- Pierantoni L, Dondini L, De Franceschi P, Musacchi S, Winkel-Shirley BJ, Sansavini S (2010) Mapping of an anthocyanin-regulating MYB transcription factor and its expression in red and green pear, *Pyrus communis*. *Plant Physiol Bioch* 48: 1020–1026
- Pirona R, Eduardo I, Pacheco I, Linge CD, Miculan M, Verde I, Tartarini S, Dondini L, Pea G, Bassi D, Rossini L (2013) Fine mapping and identification of a candidate gene for a major locus controlling maturity date in peach. *BMC Plant Biol* 13: 166
- Pfaffl MW (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* 29:e45
- Quarta R, Dettori MT, Sartori A, Verde I (2000) Genetic linkage map and QTL analysis in peach. *Acta Hort* 521:233–241
- Quilot B, Wu BH, Kervella J, Génard M, Foulongne M, Moreau K (2004) QTL analysis of quality traits in an advanced back-cross between *Prunus persica* cultivars and the wild relative species *P. davidiana*. *Theor Appl Genet* 109:884–897
- Ravaglia D, Espley RV, Henry-Kirk RA, Andreotti C, Ziosi V, Hellens RP, Costa G, Allan AC (2013) Transcriptional regulation of flavonoid biosynthesis in nectarine (*Prunus persica*) by a set of R2R3 MYB transcription factors. *BMC Plant Biol* 13(1):68
- RStudio Team (2015). RStudio: integrated development for R. RStudio, Inc., Boston, MA URL <http://www.rstudio.com/>
- Ruiz D, Egea J (2008) Phenotypic diversity and relationships of fruit quality traits in apricot (*Prunus armeniaca* L.) germplasm. *Euphytica* 163:143–158
- Ruiz D, Egea J, Gil MI, Tomás-Barberán FA (2005a) Characterization and quantification of phenolic compounds in new apricot (*Prunus armeniaca* L.) varieties. *J Agric Food Chem* 53:9544–9552
- Ruiz D, Egea J, Tomás-Barberán FA, Gil MI (2005b) Carotenoids from new apricot (*Prunus armeniaca* L.) varieties and their relationship with flesh and skin color. *J Agric Food Chem* 53: 6368–6374
- Ruiz D, Reich M, Bureau S, Renard C, Audergon JM (2008) Application of reflectance colorimeter measurements and

- infrared spectroscopy methods to rapid and nondestructive evaluation of carotenoids content in apricot (*Prunus armeniaca* L.). *J Agric Food Chem* 56(13):4916–4922
- Ruiz D, Lambert P, Audergon JM, Dondini L, Tartarini S, Adami M, Gennari F, Cervellati C, De Franceschi P, Sansavini S, Bureau S, Gouble B, Reich M, Renard CMGC, Bassi D, Testolin R (2010) Identification of QTLs for fruit quality traits in apricot. *Acta Hort* 862:587–592
- Salazar JA, Ruiz D, Egea J, Martínez-Gómez P (2013) Transmission of fruit quality traits in apricot (*Prunus armeniaca* L.) and analysis of linked quantitative trait loci (QTLs) using simple sequence repeat (SSR) markers. *Plant Mol Biol Report* 31:1506–1517
- Salazar JA, Ruiz D, Campoy JA, Sánchez-Pérez R, Crisosto CH, Martínez-García PJ, Blenda A, Jung S, Main D, Martínez-Gómez P, Rubio M (2014) Quantitative trait loci (QTL) and Mendelian trait loci (MTL) analysis in *Prunus*: a breeding perspective and beyond. *Plant Mol Biol Report* 32:1–18
- Salazar JA, Rubio M, Ruiz D, Tartarini S, Martínez-Gómez P, Dondini L (2015) SNP development for genetic diversity analysis in apricot. *Tree Genet Gen* 11:15
- Salazar JA, Ruiz D, Campoy JA, Tartarini S, Dondini L, Martínez-Gómez P (2016) Inheritance of reproductive phenology traits and related QTL identification in apricot. *Tree Genet Gen* 12(4):71
- Salazar JA, Pacheco I, Shinya P, Zapata P, Silva C, Ruiz D, Martínez-Gómez P, Infante R (2017) Genotyping by sequencing for SNP-based linkage analysis and identification of QTLs linked to fruit quality traits in Japanese plum (*Prunus salicina* Lindl.). *Front Plant Sci* 8:476
- Simpson CG, Cullen DW, Hackett CA, Smith K, Hallett PD, McNicol J, Woodhead M, Graham J (2017) Mapping and expression of genes associated with raspberry fruit ripening and softening. *Theor Appl Genet* 130:557–572
- Sims ET, Comin D (1963) Evaluation of objective maturity indices for Halehaven peaches. *Proc Am Soc Hortic Sci* 82:125–130
- Socquest-Juglard D, Christen D, Devènes G, Gessler C, Duffy B, Patocchi A (2012) Mapping architectural, phenological, and fruit quality QTLs in apricot. *Plant Mol Biol Report* 31:387–397
- Sooriyapathirana SS, Khan A, Sebolt AM, Wang D, Bushakra JM, Wang KL, Allan AC, Gardiner SE, Chagné H, Iezzoni AF (2010) QTL analysis and candidate gene mapping for skin and flesh color in sweet cherry fruit (*Prunus avium* L.). *Tree Genet Gen* 6:821–832
- Souty M, Audergon JM, Chambroy Y (1990) Abricot: les critères de qualité. *L'Arboriculture fruitière* 430:16–24
- Sugiyama A, Omura M, Shimoda T, Fuji H, Endo T, Shimizu T, Nesumi H, Nonaka K, Ikoma Y (2014) Expression quantitative trait loci analysis of carotenoid metabolism-related genes in *Citrus*. *J Japan Soc Hort Sci* 83:32–43
- Takos AM, Jaffe FW, Jacob SR, Bogs J, Robinson SP, Walker AR (2006) Light-induced expression of a MYB gene regulates anthocyanin biosynthesis in red apples. *Plant Physiol* 142:1216–1232
- Tanaka Y, Sasaki N, Ohmiya A (2008) Biosynthesis of plant pigments: anthocyanins, betalains and carotenoids. *Plant J* 54(4):733–749
- Telias A, Kui LW, Stevenson DE, Cooney JM, Hellens RP, Allan AC, Hoover EE, Bradeen JM (2011) Apple skin patterning is associated with differential expression of MYB10. *BMC Plant Biol* 11:14
- Van den Berg H, Faulks R, Fernando Granado H, Hirschberg J, Olmedilla B, Sandmann G, Southon S, Stahl W (2000) The potential for the improvement of carotenoid levels in foods and the likely systemic effects. *J Sci Food Agric* 80:880–912
- Van Ooijen JW (2006) JoinMap 4, software for the calculation of genetic linkage maps in experimental populations. Wageningen, Kyazma
- Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes." *Gen Biol* 3 (7): RESEARCH0034
- Verde I, Quarta R, Cerdrola C, Dettori MT (2002) QTL analysis of agronomic traits in a BC1 peach population. *Acta Hort* 592:291–297
- Verde I, Abbott AG, Scalabrin S, Jung S, Shu S, Marroni F et al (2013) The high-quality draft genome of peach (*Prunus persica*) identifies unique patterns of genetic diversity, domestication and genome evolution. *Nat Genet* 45(5):487–494
- Voorrips RE (2002) MapChart: software for the graphical presentation of linkage maps and QTLs. *J Hered* 93:77–78
- Wang D, Karle R, Iezzoni AF (2000) QTL analysis of flower and fruit traits in sour cherry. *Theor Appl Genet* 100:535–544
- Wang Z, Meng D, Wang A, Li T, Jiang S, Cong P, Li T (2013) The methylation of the PcMYB10 promoter is associated with green-skinned sport in Max Red Bartlett pear. *Plant Physiol* 162(2):885–896
- Xi W, Zheng H, Zhang Q, Li W (2016) Profiling taste and aroma compound metabolism during apricot fruit development and ripening. *Int J Mol Sci* 17:998
- Xie F, Xiao P, Chen D, Xu L, Zhang B (2012) miRDeepFinder: a miRNA analysis tool for deep sequencing of plant small RNAs. *Plant Mol Biol* 80(1):75–84
- Ye J, Liu P, Zhu C, Qu J, Wang X, Sun Y, Sun F, Jiang Y, Yue G, Wang C (2014) Identification of candidate genes JcARF19 and JcIAA9 associated with seed size traits in *Jatropha*. *Funct Integr Genomics* 14(4):757–766
- Zhang B, Shen JY, Wei WW, Xi WP, Xu CJ, Ferguson I, Chen KS (2010) Expression of genes associated with aroma formation derived from the fatty acid pathway during peach fruit ripening. *J Agri Food Chem* 58(10):6157–6165
- Zhou Y, Guo D, Li J, Cheng J, Zhou H, Gu C, Gardiner S, Han YP (2012) Coordinated regulation of anthocyanin biosynthesis through photorespiration and temperature in peach (*Prunus persica*). *Tree Genet Gen* 9(1):265–278
- Zhou H, Lin-Wang K, Wang H, Gu C, Dare AP, Espley RV, He H, Allan AC, Han Y (2015) Molecular genetics of blood-fleshed peach reveals activation of anthocyanin biosynthesis by NAC transcription factors. *Plant J* 82(1):105–121