

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Newly discovered associations between peritrich ciliates (Ciliophora: Peritrichia) and scale polychaetes (Annelida: Polynoidae and Sigalionidae) with a review of polychaete-peritrich epibiosis

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Mikac, B., Semprucci, F., Guidi, L., Ponti, M., Abbiati, M., Balsamo, M., et al. (2020). Newly discovered associations between peritrich ciliates (Ciliophora: Peritrichia) and scale polychaetes (Annelida: Polynoidae and Sigalionidae) with a review of polychaete-peritrich epibiosis. ZOOLOGICAL JOURNAL OF THE LINNEAN SOCIETY, 188, 939-953 [10.1093/zoolinnean/zlz111].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/708634> since: 2023-06-22

*Published:*

DOI: <http://doi.org/10.1093/zoolinnean/zlz111>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is a pre-copyedited, author-produced version of an article accepted for publication in **Zoological Journal of the Linnean Society** following peer review. The version of record:

**Barbara Mikac, Federica Semprucci, Loretta Guidi, Massimo Ponti, Marco Abbiati, Maria Balsamo, Igor Dovgal, Newly discovered associations between peritrich ciliates (Ciliophora: Peritrichia) and scale polychaetes (Annelida: Polynoidae and Sigalionidae) with a review of polychaete–peritrich epibiosis, Zoological Journal of the Linnean Society, Volume 188, Issue 4, April 2020, Pages 939–953,**

The version of record is available online at:  
<https://doi.org/10.1093/zoolinnean/zlz111>

#### Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

Journal : ZOOLIN

Article Doi : 10.1093/zoolinnean/zlz111

Article Title : Newly discovered associations between peritrich ciliates (Ciliophora: Peritrichia) and polychaetes (Polynoidae and Sigalionidae) with a review of polychaete–peritrich epibiosis

## INSTRUCTIONS

1. **Permissions:** Permission to reproduce any third party material in your paper should have been obtained prior to acceptance. If your paper contains figures or text that require permission to reproduce, please confirm that you have obtained all relevant permissions and that the correct permission text has been used as required by the copyright holders. Please contact jnls.author.support@oup.com if you have any questions regarding permissions.
2. **Author groups:** Please check that all names have been spelled correctly and appear in the correct order. Please also check that all initials are present. Please check that the author surnames (family name) have been correctly identified by a pink background. If this is incorrect, please identify the full surname of the relevant authors. Occasionally, the distinction between surnames and forenames can be ambiguous, and this is to ensure that the authors' full surnames and forenames are tagged correctly, for accurate indexing online. Please also check all author affiliations.
3. **Figures:** If applicable figures have been placed as close as possible to their first citation. Please check that they are complete and that the correct figure legend is present. Figures in the proof are low resolution versions that will be replaced with high resolution versions when the journal is printed.
4. **URLs:** Please check that all web addresses cited in the text, footnotes and reference list are up-to-date, and please provide a 'last accessed' date for each URL.
5. **Funding:** Please provide a Funding statement, detailing any funding received. Remember that any funding used while completing this work should be highlighted in a separate Funding section. Please ensure that you use the full official name of the funding body, and if your paper has received funding from any institution, such as NIH, please inform us of the grant number to go into the funding section. We use the institution names to tag NIH-funded articles so they are deposited at PMC. If we already have this information, we will have tagged it and it will appear as coloured text in the funding paragraph. Please check the information is correct. [red text to be used for suppliers who are tagging the funding]

## MAKING CORRECTIONS TO YOUR PROOF


These instructions show you how to mark changes or add notes to your proofs using Adobe Acrobat Professional versions 7 and onwards, or Adobe Reader DC. To check what version you are using go to **Help** then **About**. The latest version of Adobe Reader is available for free from [get.adobe.com/reader](http://get.adobe.com/reader).

**DO NOT OVERWRITE TEXT, USE COMMENTING TOOLS ONLY.**


### DISPLAYING THE TOOLBARS

#### Adobe Reader DC

In Adobe Reader DC, the Comment toolbar can be found by clicking 'Comment' in the menu on the right-hand side of the page (shown below).




The toolbar shown below will then display along the top.



#### Acrobat Professional 7, 8, and 9

In Adobe Professional, the Comment toolbar can be found by clicking 'Comment(s)' in the top toolbar, and then clicking 'Show Comment & Markup Toolbar' (shown below).




The toolbar shown below will then be displayed along the top.



### USING TEXT EDITS AND COMMENTS IN ADOBE ACROBAT

This is the quickest, simplest and easiest method both to make corrections, and for your corrections to be transferred and checked.



1. Click **Text Edits**
2. Select the text to be annotated or place your cursor at the insertion point and start typing.
3. Click the **Text Edits** drop down arrow and select the required action.

*You can also right click on selected text for a range of commenting options, or add sticky notes.*

#### SAVING COMMENTS

In order to save your comments and notes, you need to save the file (**File, Save**) when you close the document.


### USING COMMENTING TOOLS IN ADOBE READER

All commenting tools are displayed in the toolbar. To edit your document, use the highlighter, sticky notes, and the variety of insert/replace text options.




#### POP-UP NOTES

In both Reader and Acrobat, when you insert or edit text a pop-up box will appear. In **Acrobat** it looks like this:



In **Reader** it looks like this, and will appear in the right-hand pane:



## AUTHOR QUERY FORM

Journal : ZOOLIN

Article Doi : 10.1093/zoolinnean/zlz111

Article Title : Newly discovered associations between peritrich ciliates (Ciliophora: Peritrichia) and polychaetes (Polynoidae and Sigalionidae) with a review of polychaete–peritrich epibiosis

First Author : BARBARA MIKAC

Corr. Author : BARBARA MIKAC

### AUTHOR QUERIES - TO BE ANSWERED BY THE CORRESPONDING AUTHOR

Please ensure that all queries are answered, as otherwise publication will be delayed and we will be unable to complete the next stage of the production process.

Please note that proofs will not be sent back for further editing.

The following queries have arisen during the typesetting of your manuscript. Please click on each query number and respond by indicating the change required within the text of the article. If no change is needed please add a note saying “No change.”

AQ1	pl. provide relevant information
AQ2	Please confirm the heading levels
AQ3	Please check that the citation “Alvarez et al., 2014” not in the reference list.
AQ4	please check if this is correct
AQ5	should thi be ventral?... or ventrally?
AQ6	should this not be the other way round, i.e. redescribed then described?
AQ7	pl. check this is ok...
AQ8	interramal - ok?
AQ9	pl. reword to clarify meaning - perhaps use shorter sentences?...
AQ10	Please provide the citations for the references “Jankowski 2007, and San Martín 2003”.

# Newly discovered associations between peritrich ciliates (Ciliophora: Peritrichia) and polychaetes (Polynoidae and Sigalionidae) with a review of polychaete–peritrich epibiosis

BARBARA MIKAC<sup>1\*</sup>, FEDERICA SEMPRUCCI<sup>2</sup>, LORETTA GUIDI<sup>3</sup>, MASSIMO PONTI<sup>1,3</sup>, MARCO ABBIATI<sup>1,4</sup>, MARIA BALSAMO<sup>2</sup> and IGOR DOVGAL<sup>5</sup>

<sup>1</sup>Interdepartmental Research Centre for Environmental Sciences, University of Bologna, Via Sant’Alberto 163, 48123 Ravenna, Italy

<sup>2</sup>Department of Biomolecular Sciences, University of Urbino, Urbino, PU, Italy

<sup>3</sup>Department of Biological, Geological and Environmental Sciences, University of Bologna, Ravenna, Italy

<sup>4</sup>Department of Cultural Heritage, University of Bologna, Ravenna, Italy

<sup>5</sup>A.O. Kovalevsky Institute of Biology of the Southern Seas of RAS, Sevastopol, Russia

Received XXX; revised XXX; accepted for publication XXX

Received 31 July 2019; accepted for publication 26 August 2019

In this research, we report the presence of two ciliate protozoans of the subclass Peritrichia, *Cothurnia amphicteis* and *C. peloscolicis*, as epibionts on the chaetae of scaled polychaetes *Malmgrenia lilianae*, *M. andreapolis* (fam. Polynoidae) and *Sthenelais boa* (fam. Sigalionidae), from the north Adriatic (Mediterranean Sea). Both ciliate species are herein found for the first time after their original description and are redescribed, based on light and scanning electron microscopy analyses. This is the first record of an association between ciliates and polychaetes of the family Sigalionidae. Our results suggest that these host–epibiont relationships might be highly specific. We also present the first review of epibiosis between polychaetes and peritrich ciliates, indicating that this relationship is more diverse than previously thought. Forty taxa of peritrich ciliates from 12 genera and seven families are recorded as epibionts on polychaetes, while 48 polychaete taxa are known as their hosts. The relationship can be considered ectocommensalism, where the ciliates have the advantages of increased food availability. This association might be a more widespread phenomenon than currently known, because it could be easily overlooked or misinterpreted. It, therefore, deserves careful attention and further investigation.

**KEYWORDS:** Adriatic – Annelida – *Cothurnia* – *Malmgrenia* – Mediterranean – Protozoa – redescription – scanning electron microscopy – *Sthenelais*.

## INTRODUCTION

Epibiosis is the ecological association between a substrate organism, a basibiont, and a sessile organism, an epibiont, attached to the outer surface of the basibiont without trophic dependence on it (Wahl, 2009). Ciliate protozoans, particularly from the subclasses Peritrichia and Suctorina, are known

to establish epibiotic relationships with a variety of aquatic metazoans, such as Crustacea, Insecta, Gastropoda, Nematoda, Oligochaeta and Polychaeta (Alvarez-Campos *et al.*, 2014; Ansari *et al.*, 2017; Cabral *et al.*, 2018; Fernandez-Leborans & Tato-Porto, 2000; Sergeeva & Dovgal, 2014; Sartini *et al.*, 2018). Despite the high diversity and wide distribution of polychaetes in marine environments, records of polychaete–peritrich association are rare, and few papers deal with this relationship as their main topic (Magagnini & Verni, 1988; Alvarez *et al.*, 2014; Jankowski, 2014).

\*Corresponding author: Email: [barbara.mikac@unibo.it](mailto:barbara.mikac@unibo.it)

In this paper, we report for the first time the presence of ciliate peritrichs *Cothurnia amphicteis* Lang, 1948 and *Cothurnia peloscolicis* Precht, 1935 on scaled polychaetes *Malmgrenia lilianae* (Pettibone, 1993), *M. andreapolis* McIntosh, 1874 (fam. Polynoidae) and *Sthenelais boa* (Johnston, 1833) (fam. Sigalionidae) from the north Adriatic (Mediterranean Sea). The representatives of the protozoan genus *Cothurnia* are loricate ciliates found in fresh, brackish and marine waters, and have a cosmopolitan distribution. The lorica is attached to aquatic animals, plants, algae or inanimate substrate by a non-contractile external stalk. In many species, the stalk appears to be smooth and comparatively featureless, while others possess lines or stripes that run longitudinally down the stalk (Warren & Paynter, 1991). Transverse folds or furrows may also be present on the stalk surface. The shape of the aperture of the lorica, and sculpture of both lorica and stalk, are the principal characters used for identification of *Cothurnia* species. Most descriptions of cothurnians are based on observations from only one direction, where the shape of the aperture of the lorica is not well visible, and often only drawings are provided, which may be prone to error or misinterpretation. The two *Cothurnia* species reported, were no longer observed after their first descriptions, until the present research (Precht, 1935; Lang, 1948). Moreover, some details, especially concerning the outer sculpture of cothurnian lorica and stalk, are only visible with the use of scanning electron microscopy.

Thus, we give herein a redescription of *C. amphicteis* and *C. peloscolicis* based on light and scanning electron microscopy.

Records of ciliate epibionts may be easily overlooked in papers dealing with polychaetes, thus this association might be more common than it appears to be. In order to give a full picture of the present knowledge of polychaete–peritrich ciliate epibiosis we present the first review of this association, based on a detailed analyses of the existing literature and new data.

MATERIAL AND METHODS

RESEARCH AREA

Samples of benthos were collected by the Centre for Marine Research (Ruđer Bošković Institute, Rovinj, Croatia) at three offshore stations in the north Adriatic Sea, from 2003 to 2008. Stations SJ005 (45°18.4' N; 13°18.0' E; 31 m depth) and SJ007 (45°17.0' N; 13°16.0' E; 31 m depth) are situated on the transect Poreč (Croatia)–Venice Lido (Italy), while station SJ107 (45°02.8' N; 13°19.0' E; 37 m depth) lies on the transect Rovinj (Croatia)–Po River Delta (Italy) (Fig. 1). All three stations are characterized by silty sand substrate.

FIELD AND LABORATORY WORK

Macrofaunal samples were taken with a Van Veen grab, sieved through a 1-mm mesh and fixed in 4% buffered




Figure 1. Map of sampling area and stations (SJ005, SJ007 and SJ107).



formaldehyde–seawater solution. After sorting in the laboratory, macrobenthic organisms were preserved in 70% ethanol. Polychaetes were identified to the species level using stereo- and light-microscopes. Those with ciliate epibionts were further studied and photographed by means of light-microscope and Scanning Electron Microscope (SEM, FEI 515). Light micrographs were done under a Zeiss Axiovert 100 microscope, using a Nikon Digital sight DS-Fi2 camera and NIS-Elements D 4.30.02 64-bit programme. Measurements of ciliates were taken from light-microscopy photos. For the SEM analyses, fixed specimens were washed in 0.1 M phosphate buffer (pH 7.4), dehydrated in a graded alcohol series until 100% (5 min for each solution), then dried with hexamethyldisilazane (HMDS), according to the method of [Hochberg & Litvaitis \(2000\)](#). Dry specimens were mounted on aluminium stubs, sputter-coated with gold palladium and finally observed with a Philips 515 SEM.

In order to give a review of polychaete–peritrich ciliate epibiosis, we examined the main literature dealing with polychaetes, ciliates and their association. Classification of peritrich ciliates follows [Lynn \(2008\)](#).

## RESULTS

### ASSOCIATION OF POLYCHAETES AND CILIATES FOUND IN THIS RESEARCH

Ciliate epibionts *Cothurnia amphicteis* were observed on 18 specimens of the polychaete *Malmgrenia lilianae* and two specimens of *M. andreapolis* (fam. Polynoidae). Ciliate epibionts *Cothurnia peloscolicis* were observed only on one specimen of the polychaete *Sthenelais boa* (fam. Sigalionidae). *Cothurnia amphicteis* was found attached on the middle to distal part of *Malmgrenia* chaetae, mostly on notochaetae, more rarely on neurochaetae ([Figs 2, 3](#)). *Cothurnia peloscolicis* was found attached on the lower-basal part of the upper (simple spinous) neurochaetae of *S. boa* ([Figs 5, 6A](#)). In general, one ciliate per chaeta was found and only in a few cases two. Ciliates were never found attached on body surfaces of polychaetes. Number of epibionts per basibiont ranged from a few to about 100.

### TAXONOMIC ACCOUNT OF CILIATES

#### PHYLUM CILIOPHORA DOFLEIN, 1901

#### SUBPHYLUM INTRAMACRONUCLEATA LYNN, 1996

#### CLASS OLIGOHYMENOPHOREA DE PUYTORAC *ET AL.*, 1974

#### SUBCLASS PERITRICHIA STEIN, 1859

#### ORDER SESSILIDA KAHL, 1933

### FAMILY VAGINICOLIDAE DE FROMENTEL, 1874

#### GENUS *COTHURNIA* EHRENBERG, 1831

#### *Diagnosis*

Marine, brackish or freshwater loricate peritrichs, usually with one or two zooids per lorica. Lorica borne on a stalk and attached to aquatic animals, plants or inanimate objects. Lorica without valves or other means of closing the aperture. Inner layer or septum sometimes present enclosing a space at the posterior end of the lorica; septum connected to the base of the lorica via a mesostyle. Zooid(s) attached to the base of the lorica (or septum) directly or via an endostyle ([Warren & Paynter, 1991](#)).

#### *COTHURNIA AMPHICTEIS* LANG, 1948

([Figs 2–4](#))


#### *Material examined*

Light-microscopy measurements based on six *Cothurnia* specimens found on the polychaete *Malmgrenia lilianae*, north Adriatic Sea, **station SJ107**, 21 June 2005. Additional material examined from the polychaete *M. lilianae*, north Adriatic Sea: **station SJ005**, 30 August 2005 (three polychaetes), 18 October 2006 (one polychaete); **station SJ007**, 15 March 2005 (one polychaete; mounted for SEM), 21 June 2005 (two polychaetes), 18 October 2006 (one polychaete), 13 September 2007 (one polychaete); **station SJ107**, 21 June 2005 (one polychaete; mounted for SEM), 22 December 2005 (two polychaetes), 13 September 2007 (two polychaetes, one mounted for SEM), 12 October 2007 (one polychaete). Additional material examined from the polychaete *M. andreapolis*, north Adriatic Sea: **station SJ005**, 22 September 2006 (one polychaete, mounted for SEM); **station SJ107**, 12 October 2007 (one polychaete).

#### *Description*

Lorica conical, smooth, with thin wall, 57–77 µm long (53–56 µm after [Warren & Paynter, 1991](#)) × 35 µm wide. Aperture circular 30–33 µm (35–37 µm after [Warren & Paynter, 1991](#)) in diameter, the edge of aperture with extremely small, irregular outgrowths ([Figs 2C, 4C](#)). External stalk conical, flexed near substrate ([Fig. 4A](#)), with an annular bulge in connection with lorica base and basal disc, 18–35 µm (54–57 µm after [Warren & Paynter, 1991](#)) long, with conspicuous transverse folds ([Fig. 4D](#)). Endostyle short, broad, with longitudinal striae that are not visible in some cases. Mesostyle absent. Generally, two zooids are present ([Figs 2D, 4B](#)). Zooid conical, 85 × 40 µm, and extends between one third and





**Figure 2.** *Cothurnia amphicteis* on the chaetae of *Malmgrenia lilianae* (light microscopy). A, anterior part of *M. lilianae* showing densely attached ciliates. B, anterior chaetigers of *M. lilianae* with ciliates. C, anterior neurochaetae of *M. lilianae* with ciliates showing circular aperture of lorica. D, close-up photography of *C. amphicteis* showing two zooids. Scale bars: A = 250  $\mu$ m, B = 200  $\mu$ m and C, D = 50  $\mu$ m.

one half of its length beyond aperture (Fig. 4B). The length of contracted zooids 31–59  $\mu$ m, width 7–21  $\mu$ m. Peristomial lip well-developed, 45  $\mu$ m in diameter. Disc convex. Contractile vacuole lies just below peristome. Macronucleus elongate slightly curved anteriorly. Pellicular striations conspicuous.

#### Remarks


New specimens differ from the original description of *C. amphicteis* by the presence of two zooids in the lorica (with one exception) and the greater length of the lorica. Only contracted zooids were measured. The diagnosis of related species *C. acuta* Levander, 1915 (after Warren & Paynter, 1991) is rather similar to *C. amphicteis*. In both diagnoses, the absence of a mesostyle was mentioned. However, on Precht's

(1935) picture, *C. acuta* has the typical mesostyle with longitudinal striae. In our specimens, the mesostyle was absent, thus we consider them *C. amphicteis*. The suggested synonymy between *C. acuta* and *C. amphicteis* needs further consideration. This is the first finding of the species after its original description by Lang (1948).


#### Habitat

Marine, originally found attached to the chaetae of polychaete *Amphicteis gunneri* (M. Sars, 1835) (fam. Ampharetidae) (type host) in coastal waters of Sweden, Baltic Sea (type locality) (Lang, 1948). Other locality: north Adriatic Sea, on the chaetae of polychaetes *Malmgrenia lilianae* and *M. andreapolis* (this paper).





**Figure 3.** *Cothurnia amphiteis* on the chaetae of *Malmgrenia lilianae* (SEM). A, a part of *M. lilianae* body showing ciliates attached on the chaetae. B, close-up of one chaetiger with two specimens of *C. amphiteis* attached to the notochaeta. Arrows indicating ciliates. Scale bars: A = 280  $\mu$ m, B = 140  $\mu$ m.



**Figure 4.** *Cothurnia amphictis* on the chaetae of *Malmgrenia lilianae* (SEM). A, *C. amphictis* attached to the notochaetae, showing stalk flexed near substrate. B, one specimen of *C. amphictis* showing two zooids. C, detail of *C. amphictis* lorica showing the edge of aperture with small, irregular outgrowths. D, detail of *C. amphictis* stalk showing transverse folds. Scale bars: A = 60  $\mu$ m, B = 45  $\mu$ m, C = 20  $\mu$ m, D = 12  $\mu$ m.

*COTHURNIA PELOSCOLICIS* PRECHT, 1935

FIGS 5–6

*Material examined*


Material examined by light microscopy (measurements based on six *C. peloscolicis* specimens) and SEM microscopy from one specimen of polychaete *Sthenelais boa*, north Adriatic Sea, station SJ007, 5 March 2004.

*Description*

Lorica extended near the base and slightly converged toward aperture, smooth, not compressed, 61–86  $\mu$ m (81  $\mu$ m after Warren & Paynter, 1991)  $\times$  26–32  $\mu$ m

(27–41  $\mu$ m after Warren & Paynter, 1991) (Figs 5B, 6C). Aperture circular when viewed from above, 23–32  $\mu$ m in diameter (15  $\times$  33  $\mu$ m after Warren & Paynter, 1991), the edge of aperture without any outgrow (Fig. 6B). External stalk short, cylindrical, without folds, 10–16  $\mu$ m long (20  $\mu$ m after Warren & Paynter, 1991); mesostyle and endostyle absent (Fig. 5B). Stalk with longitudinal striae in some cases. Generally, two zooids present (Fig. 5B). Zooid 80  $\times$  18–22  $\mu$ m, extending just beyond aperture. The length of contracted zooid 26–55  $\mu$ m, width 11–18  $\mu$ m. Peristomial lip 27  $\mu$ m in diameter. Contractile vacuole small and situated in the peristomal region. Macronucleus straight, 50  $\mu$ m long. Pellicular striations inconspicuous.





**Figure 5.** *Cothurnia peloscolicis* on the chaetae of *Sthenelais boa* (light microscopy). A, one chaetiger of *S. boa* with ciliates attached on the neurochaetae. B, close-up of neurochaetae with several ciliates showing short stalk and two zooids. Scale bars: A = 200  $\mu$ m, B = 50  $\mu$ m.

#### Remarks

This is the first finding of the species after its original description by [Precht \(1935\)](#).

#### Habitat

Marine, originally found as epibiont of the oligochaete *Tubificoides benedii* (d'Udekem, 1855) [reported as *Peloscolex benedeni* (d'Udekem, 1855); type host] from the Bay of Kiel, Germany (Baltic Sea) (type locality) ([Precht, 1935](#)). Other locality: north Adriatic Sea on the chaetae of polychaete *Sthenelais boa* (this paper).

#### EPISTYLIS SP.

*Diopatra marocensis* Paxton, Fadlaoui & Lechapt, 1995 (Fam. Onuphidae) – on the gills and on the first parapodia; Bay of Biscay, Spain, north-east Atlantic ([Arias et al., 2010](#)).

#### GENUS RHABDOSTYLA KENT, 1880

*RHABDOSTYLA ARENICOLA* [FABRE-DOMERGUE, 1888](#) AQ4  
*Arenicola marina* (Linnaeus, 1758) (Fam. Arenicolidae) – on the branchial tufts; Concarneau, France, the Bay of Biscay, north-eastern Atlantic Ocean ([Fabre-Domergue, 1888](#)); on the gills, also on the body surface in the branchial region, especially ventrally, and on the slim posterior part of the body; Bay of Kiel, Germany, Baltic Sea ([Precht, 1935](#)); on the gills; Helgoland, Germany, Nord Sea ([Kahl, 1935](#)). AQ5

#### RHABDOSTYLA COMMENSALIS MOEBIUS, 1888

*Capitella capitata* (Fabricius, 1780) (Fam. Capitellidae) – on the body cuticle; Bay of Kiel, Germany, Baltic Sea ([Möbius, 1888](#)).


#### A REVIEW OF THE PERITRICH CILIATE EPIBIONTS ON POLYCHAETES

##### SUBCLASS PERITRICHIA STEIN, 1859

##### ORDER SESSILIDA KAHL, 1933

##### FAMILY EPISTYLIDIDAE [KAHL, 1935](#)

##### GENUS EPISTYLIS EHRENBERG, 1830



**Figure 6.** *Cothurnia peloscolicis* on the chaetae of *Sthenelais boa* (SEM). A, neurochaetae of *S. boa* with attached ciliates. B, close-up of ciliate *C. peloscolicis* lorica showing circular aperture. C, one *C. peloscolicis* attached to *S. boa* neurochaeta. Scale bars: A = 35  $\mu\text{m}$ , B = 15  $\mu\text{m}$  and C = 20  $\mu\text{m}$ .



*Terebellides stroemi* Sars, 1835 (Fam. Trichobranchidae) – on the cirri; Bay of Kiel, Germany, Baltic Sea (Möbius, 1888; Kahl, 1935; Precht, 1935).

*Rhabdostyla mapuche* Álvarez-Campos, Fernández-Leborans & Verdes, 2014.

*Syllis magdalena* Wesenberg-Lund, 1962 (Fam. Syllidae) – on the intersegmental furrows, close to parapodial bases and on the prostomium; Las Cruces, Central Chile, south-eastern Pacific Ocean (Álvarez-Campos *et al.*, 2014).

*Syllis* sp. 1 (Fam. Syllidae) – on the intersegmental furrows, close to parapodial bases; Las Cruces, Central Chile, south-eastern Pacific Ocean (Álvarez-Campos *et al.*, 2014).

*Syllis* sp. 2 (Fam. Syllidae) – on the intersegmental furrows, close to parapodial bases and on the prostomium; Las Cruces, Central Chile, south-eastern Pacific Ocean (Álvarez-Campos *et al.*, 2014).

*Salvatoria concinna* (Westheide, 1974) (Fam. Syllidae) – on the intersegmental furrows, close to parapodial bases; Las Cruces, Central Chile, south-eastern Pacific Ocean (Álvarez-Campos *et al.*, 2014).

*Salvatoria* sp. (Fam. Syllidae) – on the intersegmental furrows, close to parapodial bases; Las Cruces, Central Chile, south-eastern Pacific Ocean (Álvarez-Campos *et al.*, 2014).

#### *RHABDOSTYLA NEREICOLA* PRECHT, 1935

*Platynereis dumerilii* (Audouin & Milne Edwards, 1833) (Fam. Nereididae) – dorsally on the parapodia; Bay of Kiel, Germany, Baltic Sea (Precht, 1935).

*Rhabdostyla taboadai* Álvarez-Campos, Fernández-Leborans, Riesgo & Martin, 2014.

*Syllis prolifera* Krohn, 1852 (Fam. Syllidae) – on the intersegmental furrows, close to parapodial bases; Costa Brava, Spain, north-western Mediterranean coast (Álvarez-Campos *et al.*, 2014).

#### *RHABDOSTYLA VARIABILIS* DONS, 1918

*Scoloplos armiger* (Müller, 1776) (Fam. Orbiniidae) – between the parapodia and on the anterior and posterior part of parapodia; Baltic Sea, Germany (Dons, 1918; Precht, 1935).

*Phyllodoce laminosa* Savigny in Lamarck, 1818 (Fam. Phyllodocidae) – on the posterior part of parapodia; Baltic Sea, Germany (Dons, 1918); on the posterior part of the wide notopodia and on the body, positioned between the notopodia; Bay of Kiel, Germany, Baltic Sea (Precht, 1935).

#### *RHABDOSTYLA* SP. 1

*Syllis elongata* Day, 1949 (Fam. Syllidae) – on the dorsal surface, the nuchal organs, the mouth opening

and the anterior cirri; Tumbes, Peru, south-eastern Pacific Ocean (Álvarez-Campos *et al.*, 2014).

#### *RHABDOSTYLA* SP.

*Typosyllis macropectinans* Hartmann-Schröder, 1982 (Fam. Syllidae) – on the ventral surface; New South Wales, Australia (Álvarez-Campos *et al.*, 2014).

*Syllis microoculata* (Hartmann-Schröder, 1965) (Fam. Syllidae) – on the intersegmental furrows, close to parapodial bases; Maui, Hawaii, northern Pacific Ocean (Álvarez-Campos *et al.*, 2014).

*Dipolydora armata* (Langerhans, 1880) (Fam. Spionidae) – described as spermatophores attached to the chaetae of female worms; west coast of Barbados, West Indies, north-western Atlantic Ocean (Lewis, 1998).

#### *PERITRICHIA* CF. *RHABDOSTYLA* SP.

*Parapionosyllis papillosa* (Pierantoni, 1903) (Fam. Syllidae) – described as papillae placed in the interramal furrows and on the parapodia of anterior segments; Gulf of Naples, Italy, Tyrrhenian Sea, Mediterranean Sea (Pierantoni, 1903; Álvarez-Campos *et al.*, 2014).

#### FAMILY SCYPHIDIIDAE KAHL, 1933

#### GENUS *PARAVORTICELLA* KAHL, 1933

#### *PARAVORTICELLA LYCASTIS* CHAKRAVORTY, 1937

*Namalycastis indica* (Southern, 1921) (Fam. Nereididae) – on the parapodia; India (Chakravorty, 1937).

#### *PARAVORTICELLA TERESELLAE* (FAURÉ-FREMIET, 1920)

*Terebella lapidaria* Linnaeus, 1767 (Fam. Terebellidae) – forming fluffy spots on the polychaete skin; Germany (Kahl, 1935).

*Terebellides stroemii* Sars, 1835 (Fam. Trichobranchidae) – on the ventral body side, from the anterior part under gills to the mid body; Bay of Kiel, Germany, Baltic Sea (Precht, 1935).

#### GENUS *SCYPHIDIA* DUJARDIN, 1841

#### *SCYPHIDIA SPIONICOLA* PRECHT, 1935

*Pygospio elegans* Claparede, 1863 (Fam. Spionidae) – on the tentacles, the body surface, the cirri of posterior



parapodia, and between both the anterior and posterior parapodia; Bay of Kiel, Germany, Baltic Sea ([Precht, 1935](#))

*SCYPHIDIA TERESELLIDIS* [PRECHT, 1935](#)

*Terebellides stroemii* Sars, 1835 (Fam. Trichobranchidae) – on the branchiae; Bay of Kiel, Germany, Baltic Sea ([Precht, 1935](#)).

*SCYPHIDIA VARIABILIS* DONS, 1922

Terebellidae Johnston, 1846 – Norwegian coast ([Kahl, 1935](#)).

*SCYPHIDIA* SP.

*Nerilla antennata* Schmidt, 1848 (Fam. Nerillidae) – between the parapodia; Livorno coast, Italy, Tyrrhenian Sea, Mediterranean Sea ([Magagnini & Verni, 1988](#)).

FAMILY VAGINICOLIDAE DE FROMENTEL, 1874

GENUS *COTHURNIA* EHRENBERG, 1831

*COTHURNIA ACUTA* [LEVANDER, 1915](#)

*Bylgides sarsi* (Kinberg in Malmgren, 1866) (Fam. Polynoidae) – on the chaetae, particularly of the anterior parapodia; Tvärminne, Finland, Baltic Sea ([Levander, 1915](#); [Kahl, 1935](#)).

*Harmothoe imbricata* (Linnaeus, 1767) (Fam. Polynoidae) – on the chaetae; Bay of Kiel, Germany, Baltic Sea ([Precht, 1935](#)).

*COTHURNIA AMPHICTEIS* [LANG, 1948](#)

*Amphicteis gunneri* (M. Sars, 1835) (Fam. Ampharetidae) – on the chaetae; coastal waters of Sweden, Baltic Sea ([Lang, 1948](#)).

*Malmgrenia andreapolis* McIntosh, 1874 (Fam. Polynoidae) – on the chaetae; this research, north Adriatic Sea, Mediterranean Sea.

*Malmgrenia lilianae* (Pettibone, 1993) (Fam. Polynoidae) – on the chaetae; this research, north Adriatic Sea, Mediterranean Sea.

*COTHURNIA CERAMICOLA* KAHL, 1933

*Spirorbis (Spirorbis) spirorbis* (Linnaeus, 1758) (Fam. Serpulidae) – on the tube and the operculum; Bay of Kiel, Germany, Baltic Sea ([Precht, 1935](#)).

*COTHURNIA COMPLANATA* [PRECHT, 1935](#)

*Pherusa plumosa* (Müller, 1776) (Fam. Flabelligeridae) – on the chaetae; Bay of Kiel, Germany, Baltic Sea ([Precht, 1935](#)).

*Cothurnia kiwi* Álvarez-Campos, Fernández-Leborans & San Martín, 2014.

*Prosphaerosyllis magnoculata* (Hartmann-Schröder, 1986) (Fam. Syllidae) – on the intersegmental furrows on the base of parapodia; New Zealand (Álvarez-Campos *et al.*, 2014).

*COTHURNIA NEREICOLA* [PRECHT, 1935](#)

*Hediste diversicolor* (O. F. Müller, 1776) (Fam. Nereididae) – on both sides of the parapodia; Bay of Kiel, Germany, Baltic Sea ([Precht, 1935](#)).

*COTHURNIA PEDUNCULATA* DONS, 1918

*Pherusa plumosa* (Müller, 1776) (Fam. Flabelligeridae) – on the chaetae; Trøndelag, Norway, Norwegian Sea ([Dons, 1928](#); [Dons 1946](#)); Bay of Kiel, Germany, Baltic Sea ([Precht, 1935](#)).

*Remark:* [Dons \(1946\)](#) described this species as *Cothurnia trophoniae* [Dons, 1946](#), although [Dons \(1918\)](#) previously redescribed the same species as *C. pedunculata* ([Warren & Paynter, 1991](#)).

*COTHURNIA PELOSCOLICIS* [PRECHT, 1935](#)

*Sthenelais boa* (Johnston, 1833) (Fam. Sigalionidae) – on the chaetae; this research, north Adriatic Sea, Mediterranean Sea.

*COTHURNIA POLYDORICA* [JANKOWSKI, 2014](#)

*Polydora* sp. (Fam. Spionidae) – on the tips of the thick chaetae of two kinds; Sea of Japan ([Jankowski, 2014](#)).

*COTHURNIA STYLARIOIDES* [PRECHT, 1935](#)

*Pherusa plumosa* (Müller, 1776) (Fam. Flabelligeridae) – on the chaetae; Bay of Kiel, Germany, Baltic Sea ([Precht, 1935](#)).

*COTHURNIA* SP.

*Dipolydora armata* (Langerhans, 1880) (Fam. Spionidae) – on the notopodial capillary chaetae; Ibiza, Spain, Mediterranean Sea ([Bick, 2001](#)).

FAMILY VORTICELLIDAE EHRENBERG, 1838

GENUS *VORTICELLA* LINNAEUS, 1767

*VORTICELLA OBCONICA* [KAHL, 1935](#)

*SPIROBIS* SP. (FAM. SERPULIDAE) – NORWAY ([KAHL, 1935](#))

## VORTICELLA SP.

RHAMPHOBRACHIUM MACULATUM ESTCOURT, 1966  
 (FAM. ONUPHIDAE) – ON THE DORSAL SURFACE OF  
 THE ANTERIOR END OF THE BODY; NEW ZEALAND  
 (KNOX & HICKS, 1973).

GENUS PSEUDOVORTICELLA FOISSNER &  
 SCHIFFMANN, 1975

PSEUDOVORTICELLA PUNCTATA (DONS, 1918)

*Harmothoe imbricata* (Linnaeus, 1767) (Fam.  
 Polynoidae) – on the chaetae; Bay of Kiel, Germany,  
 Baltic Sea (Precht, 1935).

*Spirorbis (Spirorbis) spirorbis* (Linnaeus,  
 1758) (Fam. Serpulidae) – on the tube; Bay of Kiel,  
 Germany, Baltic Sea (Precht, 1935).

FAMILY ZOOTHAMNIIDAE SOMMER, 1951

GENUS HAPLOCAULUS WARREN, 1988

HAPLOCAULUS NICOLEAE PRECHT, 1935

*Nicolea zostericola* Örsted, 1844 (Fam. Terebellidae) –  
 on the posterior slim body part; Bay of Kiel, Germany,  
 Baltic Sea (Precht, 1935).

GENUS ZOOTHAMNIUM BORY DE ST. VINCENT, 1826

ZOOTHAMNIUM DUPLICATUM KAHL, 1933

*Spirorbis (Spirorbis) spirorbis* (Linnaeus, 1758) (Fam.  
 Serpulidae) – on the tube; Bay of Kiel, Germany, Baltic  
 Sea (Precht, 1935).

ZOOTHAMNIUM VERMICOLA PRECHT, 1935

*Lagis koreni* Malmgren, 1866 (Fam. Pectinariidae) – on  
 the tentacles, the parapodia, the branchiae, the body  
 surface; Bay of Kiel, Germany, Baltic Sea (Precht, 1935).

*Nephtys* sp. (Fam. Nephtyidae) – Bay of Kiel,  
 Germany, Baltic Sea (Precht, 1935).

*Nereimyra punctata* (Müller, 1788) (Fam.  
 Phyllodocidae) – on the whole body, including the long  
 cirri; Bay of Kiel, Germany, Baltic Sea (Precht, 1935).

*Eteone longa* (Fabricius, 1780) (Fam. Phyllodocidae)  
 – on all parts of the body; Bay of Kiel, Germany, Baltic  
 Sea (Precht, 1935).

ZOOTHAMNIUM SP.

*Pherusa plumosa* (Müller, 1776) (Fam. Flabelligeridae)  
 – on the chaetae; Bay of Kiel, Germany, Baltic Sea  
 (Precht, 1935).

## ORDER MOBILIDA KAHL, 1933

FAMILY URCEOLARIIDAE DUJARDIN, 1840

GENUS URCEOLARIA LAMARCK, 1801

URCEOLARIA CONVEXA HAIDER & DIETRICH 1977

*Phyllodoce mucosa* Örsted, 1843 (Fam. Phyllodocidae)  
 – Büsum, Germany, North Sea (Haider & Dietrich,  
 1977).

URCEOLARIA SERPULARUM (FABRE-DOMERGUE, 1888)  
 Fam. Serpulidae – on the branchial lamellae;  
 Concarneau, France, the Bay of Biscay, north-eastern  
 Atlantic Ocean (Fabre-Domergue, 1888, as *Leiotrocha*  
*serpularum*).

*Nephtys* sp. (Fam. Nephtyidae) – Bay of Kiel,  
 Germany, Baltic Sea (Precht, 1935, as *Cyclochaeta*  
*serpularum*).

*Phyllodoce laminosa* Savigny in Lamarck, 1818  
 (Fam. Phyllodocidae) – Bay of Kiel, Germany, Baltic  
 Sea (Precht, 1935, as *C. serpularum*).

*Serpula* sp. (Fam. Serpulidae) – branchiae; Germany  
 (Kahl, 1935, as *Cyclochaeta (Leiotrocha) serpularum*).  
 Remark: *Urceolaria serpularum* was originally  
 described in the monotypic genus *Leiotrocha*  
 Fabre-Domergue, 1888. Haider (1964) transferred  
*Leiotrocha serpularum* into the genus *Urceolaria*,  
 which was accepted by Xu & Song (2003), but not  
 by Lynn (2008), who insisted on validity of the  
 monotypic family Leiotrochidae Johnston, 1938.  
 Recent morphological and genetic analyses by Zhan  
 et al. (2013) support that *Leiotricha* should be placed  
 in *Urceolaria*.

URCEOLARIA SP.

*Polydora colonia* Moore, 1907 (Fam. Spionidae) – on  
 the palps, the anterior and the posterior chaetigers  
 and the pygidium; Hempstead East Marina, New  
 York, north-western Atlantic Ocean (David &  
 Williams, 2012).

*Polydora cornuta* Bosc, 1802 (Fam. Spionidae) – on  
 the body surface; Los Angeles Bay, southern California,  
 north-eastern Pacific Ocean (Douglas & Jones, 1991).

Family Trichodinidae Claus, 1951.

GENUS TRICHODINA EHRENBERG, 1830

TRICHODINA SCOLOPLONTIS PRECHT, 1935

*Scoloplos armiger* (Müller, 1776) (Fam. Orbiniidae) –  
 on the posterior end of the body; Bay of Kiel, Germany,  
 Baltic Sea (Precht, 1935).

## TRICHODINA TEREPELLIDIS PRECHT, 1935

*Terebellides stroemii* Sars, 1835 (Fam. Trichobranchidae) – on the branchiae; Bay of Kiel, Germany, Baltic Sea (Precht, 1935).

## UNCLASSIFIED PERITRICHIOUS CILIATES

*Polydora neocaeca* Williams & Radaskevsky, 1999 (Fam. Spionidae) – on the hoods of the bidentate hooded hooks; State of Rhode Island, North America, north-western Atlantic Ocean (Williams & Radaskevsky, 1999).

*Ampharete santillani* Parapar, Kongsrud, Kongshavn, Alvestad, Aneiros & Moreira, 2017 (Fam. Ampharetidae) – on the abdominal uncini, the branchial surface, the dorsolateral area behind the branchial surface and the ciliated buttons over the abdominal neuropodia; Galicia, north-western Spain and off Morocco, north-eastern Atlantic Ocean (Parapar *et al.*, 2018).

## ANALYSES OF THE POLYCHAETE–PERITRICH CILIATE ASSOCIATION

Forty taxa (30 determined to species level) of peritrich ciliates belonging to 12 genera and seven families are up to date recorded as epibionts of polychaetes. Most of them belong to the order Sessilida, while only five taxa belong to the order Mobilida. The most diverse are representatives of the genus *Cothurnia* (11 taxa, of which ten determined to species level). Forty-eight polychaete taxa (39 determined up to species level) are known as hosts for peritrich ciliate epibionts. They belong to the families Ampharetidae, Arenicolidae, Capitellidae, Flabelligeridae, Nephtyidae, Nereididae, Nerillidae, Onuphidae, Orbiniidae, Pectinariidae, Phyllodocidae, Polynoidae, Serpulidae, Sigalionidae, Spionidae, Syllidae, Terebellidae and Trichobranchidae. Most polychaete basibionts pertain to the family Syllidae (11 taxa, of which eight determined to species level).

## DISCUSSION

In this research, the peritrich ciliate *Cothurnia amphicteis* was found on polynoid polychaetes *Malmgrenia lilianae* and *M. andreapolis*, while *Cothurnia peloscolicis* was found on sigalionid polychaete *Sthenelais boa*. This is the first observation of polychaete–ciliate epibiosis from the Adriatic Sea and the first record of epibiosis between peritrich ciliates and polychaetes of the family Sigalionidae.

Other reports of Polynoidae as epibionts of polychaetes are sparse. Levander (1915) described *Cothurnia acuta* from the polychaete *Bylgides sarsi* [reported

as *Harmothoe sarsi* (Kinberg in Malmgren, 1865)] in Tvärminne (Finland, Baltic Sea), while Precht (1935) found *Cothurnia acuta* and *Pseudovorticella punctata* on *Harmothoe imbricata* in the Bay of Kiel (Germany, Baltic Sea).

The number of epibionts per host ranged from a few individuals to about 100. In all analysed specimens, ciliates were localized only on the chaetae (one ciliate per chaeta, rarely two), never on the body surface or other structures. Accordingly, in several previous studies, ciliates, particularly of the genus *Cothurnia*, were found attached on the chaetae (Levander, 1915; Dons, 1928; Kahl, 1935; Precht, 1935; Dons 1946; Lang, 1948; Lewis, 1998; Williams & Radaskevsky, 1999; Bick, 2001; Jankowski, 2014; Parapar *et al.*, 2018). Williams & Radashevsky (1999) found peritrich ciliates attached to the hoods of the bidentate hooded hooks of the spionid *Polydora neocaeca*; the stalks of the peritrich extended dorsally so that its body and oral region were positioned near the branchiae of the worm. Similarly, Bick (2001) found *Cothurnia* sp. attached on the notopodial capillaries of the postbranchiate chaetigers of the spionid *Dipolydora armata*. Ciliates were previously also found in other regions of the polychaete body, such as the body surface, the intersegmental furrows, the parapodia, the branchiae, the cirri, the tentacles, the prostomium, the mouth opening, the palps, the nuchal organs, the pygidium and the tube of the spirorbids. Parapar *et al.* (2018) found peritrich epibionts on different body parts, including the chaetae of the ampharetid *Ampharete santillani*, but noticed that their abundance was higher in ciliated body parts, such as the branchial surface, the dorsolateral area behind them and the ciliated buttons over the abdominal neuropodia.

Using fixed specimens, we could not address the ecological features of the epibiotic relationship. However, some general reflections on the consequences of this epibiosis can be retrieved from the literature. In general, an epibiotic association entails a highly complex suite of advantages and disadvantages for both partners (Wahl, 2009). The major advantage that ciliates gain from being associated with a motile substratum is the increased food availability, assured by the free transport to a variety of habitats, and by increased water flow. In fact, their frequent location on ciliated parts of the polychaete body suggests that they may take advantage of the water currents produced by the polychaete branchiae and cilia for feeding (Bick, 2001; Parapar *et al.*, 2018). Magagnini & Verni (1988) supposed that movement of the polychaete *Nerilla antennata*, determined by its ventral ciliated tract, resuspends the bottom debris containing bacteria and other microorganisms, making it available for the epizoic ciliate *Scyphidia*. In order to assess whether epibiosis interferes with the life cycle of



the polychaete host, the same authors measured for several months various parameters pertaining to the life cycle of *N. antennata*, with and without epibiotic ciliates. Their study indicated that the epibiosis did not appear to affect the life cycle of the polychaete. David & Williams (2012) reported that specimens of *Polydora colonia* hosting the ciliate *Urceolaria* in New York harbour did not appear to be negatively affected by the ciliate. In their study on epibiosis between ciliates and syllid polychaetes, Alvarez-Campos *et al.* (2014) did not notice alteration of swimming efficiency or other external harm in the specimens carrying the ciliate protozoans. Likewise, Parapar *et al.* (2018) did not observe damage caused by peritrich ciliates on the body surface of the polychaete *Ampharete santillani*. All this suggests that in polychaete–peritrich ciliate association, the advantages are limited to the ciliate, while the polychaete gets no harm from the epibiont. This association can, therefore, be considered as ectocommensalism.

Epibiotic relationships are rarely species-specific (Wahl, 2009). During the monitoring study on the three research stations, sampling was done in different seasons from 2003 to 2008; more than 21 000 individuals belonging to 230 polychaete species were carefully analysed and epibiosis was observed only between the ciliate *Cothurnia amphiteis* and the polychaetes *Malmgrenia lilianae* and *M. andreapolis*, and between *C. peloscolicis* and the polychaete *Sthenelais boa*. This suggests that these host–epibiont relationships might be species-specific. Congruently, Magagnini & Verni (1988) found the ciliate *Scyphidia* sp. associated only to the polychaete *Nerilla antennata* in benthic samples from Livorno (Italy), and concluded that the observed epibiosis is likely species-specific. In their survey of symbionts associated with spionid polychaetes from California, Douglass & Jones (1991) showed that a ciliate *Urceolaria* sp. was a specific epibiont of the polychaete *Polydora cornuta* and that its presence on the surface of the polychaete allowed the identification of the worm. They also showed that ciliates tended to have an affinity for polydorids versus other spionids.

Our research adds two more species to the ciliate fauna of the Adriatic Sea. Until today, representatives of the genus *Cothurnia* were reported from the Adriatic Sea only twice. Stiller (1968) found *Cothurnia membranoloricata* Stiller, 1968, in the vicinity of Rovinj (north Adriatic Sea) attached to the algae *Cladophora coelothrix* Kütz. and *C. laetevirens* (Dillwyn) Kütz. (Cladophoraceae). Recently, Fernandez-Leborans *et al.* (2012) reported *Cothurnia triangula* (Precht, 1935) as epibiont of the copepod *Typhlamphiascus* sp. from the Bay of Piran (Slovenia, north Adriatic Sea). *Cothurnia amphiteis* and *C. peloscolicis*, newly recorded in the northern Adriatic, were previously reported only for

the Baltic Sea. These new findings support previous observations showing that the north Adriatic Sea hosts elements of the flora and fauna with cold-temperate water affinities. In fact, the north Adriatic Sea is, together with the Gulf of Lion, the coldest sector of the Mediterranean Sea, showing ecological and biogeographical similarities with the north Atlantic Ocean (Bianchi *et al.* 2004; Boero & Bonsdorff 2007; Boero *et al.* 2008).

Records of ciliate epibionts may easily be overlooked in the extensive literature on polychaetes, as already stressed by Jankowski (2014), and this association might be even more common than it appears to be. Many papers reporting polychaete–peritrich associations are focused on polychaete taxonomy or ecology, and the presence of ciliates is barely mentioned, or eventually accompanied by an iconography and a short description of the association (Knox & Hicks, 1973; Douglas & Jones, 1991; Williams & Radashevski, 1999; Bick, 2001; Arias *et al.*, 2010; David & Williams, 2012; Parapar *et al.*, 2018). On the other hand, several peritrich ciliate species, particularly those of the genus *Cothurnia*, were originally described from the specimens found as epibionts on polychaetes (Fabre-Domergue, 1888; Möbius, 1888; Levander, 1915; Precht, 1935; Dons, 1946; Alvarez *et al.*, 2014; Jankowski, 2014). Although polychaete–peritrich association appears not to be so frequent and diversified as the epibiosis of peritrichs on some other invertebrate groups, such as Crustaceans (Fernandez-Leborans & Tato-Porto, 2000), our review has shown that it is documented in a remarkable number of reports. This epibiosis might even be a more widespread phenomenon, due to the possibility of overlooking or misinterpretations. In fact, ciliate epibionts were previously interpreted as morphological structures of the polychaete body (i.e. papillae and reproductive structures). Pierantoni (1903) described the polychaete *Pionosyllis papillosa* (today acknowledged as *Parapionosyllis papillosa*) from the Gulf of Naples (Italy), emphasizing as the principal diagnostic character of the species, the presence of a high number of large papillae of characteristic shape protruding from the polychaete skin surface of anterior body segments. ‘Papillae’ were gathered in small groups, particularly in interramal furrows and on parapodia. San Martin (2003) suggested that these papillae could actually be parasites and queried the taxonomic validity of *P. papillosa*. Later analyses by Alvarez-Campos *et al.* (2014) of the original description and drawings from Pierantoni (1903) and Fauvel (1923), as well as of the specimens identified and described by Campoy (1982), revealed that papillae were actually ciliate epibionts of the genus *Rhabdostyla*. The same authors analysed museum specimens of the species *Syllis microoculata*, originally collected in Manui, Hawaii. Structures of *S. microoculata* reported by

Hartmann-Schröder (1965) as papillae placed on intersegmental furrows, close to parapodial bases, were found to be misinterpreted ciliate epibionts of the genus *Rhabdostyla*. Alvarez-Campos *et al.* (2014) stated that, except for the ‘papillae’, *S. microoculata* is identical to the Mediterranean *S. prolifera* Krohn, 1852 but, given the distance between the Mediterranean Sea and the species type locality (Hawaii), further studies are needed to consider if they are the same species or a case of convergence. These later findings of overlooked epibiosis of ciliates on syllids have important taxonomic implications, since the papillae have been considered as a diagnostic character to distinguish among species or to erect new species (Alvarez-Campos *et al.*, 2014). The fixation method for polychaete conservation provokes contraction of the ciliate epibionts, which, together with their small size, causes difficulties in distinguishing them from papillae. Similar misinterpretations might have happened in other polychaete families bearing papillae (Alvarez-Campos *et al.*, 2014). Lewis (1998) described small, oval spermatophores produced by males and found attached by a stalk to the capillary chaetae or, occasionally, to the body wall on the genital segments of the females of the spionid polychaete *Dipolydora armata*, boring in the calcareous hydrozoans *Millepora complanata* Lamarck, 1816 on fringing reefs on the western coast of Barbados (West Indies, north-western Atlantic Ocean). Adult females commonly carried two or three, but up to a dozen of ‘spermatophores’ each. Careful analyses of the ‘spermatophores’ description, and light-microscopy photographs, revealed that they actually are peritrich ciliates, possibly belonging to *Rhabdostyla* (Jankowski, 2014).

Together with novel redescrptions of two *Cothurnia* species and new discoveries of polychaete–peritrich epibiosis, our analyses and results suggest that this association warrants further investigations, both to elucidate its real diversity and to deepen our knowledge on its ecological peculiarities.

#### ACKNOWLEDGEMENTS

Igor Dovgal’s work was made within the framework of research issue of A.O. Kovalevsky Institute of Biology of the Southern Seas of RAS “Fundamental research of population biology in marine animals, their morphological and genetic diversity” #AAAA-A19-119060690014-5.

#### REFERENCES

Álvarez-Campos P, Fernández-Leborans G, Verdes A, San Martín G, Martín D, Riesgo A. 2014. The tag-along

friendship: epibiotic protozoans and syllid polychaetes. Implications for the taxonomy of Syllidae (Annelida), and description of three new species of *Rhabdostyla* and *Cothurnia* (Ciliophora, Peritrichia). *Zoological Journal of the Linnean Society* **172**: 265–281.

Ansari KGMT, Guidi L, Dovgal I, Balsamo M, Semprucci F. 2017. Some epibiont suctorian ciliates from meiofaunal organisms of Maldivian archipelago with description of a new ciliate species. *Zootaxa* **4258**: 375–387.

Arias A, Anadón N, Paxton H. 2010. New records of *Diopatra marocensis* (Annelida: Onuphidae) from northern Spain. *Zootaxa* **2691**: 67–68.

Bianchi CN, Boero F, Fraschetti S, Morri C. 2004. *The wildlife of the Mediterranean*. In: Minelli A, Chemini C, Argano R, Ruffo S, eds. *Wildlife in Italy*. Milan: Touring Editor and Rome: Italian Ministry for the Environment and Territory, 248–335.

Bick A. 2001. The morphology and ecology of *Dipolydora armata* (Polychaeta, Spionidae) from the western Mediterranean Sea. *Acta Zoologica* **82**: 177–187.

Boero F, Bonsdorff E. 2007. A conceptual framework for marine biodiversity and ecosystem functioning. *Marine Ecology – an Evolutionary Perspective* **28**(Supplement 1): 134–145.

Boero F, Féral JP, Azzurro E, Cardin V, Riedel B, Despalatović M, Munda I, Moschella P, Zaouali J, Fonda Umani S, Theocharis A, Wiltshire K, Briand F. 2008. Climate warming and related changes in Mediterranean marine biota. In: Briand F, ed. *Climate warming and related changes in Mediterranean marine biota*, Vol. 35. Monaco: CIESM Workshop Monographs, 5–21.

Cabral AF, Dias RJP, Oliveira VC, Alves RDG, D’Agosto M. 2018. Rhabdostylid Ciliates (Ciliophora, Peritrichia, Epistylididae) as Epibionts on Chironomid Larvae: Evidence of High Specificity and Association with Organic Pollution. *Zoological Science* **35**: 514–520.

Campoy A. 1982. *Fauna de España. Fauna de anélidos poliquetos de la Península Ibérica*. Pamplona: EUNSA (Ediciones de la Universidad de Navarra), Publicaciones de Biología de la Universidad de Navarra, Serie Zoológica 7(1), 781.

Chakravorty M. 1937. On *Paravorticella lycastis* n. sp. an ectoparasitic ciliate on the parapodia of an Indian polychaete, *Lycastis indica*. *Journal of the Royal Microscopical Society* **57**(2): 71–74.

David AA, Williams, JD. 2012. Morphology and natural history of the cryptogenic sponge associate *Polydora colonia* Moore, 1907 (Polychaeta: Spionidae). *Journal of Natural History* **46**(23–24): 1509–1528.

Dons C. 1918. Neue marine Ciliaten und Suctorien. *Tromsø Museums Aarschefter* 38–39: 75–100.

Dons C. 1928. Neue und wenig bekannte Protozoen. *Det Kongelige Norske Videnskabers Selskabs Skrifter* **7**: 1–17.

Dons C. 1946. Zwei neue Cothurnien. *Det Kongelige Norske Videnskabers Selskabs* **12**: 119–121.

Douglass TG, Jones I. 1991. Parasites of California spionid polychaetes. *Bulletin of Marine Science* **48**(2): 308–317.

Fabre-Domergue P. 1888. Étude sur l’organisation des Urcéolaires et sur quelques genres d’infusoires voisins de

cette famille. *Journal de l'Anatomie et de la Physiologie Normales et Pathologiques de l'Homme et des Animaux* **23**: 214–260.

**Fauvel P. 1923.** *Faune de France 5. Polychètes errantes*. Paris: Le Chevalier, 488.

**Fernandez-Leborans G, Tato-Porto ML. 2000.** A review of the species of protozoan epibionts on crustaceans. I. Peritrich ciliates. *Crustaceana* **73**(6): 643–683.

**Fernandez-Leborans G, Chatterjee T, Grego M. 2012.** New records of epibiont ciliates (Ciliophora) on Harpacticoida (Copepoda, Crustacea) from the Bay of Piran (Gulf of Trieste, northern Adriatic). *Cahiers de Biologie Marine* **53**: 53–63.

**Haider G. 1964.** Monographie der Familie Urceolariidae (Ciliata, Peritricha, Mobilina) mit besonderer Berücksichtigung der im süddeutschen Raum vorkommenden Arten. *Parasitologische Schriftenreihe* **17**: 1–251.

**Haider G, Dietrich W. 1977.** Das mobile peritriche ciliat *Urceolaria convexa* n. sp. *Hydrobiologia* **52**(2–3): 135–141.

**Hartmann-Schröder G. 1965.** Zur Kenntnis der eulitoralen Polychaetenfauna von Hawaii, Palmira und Samoa. *Abhandlungen und Verhandlungen des Naturwissenschaftlichen Vereins in Hamburg* **9**: 81–161.

**Hochberg R, Litvaitis MK. 2000.** Phylogeny of Gastrotricha: a morphology-based framework of gastrotrich relationships. *The Biological bulletin* **198**: 299–305.

**Jankowski AW. 2007.** Review of taxa phylum Ciliophora Doflein, 1901. In: Alimov AF, ed. *Protista: handbook on zoology*. St. Petersburg: Nauka, Part 2, 415–993. [In Russian]

**Jankowski AW. 2014.** *Cothurnia polydorica* sp. nov. (Ciliophora: Peritrichia), a symbiont of shell-boring spionid polychaetes in the Sea of Japan, and a review of ciliates associated with polydorids. *Zoosystematica Rossica* **23**(2): 167–173.

**Kahl A. 1935.** Urtiere oder Protozoa I: Wimpertiere oder Ciliata (Infusoria), eine Bearbeitung der freilebenden und ectocommensalen Infusorien der Erde, unter Ausschluss der marinen Tintinnidae, 4. Peritricha und Chonotricha. In: Dahl F, ed. *Die Tierwelt Deutschlands und der angrenzenden Meeresgebiete nach ihren Merkmalen und nach ihrer Lebensweise*, 30. Jena: Verlag von Gustav Fischer, 651–886.

**Knox GA, Hicks K. 1973.** The polychaetes of New Zealand part 5. Onuphidae. *Journal of the Royal Society of New Zealand* **3**(2): 281–294.

**Lang K. 1948.** On *Rhabdostyla commensalis* Möbius and two new *Cothurnia* species (Ciliata, Peritrichia) from the west coast of Sweden. *Arkiv för Zoologi* **41-B**: 1–6.

**Levander KM. 1915.** Über eine neue marine *Cothurnia*-Art. *Meddelanden et Societas Profauna et Flora Fennica* **10**: 86–88.

**Lewis JB. 1998.** Reproduction, larval development and functional relationships of the burrowing, spionid polychaete *Dipolydora armata* with the calcareous hydrozoan *Millepora complanata*. *Marine Biology* **130**: 651–662.

**Lynn DH. 2008.** *The ciliated Protozoa: characterization, classification, and guide to the literature*. Dordrecht: Springer, 605.

**Magagnini G, Verni F. 1988.** Epibiosis of *Scyphidia* sp. (Ciliophora, Peritrichida) on *Nerilla antennata* (Archannelida, Nerillidae). *Bolletino di Zoologia* **3**: 185–189.

**Möbius K. 1888.** Bruchstücke einer Infusorienfauna der Kieler Bucht. *Archiv für Naturgeschichte* **1888**: 81–116.

**Parapar J, Kongsrud JA, Kongshavn K, Alvestad T, Aneiros F, Moreira J. 2018.** A new species of *Ampharete* (Annelida: Ampharetidae) from the NW Iberian Peninsula, with a synoptic table comparing NE Atlantic species of the genus. *Zoological Journal of the Linnean Society* **183**(3): 526–555.

**Pierantoni U. 1903.** La gestazione esterna (Contributo alla biologia ed alla embriologia dei Sillidi). *Archivio Zoologico Italiano* **1**: 231–252.

**Precht H. 1935.** Epizoen der Kieler Bucht. *Deutsche Akademie der Naturforscher Nova Acta Leopoldiana (Halle) N.F.* **3**: 405–474.

**San Martín G. 2003.** Annelida Polychaeta II: Syllidae. In: Ramos MA, Alba J, Bellés X, Gosálbez J, Guerra A, Macpherson E, Martín F, Serrano J, Templado J, EDS. *Fauna Ibérica. Vol. 21*. Madrid: Museo Nacional de Ciencias Naturales, CSIC, 1–554.

**Sartini B, Marchesini R, D'ávila S, D'Agosto M, Dias RJP. 2018.** Diversity and distribution of peritrich ciliates on the snail *Physa acuta* Draparnaud, 1805 (Gastropoda: Physidae) in a eutrophic lotic system. *Zoological Studies* **57**: 1–12

**Sergeeva N, Dovgal I. 2014.** First finding of epibiont peritrich and suctorian ciliates (Ciliophora) on oligochaetes and harpacticoid copepods from the deep-water hypoxic/anoxic conditions of the Black Sea. *Ecologica Montenegrina* **1**(1): 49–54.

**Stiller J. 1968.** Peritriche Ciliaten ökologisch verschiedener Biotope von Rovinj und Umgebung. *Acta Zoologica Academiae Scientiarum Hungaricae* **14**: 185–211.

**Wahl M. 2009.** Epibiosis: ecology, effects and defences. In: Wahl M, ed. *Marine hard bottom communities. Patterns, dynamics, diversity, and change*. *Ecological Studies* 206. Berlin, Heidelberg: Springer-Verlag, 61–72.

**Warren A, Paynter J. 1991.** A revision of *Cothurnia* (Ciliophora, Peritrichida) and its morphological relatives. *Bulletin of the British Museum (Natural History), Zoology* **57**(1): 17–59.

**Williams JD, Radashevsky VI. 1999.** Morphology, ecology and reproduction of a new *Polydora* species (Polychaeta: Spionidae) from the east coast of North America. *Ophelia* **51**(2): 115–127.

**Xu K, Song W. 2003.** Brief introduction to the mobile peritrich ciliates with revision of their systematics at the generic level. In: Song W, Zhao Y, Xu K, Hu X, Gong J, eds. *Pathogenic Protozoa in mariculture*. Beijing: Science Press, 291–314.

**Zhan Z, Xu K, Dunthorn M. 2013.** Evaluating molecular support for and against the monophyly of the Peritrichia and phylogenetic relationships within the Mobilida (Ciliophora, Oligohymenophorea). *Zoologica Scripta* **42**(2): 213–226.