This is the peer reviewed version of the following article:

F. Farabegoli, S. Nesci, V. Ventrella, A. Badiani, S. Albonetti, M. Pirini (2019). Season and cooking may alter fatty acids profile of polar lipids from blue-back fish. *Lipids*, 54: 741–753.

which has been published in final form at <https://doi.org/10.1002/lipd.12202>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

Federica Farabegoli^{1,2,*}, Salvatore Nesci², Vittoria Ventrella², Anna Badiani², Sabrina Albonetti²,
Maurizio Pirini²

Season and cooking may alter fatty acids profile of polar lipids from Blue-Back Fish

¹ ANFACO-CECOPESCA, Estrada Colexio Universitario, 16 - 30310 Vigo, Pontevedra, Spain

² Department of Veterinary Medical Science (DIMEVET), University of Bologna, Via Tolara di Sopra,
50 - 40064 Ozzano dell'Emilia (BO), Italy

*Corresponding author.

E-mail address: farabegolif@hotmail.com

Postal address: Estrada Colexio Universitario, 16 - 30310 Vigo, Pontevedra, Spain

Tel: (+34) 986 469 373

Fax: (+34) 986 469 269

Abstract

Polar lipids (PoL) represent a new promising dietary approach in the prevention and treatment of many human diseases, due to their potential nutritional value and unique biophysical properties. This study investigates the effects of catching season and oven baking on the fatty acid profiles (FAP) of PoL in four species of blue-back fish widely present in the North Adriatic Sea: anchovy (*Engraulis encrasicolus*), sardine (*Sardina pilchardus*), sprat (*Sprattus sprattus*) and horse mackerel (*Trachurus trachurus*).

PoL levels (427–652 mg/100 g flesh) varied among the four species, with no significant seasonal variations within species. FAP of raw fillets were particularly high in PUFA, especially DHA and EPA; total PUFA was constant in all species throughout the year, while n-3 PUFA rose in spring (except in sprat), especially due to the contribution of DHA. The FAP response for PoL to oven baking was species-specific and, among n-3 PUFA, DHA exhibited the greatest heat resistance; the influence of oven baking on FAP was found to be correlated with the catching season, especially for anchovy and sardine, while sprat PoL were not affected by cooking processes.

The four species analyzed in this study presented very low n-6/n-3 fatty acid ratios and highly favorable nutritional indices, emphasizing their PoL qualities and promoting their role in increasing human n-3 PUFA intake. The four species can be considered as superior sources of n-3 PUFA and can be employed as supplements in functional food manufacturing and in pharmaceutical and cosmetic industries.

Keywords

Blue-back fish · Polar lipids · Neutral lipids · Fatty acid composition · Season influence · Cooking effects · Baking · Nutritional quality

40 **Abbreviations**

41	AI	Atherogenicity index
42	ALA	α -Linolenic acid
43	ARA	Arachidonic Acid
44	DHA	Docosahexaenoic acid
45	EPA	Eicosapentaenoic acid
46	FAME	Fatty acid methyl esters
47	FAP	Fatty acid profile
48	HH	Hypocholesterolemic to hypercholesterolemic fatty acid ratio
49	LNA	Linoleic acid
50	MUFA	Monounsaturated fatty acid(s)
51	n-3 PUFA	Long-chain n-3 polyunsaturated fatty acid(s)
52	n-6 PUFA	Long-chain n-6 polyunsaturated fatty acid(s)
53	NL	Neutral lipids
54	PL	Phospholipids
55	PoL	Polar lipids
56	PUFA	Polyunsaturated fatty acid(s)
57	SFA	Saturated fatty acid(s)
58	TI	Thrombogenicity index
59	TL	Total lipids

60

Introduction

Fish is not only a valuable source of high quality animal protein, but it is also well known that fish lipids are rich in long-chain n-3 polyunsaturated fatty acids (n-3 PUFA), especially 20:5n-3 (EPA) and 22:6n-3 (DHA); these fatty acids are unanimously considered as molecules with high nutritional value for the human diet and represent a focus of nutritionists, for the prevention and treatment of obesity (Fernandes et al., 2014; Parmentier, Al Sayed Mahmoud, Linder, & Fanni, 2007; Schneedorferová, Tomčala, & Valterová, 2015). Several studies have demonstrated that n-3 PUFA may be considered as nutraceuticals, i.e. medicinal foods that play a role in maintaining well-being, enhancing health and modulating immunity and thereby able to prevent as well as to treat specific diseases such as cardiovascular diseases, rheumatoid arthritis, diabetes, ulcerative colitis, allergies, eczema, thickening of the skin, weight gain, premature labor, depression and cancer (Fernandes et al., 2014; Hossain, Hosokawa, & Takahashi, 2009; Loftsson, Ilievska, Asgrimsdottir, Ormarsson, & Stefansson, 2016; Murakawa et al., 2007; Taylor, Pletschen, Arends, Unger, & Massing, 2010).

The fatty acid profile (FAP) of marine fish can vary among species and among individuals, depending on their diet, size, age, gender, environmental conditions, season and method of capture (Fernandes et al., 2014).

Several studies have observed that the total lipid (TL) content of fish and the FAP may vary throughout the year, due to exogenous and endogenous factors, and may also be affected by processing or cooking methods (Kaçar & Başhan, 2015); in fact, cooking may significantly alter the content, composition, biological activity and nutritional value of fish lipids, due to water loss, oxidation, loss of fatty acids by leaching, and fatty acid exchange between fish and other oils (Little, Armstrong, & Bergan, 2000). The FAP of TL is influenced by the lipid classes by which they are constituted: neutral lipids (NL) and polar lipids (PoL), which are different in composition and role. NL mainly consist of triacylglycerols, storage lipids used for energy purposes, for the maturation of gametes during the breeding season and as a temporary PUFA reservoir which can be forwarded to structural lipids or directed to specific metabolic pathways (Varljen, Baticic, Sincic-Modric, Obersnel, & Kapovic, 2004). The PoL include mainly phospholipids (PL) (De Leonardis & Macciola, 2004; Fanni, Linder, & Parmentier, 2004), important structural components of cell membranes and eicosanoid precursors. The

FAP of NL seems to depend mainly on the quality and availability of dietary intake and on the reproductive status of the subject, while the PoL fraction is less representative of dietary composition and can vary mainly due to environmental conditions such as temperature and salinity (Cordier, Brichon, Weber, & Zwingelstein, 2002).

The n-3 PUFA of PoL are important precursors of beneficial prostaglandins and platelet-activating factors and have a number of interesting biological activities, ranging from involvement in inflammatory and allergic reactions and antihypertensive responses to physiological processes (reproduction, fetal development, childbirth) (Blank, Cress, Smith, & Snyder, 1992). Conversely, oxylipins formed from n-3 and n-6 PUFA have essential roles in normal physiology and function, but can also have detrimental effects (Gabbs, Leng, Devassy, Monirujjaman, & Aukema, 2015). In mammalian systems, longer-chain n-3 and n-6 PUFA are synthesized from two essential precursors: α -linolenic acid (ALA; 18:3 n-3), leading to n-3 PUFA, and linoleic acid (LNA; 18:2 n-6), resulting in n-6 PUFA. The precursors compete for the enzymatic system, thus a high intake of ALA supports the elongation of n-3 PUFA and reduces the elongation of n-6 PUFA (Brenner, 1977). The n-6/n-3 PUFA ratio is the main important index of nutritional value of fish lipids; the intake of an appropriate ratio of essential fatty acids appears to be critical for the prevention and reduction of obesity, and nutritionists recommend human intakes of PUFA with ratios below 4 (Santos-Silva, Bessa, & Santos-Silva, 2002). Currently, the n-6 PUFA content in the human diet is from 3.75 to 10 times higher than recommended value, which is strongly correlated to an increased incidence of chronic non-transmissible diseases. Western diet, composed of high contents of red meat, refined flours and industrial products (with high levels of n-6 precursors and low levels of n-3 ones), enhances the production of long-chain n-6 PUFA and suppresses the production of n-3 PUFA (Adkins & Kelley, 2010; Fernandes et al., 2014). Moreover, the amount of DHA available for the global body metabolism remains insufficient for a large proportion of consumers (Parmentier et al., 2007), so that a higher dietary intake of n-3 and a reduction of n-6 consumption is strongly recommended for the correction of an unbalanced diet (Fernandes et al., 2014).

Currently, there is increasing attention on marine PoL, especially on PL, because of their potential nutritional value and their unique biophysical properties, so that their administration seems to be a promising new dietary approach in the prevention and treatment of many human diseases. Compared to

NL, PL of marine origin show greater nutritional value, higher bioavailability and higher intestinal absorption of n-3 PUFA, properties due to the esterification in the sn-2 position of ARA, EPA and DHA (Parmentier et al., 2007); PL also present better resistance to oxidation compared to triacylglycerols from the same source (Adkins & Kelley, 2010). Fish species with a high PL content are considered valuable food products due to their high levels of PUFA. Previous studies suggested that fish meals rich in PL could be used as a valuable functional ingredient of food or nutraceuticals (Henna Lu, Nielsen, Timm-Heinrich, & Jacobsen, 2011). The demand for PUFA in health-related products is increasing; modern biotechnology and engineering make it possible to design and purify marine PL for a wide range of applications (e.g. in nutrition as dietary supplements, and in drug delivery in the form of liposomes) (Loftsson et al., 2016; Mika, Swiezewska, & Stepnowski, 2016). Despite the nutritional, pharmaceutical and cosmetic value of marine PL having been widely discussed (Burri, Hoem, Banni, & Berge, 2012), there are very limited studies on the FAP of PoL in blue-back fish, including species widely caught and commonly consumed; these small pelagic fish may provide a valuable source of n-3 PUFA due to the abundance of the stocks. The aim of this study was the market promotion of these species, currently with a low commercial value, and the divulgation of their nutritional properties. Moreover, since these species analyzed are generally consumed cooked in the area subjected to study, the effects of cooking process have been assessed. An increment in the consumption of local fish in place of imported or over-exploited fish species would represent an example of sustainable utilization of the natural marine resources, and would also have a positive environmental impact (Guo, Vikbjerg, & Xu, 2005). Therefore, the objectives of this study are: (1) characterizing the PoL content in four species of blue-back fish from the Adriatic Sea; (2) highlighting any changes in PoL content and FAP due to the seasonal period and/or cooking process; (3) providing information about the best season for catching and consuming the fish species considered.

Materials and Methods

Sample Collection and Preparation

Four fish species were included in this study: anchovy (*Engraulis encrasicolus*), sardine (*Sardina pilchardus*), sprat (*Sprattus sprattus*) and horse mackerel (*Trachurus trachurus*). Fish caught in the

North Adriatic Sea were obtained from wholesale fish markets in Cesenatico and Rimini, in three different seasons, fall (four batches for anchovy, sardine and horse mackerel) or winter (three batches for sprat), and in the following spring, as described in the study of Pirini, Testi, Ventrella, Pagliarani and Badiani (2010). Within each batch, one fillet out of each of 30 specimens was retained to constitute the pooled raw reference.

The contralateral fillets were oven-baked as described by Pirini, Testi, Ventrella, Pagliarani and Badiani (2010). Briefly, the oven temperature was set at 190°C; cooking time ranged from a minimum of 5 min for sprat to a maximum of 9 min for horse mackerel to reach a core temperature of 70 °C, as checked by an iron–constantan thermocouple connected to a digital potentiometer (Type J/K Thermometer mod. 421502, Extech Instruments Corp., Waltham, MA, USA).

Raw and cooked fillets were either skinned (sardine, sprat and horse mackerel) or left unskinned (anchovy), as customarily prepared in Italy, then cut into small pieces, homogenized using a stainless-steel meat mincer, and stored at –20 °C until lipid analysis.

Lipid Analysis

Lipid analyses were carried out in duplicate on 5 g of homogenized fillet, both raw and cooked. TL were extracted according to the method of Folch, Lees and Sloane Stanley (1957) and weighed after complete evaporation of the solvent. TL were separated into NL and PoL fractions by SPE (solid phase extraction) using Phenomenex Strata SI-1 Silica normal phase columns (55 µm, 70 Å) (Torrance, CA, USA) with a capacity of 12 mL, containing 2 g silica, according to the method described by Bayır et al. (2010), with some modifications as shown below. The sample of TL diluted in 3 mL of chloroform was applied to the column equilibrated by four aliquots of 3 mL of chloroform. The NL fraction was eluted by adding 3 mL of chloroform eight times with a flow of two drops per second; PoL were eluted by four aliquots of 3 mL of methanol and four aliquots of 3 mL of a chloroform/methanol mixture (3:7 v:v), with a flow of two drops per second. PoL content was evaluated gravimetrically after completely solvent evaporation. Colorimetric determination of phosphorus content confirmed gravimetrical results, thus confirming that the PoL fraction coincided with PL (Bayır et al, 2010). Fractions obtained were weighed, diluted in a specific volume of chloroform and methanol in the ratio 1:1, and stored at –20 °C prior to

use for further analysis. The methyl esters of the PoL fatty acids were prepared by transmethylation with acid catalyst (Christie, 1989) and analyzed on a Varian 3380 gas chromatograph equipped with a J&W Scientific DB-23 fused silica capillary column (30 m × 0.25 mm), under the conditions reported by Pirini et al. (2010).

Fatty acid identification was accomplished by comparing the retention time of unknown FAME with those of known FAME standard mixtures (Sigma-Aldrich Corp., St. Louis, MO, USA; PUFA No.1, Marine Source, and PUFA No.3, Menhaden Oil, SUPELCO, Inc., Bellefonte, PA, USA).

Lipid Quality Indices

A qualitative assessment of the nutritional profile of the PoL fraction was done through computation of the ratio between the percentages of n-6 and n-3 PUFA, often highlighted as a key element for a healthy diet (Simopoulos, 2006). Moreover, the nutritional quality of the PoL fraction was assessed by three indices: atherogenicity index (AI), thrombogenicity index (TI) and hypocholesterolemic to hypercholesterolemic fatty acid ratio (HH), as described by Ulbricht and Southgate (1991) and Santos-Silva, Bessa, & Santos-Silva (2002). AI indicates the relationship between the sum of the main saturated fatty acids (14:0 and 16:0) and that of the main classes of unsaturated fatty acids (MUFA and n-3 and n-6 PUFA); AI was calculated by applying the following equation:

$$AI = \frac{[(4 \times 14:0) + 16:0]}{MUFA + n-6 + n-3}$$

IT is expressed as the relationship between pro-thrombogenic (saturated) and anti-thrombogenic fatty acids (MUFA, n-6 and PUFA) and was calculated by applying the following equation:

$$TI = \frac{(14:0 + 16:0 + 18:0)}{(0.5 \times MUFA) + (0.5 \times n-6) + (3 \times n-3) + \frac{n-3}{n-6}}$$

The HH ratio is related to cholesterol metabolism and was calculated according to the following equation:

$$HH = \frac{[(18:1n-9 + 18:2n-6 + 18:3n-3 + 20:5n-3 + 22:5n-3 + 22:6n-3)]}{(14:0 + 16:0)}$$

Analytical Quality Assurance

The analytical quality was controlled for three times by analyzing the standard reference material “Meat Homogenate” (SRM 1546; NIST, Gaithersburg, MD) for lipid content as well as FAP, following the same analytical procedure used in this work. The means determined were always within the certified (or reference) intervals.

Statistical Analysis

For each species, data (lipid content and FAP, expressed as mole%) were arcsin-transformed and analyzed using a two way “between group/within subjects” analysis of variance (ANOVA). “Season of catch” (fall/winter vs spring) was the between-group factor, whereas “State” (raw vs cooked) was the within-subjects (repeated measures) factor. Means were separated at, or below, the 5% probability level using Tukey’s HSD *post hoc* test. All statistical computations were performed using the Statistica® software package (Release 7, 2005; StatSoft Inc., Tulsa, OK, USA).

Results

Polar Lipids Content

The percentages of PoL in raw and cooked fillets of the blue-back fish examined are shown in Table 1. Levels of PoL, calculated as g/100 g of TL (or %PoL) in fresh fillets, ranged from 4.83% (sprat in spring) to 22.8% (sardine in spring). In sardine, these values varied significantly among fishing seasons: they were higher in spring, when the lower amount of TL characterizes this species in this period. The same parameters expressed in absolute terms (mg of PoL/100 g of raw fillet) showed levels between 427 and 652 mg/100 g tissue, with no statistically significant seasonal variation within species, suggesting variability in TL profile during the year.

The cooking process significantly influenced total PoL content (expressed in absolute terms) in all species, due to the loss of water. The %PoL of anchovy increased significantly in both seasons, while in sardine and in horse mackerel this effect was only appreciable in subjects collected in fall.

Fatty Acid Profile of Polar Lipids in Raw Samples

The FAP of PoL fractions of the four species analyzed in fall/winter and in spring is shown in Tables 2 and 3. The n-3 PUFA were the major fraction, with percentages ranging between 42.40% and 57.06%, followed in decreasing order by SFA (26.70-30.85%), MUFA (8.58-13.89%) and n-6 PUFA (2.85-8.28%). The SFA were high in palmitic acid (16:0), with a peak of 35.40% in the spring in sardine; among the MUFA, oleic acid (18:1 n-9) was predominant, with values ranging between 2.90% and 6.75%; ARA (1.07-3.70%) was the most represented n-6 PUFA, while DHA (30.50-45.23%) and EPA (4.67-13.22%) were the major ones among n-3 PUFA. The species that contains the highest PUFA content is horse mackerel although sprat is richer in n-3 PUFA, followed by horse mackerel and anchovy fished in spring.

Seasonal Dynamics of Fatty Acid Profile of Polar Lipids

The SFA, MUFA and PUFA composition of PoL (Tables 2 and 3) generally showed a remarkable pattern of seasonality, reflecting fatty acid fluctuations. Almost all SFA tended to decline from fall to spring, i.e. the 14:0 in sardine (from 3.98% to 0.68%, about 1/6 of the fall value). An exception is the increase of 16:0 content in sardine (from 18.0% in fall to 31.2% in spring) and, to a lesser extent, in anchovy (from 20.39% in fall to 35.40% in spring). However, these changes had no effect on the overall value of SFA, except in sardine (from 30.85% in fall to 41.73% in spring). Significant variations in the percentages of MUFA were found in anchovy, sardine and horse mackerel caught in fall, mainly due to the increase in both 16:1n-7 and 18:1n-9. In these very same species, seasonal changes in fatty acid composition of MUFA were also recorded for NL. Furthermore, fall-caught horse mackerel showed an increase in SFA (especially 16:0), as it can be seen in Table 5 and Table 6.

The percentage of total PUFA, instead, did not change seasonally: in anchovy, sardine and horse mackerel caught in fall, the reduction in DHA content was compensated for by increases in the percentage of EPA or of n-6 PUFA. In fact, anchovy and sardine were richer in n-6 PUFA in fall, while in sprat and in horse mackerel seasonal rates remain unchanged. The levels of n-3 PUFA were significantly higher in spring, except for sprat. In anchovy, this trend reflected that of the major n-3 PUFA (EPA and DHA), while in sardine and horse mackerel it was connected to a marked increase in DHA, which compensated for the significant decline of EPA in spring. A similar pattern was observed

for NL PUFA, where higher levels of n-6 are detected in anchovy caught in fall and increased n-3 in spring horse mackerel (see Table 5 and Table 6).

Effect of Cooking on Fatty Acid Profile of Polar Lipids

Upon cooking, significant changes in percentages of fatty acids, connected to season and species, were frequently observed (see Tables 2 and 3). As was observed for raw samples, the n-3 PUFA fraction was the highest, with percentages ranging from 39.38% to 57.38%, followed by SFA (26.73-38.10%), MUFA (5.93-11.75%) and n-6 PUFA (2.69-7.99%). In particular, cooking seemed to cause a significant reduction in SFA level in spring sardine, mainly due to the remarkable decrease of 16:0; on the contrary, in fall in anchovy, and in spring in horse mackerel, the same fatty acid tended to achieve notably higher values. In sprat and horse mackerel, the quantities of SFA were less influenced by cooking and season, while in anchovy and in sardine the interaction (Sc*St) between season and status was significant. The cooking process induced an important decrease in MUFA content in sardines, particularly those collected in fall; on the contrary, an increase could be observed in anchovy. Cooking caused a significant drop in total PUFA only in the fall anchovy. For n-6 PUFA, the cooking effect seemed to consist of a small decrease in the total concentration in horse mackerel fished in spring and in anchovy and sardine fished in fall, in which the interactions Sc*St are significant for many fatty acids. The effects of the cooking process on n-3 PUFA mainly involved fall anchovy and sardine; in the former species, cooking caused significant loss of DHA and thus a significant decrease in n-3 PUFA; in the latter, all n-3 PUFA, excluding DHA, incurred significant losses, while in spring, EPA and DHA appeared to increase.

NL showed a different behavior towards cooking processes compared to PoL; in each species analyzed, SFA, MUFA and n-6 PUFA from NL are quite stable, or they can be even found concentrated in cooked fillet, due to the water loss. On the contrary, the trend of n-3 PUFA is similar to that of PoL, since drops are recorded in fall anchovy and spring sardines, mainly due to loss in DHA (see Table 5 and Table 6).

Nutritional Indices

The n-6/n-3 PUFA ratios (see Tables 2 and 3) were steadily below the value of 1 (0.05 to 0.16), given the clear predominance of n-3 PUFA. In particular, a seasonal reduction in n-6/n-3 ratio was found in raw anchovy (0.11 in fall to 0.06 in spring), while a less important decrease was shown for horse mackerel (0.16 to 0.12). Cooking finally produced slight but significant increases in n-6/n-3 ratio only in fall anchovy.

Table 4 shows the nutritional profile of PoL of the four species in the two seasonal periods in the raw and cooked states. Within each species, the comparison between seasons did not reveal significant differences in AI and in TI. The values of AI in subjects in the raw state varied from a minimum of 0.21 in horse mackerel to a maximum of 0.56 in sardine, both species caught in spring; in sprat (in both seasons) and in anchovy caught in spring, the TI were in the range 0.11–0.25, respectively. After baking, the values of AI were included in the range 0.23 to 0.59, respectively in spring sprat and in fall sardine. Within each species, comparison between seasons showed a significant decrease of AI in spring anchovy and sardine only after cooking. Finally, the highest HH value observed was 4.49 in spring horse mackerel; the lowest was 1.64 in spring sardine. In raw anchovy and sardine, HH values tended to be slightly higher in fall. In all other cases, the differences between seasons and between raw and cooked fish seemed to be minor.

Discussion

Seasonal Effect on Polar Lipids Content and Fatty Acid Profile

The proportions of PoL (g/100 g of TL) did not differ from those reported in the literature (De Leonardis & Macciola, 2004; Fanni, Linder, & Parmentier, 2004; Shahidi & Miraliakbari, 2006): the highest values of PoL were detected during the “lean period”, when the amount of NL in the flesh decreases in favor of gonadal maturation. Even the amount of PoL agrees by and large with the data from other authors (Kolakowska, Olley & Dunstan, 2006), ranging between 300 and 500 mg/100 g of flesh (see Table 1). The lack of significant seasonal variation in PoL content confirms that the variations in TL detected in the course of the year should be mostly attributable to cyclical variations of NL, as reported by other authors (Fanni, Linder & Parmentier, 2004). As far as we know, this is one of the very

few studies reporting seasonal changes of PoL fatty acids of blue-back fish from the Adriatic Sea; therefore the comparison with literature data is fragmented and incomplete.

Overall, the results of this survey indicate that the FAP of raw PoL exhibits a preponderance of PUFA, especially DHA and EPA. Total PUFA remains substantially constant in all species during the year, while the n-3 PUFA content (except in sprat) tends to rise in spring, especially due to the contribution of DHA. Furthermore, n-6 PUFA show significant seasonal variations only in anchovy and sardine, with lower values in spring compared to fall. These findings are consistent with the results obtained in the study by Passi et al. (2002), where the FAP of PoL were analyzed in sardine and horse mackerel from the central Tyrrhenian Sea, without specifying the season of capture. Other investigations on PoL from *Anguilla anguilla* reported low levels of MUFA and very high levels of PUFA, mostly n-3 (Ciappellano, Erba, Colombo, Testolin & Bolis, 1999). Similarly, in a study of seasonal fluctuations in FAP in three anadromous subspecies (*Salmo trutta caspius*, *S. t. labrax* and *S. t. macrostigma*), Bayır et al. (2010) reported PoL fractions with very high percentages of PUFA and SFA (due to the high levels of 16:0). Seasonal changes in the various groups of fatty acids varied depending on the species considered, in agreement with the lack of homogeneity occurring in changes of PoL in blue-back fish. Given the large number of variables involved, it is difficult to distinguish the origin of PoL fluctuations, especially since in aquatic animals this fraction was found to be susceptible to the dietary quality, although to a lesser extent than NL (De Souza et al., 2008; Standal, Axelson & Aursand, 2010).

On this basis, the seasonal changes we observed in PoL of blue-back fish can be attributed to the different seasonal availability of fatty acids in the aquatic food chain rather than homeoviscous adaptation. On the other hand, high levels of n-3 PUFA, typical in winter or early spring due to the adaptation of aquatic animals at low temperatures, were detected only in horse mackerel.

Effect of Cooking on Fatty Acid Profile

Important changes in the FAP of fish muscle, caused by culinary treatments, consist of considerable loss of DHA and EPA and depend on the cooking method (Larsen, Quek & Eyres, 2010; Türkkan, Cakli & Kilinc, 2008). Baking is considered by many authors the best heat treatment for preservation of all lipid features of fish meat, including the PUFA content and n-3/n-6 ratio (García-Arias et al., 2003); it

has been found to be the best preparation technique with regard to retention of EPA and DHA of TL, maintaining over 80% for both fatty acids (Al-Shagir et al., 2004; Mierke-Klemeyer et al., 2008). The great increase of PoL content in cooked fillets (see Tables 2 and 3) when compared to the raw counterpart is rationally connected to the loss of moisture and storage lipids, subjected to melting or leaching from the flesh with the so-called “cook-out” (Little, Armstrong, & Bergan, 2000).

The heat resistance of many fatty acids found in this study, including n-3 PUFA (especially in spring sardine, sprat and horse mackerel), can be related to some extent of “internal protection” of n-3 fatty acids that prevent the degradation of PUFA during heat treatments. This protective effect may be due to the presence of NL and PoL incorporated in lipoprotein structures, such as biological membranes (Küllenberg, Taylor, Schneider, Massing and Ulrich, 2012), and to the presence of high levels of natural antioxidants (vitamin E, astaxanthin) in the flesh (Gladyshev et al., 2006; Larsen, Quek & Eyres, 2010). From the results of this study (Tables 2 and 3), the response to baking is species-specific, as also reported by other authors (García-Arias et al., 2003; Schneedorferová, Tomčala & Valterová, 2015); it seems to be correlated to the catch season in anchovy and sardine: in baked anchovy caught in fall, significant increments in the percentages of SFA, MUFA and n-6 PUFA were detected, together with a decrease in the global percentage of PUFA (due to significant reductions in n-3 PUFA), and a consequent increase in n-6/n-3 ratio. In baked sardine, significant increments were detectable in the MUFA and n-6 PUFA percentages of subjects caught in fall. These findings agree with those reported in the study of Martelli et al. (2013), in which the FAP was influenced by both catch season and cooking treatments.

Nutritional Indices

The n-6/n-3 ratio has been suggested to be the best index for comparing the relative nutritional value of fish oils (Simopoulos, 1991). The typical Western diet is characterized by a high intake of n-6 PUFA and a low intake of n-3 PUFA. According to the recommendations of nutritional advisers, the attempt is to increase the levels of n-3 PUFA in the diet, such that the n-6/n-3 ratio does not exceed 4.0 (Santos-Silva, Bessa, & Santos-Silva, 2002). Analysis of PoL in the fish species included in this study reported an n-6/n-3 ratio between 0.05 and 0.17 (see Tables 2 and 3), which are values 2-15 fold lower than those reported by Simonetti et al. (2008) in a study of PoL in four freshwater species; these findings

suggest excellent nutritional quality in terms of FAP of PoL in the fish sources investigated, also confirmed by the AI and indices always lower than 1 (between 0.10 and 0.59, among raw and cooked samples). These indices (see Table 4) indicate the potential for stimulating platelet aggregation and are primarily related to a decrease in cardiovascular disease risk; thus, the smaller the values, the greater the protective potential for coronary artery disease (Turan, Sönmez & Kaya, 2007). From the findings of this study, sardine was the species with the highest index values. AI and TI indices were not influenced by seasonality or by cooking, with the exception of cooked sardine and anchovy which showed a decrease of both AI and TI indices in spring compared to the raw state.

The HH index refers to the ratio between the sum of hypercholesterolemic fatty acids and the sum of hypocholesterolemic fatty acids, and indicates the specific effects of fatty acids on the cholesterol metabolism; from a nutritional point of view, high HH values are considered beneficial to human health (Testi, Bonaldo, Gatta & Badiani, 2006). HH index values from this study (means ranging from 1.64 to 4.49, see Table 4) were higher than those obtained in similar investigations on black needle, white needle, mackerel and sardine from a Brazilian market (Fernandes et al., 2014). HH indices tended to be higher in horse mackerel and sprat and showed significant seasonal variability in raw and cooked sardine; furthermore, in every species analyzed, they exhibited a decrease after the cooking process, with the exception of sprat, for which baking increased the index.

Due to the very low n-6/n-3 fatty acid ratios and the highly favorable nutritional indices found in the blue-back fish analyzed in this study, the PoL qualities of the four species examined may play an important role in the human diet, helping to increase significantly the n-3 fatty acid intake and improving consumer health (Hossain, Hosokawa & Takahashi, 2009; Murakawa et al., 2007; Ulbricht & Southgate, 1991; Taylor et al., 2010). Furthermore, the existence of such high levels of EPA and DHA could make blue-back fish PoL biologically valuable in the food industry for enrichment and production of functional foods with higher nutritional value (Burri, Hoem, Banni, & Berge, 2012). This study contributes to finding new commercialization strategies for these species, possibly as value-added products. This study promotes the importance of marine PoL for the human diet, as a valuable source of PUFA, and highlights the nutritional quality of four popular blue-back species from the Adriatic Sea (*E. encrasicolus*, *S. pilchardus*, *S. sprattus* and *T. trachurus*). This manuscript presents, for the first time,

data on the influences of catch season and cooking treatments on the FAP of PoL, and therefore on the nutritional quality of the species analyzed. Sprat was the only species not affected by seasonal variations or by cooking processes. For the other species, especially sardine and anchovy, it is difficult to delineate the best season for capture and consumption because, although the data indicate a higher level of PUFA (in particular n-3) in the specimens caught in spring, the nutritional indices of cooked samples were generally higher in those collected in fall.

The increased accessibility of marine lipids during recent years has opened up new possibilities for the use of PoL not only as a superior nutritional source of n-3 PUFA, but also for use in the pharmaceutical, cosmetic and functional food industries (Burri, Hoem, Banni, & Berge, 2012). Since the only available source of n-3 PUFA comes at present from marine lipids, there is a need to scale-up productions of these compounds and to identify new affordable resources, natural or synthetic (Løvaas et al., 2006); future investigations should explore the PoL content in other marine species, krill, fish roe, fish by-products (viscera and skin) and even in non-animal marine sources such as microalgae, in order to meet the growing demands of the market. In addition, in order to complete the assessment of the role of PoL in food manufacturing and in pharmaceutical and cosmetic applications, other important aspects which need to be considered and deepened are the bioavailability of PoL from different sources, and the identification of reliable physiological markers to confirm and determine their bioactivity.

Acknowledgments

This study was founded by a grant from Alma Mater Studiorum – University of Bologna (“RFO” 2006). Thanks are due to Micaela Fabbri for her contribution to sample analysis.

Conflict of interest

The authors declare that there is no conflict of interest regarding the publication of this article.

References

Adkins Y. & Kelley D. S. (2010) Mechanisms underlying the cardioprotective effects of omega-3 polyunsaturated fatty acids. *J Nutr Biochem* **21**:781–792. <https://doi.org/10.1016/j.jnutbio.2009.12.004>

420 Al-Saghir S., Thurner K., Wagner K.-H., Frisch, G., Luf, W., Razzazi-Fazeli, E. & Elmadfa, I.
 421 (2004) Effects of different cooking procedures on lipid quality and cholesterol oxidation of farmed
 422 salmon fish (*Salmo salar*). J Agric Food Chem **52** (16), 5290–5296. DOI: 10.1021/jf0495946

423 Bayır A., Sirkecioğlu A. N., Aras N .M., Aksakal E., Haliloğlu, H. I. & Bayır M. (2010) Fatty acids
 424 of neutral and phospholipids of three endangered trout: *Salmo trutta caspius* Kessler, *Salmo trutta*
 425 *labrax* Pallas and *Salmo trutta macrostigma* Dumeril. Food Chem **119**:1050–1056.
 426 <https://doi.org/10.1016/j.foodchem.2009.07.064>

427 Blank M.L., Cress E.A., Smith Z.L. & Snyder F. (1992) Meats and fish consumed in the American
 428 diet contain substantial amounts of ether-linked phospholipids. J Nutr **122**:1656–1661.
 429 <https://doi.org/10.1093/jn/122.8.1656>

430 Brenner R.R. (1977) Regulatory Function of $\Delta 6$ Desaturase — Key Enzyme of Polyunsaturated
 431 Fatty Acid Synthesis. In: Bazán N.G., Brenner R.R., Giusto N.M. (eds) Function and Biosynthesis of
 432 Lipids. Advances in Experimental Medicine and Biology, vol 83. Springer, Boston, MA.
 433 https://doi.org/10.1007/978-1-4684-3276-3_8

434 Burri L., Hoem N., Banni S. & Berge K. (2012) Marine Omega-3 phospholipids: metabolism and
 435 biological activities. Int J Mol Sci **13**:15401–15419. <https://doi.org/10.3390/ijms131115401>

436 Christie W. W. (1989) Gas Chromatography and Lipids: A Practical Guide. The Oily Press Ltd,
 437 1989, The Hannah Research Institute, Ayr, Scotland KA6 5HL, vol. 1, 307 pp.

438 Ciappellano S., Erba D., Colombo A., Testolin G. & Bolis C. G. (1999) Composition and
 439 nutritional aspects of flesh eel (*Anguilla anguilla*) of different size subjects. Rivista di Scienze
 440 dell'Alimentazione **28**(4):431-437. ISSN : 0391-4887.

441 Cordier M., Brichon G., Weber J.-M. & Zwingelstein G. (2002) Changes in the fatty acid
 442 composition of phospholipids in tissues of farmed sea bass (*Dicentrarchus labrax*) during an annual
 443 cycle. Roles of environmental temperature and salinity. Comp Biochem Physiol Part B **133**:281–288.
 444 [https://doi.org/10.1016/S1096-4959\(02\)00149-5](https://doi.org/10.1016/S1096-4959(02)00149-5)

445 De Leonardis A. & Macciola V. (2004) A study on the lipid fraction of Adriatic sardine fillets
 446 (*Sardina pilchardus*). *Nahrung - Food* **48**:209–212. <https://doi.org/10.1002/food.200300408>

447 De Souza N. E., Stevanato F. B., Garcia E. E., Visentainer J. E. L., Zara R. F. & Visentainer J. V.
 448 (2008) Supplemental dietary flaxseed oil affects both neutral and phospholipid fatty acids in cultured
 449 tilapia. *Eur J Lipid Sci Technol* **110**:709–713. <https://doi.org/10.1002/ejlt.200700245>

450 Fanni J., Linder M. & Parmentier M. (2004) Lipides polaires marins. Oléagineux, Corps Gras,
 451 *Lipides* **11**:142–145. <https://doi.org/http://dx.doi.org/10.1051/ocl.2004.0142>

452 Fernandes C. E., da Silva Vasconcelos M. A., De Almeida Ribeiro M., Asfora Sarubbo L.,
 453 Alvachian Cardoso Andrade, S. & Bibiano De Melo Filho A. (2014) Nutritional and lipid profiles in
 454 marine fish species from Brazil. *Food Chem* **160**:67–71.
 455 <https://doi.org/10.1016/j.foodchem.2014.03.055>

456 Folch J., Lees M. & Sloane Stanley G. H. (1953) A simple method for the isolation and purification
 457 of total lipids from animal tissues. *J Biol Chem* **226**:497–509.

458 Gabbs M., Leng S., Devassy J. G., Monirujjaman M., & Aukema H. M. (2015) Gabbs, M.,
 459 Leng, S., Devassy, J. G., Monirujjaman, M., & Aukema, H. M. (2015). Advances in our
 460 understanding of oxylipins derived from dietary PUFAs. *Advances in Nutrition*, **6**:513–540.
 461 <https://doi:10.3945/an.114.007732>

462 García-Arias M. T., Álvarez Pontes E., García-Linares M. C., García-Fernández M. C. & Sánchez-
 463 Muniz F. J. (2003) Cooking-freezing-reheating (CFR) of sardine (*Sardina pilchardus*) fillets. Effect of
 464 different cooking and reheating procedures on the proximate and fatty acid compositions. *Food Chem*
 465 **83**:349–356. [https://doi.org/10.1016/S0308-8146\(03\)00095-5](https://doi.org/10.1016/S0308-8146(03)00095-5)

466 Gladyshev M. I., Sushchik N. N., Gubanenko G. A., Demirchieva S M. & Kalachova, G. S. (2006)
 467 Effect of way of cooking on content of essential polyunsaturated fatty acids in muscle tissue of
 468 humpback salmon (*Oncorhynchus gorbuscha*). *Food Chem* **96**:446–451.
 469 <https://doi.org/10.1016/j.foodchem.2005.02.034>

470 Guo Z., Vikbjerg A. F. & Xu X. (2005) Enzymatic modification of phospholipids for functional
 471 applications and human nutrition. *Biotechnol Adv* **23**:203–259.
 472 <https://doi.org/10.1016/j.biotechadv.2005.02.001>

473 Henna Lu F. S., Nielsen N. S. & Timm-Heinrich M, Jacobsen C (2011) Oxidative stability of
 474 marine phospholipids in the liposomal form and their applications. *Lipids* **46**:3–23.
 475 <https://doi.org/10.1007/s11745-010-3496-y>

476 Hossain Z., Hosokawa M. & Takahashi K. (2009) Growth inhibition and induction of apoptosis of
 477 colon cancer cell lines by applying marine phospholipid. *Nutr Cancer* **61**:123–130.
 478 <https://doi.org/10.1080/01635580802395725>

479 Kaçar S. & Başhan M. (2015) Seasonal variations in the fatty acid composition of phospholipid and
 480 triacylglycerol in gonad and liver of *Mastacembelus simack*. *J Am Oil Chem Soc* **92**:1313–1320.
 481 <https://doi.org/10.1007/s11746-015-2692-6>

482 Kolakowska A., Olley J. & Dunstan G.A. (2006) Fish lipids. In: Chemical, Biological, and
 483 Functional Aspects of Food Lipids, 2nd Edition, Zdzislaw Z. E. Sikorski, Anna Kolakowska. CRC
 484 Press. Chapter 15, p 273-312. ISBN 9781439802373.

485 Küllenberg D., Taylor L. A., Schneider M. & Massing U. (2012) Health effects of dietary
 486 phospholipids. *Lipids Health Dis* **11**:1–16. <https://doi.org/10.1186/1476-511X-11-3>

487 Larsen D., Quek S. Y. & Eyres L. (2010) Effect of cooking method on the fatty acid profile of New
 488 Zealand King Salmon (*Oncorhynchus tshawytscha*). *Food Chem* **119**:785–790.
 489 <https://doi.org/10.1016/j.foodchem.2009.07.037>

490 Little, S. O. , Armstrong, S. G. & Bergan J. G. (2000) Fatty acids in foods and their health
 491 implications. In: Chow CK (ed) Factors affecting stability and nutritive value of fatty acids: culinary
 492 practices, 2nd ed. Marcel Dekker Inc., New York, USA, pp 427–437

493 Loftsson T., Ilievska B., Asgrimsdottir G. M., Ormarsson O. T., Stefansson E. (2016) Fatty acids
 494 from marine lipids: Biological activity, formulation and stability. *J Drug Deliv Sci Technol* **34**:71–75.
 495 <https://doi.org/10.1016/j.jddst.2016.03.007>

496 Løvaas E. (2006) Marine phospholipids (MPL): Resources, application and markets. In: Luten JB,
 497 Jacobsen C, Bekaert K, Saebø A, Oehlenschläger (Ed.) *Seafood Research from Fish to Dish*, pp. 17-28.
 498 Wageningen Academic Publishers, The Netherland. ISBN: 978-90-8686-005-0.
 499 <https://doi.org/10.3920/978-90-8686-581-9>

500 Martelli R., Dalle Zotte A., Bonelli A., Lupi P., Franci O. & Parisi G. (2013) Macronutrient and
 501 fatty acid profiles of meagre (*Argyrosomus regius*) fillets as influenced by harvesting time and boiling.
 502 *Ital J Anim Sci* **12**:538–545. <https://doi.org/10.4081/ijas.2013.e88>

503 Mierke-Klemeyer S., Larsen R., Oehlenschläger J., Maehre H., Elvevoll E. O., Bandarra N. M.,
 504 Parreira, R., Andrade A. M., Nunes, M. L., Schram, E. & Luten, J. (2008) Retention of health-related
 505 beneficial components during household preparation of selenium-enriched African catfish (*Clarias*
 506 *gariepinus*) fillets. *Eur Food Res Technol* **227**:827–833. <https://doi.org/10.1007/s00217-007-0793-7>

507 Mika A., Swiezewska E. & Stepnowski P. (2016) Polar and neutral lipid composition and fatty
 508 acids profile in selected fish meals depending on raw material and grade of products. *LWT - Food Sci*
 509 *Technol* **70**:199–207. <https://doi.org/10.1016/j.lwt.2016.02.051>

510 Murakawa K., Fukunaga K., Tanouchi M., Hosokawa M., Hossain Z. & Takahashi K. (2007)
 511 Therapy of myeloma *in vivo* using marine phospholipid in combination with *Agaricus blazei* Murill as
 512 an immune respond activator. *J Oleo Sci* **56**:179–188. <https://doi.org/10.5650/jos.56.179>

513 Parmentier M., Al Sayed Mahmoud C., Linder M. & Fanni J. (2007) Polar lipids: n-3 PUFA
 514 carriers for membranes and brain. Nutritional interest and emerging processes. *OCL - Ol Corps Gras*
 515 *Lipides* **14**:224–229. <https://doi.org/10.1684/ocl.2007.0127>

516 Passi S., Cataudella S., Di Marco P., De Simone F. & Rastrelli L. (2002) Fatty Acid Composition
 517 and antioxidant levels in muscle tissue of different Mediterranean marine species of fish and shellfish.
 518 J Agric Food Chem **50**:7314–7322. <https://doi.org/10.1021/jf020451y>
 519

520 Pirini M., Testi S., Ventrella V., Pagliarani A. & Badiani A. (2010) Blue-back fish: Fatty acid
 521 profile in selected seasons and retention upon baking. Food Chem **123**:306–314.
 522 <https://doi.org/10.1016/j.foodchem.2010.04.036>

523 Santos-Silva J., Bessa R. J. B. & Santos-Silva F. (2002) Effect of genotype, feeding system and
 524 slaughter weight on the quality of light lambs. II. Fatty acid composition of meat. Livest Prod Sci
 525 **77**:187–194. [https://doi.org/10.1016/S0301-6226\(02\)00059-3](https://doi.org/10.1016/S0301-6226(02)00059-3)

526 Schneedorferová I., Tomčala A., Valterová I. (2015) Effect of heat treatment on the n-3/n-6 ratio
 527 and content of polyunsaturated fatty acids in fish tissues. Food Chem **176**:205–211.
 528 <https://doi.org/10.1016/j.foodchem.2014.12.058>

529 Shahidi F. & Miraliakbari H. (2006) Marine oils: compositional characteristics and health effects.
 530 In: Shahidi F. (ed) Nutraceutical and specialty lipids and their co-products. CRC Press, chapter 15, p
 531 227-247. ISBN 9781574444995.

532 Simopoulos A. P. (1991) Omega-3 fatty acids in health and disease and in growth and
 533 development. Am J Clin Nutr **54**:438–63. <https://doi.org/10.1093/ajcn/54.4.771>

534 Simopoulos A. P. (2006) Evolutionary aspects of diet, the omega-6/omega-3 ratio and genetic
 535 variation: nutritional implications for chronic diseases. Biomed Pharmacother **60**:502–507.
 536 <https://doi.org/10.1016/j.biopha.2006.07.080>

537 Simonetti M. S., Blasi F., Bosi A., Maurizi A., Cossignani L., Damiani P. (2008) Stereospecific
 538 analysis of triacylglycerol and phospholipid fractions of four freshwater fish species: *Salmo trutta*,
 539 *Ictalurus punctatus*, *Ictalurus melas* and *Micropterus salmoides*. Food Chem **110**: 199–206.
 540 <https://doi.org/10.1016/j.foodchem.2008.01.033>

541 Standal I. B., Axelson D. E. & Aursand M. (2010) ^{13}C NMR as a tool for authentication of different
542 gadoid fish species with emphasis on phospholipid profiles. Food Chem **121**:608–615.
543 <https://doi.org/10.1016/j.foodchem.2009.12.074>

544 Taylor L. A., Pletschen L., Arends J., Unger C. & Massing U. (2010) Marine phospholipids-a
545 promising new dietary approach to tumor-associated weight loss. Support Care Cancer **18**:159–170.
546 <https://doi.org/10.1007/s00520-009-0640-4>

547 Testi S., Bonaldo A., Gatta P. P. & Badiani A. (2006) Nutritional traits of dorsal and ventral fillets
548 from three farmed fish species. Food Chem **98**:104–111.
549 <https://doi.org/10.1016/j.foodchem.2005.05.053>

550 Turan H., Sönmez G. & Kaya Y. (2007) Fatty acid profile and proximate composition of the
551 thornback ray (*Raja clavata*, L. 1758) from the Sinop coast in the Black Sea. J Fish Sci **1**:97–103.
552 <https://doi.org/10.3153/jfscom.2007012>

553 Türkkan A. U., Cakli S. & Kilinc B. (2008) Effects of cooking methods on the proximate
554 composition and fatty acid composition of seabass (*Dicentrarchus labrax*, Linnaeus, 1758). Food
555 Bioprod Process **86**:163–166. <https://doi.org/10.1016/j.fbp.2007.10.004>

556 Ulbricht T. L. V. & Southgate D. A. T. (1991) Coronary heart disease: seven dietary factors.
557 Lancet **338**:985–992. [https://doi.org/10.1016/0140-6736\(91\)91846-M](https://doi.org/10.1016/0140-6736(91)91846-M)

558 Varljen J., Baticic L., Sincic-Modric G., Obersnel V. & Kapovic M. (2004) Composition and
559 seasonal variation of fatty acids of *Diplodus vulgaris* L. from the Adriatic sea. J Am Oil Chem Soc
560 **81**:759–763. <https://doi.org/10.1007/s11746-004-0975-7>

561

Table 1 Polar lipid (PoL) content in blue-back fish fillets from the four species, in the raw and cooked state, and from different seasons

Fish species	Season	PoL (g/100g TL)		PoL (mg/100g flesh)	
		Raw	Cooked	Raw	Cooked
Anchovy	<i>Fall</i>	y11.1 ± 2.25	x18.9 ± 1.61	y486 ± 24	x1028 ± 174
	<i>Spring</i>	y7.71 ± 1.31	x15.1 ± 2.12	y442 ± 56	x977 ± 81
Sardine	<i>Fall</i>	y6.70 ± 1.38b	x11.9 ± 1.46b	y486 ± 58	x827 ± 46
	<i>Spring</i>	22.8 ± 2.21a	24.9 ± 3.65a	y652 ± 68	x927 ± 71
Sprat	<i>Winter</i>	14.1 ± 4.22	25.9 ± 14.4	y482 ± 130	x967 ± 154a
	<i>Spring</i>	4.83 ± 0.43	6.08 ± 0.73	539 ± 31	662 ± 30b
Horse mackerel	<i>Fall</i>	y20.5 ± 5.41	x29.6 ± 3.58	y427 ± 34	x679 ± 122
	<i>Spring</i>	20.4 ± 4.56	25.0 ± 6.93	y595 ± 37	x845 ± 86

Values are means of four determinations ± Std Dev; values in the same row, with trait preceded by different letters (x, y), differ significantly ($p < 0.05$) for state (raw or cooked); values within a column, followed by different letters (a, b), differ significantly ($p < 0.05$) for catch season.

569 **Table 2** Content of selected fatty acids (mole%) in polar lipids from skin-on fillets of anchovy and
570 skinned sardine in the raw (R) and cooked (C) state (St)

Fatty acids ^{A, B, C}	St	Anchovy		Statistical analysis				Sardine		Statistical analysis			
		<i>Fall</i>	<i>Spring</i>	RSD	Sc	St	Sc*St	<i>Fall</i>	<i>Spring</i>	RSD	Sc	St	Sc*St
14:0	R	3.00a	1.29b	0.940	***	*	n.s.	x3.98a	0.68b	0.267	***	**	*
	C	3.30a	1.41b					y2.47a	0.65b				
16:0	R	x19.54b	25.83a	0.762	***	*	***	y20.39b	x35.40a	2.078	**	n.s.	***
	C	x23.23	24.51					x28.68	y27.95				
18:0	R	y5.46a	4.36b	0.384	***	n.s.	**	5.83	x5.26	0.451	**	*	**
	C	x6.60a	3.79b					6.21a	y3.76b				
Σ SFA^D	R	y29.62	32.65	1.253	n.s.	*	**	30.85b	x41.73a	2.773	n.s.	n.s.	**
	C	x35.10a	30.84b					38.10	y32.72				
16:1n-7	R	2.71a	1.21b	0.092	***	*	n.s.	x3.26a	0.64b	0.181	***	**	**
	C	2.98a	1.28b					y1.99a	0.62b				
18:1n-9	R	y3.71a	x2.90b	0.069	***	n.s.	***	x6.19a	3.19b	0.524	***	**	n.s.
	C	x4.16a	y2.51b					y4.16a	2.73b				
18:1n-7	R	x2.94	x3.23	0.082	n.s.	n.s.	***	x2.66	x2.85	0.234	n.s.	**	n.s.
	C	x3.26	y2.90					y2.04	y2.17				
Σ MUFA^E	R	y9.84a	x7.70b	0.167	***	n.s.	***	x13.89a	6.90b	0.901	***	**	*
	C	x11.16a	y6.93b					y8.94	5.93b				
18:2n-6	R	1.15a	0.54b	0.049	***	n.s.	n.s.	x1.29a	0.96b	0.083	*	***	**
	C	1.10a	0.53b					y0.79	0.88				
20:4n-6	R	2.20a	1.29b	0.032	***	n.s.	**	1.59a	1.07b	0.200	***	n.s.	n.s.
	C	2.12a	1.35b					1.64a	1.13b				
22:5n-6	R	x1.52a	0.96b	0.033	***	***	**	1.11	1.07	0.112	n.s.	n.s.	n.s.
	C	y1.28a	0.93b					1.12	1.31				
Σ n-6 PUFA^F	R	x5.76a	3.33b	0.053	***	***	**	x5.01a	3.78b	0.216	***	**	**
	C	y5.34a	3.29b					y4.08	3.91				
18:3n-3	R	0.66a	0.26b	0.023	***	n.s.	*	x0.80a	0.43b	0.046	**	***	***
	C	0.61a	0.29b					y0.34	0.42				
18:4n-3	R	x0.69a	0.26b	0.025	***	**	***	x1.02a	0.22b	0.084	***	***	***
	C	y0.53a	0.30b					y0.24	0.23				
20:5n-3	R	8.52b	y10.15a	0.276	***	**	***	x8.26a	y4.67b	0.525	***	n.s.	***
	C	7.79b	x12.45a					y5.78	x5.88				
22:5n-3	R	x0.98a	0.86b	0.028	*	***	**	x1.24a	0.67b	0.051	***	***	***
	C	y0.83	0.83					y0.62	0.74				
22:6n-3	R	x38.39	42.41	0.911	***	***	***	30.50	37.71	4.221	**	*	n.s.
	C	y32.74	42.71a					32.08b	47.23a				
Σ n-3PUFA^G	R	x49.49	54.00	1.140	***	*	***	42.40	44.07	4.681	*	n.s.	*
	C	y42.71	56.70a					39.38b	54.82a				
Σ PUFA^H	R	x55.76	57.56	1.221	**	**	***	47.96	48.18	4.778	*	n.s.	*
	C	y48.44b	60.18a					43.74b	58.99a				
Unknown	R	y8.53a	4.47b	0.207	***	*	*	11.88	5.18	1.699	***	n.s.	n.s.
	C	x9.48a	4.20b					11.85a	4.30b				
n-6/n-3	R	x0.12a	0.06b	0.002	***	n.s.	**	0.12	0.09	0.007	**	*	n.s.
	C	y0.13a	0.06b					0.10	0.07				

571

572 *Values are means of four determinations (each one in duplicate on pooled samples) per catch season*
573 *and state; means within a column and trait preceded by different letters (x, y) differ significantly ($p \leq$*
574 *0.05); means in the same row followed by different letters (a, b) differ significantly ($p \leq 0.05$); RSD =*
575 *residual standard deviation; Sc = catch season; A: includes 15:0 and 17:0; B: includes 20:1n-11, 20:1n-*
576 *9, 20:1n-7, 22:1n-11 and 22:1n-9; C: includes 18:3, 20:2, 20:3 and 22:4n-6; D: includes 16:4n-3 and*
577 *20:4n-3; E: includes 16:2n-4; *** $p \leq 0.001$; ** $p \leq 0.01$; * $p \leq 0.05$; n.s. = not significant.*

578

579 **Table 3** Content of selected fatty acids (mole%) in polar lipids from skinned fillets of sprat and horse
580 mackerel in the raw (R) and cooked (C) state (St)

Fatty acids ^{A,B,C}	St	Sprat		Statistical analysis				Horse mackerel		Statistical analysis			
		Winter	Spring	RSD	Sc	St	Sc*St	Autum	Spring	RSD	Sc	St	Sc*St
14:0	R	1.02	y1.31	0.074	**	**	**	0.42	0.38	0.037	n.s.	n.s.	n.s.
	C	0.97b	x1.84a					0.33	0.38				
16:0	R	22.53	22.29	1.129	n.s.	n.s.	n.s.	17.07	y16.74	1.121	n.s.	*	*
	C	22.86	21.70					17.78	x20.79				
18:0	R	3.49	x3.47	0.169	n.s.	***	**	8.51	8.67	0.558	n.s.	n.s.	n.s.
	C	3.36	y2.28					8.30	9.00				
Σ SFA^D	R	27.96	27.95	1.182	n.s.	n.s.	n.s.	27.07	26.70	1.737	n.s.	*	n.s.
	C	28.06	26.73					27.33	31.19				
16:1n-7	R	1.44	y1.78	0.072	**	*	***	1.21a	0.77b	0.077	***	n.s.	n.s.
	C	1.27b	x2.21a					1.04a	0.75b				
18:1n-9	R	4.83	y6.25	0.170	*	*	**	x6.75	5.74	0.279	n.s.	*	*
	C	4.70	x7.05					y5.80	5.59				
18:1n-7	R	1.96	x2.18	0.094	n.s.	**	***	2.35a	1.57b	0.176	***	n.s.	n.s.
	C	2.04	y1.43					2.32a	1.56b				
20: n-9	R	0.20	y0.30	0.024	n.s.	*	n.s.						
	C	0.22	x0.38										
22:1n-11	R	0.20	y0.33	0.057	n.s.	**	**						
	C	0.20	x0.58										
Σ MUFA^E	R	8.69	10.93	0.303	*	n.s.	*	11.00a	8.58b	0.521	**	n.s.	n.s.
	C	8.54	11.75					9.81	8.44				
18:2n-6	R	0.76	x0.72	0.018	n.s.	**	**	1.02	0.81	0.073	n.s.	n.s.	n.s.
	C	0.78	y0.61					0.92	0.80				
20:4n-6	R	1.15	1.11	0.072	*	n.s.	n.s.	3.70	2.87	0.098	*	n.s.	n.s.
	C	1.25a	1.08b					3.68	2.72				
22:5n-6	R	0.76	0.65	0.046	n.s.	n.s.	n.s.	1.97	2.17	0.128	n.s.	*	n.s.
	C	0.72	0.68					1.87	1.97				
Σ n-6 PUFA^F	R	3.01	2.85	0.065	n.s.	n.s.	*	8.28	x7.16	0.152	*	**	n.s.
	C	3.12	2.69					7.99	y6.76				
18:3n-3	R	0.49	x0.48	0.010	n.s.	*	*	0.22	0.20	0.027	n.s.	n.s.	n.s.
	C	0.49	y0.45					0.22	0.23				
18:4n-3	R	0.50	y0.47	0.026	n.s.	**	***	0.12	0.12	0.009	n.s.	n.s.	n.s.
	C	0.48	x0.67					0.12	0.12				
20:5n-3	R	10.62	13.22	0.675	n.s.	*	n.s.	8.38	7.43	0.302	*	n.s.	n.s.
	C	11.70	14.69					8.93	7.39				
22:5n-3	R	0.70	0.70	0.032	n.s.	n.s.	n.s.	3.00	2.54	0.114	**	*	*
	C	0.67	0.72					2.97a	2.20b				
22:6n-3	R	44.46	40.21	1.661	*	n.s.	n.s.	38.02b	45.23a	2.198	**	n.s.	n.s.
	C	43.70	39.64					38.13	41.58				
Σ n-3 PUFA^G	R	57.06	55.50	1.273	n.s.	n.s.	n.s.	49.62b	55.38a	2.490	**	n.s.	n.s.
	C	57.38	56.50					50.24	51.51				
Σ PUFA^H	R	60.48	58.72	1.273	n.s.	n.s.	n.s.	58.23	62.83	2.585	n.s.	n.s.	n.s.
	C	60.81	59.52					58.55	58.42				

Unknown	R	2.87	x2.40	0.162	n.s.	**	n.s.	3.70a	1.89b	0.599	***	n.s.	n.s.
	C	2.59	y2.00					4.30a	1.95b				
n-6/n-3	R	0.05	0.05	0.001	n.s	n.s	n.s.	0.17a	0.13b	0.004	**	n.s.	n.s.
	C	0.05	0.05					0.16	0.13				

Values are means of four determinations (each one in duplicate on pooled samples) per catch season and state; means within a column and trait preceded by different letters (x, y) differ significantly ($p \leq 0.05$); means in the same row followed by different letters (a, b) differ significantly ($p \leq 0.05$); RSD = residual standard deviation; Sc = catch season; A: includes 15:0 and 17:0; B: includes 20:1n-11, 20:1n-7 and 22:1n-9; C: includes 18:3, 20:2, 20:3 and 22:4n-6; D: includes 16:4n-3 and 20:4n-3; E: includes 16:2n-4; *** $p \leq 0.001$; ** $p \leq 0.01$; * $p \leq 0.05$; n.s. = not significant.

Table 4 Seasonal variation of nutritional quality indices AI (atherogenicity index), TI (thrombogenicity index) and HH (hypocholesterolemic/hypercholesterolemic fatty acid ratio) in polar lipids in the raw (R) and cooked (C) state (St)

Fish species	St	AI		TI		HH	
		Fall/Winter	Spring	Fall/Winter	Spring	Fall/Winter	Spring
Anchovy	R	y0.38±0.02	0.38±0.03	0.14±0.02	y0.14±0.02	x3.20±0.05	2.62±0.04
	C	x0.49±0.02a	0.36±0.04b	0.19±0.03	x0.25±0.04	y2.22±0.02	2.39±0.02
Sardine	R	y0.47±0.02	x0.56±0.04	0.17±0.02	x0.22±0.03	x2.46±0.04a	y1.64±0.04b
	C	x0.59±0.04a	y0.38±0.02b	0.17±0.02	y0.14±0.02	y1.75±0.03a	x2.53±0.02b
Sprat	R	0.31±0.02	0.31±0.02	0.11±0.02	0.11±0.02	3.25±0.06	2.97±0.05
	C	0.31±0.03	0.33±0.02	0.13±0.02	0.11±0.02	3.26±0.06	3.31±0.05
Horse mackerel	R	0.22±0.03	0.21±0.02	0.13±0.02	0.12±0.02	4.04±0.03	x4.49±0.05
	C	0.23±0.02	0.24±0.02	0.10±0.02	0.15±0.03	3.88±0.02	y3.38±0.06

Values are means of four determinations $\pm$ Std Dev. Means within a column and trait preceded by different letters (x, y) differ significantly ($p \leq 0.05$); means in the same row followed by different letters (a, b) differ significantly ($p \leq 0.05$).

597 **Table 5** Content of selected fatty acids (mole%) in neutral lipids from skin-on fillets of
598 ANCHOVY and skinned SARDINE at raw (R) and cooked (C) state.

Fatty acids ^{A,B,C}	St	Anchovy		Statistical analysis				Sardine		Statistical analysis			
		Fall	Spring	RSD	Sc	St	Sc*St	Fall	Spring	RSD	Sc	St	Sc*St
14:0	R	y6.90b	8.54a	0.257	***	n.s.	**	6.93a	4.75b	0.267	***	**	*
	C	x7.88	8.06					6.95	7.06				
16:0	R	19.39	19.68	1.316	n.s.	n.s.	n.s.	21.69	y20.89	2.078	**	n.s.	***
	C	21.62	20.24					22.96	x24.06				
18:0	R	4.97	3.91	0.334	**	*	n.s.	y5.13a	4.58b	0.451	**	*	**
	C	5.48	4.37					x6.09a	4.61b				
Σ SFA^D	R	33.66	34.07	1.729	n.s.	*	n.s.	35.86	y32.24	2.773	n.s.	n.s.	**
	C	37.71	34.56					38.31	x38.05				
16:1n-7	R	y6.08b	9.37a	0.302	***	n.s.	**	5.50a	y3.08b	0.181	***	**	**
	C	x6.89b	9.03a					5.43	x4.06				
18:1n-9	R	y4.73	5.32	0.183	n.s.	**	n.s.	8.24	7.38	0.524	***	**	n.s.
	C	x5.20	5.63					8.71	7.84				
18:1n-7	R	2.79b	3.35a	0.116	***	*	n.s.	y2.76	2.07	0.234	n.s.	**	n.s.
	C	3.01b	3.38a					x3.27a	2.08b				
Σ MUFA^E	R	y14.79b	19.52a	0.471	***	*	**	y18.38a	y14.49b	0.901	***	**	*
	C	x16.26b	19.16a					x19.33a	x16.42b				
18:2n-6	R	1.54a	1.18b	0.052	***	*	n.s.	1.61	1.64	0.083	*	***	**
	C	1.61a	1.23b					1.66	1.72				
20:4n-6	R	1.90a	0.90b	0.085	***	n.s.	n.s.	1.25	0.92	0.200	***	n.s.	n.s.
	C	1.74a	0.92b					1.21	0.76				
22:5n-6	R	x1.09a	0.43b	0.054	***	*	**	0.72	x0.83	0.112	n.s.	n.s.	n.s.
	C	y0.90a	0.45b					0.72	y0.64				
Σ n-6 PUFA^F	R	5.56a	3.13b	0.215	***	n.s.	n.s.	5.44	5.00	0.216	***	**	**
	C	5.33a	3.26b					5.51a	4.71b				
18:3n-3	R	1.28	1.11	0.038	**	n.s.	n.s.	1.21	y1.26	0.046	**	***	***
	C	1.29a	1.08b					1.12	x1.47				
18:4n-3	R	1.76b	2.66a	0.392	***	n.s.	n.s.	1.68	1.78	0.084	***	***	***
	C	1.70b	2.53a					1.46	1.86				
20:5n-3	R	12.12b	15.88a	0.606	***	*	n.s.	10.35	8.19	0.525	***	n.s.	***
	C	10.64b	15.90a					9.76a	7.14b				
22:5n-3	R	x1.27a	0.80b	0.071	***	n.s.	**	1.47	x1.11	0.051	***	***	***
	C	y1.06a	0.86b					1.44a	y0.91b				
22:6n-3	R	x20.46a	13.04b	0.995	***	**	**	17.99b	x26.60a	4.221	**	*	n.s.
	C	y16.12a	13.32b					17.52	y19.51				
Σ n-3 PUFA^G	R	x37.34	33.89	1.549	n.s.	**	**	33.42	x39.59	4.681	*	n.s.	*
	C	y31.19	34.17					31.95	y31.43				
Σ PUFA^H	R	x43.65a	37.73b	1.575	*	**	**	39.65	x45.44	4.778	*	n.s.	*
	C	y37.42	38.11					38.35	y36.92				
Unknown	R	7.74	8.36	0.007	***	*	n.s.	x6.67	8.56	0.007	**	*	n.s.
	C	8.41	8.00					y4.71b	9.20a				
n-6/n-3	R	y0.15a	0.09b	0.821	n.s.	n.s.	n.s.	0.16	y0.13	1.699	***	n.s.	n.s.
	C	x0.17a	0.10b					0.17	x0.15				

599

600 ^A Values are means of 4 determinations (each one in duplicate on pooled samples) per season
601 of catch and state; ^B Means within a column and trait preceded by different letters (x,y) differ
602 significantly ($P \leq 0.05$); means on the same row followed by different letters (a,b) differ
603 significantly ($P \leq 0.05$); ^C RSD = Residual Standard Deviation; Sc = Season of catch; St = State;
604 ^D Includes: 15:0 and 17:0; ^E Includes: 20:1n-11; 20:1n-9; 20:1n-7; 22:1n-11 and 22:1n-9; ^F
605 Includes: 18:3, 20:2, 20:3 and 22:4n-6; ^G Includes: 16:4 and 20:4n-3; ^H Includes: 16:2n-4;
606 *** $P \leq 0.001$; ** $P \leq 0.01$; * $P \leq 0.05$; n.s. = not significant.

607

608 **Table 6** Content of selected fatty acids (mole%) in neutral lipid from skinned fillets of SPRAT
609 and HORSE MACKEREL at raw (R) and cooked (C) state.

Fatty acids A,B,C	St	Sprat		Statistical analysis				Horse mackerel		Statistical analysis			
		Winter	Spring	RSD	Sc	St	Sc*St	Autum	Spring	RSD	Sc	St	Sc*St
14:0	R	5.97	5.17	0.220	n.s.	n.s.	n.s.	4.16	3.31	0.157	*	n.s.	n.s.
	C	6.41	5.16					4.15	3.62				
16:0	R	25.23	22.08	0.747	n.s.	n.s.	n.s.	28.61a	20.34b	1.043	***	n.s.	*
	C	24.92	23.12					26.64a	21.53b				
18:0	R	2.97	y2.62	0.058	n.s.	**	n.s.	8.06	7.31	0.458	n.s.	n.s.	n.s.
	C	3.05	x2.81					8.02	8.14				
Σ SFA^D	R	35.67	30.73	0.640	n.s.	n.s.	n.s.	42.76a	32.58b	1.418	***	n.s.	*
	C	35.79	32.07					40.73a	34.96b				
16:1n-7	R	7.18	6.88	0.293	n.s.	n.s.	n.s.	8.40a	4.93b	0.493	***	n.s.	n.s.
	C	7.81	6.74					8.38a	5.07b				
18:1n-9	R	16.17	16.91	0.962	n.s.	n.s.	n.s.	y16.61	15.92	0.414	n.s.	*	*
	C	17.59	17.78					x18.03	15.81				
18:1n-7	R	2.02a	y1.29b	0.078	**	**	**	y4.31	3.44	0.095	*	*	*
	C	2.07	x1.73					x4.68	3.41				
20:1 n-9	R	1.30	1.77	0.187	n.s.	n.s.	n.s.	0.39	0.74	0.051	*	n.s.	n.s.
	C	1.52	1.55					0.36	0.70				
22:1n-11	R	2.34	3.84	0.042	n.s.	n.s.	n.s.	-	0.57	-	-	-	-
	C	2.58	3.49					-	0.57				
Σ MUFA^E	R	29.59	31.31	1.887	n.s.	n.s.	n.s.	30.49	26.26	0.819	*	n.s.	n.s.
	C	32.15	31.90					32.14	26.22				
18:2n-6	R	1.00	0.90	0.028	n.s.	n.s.	n.s.	1.08	1.28	0.267	n.s.	n.s.	n.s.
	C	1.03	0.89					1.38	1.27				
20:4n-6	R	0.44	0.36	0.061	n.s.	n.s.	n.s.	1.32	1.42	0.094	n.s.	n.s.	n.s.
	C	0.35	0.36					1.34	1.54				
22:5n-6	R	0.21	0.17	0.069	n.s.	n.s.	n.s.	0.33b	0.68a	0.049	***	n.s.	n.s.
	C	0.18	0.24					0.31b	0.59°				
Σ n-6 PUFA^F	R	3.11a	2.58b	0.095	**	n.s.	n.s.	3.48	4.35	0.291	*	n.s.	n.s.
	C	3.06	2.64					3.93	4.35				
18:3n-3	R	1.09	x0.98	0.005	*	***	**	0.49	0.67	0.035	n.s.	n.s.	n.s.
	C	1.07	y0.94					0.53	0.66				
18:4n-3	R	2.34	2.06	0.045	n.s.	*	n.s.	0.67	0.91	0.040	n.s.	n.s.	*
	C	2.26	1.94					0.73	0.82				
20:5n-3	R	7.66	9.62	0.479	n.s.	*	n.s.	6.26	7.66	0.403	n.s.	n.s.	*
	C	6.64	9.23					6.88	6.81				
22:5n-3	R	0.41	0.65	0.030	*	n.s.	n.s.	1.24b	2.28a	0.178	**	n.s.	n.s.
	C	0.39	0.67					1.34	2.04				
22:6n-3	R	12.96	15.86	1.017	n.s.	n.s.	n.s.	7.11b	17.99a	1.513	***	n.s.	n.s.
	C	11.63	15.47					7.09b	15.77a				
Σ n-3 PUFA^G	R	24.81	29.62	1.463	n.s.	n.s.	n.s.	15.96b	29.97a	2.106	***	n.s.	n.s.
	C	22.32	28.71					16.83b	26.46a				
Σ PUFA^H	R	28.74	32.87	1.497	n.s.	n.s.	n.s.	20.09b	34.69a	2.198	***	n.s.	n.s.
	C	26.16	32.04					21.28b	31.18a				

Unknown	R	6.69	5.58	0.666	**	n.s.	n.s.	6.54	6.32	0.688	n.s.	n.s.	*
	C	6.42a	4.68b					5.65	7.52				
n-6/n-3	R	0.13	0.09	0.015	n.s.	n.s.	n.s.	0.22a	0.15b	0.021	**	n.s.	n.s.
	C	0.14	0.09					0.23a	0.16b				

610

611 ^A Values are means of 4 determinations (each one in duplicate on pooled samples) per season

612 of catch and state; ^B Means within a column and trait preceded by different letters (x,y) differ

613 significantly ($P \leq 0.05$); means on the same row followed by different letters (a,b) differ

614 significantly ($P \leq 0.05$); ^C RSD = Residual Standard Deviation; Sc = Season of catch; St =

615 State; ^D Includes: 15:0 and 17:0; ^E Includes: 20:1n-11; 20:1 n-7 and 22:1n-9; ^F Includes: 18:3,

616 20:2, 20:3 and 22:4n-6; ^G Includes: 16:4 and 20:4n-3; ^H Includes: 16: 2n-4; *** $P \leq 0.001$;

617 ** $P \leq 0.01$; * $P \leq 0.05$; n.s. = not significant.

618