

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

The Soap Bubble Theorem and a p -Laplacian overdetermined problem

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Colasuonno, F., Ferrari, F. (2020). The Soap Bubble Theorem and a p -Laplacian overdetermined problem. COMMUNICATIONS ON PURE AND APPLIED ANALYSIS, 19(2), 983-1000 [10.3934/cpaa.2020045].

Availability:

This version is available at: <https://hdl.handle.net/11585/714862> since: 2020-01-18

Published:

DOI: <http://doi.org/10.3934/cpaa.2020045>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Colasuonno, F., Ferrari, F. The soap bubble theorem and a p-Laplacian overdetermined problem (2020) *Communications on Pure and Applied Analysis*, 19 (2), pp. 983-1000

The final published version is available online at:
<https://dx.doi.org/10.3934/cpaa.2020045>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

1 THE SOAP BUBBLE THEOREM AND A p -LAPLACIAN 2 OVERDETERMINED PROBLEM

3 FRANCESCA COLASUONNO AND FAUSTO FERRARI

ABSTRACT. We consider the p -Laplacian equation $-\Delta_p u = 1$ for $1 < p < 2$, on a regular bounded domain $\Omega \subset \mathbb{R}^N$, with $N \geq 2$, under homogeneous Dirichlet boundary conditions. In the spirit of Alexandrov's Soap Bubble Theorem and of Serrin's symmetry result for the overdetermined problems, we prove that if the mean curvature H of $\partial\Omega$ is constant, then Ω is a ball and the unique solution of the Dirichlet p -Laplacian problem is radial. The main tools used are integral identities, the P -function, and the maximum principle.

4 1. INTRODUCTION

5 The celebrated Alexandrov's Soap Bubble Theorem [2], dated back to 1958,
6 states that if Γ is a compact hypersurface, embedded in \mathbb{R}^N , having constant mean
7 curvature, then Γ is a sphere. On the other hand, Serrin's symmetry result (1971)
8 [26] for the following overdetermined problem

$$-\Delta u = 1 \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega, \quad (1.1)$$

$$9 \quad u_\nu = c \quad \text{on } \partial\Omega, \quad (1.2)$$

10 where $\Omega \subset \mathbb{R}^N$ is a bounded domain and u_ν is the outer normal derivative, states
11 that if (1.1)–(1.2) has a solution, then Ω must be a ball, and the unique solution u
12 must be radial. It is nowadays well-known that these two results are strictly related.
13 Indeed, for his proof, Serrin adapted to the PDEs the reflection principle, a geomet-
14 rical technique introduced by Alexandrov in [2], and combined it with the maximum
15 principle, giving rise to a very powerful and versatile tool, the *moving plane method*.
16 This method is still very much used, since it can be successfully applied to a large
17 class of PDEs. Besides the common techniques used, the link between these two
18 results has been further highlighted by Reilly in [25], where the author proposed
19 an alternative proof of the Soap Bubble Theorem, considering the hypersurface Γ
20 as a level set (i.e., $\partial\Omega$) of the solution of (1.1). For his proof, Reilly found and
21 exploited a relation between the Laplacian operator and the geometrical concept of
22 mean curvature. Interestingly enough, Serrin's result for the overdetermined prob-
23 lem has been proved via a different technique by Weinberger in a two-page paper
24 [29] that was published in the same volume of the same journal as the paper by
25 Serrin [26]. Weinberger's proof is much simpler, it relies on some integral identities,
26 the maximum principle, and the introduction of an auxiliary function, the so-called
27 P -function. Even if Weinberger's technique is less flexible than the moving plane

Date: July 11, 2019.

2010 Mathematics Subject Classification. 35J92, 35B06, 35N25, 53A10, 35A23.

Key words and phrases. Alexandrov's Soap Bubble Theorem, Serrin-type result for overdeter-
mined p -Laplacian problems, p -torsional problem, P -function, Radial symmetry results.

method, it lends itself well to being re-read in quantitative terms. Recently, using this approach, Magnanini and Poggesi in [17, 18, 19, 22] proved the stability both for the Alexandrov's Soap Bubble Theorem and for Serrin's result, cf. also [21]. Also the moving plane method has been reformulated in a quantitative version to get the stability of both Serrin's result, cf. [1], and [8] where the best estimate obtained so far with this method appears, and Alexandrov's Theorem, cf. [10]. In those stability results, the idea is to measure how much Ω is close to being a ball by estimating from above the difference $r_e - r_i$ (r_e and r_i being the radii of two suitable balls such that $B_{r_e} \subset \Omega \subset B_{r_i}$) in terms of the deviation of the normal derivative u_ν from being constant on $\partial\Omega$, or in terms of the deviation of the mean curvature H from being constant on Γ . Furthermore, under more general assumptions, without any hypotheses on the domain, Ciruolo and Maggi in [7] proved a stability result for Alexandrov's Theorem using the torsion potentials. Other stability issues for the Serrin problem have been treated in [4].

Serrin's symmetry result has been extensively studied and generalized also to the case of quasilinear problems. For the p -Laplacian operator $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2} \nabla u)$, $1 < p < \infty$, it has been proved that if the following problem

$$\begin{cases} -\Delta_p u = 1 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \\ |\nabla u| = c & \text{on } \partial\Omega \text{ for some } c > 0 \end{cases} \quad (1.3)$$

admits a weak solution in the bounded domain $\Omega \subset \mathbb{R}^N$, then Ω is a ball. Garofalo and Lewis [14] proved this result via Weinberger's approach; Brock and Henrot [6] proposed a different proof via Steiner symmetrization for $p \geq 2$; Damascelli and Pacella [11] succeeded in adapting the moving plane method to the case $1 < p < 2$. Finally, Bianchini and Ciruolo in [3] proved the result for every p , in a framework which is possibly anisotropic, using Newton's inequality. Later, many other refinements and generalizations have been proposed in [13, 12, 5] for more general operators, and in [20, 9] for the case of convex cones.

In this paper, we consider the following Dirichlet p -Laplacian problem

$$\begin{cases} -\Delta_p u = 1 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.4)$$

for $1 < p < 2$. Here $\Omega \subset \mathbb{R}^N$ is a smooth bounded domain and $N \geq 2$. Due to its physical meaning, (1.4) is often referred to as p -torsion problem. For this problem, existence and uniqueness of the solution can be easily proved via the Direct Method of the Calculus of Variations and using the strict convexity of the action functional associated, see Section 2. In the spirit of Reilly's result, we regard the hypersurface Γ of Alexandrov's Theorem as the level set $\partial\Omega$ of the solution of (1.4) and we obtain, for smooth hypersurfaces, an alternative proof of the Soap Bubble Theorem. As a consequence, we prove the equivalence of the Soap Bubble Theorem to the Serrin-type symmetry result for the overdetermined problem (1.3), when $1 < p < 2$. We state here our main results.

Theorem 1.1. *Let $\Gamma \subset \mathbb{R}^N$ be a $C^{2,\alpha}$ surface which is the boundary of a bounded domain $\Omega \subset \mathbb{R}^N$, i.e. $\Gamma = \partial\Omega$, and denote by $H = H(x)$ the mean curvature of $\partial\Omega$. Suppose that $1 < p < 2$, that u solves (1.4), and that the set of critical points of u has zero measure. Then the following statements are equivalent:*

- 1 a. Ω is a ball;
- 2 b. $|u_\nu(x)|^{p-2}u_\nu(x) = -\frac{1}{NH(x)}$ for every $x \in \partial\Omega$;
- 3 c. u is radial;
- 4 d. $H(x) = H_0$ for every $x \in \partial\Omega$.

5 Moreover, if one of the previous ones holds, then

- 6 e. $|\nabla u(x)| = \left(\frac{1}{NH_0}\right)^{\frac{1}{p-1}}$ for every $x \in \partial\Omega$.

7 The implication d. \Rightarrow a. in the previous theorem is a special case of the Soap
8 Bubble Theorem of Alexandrov. We further observe that from the proof of the
9 previous theorem, cf. formula (3.4), it results that if d. holds, then Ω must be a
10 ball of radius $R_0 = 1/H_0$. Moreover, the fact that any of the statements a., b.,
11 c., or d. implies e. is a simple consequence of the previous results, but we know
12 that the converse implication e. \Rightarrow a. holds as well: as proved in [14, 13, 12], the
13 overdetermined problem (1.3) admits a solution only if Ω is a ball of radius R_0 .
14 This allows us to state the equivalence of the Soap Bubble Theorem and of the
15 Serrin-type result for the overdetermined p -Laplacian problem (1.3) under suitable
16 regularity assumptions, in the case $1 < p < 2$.

17 **Corollary 1.2.** *Under the assumptions of Theorem 1.1, statements a., b., c., d.,*
18 *and e. are all equivalent.*

19 Our proof technique takes inspiration from [17] and follows the approach of
20 Weinberger. After having introduced the P -function (2.5) in terms of the solution
21 of (1.4), we derive the integral identity (2.7) using the Divergence Theorem. The
22 identity (2.7) will be a key tool for the estimates in the rest of the paper. We
23 recall then that the p -Laplacian of a smooth function can be expressed as the trace
24 of a matrix-operator applied to the same function, cf. (2.2), and we use a simple
25 algebraic inequality (2.11) (known as Newton's inequality) to get an estimate of
26 the p -Laplacian of a function. This suggests us to introduce in (3.1) the integral
27 $\mathcal{I}_p(u)$ which will play the role of the so-called Cauchy-Schwarz deficit in [17] for the
28 linear case $p = 2$. In view of Newton's inequality, the integral $\mathcal{I}_p(u)$ has a sign, it is
29 always non-negative. Now the P -function comes into play: thanks to the fact that
30 it satisfies a maximum principle, we can prove that, when $1 < p < 2$, $\mathcal{I}_p(u)$ vanishes
31 only on radial solutions of (1.4), cf. Lemma 2.6. This, combined with the integral
32 identity (2.7), allows us to obtain an estimate from above of $\mathcal{I}_p(u)$ in terms of some
33 boundary integrals involving only the mean curvature H and the normal derivative
34 u_ν , see Theorem 3.1. Then Theorem 1.1 and Corollary 1.2 are easy consequences:
35 $\mathcal{I}_p(u)$ is zero (or equivalently the solution of (1.4) is radial) if and only if the mean
36 curvature H is constant on $\partial\Omega$ or the modulus of the gradient of u is constant on
37 $\partial\Omega$. Finally, in Corollary 3.6, we give an estimate from above of the integral $\mathcal{I}_p(u)$
38 in terms of the $L^1(\partial\Omega)$ -norm of the deviation of H from being constant and some
39 constants which only depend on the geometry of the problem, cf. (3.6).

40 The paper is organized as follows: in Section 2 we introduce some useful notation,
41 the P -function, some known results, and some preliminary lemmas. In Section 3
42 we prove Theorem 1.1 and its consequences, while in Section 4, we present some
43 comments on the stability for the p -overdetermined problem.

2. PRELIMINARIES

We first introduce the main important quantities and notation involved. Throughout the paper, with abuse of notation, we use the symbol $|\cdot|$ to denote both the N -dimensional and the $(N-1)$ -dimensional Lebesgue measures. We further denote by $\|\cdot\|$ the Frobenius matrix norm and by $\langle \cdot, \cdot \rangle$ the scalar product in \mathbb{R}^N .

The p -Laplacian on non-critical level sets of u . The p -Laplacian of a regular function v can be expressed as follows

$$\Delta_p v = |\nabla v|^{p-2} \left(\Delta v + (p-2) \frac{\langle D^2 v \nabla v, \nabla v \rangle}{|\nabla v|^2} \right), \quad (2.1)$$

where $D^2 v$ denotes the Hessian matrix of v . Moreover, we recall that, in view of (2.1), it is possible to express the p -Laplacian of any C^2 -function v as follows

$$\begin{aligned} \Delta_p v &= |\nabla v|^{p-2} \left(\Delta v + (p-2) \left\langle D^2 v \frac{\nabla v}{|\nabla v|}, \frac{\nabla v}{|\nabla v|} \right\rangle \right) \\ &= |\nabla v|^{p-2} \left(\text{Tr}(D^2 v) + \frac{p-2}{|\nabla v|^2} \sum_{i,j=1}^N \frac{\partial^2 v}{\partial x_i \partial x_j} \frac{\partial v}{\partial x_i} \frac{\partial v}{\partial x_j} \right) \\ &= |\nabla v|^{p-2} \left[\text{Tr}(D^2 v) + (p-2) \text{Tr} \left(\frac{\nabla v}{|\nabla v|} \otimes \frac{\nabla v}{|\nabla v|} \cdot D^2 v \right) \right] \\ &= \text{Tr} \left[|\nabla v|^{p-2} \left(\text{I} + (p-2) \frac{\nabla v}{|\nabla v|} \otimes \frac{\nabla v}{|\nabla v|} \right) D^2 v \right], \end{aligned} \quad (2.2)$$

where we have denoted simply by I the $N \times N$ identity matrix.

Let u be a solution of (1.4). We denote by ν the following vector field

$$\nu = -\frac{\nabla u}{|\nabla u|},$$

which coincides with the external unit normal on $\partial\Omega$, being $u|_{\partial\Omega}$ constant. The mean curvature of the regular level sets of u is given by

$$H = -\frac{1}{N-1} \text{div} \frac{\nabla u}{|\nabla u|}.$$

It is possible to see that, on non-critical level sets of u , the Laplacian of u can be expressed in terms of H as follows

$$\Delta u = u_{\nu\nu} + (N-1)Hu_\nu, \quad (2.3)$$

where $u_\nu = \nabla u \cdot \nu = -|\nabla u|$ and $u_{\nu\nu} = \langle D^2 u \nu, \nu \rangle$. Therefore, on non-critical level sets of u , we can write the p -Laplacian as

$$\Delta_p u = |u_\nu|^{p-2} [(p-1)u_{\nu\nu} + (N-1)Hu_\nu]. \quad (2.4)$$

The P -function. In terms of a solution u of (1.4), we can define the so-called P -function as

$$P := \frac{2(p-1)}{p} |\nabla u|^p + \frac{2}{N} u \quad \text{a.e. in } \Omega, \quad (2.5)$$

we refer to [27, Chapter 7, formula (7.6) with $v(q) = q^{\frac{p-2}{2}}$ and $q = |\nabla u|^2$] for its derivation. The main feature of P is that it satisfies a maximum principle, which is the starting point for finding useful bounds for the main quantities involved in this problem.

Definition 2.1. Let $\Omega \subset \mathbb{R}^N$ be a bounded domain. Ω satisfies the *interior sphere condition* if for every $x \in \partial\Omega$ there exist $x_0 \in \Omega$ and $r > 0$ such that $B_r(x_0) := \{y \in \mathbb{R}^N : |y - x_0| < r\} \subset \Omega$ and $x \in \partial B_r(x_0)$.

We recall that if Ω is a C^2 bounded domain, then it satisfies the interior sphere condition.

Lemma 2.2. Let Ω be of class $C^{1,\alpha}$ and satisfy the interior sphere condition. If u solves (1.4), then P is either constant in $\bar{\Omega}$ or it satisfies $P_\nu > 0$ on $\partial\Omega$.

Proof. The proof of this lemma is given in [13, Lemma 3.2] for a solution of the overdetermined problem (1.3); we report the outline of the proof here in order to highlight that it continues to hold even if u does not satisfy $|\nabla u| = \text{const.}$ on $\partial\Omega$.

Since u solves (1.4), then by [24, Theorem 3.2.2], $u \geq 0$ a.e. in Ω and by [16, Theorem 1], u is of class $C^{1,\alpha}(\bar{\Omega})$. Now, [28, Theorem 5] guarantees that $|\nabla u| \geq \min_{\partial\Omega} |\nabla u| > 0$ on $\partial\Omega$. By continuity, $|\nabla u| \neq 0$ in a closed neighborhood $D \subset \bar{\Omega}$ of $\partial\Omega$.

Now, suppose that P is not constant in $\bar{\Omega}$. Under this assumption, as in [13, Lemma 3.2 - Claim - Step 2], it is possible to prove that P attains its maximum on $\partial\Omega$ and that, if P also attains its maximum at a point $\bar{x} \in \Omega$, then necessarily $\nabla u(\bar{x}) = 0$. Therefore, being $D \subset \bar{\Omega}$ a closed neighborhood of $\partial\Omega$, P attains its maximum in D only on $\partial\Omega$. By the proof of [13, Lemma 3.2], we know that P satisfies in D a uniformly elliptic equation and so it satisfies the classical Hopf's lemma. Hence, $P_\nu > 0$ on $\partial\Omega$. \square

For future use, we derive here an easy identity holding true for any u solution of (1.4). By integration by parts, the Divergence Theorem, and (2.3) we get

$$\begin{aligned} \int_{\Omega} \langle |\nabla u|^{p-2} \nabla u, \nabla \Delta u \rangle dx &= - \int_{\Omega} \Delta_p u \Delta u dx + \int_{\partial\Omega} \Delta u |\nabla u|^{p-2} \nabla u \cdot \nu d\sigma \\ &= \int_{\Omega} \Delta u dx + \int_{\partial\Omega} \Delta u |u_\nu|^{p-2} u_\nu d\sigma \\ &= \int_{\partial\Omega} u_\nu d\sigma + \int_{\partial\Omega} |u_\nu|^{p-2} u_\nu [u_{\nu\nu} + (N-1)H u_\nu] d\sigma \\ &= \int_{\partial\Omega} u_\nu d\sigma - \int_{\partial\Omega} |u_\nu|^{p-1} u_{\nu\nu} d\sigma + (N-1) \int_{\partial\Omega} H |u_\nu|^p d\sigma, \end{aligned} \tag{2.6}$$

where we used that $\partial\Omega$ is a non-critical level set of u , as showed in the proof of Lemma 2.2.

Reference constant mean curvature and reference domain. We introduce here some reference geometric constants which are related to problem (1.4). These constants will be useful to compare problem (1.4) with the same problem set in a ball instead of a general domain Ω .

By Minkowski's identity, i.e.,

$$\int_{\partial\Omega} H(x) \langle x - z, \nu(x) \rangle d\sigma = |\partial\Omega| \quad \text{for any } z \in \mathbb{R}^N,$$

we get, by the Divergence Theorem and if H is constant:

$$|\partial\Omega| = H \int_{\partial\Omega} \langle x - z, \nu(x) \rangle d\sigma = H \int_{\Omega} \sum_{i=1}^N \frac{\partial(x-z)_i}{\partial x_i} dx = H |\Omega| N.$$

If H is not constant, we can take as reference constant mean curvature the quantity

$$H_0 := \frac{|\partial\Omega|}{N|\Omega|}$$

and, as reference domain, a ball of radius

$$R_0 = \frac{1}{H_0} = \frac{N|\Omega|}{|\partial\Omega|}.$$

1

Existence and uniqueness for (1.4). Problem (1.4) has a variational structure with associated action functional $I : W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$ given by

$$I(u) := \int_{\Omega} \left(\frac{1}{p} |\nabla u|^p - u \right) dx.$$

2 By strict convexity and the Direct Method of Calculus of Variations, it is possible
3 to prove that I has a unique minimizer. Hence, (1.4) has a unique weak solution
4 $u \in W_0^{1,p}(\Omega)$.

From now on in the paper, we denote by \mathcal{C} the critical set of the solution u of problem (1.4), namely

$$\mathcal{C} := \{x \in \Omega : |\nabla u(x)| = 0\}.$$

5 By [13, Lemma 3.1], we know that the solution u of (1.4) is of class $C^{2,\alpha}(\bar{\Omega} \setminus \mathcal{C})$.

6 Therefore, hereafter we assume that Ω is of class $C^{2,\alpha}$ in order to guarantee that
7 the solution u of (1.4) is of class $C^{2,\alpha}$ in a neighborhood of $\partial\Omega$ (this is a consequence
8 of the regularity of u and of the first part of the proof of Lemma 2.2).

9 **Lemma 2.3.** *Let u solve (1.4) and suppose that its critical set \mathcal{C} has zero N -
10 dimensional measure. The following identity holds*

$$\begin{aligned} \int_{\Omega} \left\{ (p-1) |\nabla u|^{p-2} \left[(p-2) \left\| D^2 u \frac{\nabla u}{|\nabla u|} \right\|^2 + \|D^2 u\|^2 + \langle \nabla u, \nabla \Delta u \rangle \right] + \frac{\Delta u}{N} \right\} dx \\ = - \int_{\partial\Omega} (N-1) \left(\frac{1}{N} u_{\nu} + H |u_{\nu}|^p \right) d\sigma \end{aligned} \quad (2.7)$$

11 *Proof.* By straightforward calculations, we get

$$P_{\nu} = \nabla P \cdot \nu = 2u_{\nu} \left((p-1) |u_{\nu}|^{p-2} u_{\nu\nu} + \frac{1}{N} \right), \quad (2.8)$$

12 cf. [27, formula (7.7)] with $f \equiv w \equiv 1$, $\alpha = 2/N$, $q = |\nabla u|^2$, and $v(q) = q^{(p-2)/2}$.

13 By taking into account (2.3), (2.4), and the equation in (1.4), we can rewrite P_{ν} as

14

$$\begin{aligned} P_{\nu} &= 2u_{\nu} \left(\Delta_p u - (N-1) H |u_{\nu}|^{p-2} u_{\nu} + \frac{1}{N} \right) \\ &= -2(N-1) \left(\frac{1}{N} u_{\nu} + H |u_{\nu}|^p \right). \end{aligned} \quad (2.9)$$

1 Moreover,

$$\Delta P = 2 \left\{ (p-1) |\nabla u|^{p-2} \left[(p-2) \left\| D^2 u \frac{\nabla u}{|\nabla u|} \right\|^2 + \|D^2 u\|^2 + \langle \nabla u, \nabla \Delta u \rangle \right] + \frac{\Delta u}{N} \right\} \quad (2.10)$$

cf. [27, formula (7.9)]. The conclusion then follows, since $\int_{\Omega} \Delta P dx = \int_{\partial\Omega} P_{\nu} d\sigma$, by the Divergence Theorem. \square

Proposition 2.4 (Newton's inequality). *Let $n \in \mathbb{N}$ and A be a $(n \times n)$ -matrix, then*

$$\|A\|^2 \geq \frac{(\text{Tr}(A))^2}{n}, \quad (2.11)$$

where $\text{Tr}(\cdot)$ denotes the trace of a matrix. Furthermore, the equality holds in (2.11) if and only if $A = kI_n$ for some constant k .

Proof. The proof is standard, but we report it here for the sake of completeness. The statement is trivial for $n = 1$. We proceed by induction on $n \geq 2$. If we denote by a_{ij} the elements of the matrix A , we obtain for $n = 2$ that

$$(\text{Tr}(A))^2 = (a_{11} + a_{22})^2 = a_{11}^2 + a_{22}^2 + 2a_{11}a_{22} \leq 2(a_{11}^2 + a_{22}^2) \leq 2\|A\|^2, \quad (2.12)$$

where we have used that $2a_{11}a_{22} \leq a_{11}^2 + a_{22}^2$, being $(a_{11} - a_{22})^2 = a_{11}^2 + a_{22}^2 - 2a_{11}a_{22} \geq 0$. As a consequence, we observe that (2.12) holds with the equality signs if and only if $a_{11} = a_{22}$ and $a_{12} = a_{21} = 0$. We now assume that (2.11) holds true for n and we prove it for $n + 1$. Indeed,

$$\begin{aligned} (\text{Tr}(A))^2 &= \left(\sum_{i=1}^{n+1} a_{ii} \right)^2 = \left(\sum_{i=1}^n a_{ii} + a_{n+1,n+1} \right)^2 \\ &= \left(\sum_{i=1}^n a_{ii} \right)^2 + 2 \left(\sum_{i=1}^n a_{ii} \right) a_{n+1,n+1} + a_{n+1,n+1}^2 \\ &\leq n \sum_{i=1}^n a_{ii}^2 + n \sum_{\substack{i,j=1 \\ i \neq j}}^n a_{ij}^2 + 2 \left(\sum_{i=1}^n a_{ii} \right) a_{n+1,n+1} + a_{n+1,n+1}^2. \end{aligned} \quad (2.13)$$

Now, as above, we can estimate

$$\begin{aligned} 2 \left(\sum_{i=1}^n a_{ii} \right) a_{n+1,n+1} &= \sum_{i=1}^n 2a_{ii}a_{n+1,n+1} \\ &\leq \sum_{i=1}^n (a_{ii}^2 + a_{n+1,n+1}^2) = na_{n+1,n+1}^2 + \sum_{i=1}^n a_{ii}^2, \end{aligned}$$

where the equality is achieved only for $a_{ii} = a_{n+1,n+1}$ for every $i = 1, \dots, n$. Therefore, combining this estimate with (2.13), we obtain

$$\begin{aligned} (\text{Tr}(A))^2 &\leq n \sum_{i=1}^n a_{ii}^2 + a_{n+1,n+1}^2 + na_{n+1,n+1}^2 + \sum_{i=1}^n a_{ii}^2 + n \sum_{\substack{i,j=1 \\ i \neq j}}^n a_{ij}^2 \\ &= (n+1) \sum_{i=1}^n a_{ii}^2 + (n+1)a_{n+1,n+1}^2 + n \sum_{\substack{i,j=1 \\ i \neq j}}^n a_{ij}^2 \\ &= (n+1) \sum_{i=1}^{n+1} a_{ii}^2 + n \sum_{\substack{i,j=1 \\ i \neq j}}^n a_{ij}^2 \leq (n+1) \sum_{i,j=1}^{n+1} a_{ij}^2, \end{aligned}$$

where the equalities hold only when $A = kI_{n+1}$ for some constant k , and the proof is complete. \square

Corollary 2.5. *Let v be any C^2 -function, then the following inequality holds*

$$(\Delta_p v)^2 \leq N |\nabla v|^{2(p-2)} \left\| \left(I + (p-2) \frac{\nabla v}{|\nabla v|} \otimes \frac{\nabla v}{|\nabla v|} \right) D^2 v \right\|^2. \quad (2.14)$$

Proof. Taking into account (2.2), it is enough to apply Proposition 2.4 with $n := N$ and $A := |\nabla v|^{p-2} \left(I + (p-2) \frac{\nabla v}{|\nabla v|} \otimes \frac{\nabla v}{|\nabla v|} \right) D^2 v$. \square

For every $z \in \mathbb{R}^N$ and $r > 0$, we introduce the function

$$w_r(x) := -\frac{p-1}{pN^{\frac{1}{p-1}}} \left(|x-z|^{\frac{p}{p-1}} - r \right) \quad \text{for every } x \in \Omega. \quad (2.15)$$

We observe that, if $z \in \Omega$ and $p > 2$, then $w_r \notin C^2$. Clearly, w_r is radial about z , and, if $\Omega = B_r(z)$, it solves (1.4). Indeed, by straightforward calculations we get

$$\begin{aligned} \nabla w_r &= -N^{-\frac{1}{p-1}} |x-z|^{\frac{p}{p-1}-2} (x-z), \\ |\nabla w_r|^{p-2} \nabla w_r &= -\frac{1}{N} (x-z), \end{aligned}$$

and so

$$\Delta_p w_r = \operatorname{div} \left(-\frac{1}{N} (x-z) \right) = -1.$$

We are now ready to prove the following result.

Lemma 2.6. *Let $1 < p < 2$, then the following statements hold true.*

- (i) *Let w_r be defined as in (2.15), then for $v := w_r$ the equality holds in (2.14).*
- (ii) *Let u solve (1.4). Suppose that the critical set C of u has zero N -dimensional measure and that for $v := u$ the equality holds in (2.14) for every $x \in \Omega \setminus C$. Then u is radial.*

Proof. (i) Since

$$\frac{\partial^2 w_r}{\partial x_i \partial x_j} = -N^{-\frac{1}{p-1}} \left[\frac{2-p}{p-1} |x-z|^{\frac{p}{p-1}-4} (x_j - z_j)(x_i - z_i) + \delta_{ij} |x-z|^{\frac{p}{p-1}-2} \right],$$

the Hessian of w_r has the following expression

$$D^2 w_r = -N^{-\frac{1}{p-1}} |x-z|^{\frac{2-p}{p-1}} \left(\frac{2-p}{p-1} \cdot \frac{x-z}{|x-z|} \otimes \frac{x-z}{|x-z|} + I \right).$$

By

$$\left(\frac{x-z}{|x-z|} \otimes \frac{x-z}{|x-z|} \right)^2 = \frac{x-z}{|x-z|} \otimes \frac{x-z}{|x-z|} \quad \text{and} \quad \frac{\nabla w_r}{|\nabla w_r|} = \frac{x-z}{|x-z|},$$

we get

$$\begin{aligned} & |\nabla w_r|^{p-2} \left(I + (p-2) \frac{\nabla w_r}{|\nabla w_r|} \otimes \frac{\nabla w_r}{|\nabla w_r|} \right) D^2 w_r \\ &= -\frac{|x-z|^{\frac{2-p}{p-1} + (\frac{p}{p-1}-1)(p-2)}}{N} \left[I + \left(\frac{2-p}{p-1} - \frac{(p-2)^2}{p-1} + p-2 \right) \frac{x-z}{|x-z|} \otimes \frac{x-z}{|x-z|} \right] \\ &= -\frac{1}{N} I. \end{aligned}$$

2 Hence, by Proposition 2.4, (2.14) holds with the equality sign for $v := w_r$.

(ii) By Proposition 2.4, we know that the equality holds in (2.14) if and only if

$$|\nabla u|^{p-2} \left(\mathbf{I} + (p-2) \frac{\nabla u}{|\nabla u|} \otimes \frac{\nabla u}{|\nabla u|} \right) D^2 u = k \mathbf{I}$$

for some constant k . By $\left\| (2-p) \frac{\nabla u}{|\nabla u|} \otimes \frac{\nabla u}{|\nabla u|} \right\| = |2-p| < 1$,

$$\det \left(\mathbf{I} - (2-p) \frac{\nabla u}{|\nabla u|} \otimes \frac{\nabla u}{|\nabla u|} \right) \neq 0,$$

and

$$\left(\frac{x}{|x|} \otimes \frac{x}{|x|} \right)^i = \frac{x}{|x|} \otimes \frac{x}{|x|} \quad \text{for all } x \in \mathbb{R}^N \text{ and all } i \in \mathbb{N},$$

3 we get on $\Omega \setminus \mathcal{C}$

$$\begin{aligned} D^2 u &= \frac{k}{|\nabla u|^{p-2}} \left(\mathbf{I} - (2-p) \frac{\nabla u}{|\nabla u|} \otimes \frac{\nabla u}{|\nabla u|} \right)^{-1} \\ &= \frac{k}{|\nabla u|^{p-2}} \sum_{i=0}^{\infty} (2-p)^i \left(\frac{\nabla u}{|\nabla u|} \otimes \frac{\nabla u}{|\nabla u|} \right)^i \\ &= \frac{k}{|\nabla u|^{p-2}} \left(\mathbf{I} + \frac{\nabla u}{|\nabla u|} \otimes \frac{\nabla u}{|\nabla u|} \sum_{i=1}^{\infty} (2-p)^i \right) \\ &= \frac{k}{|\nabla u|^{p-2}} \left[\mathbf{I} + \frac{\nabla u}{|\nabla u|} \otimes \frac{\nabla u}{|\nabla u|} \left(\frac{1}{1-(2-p)} - 1 \right) \right] \\ &= \frac{k}{|\nabla u|^{p-2}} \left(\mathbf{I} - \frac{p-2}{p-1} \frac{\nabla u}{|\nabla u|} \otimes \frac{\nabla u}{|\nabla u|} \right). \end{aligned} \tag{2.16}$$

Namely, for $i, j = 1, \dots, N$

$$\partial_{ij}^2 u = \frac{k}{|\nabla u|^{p-2}} \left(\delta_{ij} - \frac{p-2}{p-1} \frac{\partial_i u \partial_j u}{|\nabla u|^2} \right).$$

1 Hence, in particular,

$$\Delta u = \frac{k}{|\nabla u|^{p-2}} \sum_{i=1}^N \left(1 - \frac{p-2}{p-1} \frac{(\partial_i u)^2}{|\nabla u|^2} \right) = \frac{k}{|\nabla u|^{p-2}} \left(N - \frac{p-2}{p-1} \right). \tag{2.17}$$

Furthermore, since u solves (1.4), then by (2.16), (2.17), and (2.1), we have

$$\begin{aligned} -1 &= |\nabla u|^{p-2} \left(\Delta u + (p-2) \left\langle D^2 u \frac{\nabla u}{|\nabla u|}, \frac{\nabla u}{|\nabla u|} \right\rangle \right) \\ &= k \sum_{i=1}^N \left(1 - \frac{p-2}{p-1} \frac{(\partial_i u)^2}{|\nabla u|^2} \right) + (p-2) \left\langle |\nabla u|^{p-2} D^2 u \frac{\nabla u}{|\nabla u|}, \frac{\nabla u}{|\nabla u|} \right\rangle \\ &= k \sum_{i=1}^N \left(1 - \frac{p-2}{p-1} \frac{(\partial_i u)^2}{|\nabla u|^2} \right) + (p-2) \left\langle k \left(\mathbf{I} - \frac{p-2}{p-1} \frac{\nabla u}{|\nabla u|} \otimes \frac{\nabla u}{|\nabla u|} \right) \frac{\nabla u}{|\nabla u|}, \frac{\nabla u}{|\nabla u|} \right\rangle \\ &= k \left[N - \frac{p-2}{p-1} + (p-2) \left(1 - \frac{p-2}{p-1} \left\langle \frac{\nabla u}{|\nabla u|} \otimes \frac{\nabla u}{|\nabla u|} \frac{\nabla u}{|\nabla u|}, \frac{\nabla u}{|\nabla u|} \right\rangle \right) \right] \\ &= k \left[N - \frac{p-2}{p-1} + (p-2) \left(1 - \frac{p-2}{p-1} \right) \right] = kN, \end{aligned}$$

where in the last equality, we have used that

$$\frac{x}{|x|} \otimes \frac{x}{|x|} \frac{x}{|x|} = \frac{x}{|x|} \quad \text{for all } x \in \mathbb{R}^N.$$

2 Hence, $k = -\frac{1}{N}$.

Now, by the equation in (1.4), (2.17), and (2.3), we get on non-critical level sets of u

$$\begin{aligned} |u_\nu|^{p-2} [(p-1)u_{\nu\nu} + (N-1)Hu_\nu] &= -1, \\ u_{\nu\nu} + (N-1)Hu_\nu &= \left(\frac{p-2}{N(p-1)} - 1 \right) \frac{1}{|u_\nu|^{p-2}}, \end{aligned}$$

being $u_\nu = -|\nabla u|$. These two identities give

$$|u_\nu|^{p-2} u_{\nu\nu} = -\frac{1}{N(p-1)}$$

3 and consequently

$$H = \frac{1}{N|u_\nu|^{p-1}} \quad \text{on } \partial\Omega. \quad (2.18)$$

Now, by Lemma 2.2, we know that either P is constant on $\bar{\Omega}$, or $P_\nu > 0$ on $\partial\Omega$. If the first case occurs, then it is possible to see that all level sets of u are isoparametric surfaces. In particular, since u satisfies homogeneous Dirichlet boundary conditions, all level sets must be concentric spheres and so u is radial, cf. [13, Remark 5.5] and [15, Theorem 5]. If the second case occurs, then by (2.9),

$$\frac{1}{N}u_\nu + H|u_\nu|^p < 0 \quad \text{on } \partial\Omega,$$

therefore, by (2.18),

$$0 = \frac{1}{N}(u_\nu - u_\nu) = \frac{u_\nu}{N} + \frac{|u_\nu|}{N} < 0 \quad \text{on } \partial\Omega.$$

4 This is impossible and concludes the proof. \square

5 3. PROOF OF THE MAIN RESULTS

6 Let u solve (1.4) and suppose that its critical set \mathcal{C} has zero N -dimensional
7 measure. We introduce the following integral

$$\mathcal{I}_p(u) := \int_{\Omega} \left[|\nabla u|^{(p-2)} \left\| \left(I + (p-2) \frac{\nabla u}{|\nabla u|} \otimes \frac{\nabla u}{|\nabla u|} \right) D^2 u \right\|^2 - \left(\frac{\Delta_p u}{N^{1/2} |\nabla u|^{\frac{p-2}{2}}} \right)^2 \right] dx. \quad (3.1)$$

8 **Theorem 3.1.** *Let $1 < p < 2$ and $\partial\Omega$ be a $C^{2,\alpha}$ bounded domain of \mathbb{R}^N . If u solves
9 (1.4) and has $|\mathcal{C}| = 0$, then*

- 10 (i) $\mathcal{I}_p(u) \geq 0$ and $\mathcal{I}_p(u) = 0$ if and only if u is radial;
- 11 (ii) $\mathcal{I}_p(u) \leq -\frac{p(N-1)}{p-1} \int_{\partial\Omega} \left(\frac{1}{N}u_\nu + H|u_\nu|^p \right) d\sigma$;
- 1 (iii) $\mathcal{I}_p(u) \leq \frac{p(N-1)}{p-1} \int_{\partial\Omega} |u_\nu|^p (H_0 - H) d\sigma$.

- 2 *Proof of Theorem 3.1.* (i) By (2.14), we know that $\mathcal{I}_p(u) \geq 0$ and, by Lemma 2.6,
 3 we know that $\mathcal{I}_p(u) = 0$ if and only if u is radial.
 (ii) First, we observe that a.e. in Ω we have

$$\begin{aligned}
 & \left\| \left(I + (p-2) \frac{\nabla u}{|\nabla u|} \otimes \frac{\nabla u}{|\nabla u|} \right) D^2 u \right\|^2 = \sum_{i,j=1}^N \left(\partial_{ij}^2 u + (p-2) \sum_{k=1}^N \frac{\partial_i u}{|\nabla u|} \frac{\partial_k u}{|\nabla u|} \partial_{kj}^2 u \right)^2 \\
 & = \|D^2 u\|^2 + 2(p-2) \sum_{i,j=1}^N \partial_{ij}^2 u \frac{\partial_i u}{|\nabla u|} \sum_{k=1}^N \frac{\partial_k u}{|\nabla u|} \partial_{kj}^2 u + (p-2)^2 \sum_{i,j=1}^N \left(\sum_{k=1}^N \frac{\partial_i u}{|\nabla u|} \frac{\partial_k u}{|\nabla u|} \partial_{kj}^2 u \right)^2 \\
 & = \|D^2 u\|^2 + 2(p-2) \sum_{j=1}^N \left(\sum_{i=1}^N \frac{\partial_i u}{|\nabla u|} \partial_{ij}^2 u \right)^2 + (p-2)^2 \left\| D^2 u \frac{\nabla u}{|\nabla u|} \right\|^2 \\
 & = \|D^2 u\|^2 + p(p-2) \left\| D^2 u \frac{\nabla u}{|\nabla u|} \right\|^2.
 \end{aligned}$$

Furthermore, by (2.7), we get

$$\begin{aligned}
 p(p-2) \int_{\Omega} |\nabla u|^{p-2} \left\| D^2 u \frac{\nabla u}{|\nabla u|} \right\|^2 dx &= -p \int_{\Omega} \left[|\nabla u|^{p-2} (\|D^2 u\|^2 + \langle \nabla u, \nabla \Delta u \rangle) + \frac{\Delta u}{N(p-1)} \right] dx \\
 &\quad - p \frac{N-1}{p-1} \int_{\partial\Omega} \left(\frac{1}{N} u_{\nu} + H|u_{\nu}|^p \right) d\sigma.
 \end{aligned}$$

Hence, using these last two identities, we can rewrite $\mathcal{I}_p(u)$ as

$$\begin{aligned}
 \mathcal{I}_p(u) &= \int_{\Omega} \left\{ |\nabla u|^{p-2} [-(p-1)\|D^2 u\|^2 - p\langle \nabla u, \nabla \Delta u \rangle] - \frac{p}{N(p-1)} \Delta u - \frac{(\Delta_p u)^2}{N|\nabla u|^{p-2}} \right\} dx \\
 &\quad - \frac{p(N-1)}{p-1} \int_{\partial\Omega} \left(\frac{1}{N} u_{\nu} + H|u_{\nu}|^p \right) d\sigma.
 \end{aligned}$$

On the other hand, by (2.1), the $C^{2,\alpha}$ regularity of u in a neighborhood of $\partial\Omega$, and the Divergence Theorem

$$\begin{aligned}
 & -p \int_{\Omega} \left(|\nabla u|^{p-2} \langle \nabla u, \nabla \Delta u \rangle + \frac{1}{N(p-1)} \Delta u \right) dx \\
 &= \int_{\Omega} -p \left(1 + \frac{1}{N(p-1)} \right) \Delta u dx + p \int_{\partial\Omega} |\nabla u|^{p-1} \Delta u d\sigma \\
 &= p \int_{\partial\Omega} \left(1 + \frac{1}{N(p-1)} \right) |\nabla u| (1 + |\nabla u|^{p-2} \Delta u) d\sigma \\
 &= -p(p-2) \left(1 + \frac{1}{N(p-1)} \right) \int_{\partial\Omega} |\nabla u|^{p-1} \langle D^2 u \frac{\nabla u}{|\nabla u|}, \frac{\nabla u}{|\nabla u|} \rangle d\sigma.
 \end{aligned}$$

1 Hence,

$$\begin{aligned}
 \mathcal{I}_p(u) &= \int_{\Omega} \left\{ -(p-1)|\nabla u|^{p-2} \|D^2 u\|^2 - \frac{(\Delta_p u)^2}{N|\nabla u|^{p-2}} \right\} dx \\
 &\quad - \frac{p(N-1)}{p-1} \int_{\partial\Omega} \left(\frac{1}{N} u_{\nu} + H|u_{\nu}|^p \right) d\sigma \\
 &\quad - p(p-2) \left(1 + \frac{1}{N(p-1)} \right) \int_{\partial\Omega} |\nabla u|^{p-1} \langle D^2 u \frac{\nabla u}{|\nabla u|}, \frac{\nabla u}{|\nabla u|} \rangle d\sigma.
 \end{aligned} \tag{3.2}$$

In order to estimate from above $\mathcal{I}_p(u)$, we want to determine the sign of the last term in (3.2). By Lemma 2.2, we know that either $P_\nu > 0$ on $\partial\Omega$ or P is constant in $\bar{\Omega}$. If the second case occurs, then, as in the proof of Lemma 2.6-(ii), all level sets of u are concentric spheres, and in particular Ω is a ball. Without loss of generality we can suppose Ω to be a ball centered in the origin B_r , thus, the unique solution of (1.4) is w_r , given in (2.15), with $z = 0$. Then, by straightforward calculations, we have for every $x \in \partial B_r$

$$H(x) = -\frac{1}{N-1} \operatorname{div} \frac{\nabla w_r}{|\nabla w_r|} = \frac{1}{N-1} \sum_{i=1}^N \left(\frac{1}{|x|} - \frac{x_i^2}{|x|^3} \right) = \frac{1}{r}$$

and

$$(w_r)_\nu(x) = -|\nabla w_r(x)| = -\frac{1}{N^{\frac{1}{p-1}}} r^{\frac{1}{p-1}}.$$

Hence,

$$\frac{1}{N} (w_r)_\nu(x) + H(x) |(w_r)_\nu(x)|^p = 0 \quad \text{for every } x \in \partial B_r$$

and the inequality in (ii) is satisfied with the equality sign and we are done. We consider now the remaining case $P_\nu > 0$ on $\partial\Omega$. In this case

$$(p-1)|u_\nu|^{p-2}u_{\nu\nu} + \frac{1}{N} < 0 \quad \text{on } \partial\Omega$$

(cf. (2.8) and remember that $u_\nu < 0$ on $\partial\Omega$), or equivalently

$$u_{\nu\nu} < -\frac{|u_\nu|^{2-p}}{N(p-1)} \quad \text{on } \partial\Omega.$$

Hence, $u_{\nu\nu} < 0$ on $\partial\Omega$, and so, when $1 < p < 2$, we get

$$\mathcal{I}_p(u) \leq -\frac{p(N-1)}{p-1} \int_{\partial\Omega} \left(\frac{1}{N} u_\nu + H|u_\nu|^p \right) d\sigma.$$

(iii) Since u is a solution of (1.4), by Divergence Theorem and Hölder's inequality we have

$$\begin{aligned} |\Omega| &= \int_{\Omega} -\Delta_p u dx = - \int_{\Omega} \operatorname{div}(|\nabla u|^{p-2} \nabla u) dx = - \int_{\partial\Omega} |\nabla u|^{p-2} \nabla u \cdot \nu d\sigma \\ &= \int_{\partial\Omega} |u_\nu|^{p-1} d\sigma \leq \left(\int_{\partial\Omega} |u_\nu|^p d\sigma \right)^{\frac{p-1}{p}} |\partial\Omega|^{\frac{1}{p}}. \end{aligned}$$

By using the definition of H_0 , the previous estimate reads as

$$\left(\int_{\partial\Omega} |u_\nu|^p d\sigma \right)^{\frac{1}{p'}} \geq \frac{|\Omega|}{|\partial\Omega|^{\frac{1}{p}}} = \frac{|\partial\Omega|^{\frac{1}{p'}}}{NH_0}.$$

Consequently, by Hölder's inequality,

$$- \int_{\partial\Omega} u_\nu d\sigma \leq \|u_\nu\|_{L^p(\partial\Omega)} |\partial\Omega|^{\frac{1}{p'}} \leq NH_0 \left(\int_{\partial\Omega} |u_\nu|^p d\sigma \right)^{\frac{1}{p} + \frac{1}{p'}} = NH_0 \int_{\partial\Omega} |u_\nu|^p d\sigma.$$

1 By using this inequality, the right-hand side of (2.7) can be estimated as

$$-(N-1) \int_{\partial\Omega} \left(\frac{1}{N} u_\nu + H|u_\nu|^p \right) d\sigma \leq (N-1) \int_{\partial\Omega} |u_\nu|^p (H_0 - H) d\sigma. \quad (3.3)$$

Therefore, in view of part (ii) of the present theorem, we have for $1 < p < 2$

$$\mathcal{I}_p(u) \leq \frac{p(N-1)}{p-1} \int_{\partial\Omega} |u_\nu|^p (H_0 - H) d\sigma.$$

2 This concludes the proof. \square

Remark 3.2. From parts (i) and (iii) of the previous theorem, since $|u_\nu|^p$ is bounded on $\partial\Omega$, we have the following upper bound for the L^1 -norm of the mean curvature H of $\partial\Omega$

$$\int_{\partial\Omega} H d\sigma \leq H_0 |\partial\Omega| = \frac{|\partial\Omega|^2}{N|\Omega|}.$$

3 The previous theorem allows us to give an alternative proof of the Soap Bubble
4 Theorem in the case in which the hypersurface is a level set of the solution of
5 problem (1.4).

6 *Proof of Theorem 1.1.* The scheme of the proof is the following: a. \Rightarrow c. \Rightarrow b. \Rightarrow
7 c. \Rightarrow a., this proves that a., b. and c. are all equivalent; then we will prove that a.
8 \Rightarrow d. \Rightarrow c., and finally b. \Rightarrow e.

9 a. \Rightarrow c. If $\Omega = B_r$, the only solution of (1.4) is the radial function w_r defined in
10 (2.15).

11 c. \Rightarrow b. As in the proof of Theorem 3.1-(ii), if the solution of (1.4) is radial,
12 $\Omega = B_r$ for some $r > 0$, and so $u = w_r$. Hence, by strighforward calculations, b.
13 holds true.

14 b. \Rightarrow c. By Theorem 3.1-(ii), we get $\mathcal{I}_p(u) = 0$, which in turn implies that u is
15 radial, by Lemma 2.6.

16 c. \Rightarrow a. If u is radial, then $\Gamma = \partial\Omega$, being a level set of u , is a sphere, and so Ω
17 is a ball.

18 a. \Rightarrow d. If $\Omega = B_r$ for some $r > 0$, then $u = w_r$ and so, for every $x \in \partial\Omega$

$$H(x) = -\frac{1}{N-1} \operatorname{div} \frac{\nabla w_r}{|\nabla w_r|} = \frac{1}{N-1} \sum_{i=1}^N \left(\frac{1}{|x|} - \frac{x_i^2}{|x|^3} \right) = \frac{1}{r} = \frac{|\partial B_r|}{N|B_r|} = H_0. \quad (3.4)$$

19

20 d. \Rightarrow c. By Theorem 3.1-(iii), we get $\mathcal{I}_p(u) = 0$, which in turn implies that u is
21 radial, by Lemma 2.6.

b. \Rightarrow e. Up to now, we have proved that a., b., c. and d. are equivalent. Thus,
if b. holds, we have by d.

$$|u_\nu|^{p-2} u_\nu = -\frac{1}{NH_0} \quad \text{on } \partial\Omega.$$

We recall that, on $\partial\Omega$, $\nu = -\frac{\nabla u}{|\nabla u|}$ and consequently $u_\nu = \nabla u \cdot \nu = -|\nabla u|$. Therefore,

$$|u_\nu|^{p-2} u_\nu = -|\nabla u|^{p-1} = -\frac{1}{NH_0} \quad \text{on } \partial\Omega,$$

22 which gives e. \square

23 In the remaining part of this section, we give an upper bound of the integral
24 $\mathcal{I}_p(u)$ in terms of the $L^1(\partial\Omega)$ -norm of the difference between the mean curvature
1 of $\partial\Omega$ and the reference constant H_0 . We start with some preliminary results.

Lemma 3.3. *Let $\Omega = \mathcal{A}(R_1, R_2)$ be an annulus of radii $0 < R_1 < R_2$, then there exists a unique $\bar{R} \in (R_1, R_2)$ such that the positive radial function*

$$u_{\mathcal{A}}(r) := \begin{cases} \int_{R_1}^r \left(\frac{\bar{R}^N}{N\tau^{N-1}} - \frac{\tau}{N} \right)^{\frac{1}{p-1}} d\tau & \text{for every } r \in [R_1, \bar{R}], \\ \int_r^{R_2} \left(\frac{\tau}{N} - \frac{\bar{R}^N}{N\tau^{N-1}} \right)^{\frac{1}{p-1}} d\tau & \text{for every } r \in (\bar{R}, R_2] \end{cases} \quad (3.5)$$

is of class $C^1([R_1, R_2])$ and solves (1.4). Furthermore, $u_{\mathcal{A}}$ achieves its maximum at \bar{R} , where with abuse of notation we have written $u_{\mathcal{A}}(x) = u_{\mathcal{A}}(r)$ for $|x| = r$.

Proof. Suppose first that such \bar{R} exists and belongs to (R_1, R_2) . In this case, it is straightforward to verify that the function $u_{\mathcal{A}}$ given in (3.5) solves problem (1.4), which can be written in radial form as

$$\begin{cases} |u'_{\mathcal{A}}|^{p-2} [(p-1)u''_{\mathcal{A}} + \frac{N-1}{r}u'_{\mathcal{A}}] = -1 & \text{in } (R_1, R_2), \\ u_{\mathcal{A}}(R_1) = u_{\mathcal{A}}(R_2) = 0, \end{cases}$$

where the symbol $'$ denotes the derivative with respect to r .

Now, if we consider the two functions

$$\begin{aligned} F_1 : \rho \in [R_1, R_2] &\mapsto \int_{R_1}^{\rho} \left(\frac{\rho^N}{N\tau^{N-1}} - \frac{\tau}{N} \right)^{\frac{1}{p-1}} d\tau \in \mathbb{R}, \\ F_2 : \rho \in [0, R_2] &\mapsto \int_{\rho}^{R_2} \left(\frac{\tau}{N} - \frac{\rho^N}{N\tau^{N-1}} \right)^{\frac{1}{p-1}} d\tau \in \mathbb{R}, \end{aligned}$$

they have the following properties:

$$\begin{aligned} F_1(R_1) &= F_2(R_2) = 0, \\ 0 < F_1(\rho) < +\infty &\text{ for every } \rho \in (R_1, R_2], \quad 0 < F_2(\rho) < +\infty \text{ for every } \rho \in [0, R_2), \\ F_1'(\rho) &= \frac{1}{p-1} \int_{R_1}^{\rho} \left(\frac{\rho^N}{N\tau^{N-1}} - \frac{\tau}{N} \right)^{\frac{2-p}{p-1}} \left(\frac{\rho}{\tau} \right)^{N-1} d\tau > 0 \text{ for every } \rho \in (R_1, R_2], \\ F_2'(\rho) &= -\frac{1}{p-1} \int_{\rho}^{R_2} \left(\frac{\tau}{N} - \frac{\rho^N}{N\tau^{N-1}} \right)^{\frac{2-p}{p-1}} \left(\frac{\rho}{\tau} \right)^{N-1} d\tau < 0 \text{ for every } \rho \in [0, R_2). \end{aligned}$$

Therefore, there exists a unique $\rho = \bar{R} \in (R_1, R_2)$ for which $F_1(\bar{R}) = F_2(\bar{R})$. This concludes the proof. \square

Definition 3.4. A domain $\Omega \subset \mathbb{R}^N$ satisfies the *uniform interior and exterior touching sphere conditions*, and we denote with ρ_i and ρ_e the optimal interior and exterior radii respectively, if for any $x_0 \in \partial\Omega$ there exist two balls $B_{\rho_i}(c^-) \subset \Omega$ and $B_{\rho_e}(c^+) \subset \mathbb{R}^N \setminus \bar{\Omega}$ such that $x_0 \in \partial B_{\rho_i}(c^-) \cap \partial B_{\rho_e}(c^+)$. We call *optimal radius* the minimum between the interior and the exterior radius, $\rho := \min\{\rho_i, \rho_e\}$.

We observe that if Ω is of class C^2 , then it satisfies the uniform interior and exterior touching sphere conditions.

Proposition 3.5. *Let $\Omega \subset \mathbb{R}^N$ be a bounded domain of class C^2 and $u \in C^1(\bar{\Omega})$ be a solution of (1.4) in Ω . Then*

$$\left(\frac{\rho_i}{N} \right)^{\frac{1}{p-1}} \leq |\nabla u| \leq \left[\frac{(\text{diam}(\Omega) + \rho_e)^N}{N\rho_e^{N-1}} - \frac{\rho_e}{N} \right]^{\frac{1}{p-1}} \quad \text{on } \partial\Omega.$$

Proof. We follow the ideas in [17, Theorem 3.10]. Let x_0 be any point on the boundary $\partial\Omega$. Without loss of generality, we can place the origin at c^- . Thus, the function

$$u_{\rho_i} := -\frac{p-1}{pN^{\frac{1}{p-1}}} \left(|x|^{\frac{p}{p-1}} - \rho_i^{\frac{p}{p-1}} \right)$$

is the solution of (1.4) in B_{ρ_i} . Now, being by definition $B_{\rho_i} \subset \Omega$,

$$\begin{cases} -\Delta_p u_{\rho_i} = -\Delta_p u & \text{in } B_{\rho_i}, \\ u_{\rho_i} \leq u & \text{on } \partial B_{\rho_i}, \end{cases}$$

and so, by comparison [13, Lemma 3.7], $u_{\rho_i} \leq u$ in B_{ρ_i} . Since $u_{\rho_i}(x_0) = u(x_0)$, we have $\partial_\nu(u_{\rho_i} - u)(x_0) > 0$, where ν is the external unit normal to B_{ρ_i} . This gives the first inequality in the statement, namely

$$|\nabla u(x_0)| \geq \left(\frac{\rho_i}{N} \right)^{\frac{1}{p-1}}.$$

On the other hand, let $\mathcal{A} := \mathcal{A}(\rho_e, \text{diam}(\Omega) + \rho_e)$ be the annulus centered at c^+ . By definition, $\Omega \subset \mathcal{A}$. Again, without loss of generality, we can place the origin at c^+ and consider the function $u_{\mathcal{A}}$ whose expression is given by (3.5) with $R_1 := \rho_e$ and $R_2 := \text{diam}(\Omega) + \rho_e$. Reasoning as above we have

$$\begin{cases} -\Delta_p u_{\mathcal{A}} = -\Delta_p u & \text{in } \Omega, \\ u_{\mathcal{A}} \geq u & \text{on } \partial\Omega, \end{cases}$$

and so $u_{\mathcal{A}} \geq u$ in Ω . Therefore, $\partial_\nu(u_{\mathcal{A}} - u)(x_0) \leq 0$, being ν the external unit normal to \mathcal{A} . This finally gives

$$|\nabla u(x_0)| \leq \left(\frac{\bar{R}^N}{N\rho_e^{N-1}} - \frac{\rho_e}{N} \right)^{\frac{1}{p-1}} \leq \left(\frac{(\rho_e + \text{diam}(\Omega))^N}{N\rho_e^{N-1}} - \frac{\rho_e}{N} \right)^{\frac{1}{p-1}}$$

2 and concludes the proof. \square

3 Combining together the results in Proposition 3.5 and Theorem 3.1, we get the
4 following corollary.

5 **Corollary 3.6.** *Let $1 < p < 2$ and $\Omega \subset \mathbb{R}^N$ be a $C^{2,\alpha}$ bounded domain. If u solves
6 (1.4) and has $|\mathcal{C}| = 0$, the following chain of inequalities holds*

$$0 \leq \mathcal{I}_p(u) \leq \frac{p(N-1)}{p-1} \left[\frac{(\text{diam}(\Omega) + \rho_e)^N}{N\rho_e^{N-1}} - \frac{\rho_e}{N} \right]^{\frac{p}{p-1}} \|H_0 - H\|_{L^1(\partial\Omega)}. \quad (3.6)$$

7 4. SOME COMMENTS ON THE STABILITY

8 With reference to the result given in Corollary 3.6, we observe that, while $\mathcal{I}_p(u)$
9 is related to the solution of problem (1.4), the constant that bounds from above
10 $\mathcal{I}_p(u)$ in (3.6) depends only on the geometry of the problem. In particular, the
11 non-negative integral $\mathcal{I}_p(u)$ that vanishes only on radial functions, goes to zero as
12 $H \rightarrow H_0$ in $L^1(\partial\Omega)$. In view of Corollary 3.6, this suggests, at least qualitatively, a
13 sort of stability of the Serrin-type result for the overdetermined problem with the
14 p -Laplacian.

15 In [10], Ciruolo and Vezzoni obtained the following stability result for the Soap
1 Bubble Theorem by Alexandrov.

Theorem 4.1 (Theorem 1.1 of [10]). *Let $\partial\Omega$ be a C^2 -regular, connected, and closed hypersurface embedded in \mathbb{R}^N . If*

$$\|H - H_0\|_{L^\infty(\partial\Omega)} < \varepsilon$$

for some $\varepsilon > 0$ depending only on N , $|\partial\Omega|$, and upper bounds on the inverse of the optimal radius (cf. Definition 3.4) ρ^{-1} of $\partial\Omega$, then $\partial\Omega \subset \bar{B}_{r_e} \setminus B_{r_i}$, with

$$0 < r_e - r_i \leq C\varepsilon,$$

2 where $C > 0$ depends on N , $|\partial\Omega|$, and upper bounds on the inverse of the optimal
3 radius ρ^{-1} of $\partial\Omega$.

4 This result gives an estimate of $r_e - r_i$ in terms of the $L^\infty(\partial\Omega)$ -norm of $H - H_0$.
Furthermore, as a consequence, for every $1 < p < \infty$, it is possible to compare
the solution u of (1.4) with the radial solutions

$$u_e(x) := -\frac{p-1}{pN^{\frac{1}{p-1}}} \left(|x|^{\frac{p}{p-1}} - (r_e)^{\frac{p}{p-1}} \right) \quad \text{for every } x \in B_{r_e}$$

and

$$u_i(x) := -\frac{p-1}{pN^{\frac{1}{p-1}}} \left(|x|^{\frac{p}{p-1}} - (r_i)^{\frac{p}{p-1}} \right) \quad \text{for every } x \in B_{r_i}$$

of

$$\begin{cases} -\Delta_p u_e = 1 & \text{in } B_{r_e}, \\ u_e = 0 & \text{on } \partial B_{r_e}, \end{cases} \quad \text{and} \quad \begin{cases} -\Delta_p u_i = 1 & \text{in } B_{r_i}, \\ u_i = 0 & \text{on } \partial B_{r_i}, \end{cases}$$

respectively. Indeed, by the weak comparison principle [13, Lemma 3.7], we easily
get

$$u \geq u_i \text{ in } B_{r_i} \quad \text{and} \quad u \leq u_e \text{ in } \Omega,$$

giving in particular the following estimate of u in terms of the radial solutions u_i
and u_e on the interior ball B_{r_i}

$$-\frac{p-1}{pN^{\frac{1}{p-1}}} \left(|x|^{\frac{p}{p-1}} - (r_i)^{\frac{p}{p-1}} \right) \leq u(x) \leq -\frac{p-1}{pN^{\frac{1}{p-1}}} \left(|x|^{\frac{p}{p-1}} - (r_e)^{\frac{p}{p-1}} \right) \quad \text{in } B_{r_i}.$$

5 It is quite challenging to obtain an estimate from below of $\mathcal{I}_p(u)$ in terms of some
6 increasing function of $r_e - r_i$. This would allow to improve—at least in some relevant
7 cases—the stability result in Theorem 4.1, getting a stability result in terms of the
8 $L^1(\partial\Omega)$ -norm, instead of the $L^1(\partial\Omega)$ -norm, of $H - H_0$. This approach was proposed
9 by Magnanini and Poggesi for the case $p = 2$ in [17], where the authors used in
10 a very clever way the mean value property for harmonic functions. Nevertheless,
11 their method works well only in the linear case and seems very difficult to generalize
12 it to the case $p \neq 2$. Some other issues related to the stability of the symmetry
13 result for the overdetermined p -Laplacian problem are treated in [23].

14

ACKNOWLEDGMENTS

15 The authors were supported by the INdAM - GNAMPA Project 2017 “Regolarità
16 delle soluzioni viscosse per equazioni a derivate parziali non lineari degeneri” and by
17 University of Bologna. F.C. acknowledges also the support of the projects “Ricerca
18 Locale 2018 Linea B - Problemi non lineari” from University of Turin and INdAM
19 - GNAMPA Project 2019 “Il modello di Born-Infeld per l’elettromagnetismo non-
20 lineare: esistenza, regolarità e molteplicità di soluzioni”. F.F. was also supported
1 by INdAM - GNAMPA Project 2019 “Proprietà di regolarità delle soluzioni viscosse

con applicazioni a problemi di frontiera libera”. The authors wish to thank Professors G. Ciraolo and R. Magnanini, whose comments were crucial to improve the bibliography of the present paper.

REFERENCES

- [1] A. Aftalion, J. Busca, and W. Reichel. Approximate radial symmetry for overdetermined boundary value problems. *Adv. Differential Equations*, 4(6):907–932, 1999.
- [2] A. D. Alexandrov. A characteristic property of spheres. *Ann. Mat. Pura Appl. (4)*, 58:303–315, 1962.
- [3] C. Bianchini and G. Ciraolo. Wulff shape characterizations in overdetermined anisotropic elliptic problems. *Comm. Partial Differential Equations*, 43(5):790–820, 2018.
- [4] B. Brandolini, C. Nitsch, P. Salani, and C. Trombetti. On the stability of the Serrin problem. *J. Differential Equations*, 245(6):1566–1583, 2008.
- [5] F. Brock. Symmetry for a general class of overdetermined elliptic problems. *Nonlinear Differential Equations and Applications NoDEA*, 23(3):36, 2016.
- [6] F. Brock and A. Henrot. A symmetry result for an overdetermined elliptic problem using continuous rearrangement and domain derivative. *Rend. Circ. Mat. Palermo (2)*, 51(3):375–390, 2002.
- [7] G. Ciraolo and F. Maggi. On the shape of compact hypersurfaces with almost-constant mean curvature. *Comm. Pure Appl. Math.*, 70(4):665–716, 2017.
- [8] G. Ciraolo, R. Magnanini, and V. Vespi. Hölder stability for Serrin’s overdetermined problem. *Ann. Mat. Pura Appl. (4)*, 195(4):1333–1345, 2016.
- [9] G. Ciraolo and A. Roncoroni. Serrin’s type overdetermined problems in convex cones. *ArXiv preprint arXiv:1806.08553*, 2018.
- [10] G. Ciraolo and L. Vezzoni. A sharp quantitative version of Alexandrov’s theorem via the method of moving planes. *ArXiv preprint arXiv:1501.07845*, 2015.
- [11] L. Damascelli and F. Pacella. Monotonicity and symmetry results for p -Laplace equations and applications. *Adv. Differential Equations*, 5(7-9):1179–1200, 2000.
- [12] A. Farina and B. Kawohl. Remarks on an overdetermined boundary value problem. *Calculus of Variations and Partial Differential Equations*, 31(3):351–357, 2008.
- [13] I. Fragalà, F. Gazzola, and B. Kawohl. Overdetermined problems with possibly degenerate ellipticity, a geometric approach. *Mathematische Zeitschrift*, 254(1):117–132, 2006.
- [14] N. Garofalo and J. L. Lewis. A symmetry result related to some overdetermined boundary value problems. *American Journal of Mathematics*, 111(1):9–33, 1989.
- [15] B. Kawohl. Symmetrization—or how to prove symmetry of solutions to a pde. *Partial differential equations (Prague, 1998)*, 406:214–229, 1999.
- [16] G. M. Lieberman. Boundary regularity for solutions of degenerate elliptic equations. *Nonlinear Anal.*, 12(11):1203–1219, 1988.
- [17] R. Magnanini and G. Poggesi. On the stability for Alexandrov’s soap bubble theorem. *ArXiv preprint arXiv:1610.07036*, 2016.
- [18] R. Magnanini and G. Poggesi. Serrin’s problem and Alexandrov’s soap bubble theorem: enhanced stability via integral identities. *ArXiv preprint arXiv:1708.07392*, 2017.
- [19] R. Magnanini and G. Poggesi. Nearly optimal stability for serrin’s problem and the soap bubble theorem. *ArXiv preprint arXiv:1903.04823*, 2019.
- [20] F. Pacella and G. Tralli. Overdetermined problems and constant mean curvature surfaces in cones. *ArXiv preprint arXiv:1802.03197*, 2018.
- [21] L. E. Payne and P. W. Schaefer. Duality theorems in some overdetermined boundary value problems. *Math. Methods Appl. Sci.*, 11(6):805–819, 1989.
- [22] G. Poggesi. *The Soap Bubble Theorem and Serrin’s problem: quantitative symmetry*. ArXiv:1902.08584, 2019.
- [23] C. Pucci and A. Colesanti. A symmetry result for the p -laplacian equation via the moving planes method. *Applicable Analysis*, 55(3-4):207–213, 1994.
- [24] P. Pucci and J. Serrin. *The maximum principle*, volume 73 of *Progress in Nonlinear Differential Equations and their Applications*. Birkhäuser Verlag, Basel, 2007.
- [25] R. C. Reilly. Mean curvature, the laplacian, and soap bubbles. *The American Mathematical Monthly*, 89(3):180–198, 1982.

- 2 [26] J. Serrin. A symmetry problem in potential theory. *Arch. Rational Mech. Anal.*, 43:304–318,
3 1971.
- 4 [27] R. P. Sperb. *Maximum principles and their applications*. Elsevier, 1981.
- 5 [28] J. L. Vázquez. A strong maximum principle for some quasilinear elliptic equations. *Appl.*
6 *Math. Optim.*, 12(3):191–202, 1984.
- 7 [29] H. F. Weinberger. Remark on the preceding paper of Serrin. *Arch. Rational Mech. Anal.*,
8 43:319–320, 1971.
- 9 FRANCESCA COLASUONNO
10 DIPARTIMENTO DI MATEMATICA “GIUSEPPE PEANO”
11 UNIVERSITÀ DEGLI STUDI DI TORINO
12 VIA CARLO ALBERTO, 10
13 10123 TORINO, ITALY
14 *Email address:* francesca.colasuonno@unito.it
- 15 FAUSTO FERRARI
16 DIPARTIMENTO DI MATEMATICA
17 ALMA MATER STUDIORUM UNIVERSITÀ DI BOLOGNA
18 PIAZZA DI PORTA S. DONATO, 5
19 40126 BOLOGNA, ITALY
364 *Email address:* fausto.ferrari@unibo.it