

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

		c
Mediterranean	anrannianc	tiahtina
Mediterraniean	quiquillalis	HIGHLING

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Eva Turicchia, M.A. (2020). Mediterranean gorgonians fighting. MARINE BIODIVERSITY, 50(3), 1-2 [10.1007/s12526-020-01064-w].

Availability:

This version is available at: https://hdl.handle.net/11585/757535 since: 2021-10-30

Published:

DOI: http://doi.org/10.1007/s12526-020-01064-w

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Turicchia E, Abbiati M, Ponti M (2020) Mediterranean gorgonians fighting. Marine Biodiversity 50: 33

The final published version is available online at https://dx.doi.org/10.1007/s12526-020-01064-w

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/)

When citing, please refer to the published version.

Mediterranean gorgonians fighting

2

1

- 3 Eva Turicchia^{1,*}, Marco Abbiati², Massimo Ponti³
- 4 CIRI FRAME, University of Bologna, Via Sant'Alberto 163, 48123 Ravenna, Italy
- 5 e-mail: eva.turicchia2@unibo.it
- 6 ² DBC Department, University of Bologna, Via degli Ariani 1, 48121 Ravenna, Italy
- ³ BiGeA Department, University of Bologna, Via Sant'Alberto 163, 48123 Ravenna, Italy
- 8 * Corresponding author

9

26

- 10 The long-lived, slow-growing, and low-resilience sea fans *Paramuricea clavata* (Risso, 1826) and 11 Eunicella cavolini (Koch, 1887) are keystone species of the Mediterranean coralligenous 12 assemblages and are affected by anthropogenic disturbances and mass mortality events triggered by 13 marine heatwaves (Turicchia et al. 2018; Garrabou et al. 2019). The frequent closeness between 14 colonies of these two sympatric species throughout all the Mediterranean Sea suggests weak 15 interspecific competition, at least at low colony densities, despite a large overlap in their 16 bathymetric distribution (Di Camillo et al. 2018; Ponti et al. 2018). However, a direct interaction 17 between their colonies has never been reported. We photographically documented the accidental 18 contact, maybe caused by basal rocks' movement, between branches of P. clavata and E. cavolini at 19 28 m depth at Capo Calvo, Elba Island, Italy (42.7350° N, 10.4342° E, Geodetic Datum WGS84), the May 25th, 2019 (Fig. 1a). The branch portions of *P. clavata* in touch with those of *E. cavolini* 20 21 showed depigmented coenenchyme and bare skeletons, while the E. cavolini ones appeared healthy 22 (Fig. 1a, b). 23 The interspecific interactions by contact with neighbouring specimens, although well documented
- for many scleractinians, are still little investigated in gorgonian octocorals. For the purpose, some

corals develop specialised (i.e. sweeper and thread-like) tentacles. Around the contact area, there

were thread-like tentacles in polyps of *P. clavata*, but not in those of *E. cavolini* (Fig. 1c). In *P.*

27	clavata, thread-like tentacles, with high densities of nematocysts, have been reported to be involved
28	in feeding activity (Lopez-Gonzalez et al. 2018). Although these modified tentacles could also have
29	a defensive function, they seem poorly effective weapon against E. cavolini, which appeared to be
30	the winner in direct fighting.
31	
32	Acknowledgements: we are grateful to Biodivers Elba Sea Academy for the valuable diving
33	assistance.
34	
35	Funding: This study was partially funded by Reef Check Italia onlus, which covered expedition
36	direct costs.
37	
38	Conflict of Interest: The authors declare that they have no conflict of interest.
39	
40	Ethical approval: No animal testing was performed during this study.
41	
42	Sampling and field studies: The study is compliant with CBD and Nagoya protocols.
43	
44	Author Contribution Statement: ET and MP conceived research and conducted field
45	observations. ET wrote the manuscript. All authors read and approved the final manuscript.
46	
47	References
48	Di Camillo CG, Ponti M, Bavestrello G, Krzelj M, Cerrano C (2018) Building a baseline for
49	habitat-forming corals by a multi-source approach, including Web Ecological Knowledge.
50	Biodivers Conserv 27:1257-1276. https://doi.org/10.1007/s10531-017-1492-8
51	Garrabou J et al. (2019) Collaborative database to track mass mortality events in the mediterranean
52	sea. Front Mar Sci 6:707. https://doi.org/10.3389/fmars.2019.00707

Lopez-Gonzalez PJ, Bramanti L, Escribano-Alvarez P, Benedetti MC, Martinez-Baraldes I, Megina C (2018) Thread-like tentacles in the Mediterranean corals *Paramuricea clavata* and *Corallium rubrum*. Mediterr Mar Sci 19:394-397. https://doi.org/10.12681/mms.13807
Ponti M, Turicchia E, Ferro F, Cerrano C, Abbiati M (2018) The understorey of gorgonian forests in mesophotic temperate reefs. Aquat Conserv 28:1153–1166. https://doi.org/10.1002/aqc.2928
Turicchia E, Abbiati M, Sweet M, Ponti M (2018) Mass mortality hits gorgonian forests at Montecristo Island. Dis Aquat Org 131:79-85. https://doi.org/10.3354/dao03284




Fig. 1 Colonies of *Paramuricea clavata* and *Eunicella cavolini* in direct contact: **a)** contact points (scale bar 5 cm); **b)** close up of depigmented coenenchyme and bare skeleton of *P. clavata* (scale bar 2 cm); and **c)** particular of thread-like tentacles in polyps of *P. clavata* around the contact area (scale bar 1 cm).