



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

On the refinement matrix mask of interpolating Hermite splines

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Romani, L., Viscardi, A. (2020). On the refinement matrix mask of interpolating Hermite splines. *APPLIED MATHEMATICS LETTERS*, 109, 1-8 [10.1016/j.aml.2020.106524].

Availability:

This version is available at: <https://hdl.handle.net/11585/760551> since: 2021-02-21

Published:

DOI: <http://doi.org/10.1016/j.aml.2020.106524>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Lucia Romani, Alberto Viscardi: On the refinement matrix mask of interpolating Hermite splines, Applied Mathematics Letters, Volume 109, 2020, 106524

The final published version is available online at:

<https://doi.org/10.1016/j.aml.2020.106524>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

On the refinement matrix mask of interpolating Hermite splines

Lucia Romani^a, Alberto Viscardi^{a,*}

^a*Dipartimento di Matematica, Alma Mater Studiorum Università di Bologna,
Piazza di Porta San Donato 5, 40126 Bologna, Italy*

Abstract

We propose a new computational approach for constructing the refinement matrix mask of interpolating Hermite splines of any order and with general dilation factor. Our strategy exploits the refinability properties of cardinal B-splines with simple knots and simplifies the constructive procedures proposed so far.

Keywords: Cardinal Hermite Spline Interpolation; Cardinal B-Splines; Vector Subdivision; Refinement Matrix Mask; Arbitrary Dilation Factor

1. Introduction

Hermite subdivision schemes with dilation factor $m \in \mathbb{N} \setminus \{1\}$ form a special subclass of vector subdivision schemes which started to be investigated in the early 90's (see, e.g., [3, 4, 7, 8]) but is still subject of recent research [1, 2, 5, 6, 10]. The distinctive feature of Hermite schemes is the fact that the sequence of vector data $\mathbf{p}^{[j+1]} = \{ \mathbf{p}_k^{[j+1]} \}_{k \in \mathbb{Z}}$, generated at refinement level $j+1$ ($j \in \mathbb{N}_0$) by the application of a *subdivision matrix* $\mathbf{A}^{[j]}$ to the previous data $\mathbf{p}^{[j]}$ so that $\mathbf{p}^{[j+1]} = \mathbf{A}^{[j]} \mathbf{p}^{[j]}$, contains vectors $\mathbf{p}_k^{[j+1]} = [p_{k,0}^{[j+1]}, \dots, p_{k,n}^{[j+1]}]^T \in \mathbb{R}^{n+1}$ ($n \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}$) whose components correspond to the evaluation at $m^{-(j+1)}\mathbb{Z}$ of a certain function and its derivatives up to order n (see, e.g., Figure 1).

In the present paper we investigate Hermite subdivision schemes capable of generating, in the limit, interpolating Hermite splines. Precisely, for any given initial sequence of vector data $\{ \mathbf{p}_k^{[0]} \in \mathbb{R}^{n+1} \}_{k \in \mathbb{Z}}$, the vector-valued limit function $\mathbf{f} = [f_0, \dots, f_n]^T \in \mathcal{C}^0(\mathbb{R}, \mathbb{R}^{n+1})$, linear combination of the shifts of the basic limit functions $\{ \phi_{0,s}^{[0]} \}_{s=0}^n$ of the scheme, is formed by a first scalar-valued function $f_0 \in \mathcal{C}^n(\mathbb{R})$ which is a polynomial spline of degree $2n+1$ satisfying $f_s(k) = f_0^{(s)}(k) = p_{k,s}^{[0]}$, for all $k \in \mathbb{Z}$ and $s \in \{0, \dots, n\}$. For any arbitrary dilation factor m and order n , such a Hermite scheme is described by the finite matrix sequence $\{ \mathbf{A}_k \in \mathbb{R}^{(n+1) \times (n+1)} \}_{k=1-m}^{m-1}$ which is called the *refinement matrix mask* and is the building block to define the subdivision matrices $\{ \mathbf{A}^{[j]} \}_{j \in \mathbb{N}_0}$.

The construction of such a refinement matrix mask is already known in the literature for $m=2$, $n \in \mathbb{N}_0$. The authors in [7] constructed the matrices $\{ \mathbf{A}_k \}_{k=-1,0,1}$ by introducing the transformation matrices \mathbf{H}_L and \mathbf{H}_R yielding $\{ \mathbf{A}_k = \mathbf{H}_L \tilde{\mathbf{A}}_k \mathbf{H}_R \}_{k=-1,0,1}$ with $\{ \tilde{\mathbf{A}}_k \}_{k=-1,0,1}$ the refinement matrix mask of degree- $(2n+1)$ B-splines with knots of multiplicity $n+1$ derived in [9]. An alternative approach, proposed more recently in [10], constructs explicitly the degree- $(2n+1)$ polynomials defining the basic limit functions $\{ \phi_{0,s}^{[0]} \}_{s=0}^n$ on the intervals $[-1, 0]$ and $[0, 1]$ which cover their supports. The refinement mask $\{ \mathbf{A}_k \}_{k=-1,0,1}$ is then obtained from the refinement equation, exploiting the evaluation of those polynomials.

The goal of our work is to provide an easier construction of the refinement matrix mask of interpolating Hermite splines for any arbitrary order and dilation factor, by exploiting the refinability properties of degree- $(2n+1)$ cardinal B-splines with simple knots. In Section 2 we recall basic facts about vector

*Corresponding author

Email addresses: lucia.romani@unibo.it (Lucia Romani), alberto.viscardi@unibo.it (Alberto Viscardi)

The vector scheme in (1) is said to be *convergent* if and only if there exist continuous vector-valued functions

$$\begin{aligned} \phi_{k,s}^{[j]} : \mathbb{R} &\longrightarrow \mathbb{R}^{n+1} \\ x &\longmapsto \begin{bmatrix} \varphi_{0,s}^{[j]}(m^j x - k) \\ \vdots \\ \varphi_{n,s}^{[j]}(m^j x - k) \end{bmatrix}, \quad s \in \{0, \dots, n\}, j \in \mathbb{N}_0, k \in \mathbb{Z}, \end{aligned}$$

that, by introducing the matrix notation $\Phi_k^{[j]}(x) := [\phi_{k,0}^{[j]}(x), \dots, \phi_{k,n}^{[j]}(x)]$, satisfy the *refinement equation*

$$\Phi_h^{[j]}(x) = \sum_{k \in \mathbb{Z}} \Phi_k^{[j+1]}(x) \mathbf{A}_{k-mh}^{[j]}, \quad h \in \mathbb{Z}. \quad (4)$$

For any $j \in \mathbb{N}_0$, (4) can be rewritten in matrix form as

$$\Phi^{[j]}(x) = \Phi^{[j+1]}(x) \mathbf{A}^{[j]} \quad \text{with} \quad \Phi^{[j]}(x) := [\dots, \Phi_{-1}^{[j]}(x), \Phi_0^{[j]}(x), \Phi_1^{[j]}(x), \dots]. \quad (5)$$

For any input vector sequence $\mathbf{p}^{[0]}$, a convergent vector scheme generates in the limit a continuous vector-valued function $\mathbf{f} = [f_0, \dots, f_n]^T$ that, in light of (2) and (5), fulfils $\mathbf{f}(x) = \Phi^{[j]}(x) \mathbf{p}^{[j]}$, for all $j \in \mathbb{N}_0$. In particular, for $j = 0$

$$\mathbf{f}(x) = \sum_{k \in \mathbb{Z}} \Phi_k^{[0]}(x) \mathbf{p}_k^{[0]} = \sum_{s=0}^n \sum_{k \in \mathbb{Z}} p_{k,s}^{[0]} \phi_{k,s}^{[0]}(x) \quad \text{with} \quad f_r(x) = \sum_{s=0}^n \sum_{k \in \mathbb{Z}} p_{k,s}^{[0]} \varphi_{r,s}^{[0]}(x - k), \quad r = 0, \dots, n. \quad (6)$$

\mathbf{f} has the regularity of the least regular function among its components f_0, \dots, f_n . Due to (6), the $n+1$ vector-valued functions $\phi_{0,s}^{[0]}(x) = [\varphi_{0,s}^{[0]}(x), \dots, \varphi_{n,s}^{[0]}(x)]^T$, $s \in \{0, \dots, n\}$ are called the *basic limit functions* of the vector scheme and each of them is obtained by refining the initial sequence of vector data defined by $p_{k,r}^{[0]} = \delta_{k,0} \delta_{r,s}$, $r \in \{0, \dots, n\}$, $k \in \mathbb{Z}$.

Note that, when $n = 0$, the above description recovers the known results on scalar subdivision with dilation factor m .

3. A short review of the subclass of interpolating Hermite spline subdivision schemes

The interpolating Hermite spline scheme of order $n \in \mathbb{N}_0$ is a special instance of convergent vector subdivision scheme that, for any choice of $m \in \mathbb{N} \setminus \{1\}$, generates interpolating Hermite splines of degree $2n+1$. The following properties [5, 7] make it a special member of the class of vector subdivision schemes.

i) For all $j \in \mathbb{N}_0$, each matrix in the j th level refinement mask $\{\mathbf{A}_k^{[j]} \in \mathbb{R}^{(n+1) \times (n+1)}\}_{k \in \mathbb{Z}}$ satisfies

$$\mathbf{A}_k^{[j]} = \mathbf{D}^{-j-1} \mathbf{A}_k \mathbf{D}^j \quad \text{with} \quad \mathbf{D} = \begin{bmatrix} 1 & & & \\ & m^{-1} & & \\ & & \ddots & \\ & & & m^{-n} \end{bmatrix};$$

moreover, the refinement mask is compactly supported on $[1-m, m-1] \cap \mathbb{Z}$, i.e. \mathbf{A}_k is the zero matrix for $k > m-1$ and $k < 1-m$.

ii) For all $r, s \in \{0, \dots, n\}$,

$$\varphi_{r,s}^{[0]}(x) = \frac{d^r}{dx^r} \varphi_{0,s}^{[0]}(x) \quad \text{and} \quad \varphi_{r,s}^{[j]}(k) = \delta_{k,0} \delta_{r,s}, \quad \forall j \in \mathbb{N}_0, k \in \mathbb{Z}.$$

In particular, the latter requires, for every $j \in \mathbb{N}_0$, $\mathbf{A}_0^{[j]}$ to be the $(n+1)$ -dimensional identity matrix which implies $\mathbf{A}_0 = \mathbf{D}$ due to i).

iii) For all $s \in \{0, \dots, n\}$, $\varphi_{0,s}^{[0]} \in \mathcal{C}^n(\mathbb{R})$ is a polynomial spline of degree $d = 2n + 1$.

Remark 3.1. Condition i) implies,

$$\text{supp}(\varphi_{r,s}^{[0]}) \subseteq [-1, 1], \quad \forall r, s \in \{0, \dots, n\}. \quad (7)$$

Indeed, consider, for every $j \in \mathbb{N}_0$, $k_L^{[j]}, k_R^{[j]} \in \mathbb{Z}$ such that

$$\mathbf{p}_{k_L^{[j]}}, \mathbf{p}_{k_R^{[j]}} \neq \mathbf{0} \quad \text{and} \quad \mathbf{p}_k^{[j]} = \mathbf{0} \quad \text{for} \quad k < k_L^{[j]} \vee k > k_R^{[j]}.$$

From (1) and i) one gets, for every $j \in \mathbb{N}_0$,

$$m(k_L^{[j]} - 1) + 1 \leq k_L^{[j+1]} \quad \text{and} \quad k_R^{[j+1]} \leq m(k_R^{[j]} + 1) - 1. \quad (8)$$

Since the vectors $\{\mathbf{p}_k^{[j]}\}_{k \in \mathbb{Z}}$ are attached to $m^{-j}\mathbb{Z}$, using (8) we have that, for \mathbf{f} in (6),

$$\bigcup_{r=0}^n \text{supp}(f_r) = \lim_{j \rightarrow \infty} \left[\frac{k_L^{[j+1]} - 1}{m^{j+1}}, \frac{k_R^{[j+1]} + 1}{m^{j+1}} \right] \subseteq \lim_{j \rightarrow \infty} \left[\frac{k_L^{[j]} - 1}{m^j}, \frac{k_R^{[j]} + 1}{m^j} \right] \subseteq [k_L^{[0]} - 1, k_R^{[0]} + 1].$$

As previously observed, for every $s \in \{0, \dots, n\}$, $\mathbf{f} = \phi_{0,s}^{[0]}$ is obtained by $\mathbf{p}^{[0]}$ satisfying $p_{k,r}^{[0]} = \delta_{k,0}\delta_{r,s}$, $r \in \{0, \dots, n\}$, $k \in \mathbb{Z}$, for which $k_L^{[0]} = k_R^{[0]} = 0$.

Remark 3.2. In light of ii) and iii), for any initial data $\mathbf{p}^{[0]}$, f_0 in (6) is a \mathcal{C}^n spline of degree d and, for all $r \in \{0, \dots, n\}$,

$$f_r(x) = \frac{d^r}{dx^r} f_0(x) \quad \text{with} \quad f_r(m^{-j}k) = p_{k,r}^{[j]}, \quad j \in \mathbb{N}_0, k \in \mathbb{Z}. \quad (9)$$

4. Constructing the mask of Hermite splines from cardinal B-splines with simple knots

Let $N_d \in \mathcal{C}^{d-1}(\mathbb{R})$ be the cardinal B-spline of degree $d = 2n + 1$ supported on $[-n - 1, n + 1]$ and defined over the knot vector $\{-n - 1, -n, \dots, 0, \dots, n, n + 1\}$ with all simple knots. N_d satisfies the refinement equation (see, e.g., [11])

$$N_d(x) = \sum_{k=-(m-1)(n+1)}^{(m-1)(n+1)} b_k N_d(mx - k) \quad (10)$$

$$\text{where} \quad \frac{1}{m} \sum_{k=-(m-1)(n+1)}^{(m-1)(n+1)} b_k z^k = z^{-(m-1)(n+1)} \left(\frac{1 - z^m}{m(1 - z)} \right)^{d+1}, \quad z \in \mathbb{C}, |z| = 1.$$

In light of (10), for any given sequence of control points $\mathbf{c}^{[0]} = \{c_k^{[0]} \in \mathbb{R}\}_{k \in \mathbb{Z}}$, we can express the associated spline as

$$\begin{aligned} f(x) &= \sum_{k \in \mathbb{Z}} c_k^{[0]} N_d(x - k) = \sum_{k \in \mathbb{Z}} c_k^{[j]} N_d(m^j x - k), \quad j \in \mathbb{N}_0, \\ \text{where} \quad c_h^{[j+1]} &= \sum_{k \in \mathbb{Z}} b_{h-mk} c_k^{[j]}, \quad h \in \mathbb{Z}. \end{aligned} \quad (11)$$

Thus, for all $j \in \mathbb{N}_0$, $s \in \{0, \dots, n\}$, $y \in \mathbb{R}$,

$$\frac{1}{m^{sj}} f^{(s)}(m^{-j}y) = \frac{1}{m^{sj}} \frac{d^s}{dx^s} f(x) \Big|_{x=m^{-j}y} = \sum_{k \in \mathbb{Z}} c_k^{[j]} N_d^{(s)}(y - k),$$

Proof. $\mathbf{A}_0 = \mathbf{D}$ is already asked in condition ii). For the remaining matrices, we observe that the block $\overline{\mathbf{A}}$ can be extracted from the bigger matrix in (14) with row indices from $n + 1$ to $n + (n + 1)(m - 1)$ and column indices from 0 to $2n + 1 = d$. Thus, on the right-hand side, we need only to consider the rows of \mathbf{N} with indices from 0 to d . Since, for every $s \in \{0, \dots, n\}$, $\text{supp}(N_d^{(s)}) = [-n - 1, n + 1]$, due to (15), we can also cut the columns of \mathbf{N} considering only indices from $-n$ to $n + 1$, leading to

$$\mathbf{N}_R = \left[\begin{array}{c|c|c|c|c} N_d(-n) & N_d(1-n) & \cdots & N_d(n) & 0 \\ N_d^{(1)}(-n) & N_d^{(1)}(1-n) & \cdots & N_d^{(1)}(n) & 0 \\ \vdots & \vdots & & \vdots & 0 \\ N_d^{(n)}(-n) & N_d^{(n)}(1-n) & \cdots & N_d^{(n)}(n) & 0 \\ \hline 0 & N_d(-n) & \cdots & N_d(n-1) & N_d(n) \\ 0 & N_d^{(1)}(-n) & \cdots & N_d^{(1)}(n-1) & N_d^{(1)}(n) \\ 0 & \vdots & & \vdots & \vdots \\ 0 & N_d^{(n)}(-n) & \cdots & N_d^{(n)}(n-1) & N_d^{(n)}(n) \end{array} \right] \in \mathbb{R}^{(d+1) \times (d+1)}. \quad (17)$$

In a similar fashion we can cut the matrices on the left-hand side considering only the rows of \mathbf{N} with indices from $n + 1$ to $n + (n + 1)(m - 1)$ and the columns of \mathbf{B} with indices from $-n$ to $n + 1$. The only non-zero entries of \mathbf{N} contained in those rows are the ones with column index between $1 - n$ and $m + n - 1$, extrema included. For \mathbf{B} the only non-zero entries contained in the considered columns lie on the rows with indices from $-(m - 1)(n + 1) - mn$ to $(m - 1)(n + 1) + m(n + 1)$. Thus the useful indices for the cut are the ones between α and β where

$$\begin{aligned} \alpha &= \max_{m \geq 2, n \geq 0} \{ 1 - n, -(m - 1)(n + 1) - mn \} = 1 - n, \\ \beta &= \min_{m \geq 2, n \geq 0} \{ m + n - 1, (m - 1)(n + 1) + m(n + 1) \} = m + n - 1. \end{aligned}$$

From (17), it is clear that $\det(\mathbf{N}_R) \neq 0$. Thus \mathbf{N}_R^{-1} in (16) is well-defined as well as the matrices $\{\mathbf{A}_k\}_{k=1-m}^{m-1}$. Since all the matrices involved in (14) are properly slanted, (14) holds for every choice of the sequence $\mathbf{c}^{[j]}$.

Now we only have to prove that for every initial sequence $\mathbf{p}^{[0]}$, the scheme produces a \mathcal{C}^n polynomial spline of degree d and its derivatives. In view of (2) and (14), we observe that, after a subdivision step, each of the added new control vectors depends only on two consecutive control vectors from the previous step. Since the refinement rules have been chosen to reproduce degree- d cardinal B-splines and thus polynomials up to degree d , the first component of the vector-valued limit function will be a polynomial of degree d on every unitary interval with integer extrema. The fact that we are interpolating the first n derivatives yields \mathcal{C}^n regularity, so concluding the proof. \square

Remark 4.2. When $m = 3$ one has that $\mathbf{N}_L = \mathbf{N}_R$, for every $n \in \mathbb{N}_0$.

Since the entries of \mathbf{B} are given in (10) (see (15)) and the entries of \mathbf{N} can be easily obtained by recalling the properties of cardinal B-splines and their derivatives, Theorem 4.1 gives us an easy-to-implement method for constructing the refinement mask $\{\mathbf{A}_k \in \mathbb{R}^{(n+1) \times (n+1)}\}_{k=1-m}^{m-1}$ for any order $n \in \mathbb{N}_0$ and dilation factor $m \in \mathbb{N} \setminus \{1\}$. We conclude providing some illustrative examples.

Example 4.1 ($m = 2, n = 1, d = 3$). Figure 1 shows the result of its application to some initial data.

$$\{b_k\}_{k=-2}^2 = \{1, 4, 6, 4, 1\} / 8, \quad \mathbf{N}_L = \left[\begin{array}{c|c|c} 1/6 & 2/3 & 1/6 \\ \hline -1/2 & 0 & 1/2 \end{array} \right], \quad \mathbf{N}_R = \left[\begin{array}{c|c} \mathbf{N}_L & \begin{array}{c} 0 \\ 0 \end{array} \\ \hline 0 & \mathbf{N}_L \end{array} \right],$$

$$\mathbf{A}_0 = \begin{bmatrix} 1 & 0 \\ 0 & 1/2 \end{bmatrix}, \quad \overline{\mathbf{A}} = [\mathbf{A}_1 \quad \mathbf{A}_{-1}] = \left[\begin{array}{cc|cc} 1/2 & 1/8 & 1/2 & -1/8 \\ \hline -3/4 & -1/8 & 3/4 & -1/8 \end{array} \right].$$

Example 4.2 ($m = 2, n = 2, d = 5$).

$$\{b_k\}_{k=-3}^3 = \{1, 6, 15, 20, 15, 6, 1\} / 32,$$

$$\mathbf{N}_L = \left[\begin{array}{c|c|c|c|c} 1/120 & 13/60 & 11/20 & 13/60 & 1/120 \\ \hline -1/24 & -5/12 & 0 & 5/12 & 1/24 \\ \hline 1/6 & 1/3 & -1 & 1/3 & 1/6 \end{array} \right], \quad \mathbf{N}_R = \left[\begin{array}{c|c} \mathbf{N}_L & \begin{array}{c} 0 \\ 0 \\ 0 \end{array} \\ \hline \begin{array}{c} 0 \\ 0 \\ 0 \end{array} & \mathbf{N}_L \end{array} \right],$$

$$\mathbf{A}_0 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1/2 & 0 \\ 0 & 0 & 1/4 \end{bmatrix}, \quad \bar{\mathbf{A}} = [\mathbf{A}_1 \quad \mathbf{A}_{-1}] = \left[\begin{array}{ccc|ccc} 1/2 & 5/32 & 1/64 & 1/2 & -5/32 & 1/64 \\ \hline -15/16 & -7/32 & -1/64 & 15/16 & -7/32 & 1/64 \\ \hline 0 & -3/8 & -1/16 & 0 & 3/8 & -1/16 \end{array} \right].$$

Example 4.3 ($m = 3, n = 2, d = 5$).

$$\{b_k\}_{k=-6}^6 = \{1, 6, 21, 50, 90, 126, 141, 126, 90, 50, 21, 6, 1\} / 243,$$

$$\mathbf{N}_L = \mathbf{N}_R = \left[\begin{array}{c|c|c|c|c|c|c} 1/120 & 13/60 & 11/20 & 13/60 & 1/120 & 0 & 0 \\ \hline -1/24 & -5/12 & 0 & 5/12 & 1/24 & 0 & 0 \\ \hline 1/6 & 1/3 & -1 & 1/3 & 1/6 & 0 & 0 \\ \hline 0 & 1/120 & 13/60 & 11/20 & 13/60 & 1/120 & 0 \\ \hline 0 & -1/24 & -5/12 & 0 & 5/12 & 1/24 & 0 \\ \hline 0 & 1/6 & 1/3 & -1 & 1/3 & 1/6 & 0 \end{array} \right],$$

$$\mathbf{A}_0 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1/3 & 0 \\ 0 & 0 & 1/9 \end{bmatrix}, \quad \bar{\mathbf{A}} = [\mathbf{A}_1 \quad \mathbf{A}_{-2}] = \left[\begin{array}{cc|ccc} 64/81 & 16/81 & 4/243 & 17/81 & -2/27 & 2/243 \\ \hline -40/81 & 0 & 2/243 & 40/81 & -13/81 & 4/243 \\ \hline -40/81 & -32/81 & -10/243 & 40/81 & -8/81 & 1/243 \\ \hline 17/81 & 2/27 & 2/243 & 64/81 & -16/81 & 4/243 \\ \hline -40/81 & -13/81 & -4/243 & 40/81 & 0 & -2/243 \\ \hline 40/81 & 8/81 & 1/243 & -40/81 & 32/81 & -10/243 \end{array} \right].$$

Acknowledgements. This research has been accomplished within RITA (Research Italian network on Approximation). The authors are members of the INdAM Research group GNCS which has partially supported this work.

References

- [1] C. Conti, M. Cotronei, T. Sauer, *Factorization of Hermite subdivision operators preserving exponentials and polynomials*, Adv. Comput. Math. 42 (2016), 1055–1079.
- [2] M. Cotronei, C. Moosmüller, T. Sauer, N. Sissouno, *Level-dependent interpolatory Hermite subdivision schemes and wavelets*, Constr. Approx. 50 (2019), no. 2, 341–366.
- [3] N. Dyn, D. Levin, *Analysis of Hermite-interpolatory subdivision schemes*, Spline functions and the theory of wavelets (Montreal, PQ, 1996), 105–113, CRM Proc. Lecture Notes, 18, Amer. Math. Soc., Providence, RI, 1999.
- [4] N. Dyn, D. Levin, *Subdivision schemes in geometric modelling*, Acta Numer. 11 (2002), 73–144.
- [5] J. Fageot, S. Aziznejad, M. Unser, V. Uhlmann, *Support and approximation properties of Hermite splines*, J. Comput. Appl. Math. 368 (2020), 112503, 15 pp.
- [6] B. Jeong, J. Yoon, *Construction of Hermite subdivision schemes reproducing polynomials*, J. Math. Anal. Appl. 451 (2017), no. 1, 565–582.
- [7] B. Jüttler, U. Schwanecke, *Analysis and design of Hermite subdivision schemes*, Visual Comp 18, 326–342 (2002).
- [8] J.-L. Merrien, *A family of Hermite interpolants by bisection algorithms*, Numer. Algorithms 2 (1992), no. 2, 187–200.
- [9] G. Plonka, *Two-scale symbol and autocorrelation symbol for B-splines with multiple knots*, Adv. Comput. Math. 3 (1995), no. 1-2, 1–22.
- [10] D. Ranirina, J. de Villiers, *On Hermite vector splines and multi-wavelets*, J. Comput. Appl. Math. 349 (2019), 366–378.
- [11] M. Sabin, *Subdivision of Box-splines*. In Armin Iske, Ewald Quak and Michael S. Floater (eds.), *Tutorials on Multiresolution in Geometric Modelling*, Mathematics and Visualization, pages 3–23. Springer, 2002.