

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Characterisation of microsatellite loci in Sardinian pears (*Pyrus communis* L. and *P. spinosa* Forssk.)

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Sau S., Pastore C., D'hallewin G., Dondini L., Bacchetta G. (2020). Characterisation of microsatellite loci in Sardinian pears (*Pyrus communis* L. and *P. spinosa* Forssk.). *SCIENTIA HORTICULTURAE*, 270(25 August 2020), 1-9 [10.1016/j.scienta.2020.109443].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/761020> since: 2020-08-20

*Published:*

DOI: <http://doi.org/10.1016/j.scienta.2020.109443>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

# 1 **Characterisation of Microsatellite Loci in Sardinian Pears (*Pyrus*** 2 ***communis* L. and *P. spinosa* Forssk.).**

3

4 Silvia Sau <sup>a,b\*</sup>, Chiara Pastore<sup>c</sup>, Guy D'hallewin<sup>d</sup>, Luca Dondini<sup>c</sup>, Gianluigi Bacchetta<sup>a,b</sup>

5

6 <sup>a</sup> Banca del Germoplasma della Sardegna (BG-SAR), Hortus Botanicus Karalitanus (HBK), Università degli Studi di  
7 Cagliari, Viale Sant'Ignazio da Laconi, 9-11, 09123 Cagliari, Italy.

8 <sup>b</sup> Centro Conservazione Biodiversità (CCB), Dipartimento di Scienze della Vita e dell'Ambiente (DISVA), Università  
9 degli Studi di Cagliari, Viale Sant'Ignazio da Laconi, 13, 09123 Cagliari, Italy.

10 <sup>c</sup> Dipartimento di Scienze e Tecnologie Agro-Alimentari (DISTAL), Università di Bologna, Viale Fanin, 44, 40127,  
11 Bologna, Italy.

12 <sup>d</sup> Istituto di Scienze delle Produzioni Alimentari (ISPA), Consiglio Nazionale della Ricerche (CNR), ~~Sassari~~. Traversa  
13 la Crucca, 3 Loc. Baldinca – 07040 Sassari, Italy.

14

15 \*Corresponding author Silvia Sau email: sausilvia85@gmail.com

16

## 17 **Abstract**

18 The Sardinian pear germplasm is an important resource of genetic diversity that can be used for increasing data on  
19 European pear germplasm, to optimise the efficiency of the association studies within the genome and to identify genomic  
20 regions that control the main horticultural traits. A set of 109 Sardinian pear genotypes, composed of 82 *Pyrus communis*  
21 cultivars, 24 wild populations of *P. spinosa* and three international reference cultivars, was analysed using nine SSR  
22 markers to assess the genetic diversity of Sardinian pears, determine their genetic structure and study the cases of  
23 synonymies and homonymies. The comparison of SSR profiles indicated four groups of diploid accessions with the same  
24 SSR profile. The alignment with the Sardinian dataset, pointed out a clear genotype distinctiveness. For all studied SSR  
25 loci, 15 specific rare alleles were identified, with a minimum of two alleles found in the database of analysed accessions  
26 for the SSRs EMPc11 and EMPc117. The overall allelic diversity revealed a high polymorphism in the analysed Sardinian  
27 germplasm. The structure analysis allowed us to identify four gene pool groups (Sardinian cultivars, Japanese cultivars,  
28 late-ripening cultivars and, the most famous, standard cultivars). These results were confirmed by Evanno's  $\Delta k$  statistical

analysis which has shown unequivocally that  $k = 4$  ( $\Delta k = 150$ ) is the most likely stratification level of the cluster. The  $Q$  values of *P. communis* and *P. spinosa* accessions have confirmed an allelic interchange between wild and cultivated genotypes.

**Keywords:** Sardinia; pear; molecular marker; genetic structure.

## 1. Introduction

The husbandry of pear trees in Sardinia has a long history, which involves domestication processes of autochthonous species (*Pyrus communis* L. and *P. spinosa* Forssk.). Indeed, literature dates back to Aristotle (384-322 BC), who exalted the island of Ichnusa (actual Sardinia) for its fruits, and to a description of pear cultivars by Pliny (AD 23-79). Then, particularities of pear cultivars were reported in the scripts of Pausania, Polibio and Diodoro (Agabbio, 1994). In Sardinia, over a hundred pear accessions with different morphological characteristics of local origin have been recently examined, highlighting the literature on the rich heritage of ancient cultivars (Agabbio et al., 1986, 2015). Still, lifestyle changes and the, agronomic and productive characteristics of new breeds foster genetic erosion of these ancient cultivars in that they do not ensure comparable income with other economic activities (Muresu et al., 1997). However, in the present scenario, rather than the sole economic aspect, evidence is provided evermore on the key value of genetic diversity. Indeed, ancient local cultivars own a gene pool gathered and preserved over years under a broad range of different environmental conditions (Hammer et al., 2003). Since current breeding programs have not yet exploited this high variability for many traits, as already in progress for apple and other species, shedding light on Sardinian *Pyrus* genotype distinctiveness is pivotal (Liang et al., 2015).

Guided breeding programs start from an efficient molecular characterisation of known cultivars aimed at managing the genetic heritage by identifying clonal relationships, synonyms and homonyms. In fact, the assessment of genetic and phenotypic diversity is the starting point of screening germplasm collections for valuable characteristics useful in breeding programs. Following the marker era take-off, an increasing number of papers described the relationships among pear cultivars by using molecular markers: RFLPs (Iketani et al., 1998), RAPDs (Oliveira et al., 1999), and AFLPs (Monte-Corvo et al., 2000; Dolatowski et al., 2004; Bao et al., 2008). In eukaryotic genomes, microsatellites or simple sequence repetitions (SSR) are ubiquitous and easily employed as markers for both animal and plant species. Therefore, these approaches are highly reproducible and are of multi-allelic nature, codominant inheritance, and relative abundance and provide good genome coverage (Weber and Mat, 1989; Powell et al., 1996).

In recent years, main investigations on SSR primers of the genus *Pyrus* L. have focused on the varietal heritage of the Japanese *P. pyrifolia* Nak., and several studies were also conducted on the European *P. communis* (Yamamoto et al., 2002

a, b; Fernández-Fernández et al., 2006; Bao et al., 2007). Evans et al. (2007) standardised some SSRs markers useful to compare *P. communis* accessions within European germplasm collections. Genetic diversity analysis in pear by using SSR markers is widely reported in literature and then successfully used in Italy (Martinelli et al., 2009; Ferradini et al., 2017; Bennici et al., 2018, Baccichet et al., 2020), Spain (Miranda et al., 2010; Ferreira dos Santos et al., 2011; Urrestarazu et al., 2015), Germany and Romania (Puskas et al 2015), Portugal (Queiroz et al., 2015, 2019), Bosnia (Gasi et al., 2013), and Slovenia (Sisko et al., 2009). Additional noteworthy characterisations of pear genetic variability were performed in Europe (Wünsch and Hormaza, 2007; Sehic et al., 2012), in Africa (Tunisia: Brini et al., 2008; Ouni et al 2020) and in Asia (Yue et al., 2018), including some specific Asian regions (Ahmed et al., 2010; Rana et al., 2015).

In Italy, recent characterisation of the Mount Etna pear germplasm aimed to introduce specific traits into breeding programmes by exploiting genetic variability related to drought resistance, low chilly unit requirement, adaptation to daily temperature fluctuation and low pest and disease incidence (Bennici et al., 2018).

At present, a set of useful SSR markers for pear genetic screening has been approved by the European Cooperative Program for Plant Genetic Resources (ECPGR), and following the successful use of screening pear accessions of the Brogdale National Fruit Collection (NFC) (Fernández-Fernández, 2010), a complete dataset is now available at the Institute website (<http://www.emr.ac.uk/SPFeliFernandez.htm>). The availability of datasets describing the allelic pattern of hundreds of accessions combined with the use of common sets of markers, was employed to define gene flows among local germplasms of *Malus domestica* Borkh. (Urrestarazu et al., 2016). The same approach provided evidence of allelic interchange between wild populations and cultivated pear germplasm of the Mount Etna (Bennici et al., 2018). This result supports previous findings on Iranian *P. communis* germplasm genotypes, where gene transfer was highlighted by the presence of new S-alleles possibly introduced from other domesticated or wild pear species as corroborated by plastidial DNA analysis of the same genotypes (Gharehaghaji et al., 2014a). According to Gharehaghaji et al. (2014b), these results and the cross-compatibility of wild and domesticated *Pyrus* taxa might have left a genetic trace on the traditional Iranian pear germplasm. These interesting findings along with the great number of Sardinian pear genotypes, both domesticated and wild ones, are the catalyst of the present work. This research aims to, first estimate the genetic relationship between wild and local cultivated germplasm pear accessions and, second, extend this investigation to nationally and internationally known cultivars. The outcomes will shed light on potential flows among the analysed gene pools. To reach this goal, the constructed allele dataset of Sardinian germplasm pear accessions will be aligned with the one reported by Sehic et al. (2012). This approach aims to increase *Pyrus* molecular marker datasets and sheds light on other homonymous and/or synonymous genotypes difficult to distinguish using standard morphological descriptors.

## 2. Materials and Methods

### 2.1. Germplasm material, DNA extraction and SSR genotyping

The list of 108 analysed accessions is provided in Table 1. This inventory includes 76 *P. communis* cultivars from the Sardinian germplasm conserved in the CNR-ISPA catalogue field (39°56'04.2" N and 8°35'46" E, 7 m a.s.l.), 5 from the Fo.Re.S.T.A.S catalogue field (40°25'55" N and 9°01'43" E, 540 m a.s.l.), 24 representative accessions of *P. spinosa* collected all over Sardinia and three international reference cultivars ('Williams', 'Abate' and 'Keiser'). *P. communis* cultivars reported in this list and used throughout the work are those given by local communities and adopted in the catalogue fields and, for *P. spinosa*, the geographical area where the mother plants are localised.

For each of them, genomic DNA was extracted from 50 mg of young freeze-dried leaves following the standard CTAB protocol (Maguire et al., 1994). Genomic DNA was quantified by Nanodrop™ ND-1000 Spectrophotometer (Thermo Scientific, Wilmington, DE, USA) and diluted to 10 ng/μL.

A set of nine SSRs primers (CH01d09, CH02b10, CH01f07a, CH03d12, CH04e03, CH05c06, EMPc11, EMPc117 and GD147), chosen among the ECPGR approved ones, was used for pear genotyping (Evans et al., 2007; Sehic et al., 2012). Primer sequences are also available on the HiDRAS (High-quality Disease Resistant Apples for Sustainable Agriculture) website (<http://users.unimi.it/hidras/>). Forward SSR primers were labelled with FAM or HEX fluorescent dyes. The nine SSR markers used for the analyses were selected according to the work carried out by Sehic et al. (2012), in which ten apple markers for pears were used. The high transferability of SSRs among related species, such as apple and pear, was abundantly demonstrated (since Pierantoni et al., 2004).

The amplification of markers was performed using a PCR protocol for apple as described by Liang et al. (2015). In short, PCR was performed in a 10 μL reaction mixture containing 1× reaction buffer (Applied Biosystems, Foster City, CA, USA), 0.6 mM MgCl<sub>2</sub>, 0.4 mM dNTPs (Fermentas, Lithuania), 1 nM of each SSR locus-specific primer, 0.125 μL Taq (5u/μL, Fisher Molecular Biology, Rome, Italy) and 10 ng genomic DNA.

PCR reactions were carried out in a 2720 thermal cycler (Applied Biosystems) with the following amplification protocol: initial denaturation step at 94°C for three min, followed by 32 cycles at 94°C for 30 sec, at 58°C for one min and at 72°C for one min and a final extension step at 72°C for 10 min.

Multi-pooling groups (MPG) of SSRs labelled with the two different fluorescent dyes were designed for SSR genotyping on an ABI PRISM 3730 DNA analyser. SSRs were pooled by mixing PCR products labelled with different dyes in a ratio of 1:1 for HEX:FAM; 3μl of the PCR products mixture was added to 7 μl of formamide containing 0.2 μL of GeneScan 500 LIZ size standard (Applied Biosystems). Fragments were visually analysed and scored using Peak Scanner v.1.0 (Applied Biosystems). To monitor the reproducibility in different amplifications, three reference cultivars ('Abate', 'Kaiser' and 'William', were included in each single run.

121

## 122 2.2. SSR polymorphism, cluster and structure analysis

123 SSR profiles were used to obtain a large dataset for the 109 pear genotypes used in this study. The collected data were  
124 organised in a square matrix in which code '0' and '1' were used for allele absence and presence, respectively (code for missing  
125 data was 9).

126 The genetic distance between cultivar and wild species was then calculated through the DICE coefficient (Dice, 1945) using  
127 the Similarity for Qualitative data (SimQual) procedure of NTSyS-p.c. version 2.0 (Rohlf, 1998). The dendrogram was  
128 constructed using the unweighted pair group method of arithmetic average (UPGMA) clustering and drawn with the NTSyS-p.c.  
129 version 2.0 program (Rohlf, 1998).

130 Considering that pear accessions can be polyploid, the software SPAGeDi v.1.5 (Hardy and Vekemans, 2002) was used to  
131 compute genetic information statistics, as this software supports analyses of datasets containing individuals with different ploidy  
132 levels. Genetic statistics was performed on unique genotypes, excluding reference cultivars and duplicates and included number  
133 of alleles per locus, effective number of allele, number of rare alleles per locus (alleles whose frequency in complex resulted  
134  $<0.05$ ), number of genotype-specific alleles (unique alleles, present in only one accession), observed heterozygosity ( $H_o$ ),  
135 expected heterozygosity ( $H_e = 1 - \sum p_i^2$ , where  $p_i$  is the frequency of the  $i^{th}$  allele) and Wright's fixation index ( $F_i$ ). The probability  
136 of identity (PI) was computed as defined by Paetkau et al. (1995),  $PI = 2(\sum p_i^2)^2 - \sum p_i^4$ , where  $p_i$  is the frequency of the  $i^{th}$  allele.

137 Then triploid accessions and duplicated genotypes were removed before performing the structure analysis using  
138 STRUCTURE version 2.3.4 (Pritchard et al., 2000). For this analysis, another dataset was created, which included 454 unique  
139 pear cultivars analysed by Fernández-Fernández (2010), as detailed in the publication by Ordidge et al., 2018 and 23 unique,  
140 diploid Sardinian pear cultivars. To exclude further the duplicated genotypes from the analysis this dataset was previously  
141 validated by repeating the NTSyS-p.c. cluster analysis. To investigate the structure using the genotype data of our dataset, a  
142 Bayesian clustering method (Pritchard et al., 2000a; Falush et al., 2003) was applied using the STRUCTURE software. Previous  
143 information was not used to define clusters. Independent runs were done by setting the number of clusters (k) from 2 to 10. Each  
144 run included a burn-in period length set at 10,000 followed by 100,000 Markov Chain Monte Carlo (MCMC) number of repeats  
145 under the admixture model. The most likely number of sub-clusters (k) was identified with STRUCTURE HARVESTER (Earl  
146 and von Holdt, 2012) using the  $\Delta k$  described by Evanno et al. (2005). Genotypes were assigned to the group for which they  
147 showed the highest membership coefficient, considering an accession strongly assigned to each partitioning level if its proportion  
148 of ancestry (Q) was  $\geq 0.80$ , otherwise they were considered as "admixed", according to Urrestarazu et al. (2012). The same  
149 Structure parameters have been used for the Nested approach.

150

### 3. Results

As a result, the nine SSR markers employed showed a clear and easily readable peak amplification of the Sardinian pears dataset composed of 82 local *P. communis* cultivars, 24 wild populations of *P. spinosa* and three international reference cultivars (Table 1). Following the statistical analysis of the nine polymorphic loci, 101 unique genotypes were detected within our dataset and 162 total alleles were found. The primer CH04e03 amplified a minimum of 12 different alleles while a maximum of 30 alleles was found for the CH01f07a primer, as reported in Table 2. Furthermore, the average number of alleles per locus found in the present research (Table 2) was notably different with respect to other large-scale studies (9.4) on pear genetic diversity (Fernández-Fernández et al., 2006). The effective number of alleles varied between 1.66 (CH04e03) and 15.47 (CH01f07a), with an average of 8.45. The allele frequencies of the examined loci ranged between 0.002 and 0.775 (Supplementary Table 1) and for all studied SSR loci, it was possible to detect 26 specific rare alleles, nine of which were found in the whole panel of analysed accessions for the SSRs EMPc11 and CH01f07a (Table 2). The observed heterozygosity ( $H_o$ ) ranged from 0.667 (CH04e03) to 1.00 (CH01d09 and EMPC11), with an average of 0.83 across loci, while the expected heterozygosity ( $H_e$ ) ranged from 0.396 (CH04e03) to 0.935 (CH01f07a). Higher values of  $H_o$  than  $H_e$  observed in eight loci out of nine resulted in negative Wright's fixation index ( $F_i$ ) values, indicating a slight excess of heterozygosity across these eight loci. For the remaining one locus (CH02b10), the  $F$  value was positive, meaning a deficit of heterozygosity (Table 2). However, the average  $F_i$  ( $-0.101$ ) was close to zero, suggesting that Sardinian pear germplasm was comparable to an almost random mating unit. The highest probability of identity (PI) value (0.374) was at CH04e03 locus and the lowest (0.009) at CH01f07a. The cumulative PI was  $4.04 \times 10^{-14}$ , indicating the possibility that two randomly chosen individuals have the same SSR profile is not realistic.

Among accessions, 78 polyploids were identified (Table 3): 45 showed three alleles in more than one locus, of which 20 displayed three alleles in two loci, eight displayed three alleles in three loci, eight displayed three alleles in four loci, eight displayed three alleles in five loci and two displayed three alleles in six loci. The remaining 33 accessions presented only one locus with three alleles (Table 3).

The comparison of SSR profiles indicated four groups of diploid accessions with the same profile (Figure 1) allowing to identify cases of possible synonymy such as 'Enosa' with 'E' Donna', as well as cases of misnomer such as 'Cozzon'e AINU' with 'Bragamotta', 'Alveghina' with 'Bau' and 'Reale' with 'William' (Fig. 1). Furthermore, the accessions 'Vacchesa' and 'Pira Ortine', 'Laceni 2' and 'Camusina Precoce', 'Bianca' and 'Mamoi' and 'Armungesa' and 'Mulargia' showed very similar profile and have been discriminated for very small allelic differences at different loci (Supplementary Table 2). In addition, the data set analysis showed that the *P. communis* group is phylogenetically separated from the *P.*

180 *spinosa* one, although some *P. communis* accessions, (e.g. 'Pira Cona Arrubia' and 'Pira di Urzulè'), revealed a closer  
181 genetic distance to *P. spinosa* than to *P. communis* (Fig. 1).

182 To better identify the presence of sub-clusters in this large dataset a structure analysis was performed (Fig. 3). Evanno's  
183  $\Delta k$  statistical analysis has shown unequivocally  $k = 4$  ( $\Delta k = 150$ ) as the most likely stratification level of the clusters (Fig.  
184 2), which corresponded to the Sardinian pear gene pool (G1; yellow), the Japanese one (G2; blue), a group of late ripening  
185 cultivars (G3; green) and a group with the most famous standard cultivars (G4; red, Supplementary Table 3).

186 As far as *He*, the highest values were reached in G1 (0.834), followed by G3 having a value of 0.7069 while, G2 and  
187 G4 had similar values (0.6305 and 0.6308, respectively).

188 On behalf of the colour bar-plot it is possible to clearly evidence the four distinct gene groups (Fig. 3, Supplementary  
189 Table 3), supporting the  $k = 4$  hypothesis. The fixation index (*Fst*) which, ranging from 0 to 1, could indicate no genetic  
190 divergence within the populations or complete extreme division respectively, was calculated for each gene group, and the  
191 values were very low ranging from 0.0015 (G1) to 0.3981 (G2) with intermediate values for G3 (0.1232) and G4 (0.2322).

192 In more detail, 105 accessions from G1 (including all Sardinian ones), 12 belonging to G2, 120 within G3 and 98 from  
193 G4 presented *Q* values higher than the 0.8 threshold value. The remaining 103 accessions had values lower than 0.8;  
194 therefore, they were considered as admixed (Supplementary Table 3). Within this large number of admixed accessions, 29  
195 fit to G1, one to G2 while 43 and 30 were included in G3 and G4, respectively (Supplementary Table 3).

196 A restricted analysis of the 23 accessions belonging to Sardinian collections showed that the average of their *Q* values  
197 was 0.95 for G1, 0.01 for G2 (mainly composed of Japanese cultivars of *P. pyrifolia*) and 0.02 for both G3 and G4 (mainly  
198 composed of late cultivars of the centre of Europe and the classical pear cultivars used for breeding, respectively; Fig. 4).

199 In order to identify the presence of a possible gene flow between *P. spinosa* and *P. communis* a further structure analysis  
200 (nested approach) was conducted both on the cluster G1 dataset and on the panel of Sardinian accessions. The results of  
201 this analysis revealed that the two species shared the same gene pool and did not separate in different clusters (data not  
202 shown).

203

#### 204 4. Discussion

205 Sardinia possesses a great plant diversity with numerous local cultivars and wild pear that are likely differentiated  
206 according to soil and climatic variable conditions as well as by anthropic selection (Agabbio et al., 1986, 2015). Indeed,  
207 local cultivars own a high variability for several agronomic traits such as fruit size, ripening time, soil adaptability and  
208 resilience to environmental conditions, characterised by high temperatures and scarce water availability during the summer  
209 period (from June to September). Modern breeding programs have never exploited this genetic diversity. In this work,  
210 relationships among the local, old, safeguarded cultivars and wild related species were inferred by SSR marker analysis.



211 Reproducibility and coherence in amplified peak readings in the references ‘William’, ‘Kaiser’ and ‘Abate’ have  
212 allowed us to make sure that the SSRs used in this work are valuable markers to define pear germplasm diversity.

213 The identification of one synonymy (‘Enosa’ with ‘È Donna’) in the Sardinian germplasm emphasised the importance  
214 of linking referee collections with powerful tools such as molecular markers aimed at avoiding repetitions and promoted  
215 the withdrawal of duplicates.

216 The use of reliable SSRs is essential for the effective differentiation of cultivars and the establishment of genetic  
217 relationships such as homonymies and synonymies. Three putative misnomers are between ‘Reale’ and ‘William’,  
218 ‘Alveghina’ and ‘Bau’ and ‘Bragamotta’ and ‘Cozzonn’e AINU’ that are known to be different cultivars from their  
219 descriptors.

220 A high similarity was found between the Sardinian cultivars ‘San Domenico’ and ‘Beurré Giffard’ an old French  
221 summer pear that is usually picked at the end of July or at the beginning of August. ‘San Domenico’ is also an early-  
222 ripening pear starting from June up to mid-July, with a buttery melting flesh similar to ‘Beurré Hardy’.

223 The overall diversity analysis proved the allelic interchange between accessions of *P. communis* and *P. spinosa* or vice  
224 versa. The alignment of the two datasets confirmed the phylogenetical rift of *P. spinosa*, and on behalf of the allelic  
225 frequency analysis, highlighted the magnitude of shared alleles. After dataset analysis, ‘De Puleu’, ‘Buttidu de Austu’ and  
226 ‘Mela’ resulted in four alleles in a single locus. In the present work, the number of alleles in each locus and the range of  
227 achieved values diverge from the data available in the literature (Table 2). The overall allelic diversity evidenced by  
228 employing nine SSRs revealed a high polymorphism in the analysed Sardinian germplasm.

229 The phylogenetic proximity between the accessions ‘Pira di Urzulè’ and ‘Pira Cona Arrubia’ with the wild populations  
230 of *P. spinosa* can be explained by the fact that these cultivars, according also to morphological and chemical similarities,  
231 may be due to of hybridisation between *Pyrus* species (*P. spinosa* and *P. communis*) as reported in other geographical  
232 regions rich in the genus *Pyrus*. A botanical hybrid between *P. communis* and *P. spinosa* = *P. × jordanovii* has been  
233 discovered in Pirin Mountains of Bulgaria (Dostálek, 1984) and other hybridisations between these two species have been  
234 already described (Vincent et al., 2013). This phenomenon was also recently evidenced in the pear germplasm of the Mount  
235 Etna in which an allelic interchange was found between *P. communis*, *P. pyraeaster* and *P. spinosa* (Bennici et al., 2018).

236 The high number of putative triploids in Sardinian cultivars can be explained by the fact that anthropic selection pressure  
237 has sorted out the triploid accessions because of the generally larger size of the fruit. It is known that, generally, the triploid  
238 apple and pear accessions bear larger fruits (Ashton and Spigel-Roy, 1985). This phenomenon could explain the relatively  
239 high percentage of triploids among the pear old varieties and landraces. A percentage of 15% of triploids was recently  
240 described in the pear local germplasm of Friuli Venezia Giulia, a region in the North-eastern of Italy (Baccichet et al., 2020)  
241 while the very high percentage of 24.2% was described in a panel of German and Romanian accessions (Puskas et al.,

242 2015). Analogously to pear, a high number of triploid accessions (about 10% on more than 2000 analysed samples) was  
243 also found to be present in apple landraces and local germplasms at European level (Urrestarazu et al., 2016). A detailed  
244 study in *Pyrus* species revealed that all species are diploid, with  $2n = 34$ , although different cultivars and variants are triploid  
245 (Zielinski et al. 1967), so, in several cases, the marker might have amplified another locus of the homologous genome  
246 (Pierantoni et al. 2004).

247 Thanks to the existence of a large dataset created by Fernández-Fernández et al. (2006), which covers *Pyrus*'s  
248 international and local cultivars, comparing these cultivars with our dataset has been possible, thus increasing the  
249 availability of useful data for future analysis. In fact, the use of the same markers in both analyses made it possible to align  
250 the two datasets.

251 The SSR alleles used in the structure analysis (Fig 2) allowed us to identify four groups. The Sardinian accessions  
252 belong to G1, together with a large group of the Italian cultivars that have been analysed in the Fernández-Fernández dataset  
253 (Fernández-Fernández et al., 2006, Sehic et al., 2010) and French early ripening cultivars. These results agree with the  
254 observations of Bennici et al. (2018) that showed a notable contribution to the genetic makeup of many Italian cultivars,  
255 especially ancient ones. The relationship found among pear genotypes from Sardinia, Italy and France may have historical  
256 origin by vegetative propagation of plant material exchanged between Italy and France mainly during the Sardinian  
257 Piedmont reign (1720-1861; Agabbio et al., 2015).

258 The Q values of *P. communis* and *P. spinosa* accessions are very close as confirmation of an allelic interchange between  
259 the wild and the cultivated genotypes, to support the idea that no significant gene flow exists between the Sardinian gene  
260 pool G1 and the ones of G2, G3 and G4 (Fig. 4).

261 The very low *Fst*, observed in all four groups indicates that genetic variability is not massively linked to the structure  
262 of the population analysed. In fact, although a difference exists between the statistically valid groups, we are in a fairly  
263 homogeneous gene pool.

264

## 265 5. Conclusion

266 This document describes, for the first time, the genetic diversity of domesticated and wild Sardinian pears by considering  
267 a sample of over 100 accessions, to characterise the cultivars' genetic structure and studying the influence of wild species  
268 on specific local germplasm.. The development of a large dataset allowed us to use reliable classification criteria and  
269 estimate genetic distances among cultivars. The availability of a large pear dataset in literature allows the identification or  
270 confirmation of homonymies and synonymies that could occur in different germplasm accessions. This dataset provides an  
271 overview of the history of diversification of the pear germplasm in Sardinia, diversification that was steered by man and

272 influenced by the high environmental and climatic variability present on the island. A clear allelic interchange between *P.*  
273 *communis* and *P. spinosa* accessions was detected. This Sardinian germplasm represents a very important source of genetic  
274 diversity that was never investigated at the molecular level and that could be exploited in new breeding programs aimed at  
275 improving the resilience of the new pear cultivars in a new context characterised by deep climatic changes.

276

## 277 **Acknowledgements**

278 The authors thank Mariano Ucchesu for his help during the collection of the germplasm and Giulia Pagliarani for her  
279 support during laboratory work. Thanks to Massimiliano Orsini for his suggestion and the revision of the manuscript. This  
280 research was conducted and funded by a doctoral grant of Silvia Sau from the University of Cagliari Ministero  
281 dell'Università e della Ricerca (MIUR). The authors also thanks Proof-reading-service for the English revision  
282 professionally proofread.

283

## 284 **Figure and table captions**

285 **Table 1.** In uppercase cultivars of *P. communis* (*P.c*) and (*P.c\**) and in lowercase accessions of *P. spinosa* (*P.s*) analysed  
286 in this study. In bold, reference cultivars.

287 **Table 2.** Characteristics of the nine SSRs studied in Sardinian *P. communis* and *P. spinosa* accessions.

288 **Table 3.** Triploids and tetraploids set found in the Sardinian *P. communis* cultivars (in uppercase) and in *P. spinosa* wild  
289 populations (in lowercase).

290 **Fig. 1.** Dendrogram of 109 pear cultivars from the Sardinian germplasm based on DICE coefficient calculated from the  
291 allele frequencies of 194 alleles found in nine SSR loci. The *P. communis* is in uppercase letters, while the *P. spinosa*  
292 wild population is in lowercase. In the black circle the misnomer found in the Sardinian *Pyrus* accessions. In the black  
293 rectangles are the cultivars that are differentiated for less than 10 alleles. In bold are the reference cultivars.

294 **Fig. 2.** Slope rate change estimates of the log probability curve ( $\Delta K$ ) calculated according to Evanno  
295 et al. (2005) plotted against number of cluster (*K*).

296 **Fig. 3.** Clusters identified in the first round structure analysis: Sardinian (yellow), Japanese (blue); late cultivars (green);  
297 the most famous standard cultivars (red).

298 **Fig. 4.** Genetic composition of the Sardinian gene pool for K=4 groups inferred with Structure and described by the mean  
299 Q value distribution. Sardinian (G1); Japanese (G2); late ripening (G3); the most famous standard cultivars (G4).

300 Supplementary Material

301 **Supplementary Table 1:** Allele frequency of nine SSRs in Sardinian *Pyrus* germplasm. Rare alleles are indicated in bold  
302 and unique alleles are evidenced by underling.

303 **Supplementary Table 2:** Sardinian pear cultivars with similar genotypes. In bold, allele that differentiates the genotype.

304 **Supplementary Table 3:** Classification of 454 genotypes including Sardinian *Pyrus* germplasm and pear cultivars  
305 analysed by Fernández-Fernández et al. (2006) by structure using 9 SSR loci in K = 4 reconstructed populations.

306

307

## 308 **References**

309 Agabbio, M., 1994. Le vecchie varietà della Sardegna: patrimonio genetico di specie arboree da frutto. Carlo Delfino  
310 Editore, Sassari.

311 Agabbio, M., Ansaldi, N., Beccaro, G.L., Berra, L., Botta, R., Brunu, A., 2015. Frutti dimenticati e biodiversità  
312 recuperata. Il germoplasma frutticolo e viticolo delle agricolture tradizionali italiane. Casi di studio: Piemonte e Sardegna.  
313 Quaderni Natura e Biodiversità. 7, 123-133.

314 Agabbio, M.C.S., Suelzu, R., Mulas, M., Mannoni, G., 1986. Patrimonio genetico delle pomacee in Sardegna: il pero:  
315 cenni storici e cultivar individuate. Studi sassaresi: organo ufficiale della Società sassarese di Scienze mediche e naturali.  
316 Sez. 3: Annali della Facoltà di Agraria dell'Università di Sassari. 32, 120–139.

317 Ahmed, M., Anjum, M.A., Khan, M.Q., Ahmed, M.J., Pearce, S., 2010. Evaluation of genetic diversity in *Pyrus*  
318 germplasm native to Azad Jammu and Kashmir Northern Pakistan revealed by microsatellite markers. Afr. J. Biotechnol.  
319 9, 8323–8333.

320 Alston, F.H., Spiegel-Roy, P., 1985. Fruit tree breeding: strategies, achievements and constraints. In: Attributes of Trees  
321 as Crop Plants. M.G.R. Cannell and J.E. Jackson (eds). Institute of Terrestrial Ecology, Monks Wood Experimental Station,  
322 Huntingdon UK, pp. 49–67.

323 Baccichet, I., Foria, S., Messina, R., Peccol, E., Losa, A., Fabro, M., Gori, G., Zandigiacomo, P., Cipriani, G., Testolin,  
 324 R., 2020. Genetic and ploidy diversity of pear (*Pyrus* spp.) germplasm of Friuli Venezia Giulia, Italy. *Genet Resour Crop*  
 325 *Evol.* 67, 83–96.

326 Bao, L., Chen, K., Zhang, D., Cao, Y., Yamamoto, T., Teng, Y., 2007. Genetic diversity and similarity of pear *Pyrus*  
 327 *L.* cultivars native to East Asia revealed by SSR simple sequence repeat markers. *Genet. Resour. Crop. Evol.* 54, 959–971.

328 Bao, L., Chen, K., Zhang, D., Li, X., Teng, Y., 2008. An assessment of genetic variability and relationships within  
 329 Asian pears based on AFLP Amplified Fragment Length Polymorphism markers. *Sci. Hortic.* 116, 374–380.

330 Bennici, S., Las Casas, G., Distefano, G., Di Guardo, M., Continella, A., Ferlito, F., Gentile, A., La Malfa, S., 2018.  
 331 Elucidating the contribution of wild related species on autochthonous pear germplasm: A case study from Mount Etna.  
 332 *PLoS ONE.* 13, 6.

333 Brini, W., Mars, M., Hormaza, J.I., 2008. Genetic diversity in local Tunisian pears *Pyrus communis* L. studied with  
 334 SSR markers. *Sci Hortic.* 115, 337–341.

335 Dice, L.R., 1945. Measures of the amount of ecologic association between species. *Ecology.* 26, 297–302.

336 Dostálek, J. 1984. Hybriden von *Pyrus spinosa*. *Folia Geobot. Phytotax.* 19, 89-93.

337 Earl, D.A., Von Holdt, B.M., 2012. Structure Harvester: a website and program for visualizing Structure output and  
 338 implementing the Evanno method. *Conserv. Genet. Resour.* 4, 359–361.

339 Evanno, G., Regnaut, S., Goudet, J., 2005. Detecting the number of clusters of individuals using the software  
 340 STRUCTURE: a simulation study. *Mol. Ecol.* 14, 2611–2620.

341 Evans, K.M., Fernández-Fernández, F., Govan, C., 2007. Harmonising fingerprinting protocols to allow comparisons  
 342 between germplasm collections-Pyrus. *Acta Hort.* 814, 103-106.

343 Falush, D., Stephens, M., Pritchard, J.K., 2003. Inference of population structure using multilocus genotype data: linked  
 344 loci and correlated allele frequencies. *Genetics.* 164, 1567–1587.

345 Fernández-Fernández, F., Harvey, N.G., James, C.M., 2006. Isolation and characterization of polymorphic  
 346 microsatellite markers from European pear *Pyrus communis* L. *Mol. Ecol. Resour.* 6, 1039–1041.

347 Ferradini, N., Lancioni, H., Torricelli, R., Russi, L., Dalla Ragione, I., Cardinali, I., Marconi, G., Gramaccia, M.,  
 348 Concezzi, L., Achilli, A., Veronesi, F., and Albertini, E., 2017. Characterization and phylogenetic analysis of ancient Italian  
 349 landraces of pear. *Front. Plant Sci.* 8, 751.

350 Ferreira dos Santos, A.R., Ramos–Cabrer, A.M., Díaz–Hernández, M.B., Pereira–Lorenzo, S., 2011. Genetic variability  
 351 and diversification process in local pear cultivars from northwestern Spain using microsatellites. *Tree Genet. Genomes*. 7,  
 352 1041–1056.

353 Gasi, F., Kurtovic, M., Kalamujic, B., Pojskic, N., Grahic, J., Kaiser, C., & Melan, M., 2013. Assessment of European  
 354 pear *Pyrus communis* L. genetic resources in Bosnia and Herzegovina using microsatellite markers. *Sci. Hortic*. 157, 74–  
 355 83.

356 Gharehaghaji, A.N., Arzani, K., Abdollahi, H., Shojaeiyan, A., Dondini, L., & De Franceschi, P., 2014a. Genomic  
 357 characterization of self–incompatibility ribonucleases in the Central Asian pear germplasm and introgression of new alleles  
 358 from other species of the genus *Pyrus*. *Tree Genet. Genomes*. 10, 411–428.

359 Gharehaghaji, A.N., Arzani, K., Abdollahi, H., Shojaeiyan, A., Henareh, M., De Franceschi, P., & Dondini, L., 2014b.  
 360 Chloroplast genome diversity of the *Pyrus* genus from Iranian and European wild pear species to the cultivated cultivars  
 361 *Acta Hortic* 1032, 151–158.

362 Hammer, K., Arrowsmith, N., Gladis, T., 2003. Agrobiodiversity with emphasis on plant genetic resources.  
 363 *Naturwissenschaften*. 90, 241–250.

364 Hardy O.J., Vekemans X. 2002. SPAGeDi: a versatile computer program to analyse spatial genetic structure at the  
 365 individual or population levels. *Mol. Ecol. Notes*. 2, 618–620.

366 Iketani, H., Manabe, T., Matsuta, N., Akihama, T., and Hayashi, T., 1998. Incongruence between RFLPs of chloroplast  
 367 DNA and morphological classification in East Asia pear *Pyrus* spp.. *Genet. Resour. Crop Evol*. 45, 533–539.

368 Kalinowski, S.T., Taper, M.L., Marshall, T.C., 2007. Revising how the computer program CERVUS accommodates  
 369 genotyping error increases success in paternity assignment. *Mol. Ecol*. 16, 1099–1106.

370 Liang, W., Dondini, L., De Franceschi, P., Paris, R., Sansavini, S., Tartarini, S., 2015. Genetic diversity, population  
 371 structure and construction of a core collection of apple cultivars from Italian germplasm. *Plant Mol. Biol. Rep*. 33, 458–  
 372 473.

373 Maguire, T.L., Collins, G.G., Sedgley, M., 1994. A modified CTAB DNA extraction procedure for plants belonging to  
 374 the family Proteaceae. *Plant Mol. Biol. Rep*. 12, 106–109.

375 Martinelli, F.M., Busconi, M., Camangi, F., Fogher, C., Stefani, A., Sebastiani, L., 2008. Ancient Pomoideae *Malus*  
 376 *domestica* Borkh. and *Pyrus communis* L. cultivars in “Appenino Toscano” Tuscany, Italy: molecular SSR and  
 377 morphological characterization. *Caryologia*, 61, 320–331.

378 Martinelli, F., Busconi, M., Fogher, C., and Sebastiani, L., 2009. Development of an efficient regeneration protocol for  
 379 pear rootstock Pyrodwarf and assessment of SSR variability in regenerating shoots. *Caryologia* 62, 62–68.

380 Miranda, C., Urrestarazu, J., Santesteban, L.G., Royo, J.B., Urbina, V., 2010. Genetic diversity and structure in a  
 381 collection of ancient Spanish pear cultivars assessed by microsatellite markers. *J. Am. Soc. Hortic. Sci.* 135, 428–437.

382 Monte–Corvo, L., Cabrita, L., Oliveira, C., and Leitão, J., 2000. Assessment of genetic relationships among *Pyrus*  
 383 species and cultivars using AFLP and RAPD markers. *Genet. Resour. Crop. Evol.* 47, 257–265.

384 Muresu, E., Castiglia, P., Piana, A., Masia, M.D., Ciappeddu, P.L., 1997. Caratterizzazione di prodotti alimentari  
 385 dell'area sardo-corsa: analisi chimico-fisica e batteriologica. *Atti Workshop Caratterizzazione di prodotti alimentari*  
 386 dell'area sardo-corsa. Sassari 11 aprile 1997. pp. 5–18.

387 Oliveira, C.M., Mota, M., Monte–Corvo, L., Goulão, L., and Silva, D.M., 1999. Molecular typing of *Pyrus* based on  
 388 RAPD markers. *Sci. Hortic.* 79, 163–174.

389 Ordidge, M., & Venison, E., 2018. Simple Sequence Repeat (SSR) Fingerprint data for apples from the National Fruit  
 390 Collection as used in Ordidge et al PLOS One 2018. University of Reading. Dataset. <http://dx.doi.org/10.17864/1947.163>

391 Ouni, R., Zborowska, A., Sehic, J., Choulak, S., Hormaza, J.I., Garkava-Gustavsson, L., Mars, M., 2020. Genetic  
 392 diversity and structure of Tunisian local pear germplasm as revealed by SSR markers. *Hortic. Plant J.* pp. 74–83.

393 Paetkau D., Calvert W., Stirling I., Strobeck C. 1995. Microsatellite analysis of population structure in Canadian polar  
 394 bears. *Mol. Ecol.* 4, 347–354.

395 Pierantoni, L., Cho, K.H., Shin, I.S., Chiodini, R., Tartarini, S., Dondini, L., Sansavini, S., 2004. Characterisation and  
 396 transferability of apple SSRs to two European pear F1 populations. *Theor Appl Genet.* 109, 1519–1524.

397 Powell, W., Machray, G.C., Provan, J., 1996. Polymorphism revealed by simple sequence repeats. *Trends Plant Sci.* 1,  
 398 215–222.

399 Pritchard, J.K., Stephens, M., Donnelly, P., 2000. Inference of population structure using multilocus genotype data.  
 400 *Genetics.* 155, 945–959.

401 Puskas, M., Hofer, M., Sestras, R.E., Peil, A., Sestras, A.F., Hanke, M.V., Flachowsky, E., 2015. Molecular and flow  
 402 cytometric evaluation of pear (*Pyrus communis* L.) genetic resources of the German and Romanian national fruit  
 403 collections. Genet. Res. Crop Evol. 63, 1023–1033.

404 Queiroz, A., Assunção, A., Ramadas, I., Viegas, W., Veloso, M.M., 2015. Molecular characterization of Portuguese  
 405 pear landraces (*Pyrus communis* L.) using SSR markers. Sci. Hortic. 183, 72–76.

406 Queiroz, A., Bagoin Guimarães, J., Sánchez, C., Simões, F., Maia de Sousa, R., Viegas, W., Veloso, M.M., 2019.  
 407 Genetic diversity and structure of the Portuguese pear (*Pyrus communis* L.) germplasm. Sustainability, 11, 5340.

408 Rana, J.C., Chahota, R.K., Sharma, V., Rana, M., Verma, N., Verma, B., Sharma, T.R., 2015. Genetic diversity and  
 409 structure of *Pyrus* accessions of Indian Himalayan region based on morphological and SSR markers. Tree Genet. Genomes.  
 410 11, 821.

411 Rohlf, F.J., 1998. NTSyS–p.c. Numerical taxonomy and multivariate analysis system, version 2.0. Exeter Software  
 412 Publishers Ltd., Setauket.

413 Sehic, J., Garkava–Gustavsson, L., Fernández–Fernández, F., Nybom, H., 2012. Genetic diversity in a collection of  
 414 European pear *Pyrus communis* cultivars determined with SSR markers chosen by ECPGR. Sci. Hortic. 145, 39–45.

415 Sisko, M., Javornik, B., Siftar, A., and Ivancic, A., 2009. Genetic relationships among Slovenian pears assessed by  
 416 molecular markers. J. Am. Soc. Hortic. Sci. 134, 97–108.

417 Urrestarazu, J., Denancé, C., Ravon, E., Guyader, A., Guisnel, R., Feugey, L., Poncet, C., Lateur, M., Houben, P.,  
 418 Ordidge, M., Fernandez-Fernandez, F., Evans, K.M., Paprstein, F., Sedlak, J., Nybom, H., Garkava-Gustavsson, L.,  
 419 Miranda, C., Gassmann, J., Kellerhals, M., Suprun, I., Pikunova, A.V., Krasova, N.G., Torutaeva, E., Dondini, L., Tartarini,  
 420 S., Laurens, F., Durel, C.E., 2016. Analysis of the genetic diversity and structure across a wide range of germplasm reveals  
 421 prominent gene flow in apple at the European level. BMC Plant Biol. 16-130.

422 Urrestarazu, J., Miranda, C., Santesteban, L.G., Royo, J.B., 2012. Genetic diversity and structure of local apple cultivars  
 423 from northeastern Spain assessed by microsatellite markers. Tree Genet. Genomes. 8, 1163–1180.

424 Urrestarazu, J., Royo, J.B., Santesteban, L.G., Miranda, C., 2015. Evaluating the influence of the microsatellite marker  
 425 set on the genetic structure inferred in *Pyrus communis* L. PloS ONE. 10, 9.

426 Vincent, H., Wiersema, J., Kell, S., Fielder, H., Dobbie, S., Castañeda–Álvarez, N.P., Maxted, N., 2013. A prioritized  
 427 crop wild relative inventory to help underpin global food security. Biol. Conserv. 167, 265–275.



428 Weber, J.L., & May, P.E., 1989. Abundant class of human DNA polymorphisms which can be typed using the  
 429 polymerase chain reaction. *Am. J. Hum. Genet.* 44, 388-396.

430 Wünsch, A., & Hormaza, J. I. (2007). Characterization of variability and genetic similarity of European pear using  
 431 microsatellite loci developed in apple. *Sci. Hortic.* 113, 37–43.

432 Yamamoto, T., Kimura, T., Shoda, M., Ban, Y., Hayashi, T., and Matsuta, N., 2002a. Development of microsatellite  
 433 markers in Japanese pear *Pyrus pyrifolia* Nakai. *Mol. Ecol. Notes.* 2, 14–16.

434 Yamamoto, T., Kimura, T., Shoda, M., Imai, T., Saito, T., Sawamura, Y., Kotobuki, K., Hayashi, T., Matsuta, N., 2002b.  
 435 Genetic linkage maps constructed by using an interspecific cross between Japanese and European pears. *Theor. Appl.*  
 436 *Genet.* 106, 9–18.

437 Yue, X., Zheng, X., Zong, Y., Jiang, S., Hu, C., Yu, at al., (2018). Combined analyses of chloroplast DNA haplotypes  
 438 and microsatellite markers reveal new insights into the origin and dissemination route of cultivated pears native to East  
 439 Asia. *Frontiers in plant science*, 9, 591.

440 Zielinski, Q.B. & Thompson M., 1967. Speciation in *Pyrus*: chromosome number and meiotic behavior. *Bot. Gaz.* 128,  
 441 109–112.