

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Study of the efficacy of Aureobasidium strains belonging to three different species: *A. pullulans*, *A. subglaciale*, and *A. melanogenum* against *Botrytis cinerea* of tomato

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Alessandra Di Francesco, M.D.F. (2020). Study of the efficacy of Aureobasidium strains belonging to three different species: *A. pullulans*, *A. subglaciale*, and *A. melanogenum* against *Botrytis cinerea* of tomato. ANNALS OF APPLIED BIOLOGY, 177(2), 266-275 [10.1111/aab.12627].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/766018> since: 2024-05-07

*Published:*

DOI: <http://doi.org/10.1111/aab.12627>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

# Study of the efficacy of *Aureobasidium* strains belonging to three different species: *A. pullulans*, *A. subglaciale*, and *A. melanogenum* against *Botrytis cinerea* of tomato

Alessandra Di Francesco<sup>1,2</sup> | Michele Di Foggia<sup>3</sup> | Janja Zajc<sup>4</sup> | Nina Gunde-Cimerman<sup>5</sup> | Elena Baraldi<sup>1,2</sup>

<sup>1</sup>CRIOF–Department of Agricultural Sciences, University of Bologna, Via Gandolfi, 19, 40057 Cadriano, Bologna, Italy.

<sup>2</sup>Department of Agricultural and Food Sciences, University of Bologna, Viale Fanin, 42, 40127 Bologna, Italy.

<sup>3</sup>Department of Biomedical and Neuromotor Sciences, University of Bologna, Via Belmeloro, 8/2, 40126 Bologna, Italy.

<sup>4</sup>Plant Protection Department, Agricultural Institute of Slovenia, Hacquetova ulica 17, SI-1000 Ljubljana, Slovenia.

<sup>5</sup>Department of Biology, Biotechnical Faculty, University of Ljubljana, Jamnikarjeva 101, SI-1000 Ljubljana, Slovenia.

## Correspondence

Alessandra Di Francesco, CRIOF–Department of Agricultural Sciences, University of Bologna, Via Gandolfi, 19, 40057 Cadriano, Bologna, Italy. Email: [alessand.difrancesc3@unibo.it](mailto:alessand.difrancesc3@unibo.it)

Received: 7 April 2020; revised manuscripts received: 8 and 25 June 2020; accepted: 26 June 2020.

## Abstract

The difference in antagonistic activity against the causal agent of grey mould (*Botrytis cinerea*) of tomato between *Aureobasidium* strains belonging to three different species, namely *A. pullulans*, *A. melanogenum*, and *A. subglaciale*, was evaluated by *in vitro* and *in vivo* assays. In the yeast-pathogen direct interaction experiment, all the strains significantly reduced *B. cinerea* growth, with *A. melanogenum* the least efficient species (17.8% of reduction) compared to *A. pullulans* and *subglaciale* (22% and 27.8%). The non-volatile metabolites produced by all three species reduced mycelial growth between 95–100%. These metabolites were characterized by FT-IR spectroscopy as polysaccharides, lytic enzymes, siderophores, and antibiotics. The inhibitory effect of *Aureobasidium* strains on pathogenic enzymes such as xylanase, polygalacturonase, and pectinase was measured showing *A. pullulans* strains as capable of strong inhibition of xylanase, an enzyme

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/aab.12627

directly related to the virulence of necrotrophic pathogens such as *B. cinerea*. Our data demonstrate that the different species of *Aureobasidium* isolated from a range of non-conventional environments exerted variable efficacy against *B. cinerea*, with *A. pullulans* as the most active species followed by *A. subglaciale*, and *A. melanogenum* as ineffective and not suitable for biocontrol applications.

## **Keywords**

Yeasts, Non Volatile Metabolites, Enzymes, Bio Control

## **1 | Introduction**

*Aureobasidium pullulans* (De Bary) G. Arnaud is a former species complex of black yeast-like fungi, particularly known for their biotechnological significance as producers of an extracellular polysaccharide known as ‘pullulan’, a promising biomaterial for the packaging of food (Rekha & Chandra, 2007; Singh *et al.*, 2008). Importantly, in the last years, *A. pullulans* has been considered as one of the most efficient microbial species to control fungal diseases of fruits and vegetables mainly during the post-harvest phase, used as a biocontrol agent (BCA) alone or associated with sustainable physical methods (Zhang *et al.*, 2010; Di Francesco & Mari, 2014; Di Francesco *et al.*, 2018a, 2020).

The multiloci DNA analysis distinguished four varieties of the species *A. pullulans* – var. *pullulans*, *melanogenum*, *subglaciale*, and *namibiae* (Zalar *et al.*, 2008), and later on distinguished into four well-defined species (Gostinčar *et al.*, 2014).

*Aureobasidium pullulans* is most frequently found in the phyllosphere and carposphere of various fruits and vegetables and is associated with the endophyte population of many plant species (Bozoudi & Tsaltas, 2018). Moreover, its ecology is highly diverse and cosmopolitan, as it can be found in environments ranging from hypersaline water of salterns, glacial ice, polluted water, frozen and salt-preserved food, household surfaces and house dust, synthetic polymers, and aviation fuel

tanks (Gostinčar *et al.*, 2019). The species *A. melanogenum* is mainly found in oligotrophic and aqueous environments. Importantly, it grows at 37°C, when other species of the genus cannot, and is considered as an emerging human pathogen (Chan *et al.*, 2011). Conversely, *A. subglaciale* shows a psychrotolerant nature (it grows at 4°C) and occurs mainly in glacial habitats. Finally, *A. namibiae* was named based on a single isolate from Namib Desert marble, able to tolerate up to 10% of NaCl and grows between 10°C and 30°C (Zalar *et al.*, 2008; Gostincar *et al.*, 2014).

In the most recent studies, *A. pullulans* was shown to have substantial biocontrol efficacy for the control of postharvest pathogens such as *Monilinia* spp. of stone fruits (Di Francesco *et al.*, 2020), *Botrytis cinerea* of grape, strawberry, and kiwifruit (Parafati *et al.*, 2015, 2017; Di Francesco *et al.*, 2018b) and *Penicillium expansum* of apple (Mari *et al.*, 2012); this efficacy can be attributed to different mechanisms of action such as the production of antifungal compounds (volatile and non-volatile) and cell wall degrading enzymes (Parafati *et al.*, 2015; Zhang *et al.*, 2010), the competition for nutrients and space (Janisiewicz *et al.*, 2000, Di Francesco *et al.*, 2018b; Klein & Kupper, 2018), induction of resistance (Madhupani & Adikaram, 2017; Di Francesco *et al.*, 2017), and mycoparasitism (Klein & Kupper, 2018). In fact, the application of BCAs, such as *A. pullulans*, is considered a safe and environmentally friendly alternative (Zhang *et al.*, 2010) to synthetic fungicides, considering their massive use in crop protection with possible harmful effects on human health, and the emergence of pathogen resistance to single-site mode of action fungicides (Eckert & Ogawa, 1988).

No studies have been conducted to explore the efficacy of utilizing *Aureobasidium* spp. for the control of postharvest fungal pathogens. The objective of this study was to evaluate the principal differences of efficacy between the strains belonging to the three species: *A. pullulans*, *A. subglaciale*, and *A. melanogenum* by testing their ability to inhibit *B. cinerea*, one of the most important pre-harvest and post-harvest fungal pathogen on fresh-market tomatoes (Mari *et al.*, 1996).

This research was conducted through i) the evaluation of the pathogen mycelial growth inhibition in co-culturing; ii) the activity of non-volatile metabolites; iii) the analysis of the non-volatile compounds by FT-IR spectroscopy; iv) the evaluation of the inhibitory effect of *Aureobasidium* species on the fungal pathogenic enzymes; v) and the reduction in grey mould incidence on tomato fruit.

## 2 | Materials and methods

### 2.1 | *Aureobasidium pullulans* strains

The selected *Aureobasidium* spp. strains (Table 1) were obtained from the Culture Collection Ex Infrastructural Centre Mycosmo, (Department of Biology, Biotechnical Faculty, University of Ljubljana, Ljubljana, Slovenia), and the Westerdijk Fungal Biodiversity Institute (Utrecht, The Netherlands). Strains were maintained on nutrient yeast dextrose agar (NYDA: 8 g of nutrient broth, 5 g of yeast extract, 10 g of dextrose, and 15 g of technical agar, Oxoid, Basingstoke - UK) in 1 L of distilled water) and stored at 4 °C until use. Two days before the experiments, each antagonist was grown on NYDA at 25°C, and yeast cells were collected in sterile distilled water (SDW) containing 0.05% (v/v) Tween 80 (Sigma-Aldrich, St. Louis, MO, USA) then adjusted to a final concentration of  $10^8$  cells mL<sup>-1</sup>.

### 2.2 | Pathogen

*Botrytis cinerea* (CRIOF-DipSA collection), derived from a diseased tomato, was grown on oatmeal agar (60 g of oatmeal, 10 g of sodium nitrate, 30 g of sucrose, 12 g of agar per 1 L of SDW, Oxoid, Basingstoke - UK) and incubated at 25°C for 10 days. Pathogen conidia were collected and suspended in SDW containing 0.05% (v/v) Tween 80 and the suspension was adjusted to  $10^5$  conidia mL<sup>-1</sup>, with a haemocytometer.

## 2.3 | Tomato

“Datterini” tomatoes (*Solanum lycopersicum*, L.) obtained from an organic orchard located in Cesena (Italy) were used. Tomatoes were harvested at commercial maturity and immediately used. Fruits were wounded with a sterile nail (3 mm × 3 mm × 3 mm) on the middle line (one wound per fruit) and inoculated with yeast strains cell suspensions and pathogen conidial suspension.

## 2.4 | *In vitro* assays

In the co-culturing assay, *B. cinerea* was separately co-cultured with each yeast strain on potato dextrose agar (PDA: 39 g per 1 L of SDW, Oxoid, Basingstoke - UK) plates. In each plate, a mycelial plug (6 mm in diameter) derived from a 6-day-old colony of *B. cinerea* was inoculated 30 mm from the plate edge and *A. pullulans* cells, from a 48 h-old colony grown on NYDA, were streaked by a sterile loop 30 mm from the fungal plug (Di Francesco *et al.*, 2017). In control plates, only *B. cinerea* plugs were inoculated. Plates were incubated at 25°C in the dark for 5 days, and the radius of the fungal colony was measured on a line from the centre of the plug to the yeast swipe.

For the No-VOCs (Non-Volatile Organic Compounds) assay, cells suspension (100 µL; 10<sup>8</sup> cells mL<sup>-1</sup>) of each *Aureobasidium* spp. strain obtained from 48 h-old colonies grown on NYDA, was spread on a sterile cellophane layer (Safta, PC, Italy) previously positioned on NYDA plate. In control plates, 100 µL of SDW were used.

After 48-h of incubation at 25°C, cellophane was removed, and a mycelial plug (6 mm diameter) of *B. cinerea* was inoculated in the centre of the plate. The plates were incubated at 25°C and the colony diameter was measured after 5 days from the inoculation.

In all experiments, five plates (replicates) were used for each combination and the controls. The experiments were conducted twice.

## 2.5 | Antagonists metabolites influence on *Botrytis cinerea* pathogenic enzymes

To verify the effect of secondary metabolites produced by *Aureobasidium* spp. strains on the pathogenic enzymes activity of *B. cinerea*, tissue culture plates (Costar, Corning Inc., Corning, NY) and culture plate inserts Millicell-CM (Millipore Corp., Bedford, MA) were used as reported by Janisiewicz *et al.*, (2000). The system allows for the interchange of metabolites without physical contact between antagonists cells and pathogen conidia. Aliquots (120  $\mu$ L) of fungal suspension ( $10^5$  conidia  $\text{mL}^{-1}$ ) derived from 6 day-old colony or water (control) were dispensed in the wells of culture plates, while the same aliquots of *Aureobasidium* spp. cells suspensions ( $10^8$  cells  $\text{mL}^{-1}$ ) were dispensed inside the cylinder inserts. The plates with the cylinders were placed at 25°C on a rotary shaker at 50 rpm; after 6 h of incubation, cylinders were removed from the wells and fungal conidia aliquots were used for the enzymes assays. For each *Aureobasidium* spp. strain three wells of culture plate were used. The experiment was performed twice.

## 2.6 | Enzyme assays

Xylanase, polygalacturonase, and pectinase assays were performed in Petri dishes containing specific agar medium previously described in Di Francesco *et al.* (2018c). For the xylanase assay, the medium contained 0.5% beechwood xylan (Sigma-Aldrich) in a minimal medium consisting of  $\text{NaNO}_3$  0.3%;  $\text{KH}_2\text{PO}_4$  0.1%;  $\text{MgSO}_4$  0.05%; yeast extract 0.1%; agar 1.2% (St Leger *et al.*, 1997).

Polygalacturonase activity was determined through the Eriksson & Pettersson culture medium (Eriksson & Pettersson, 1975) enriched with the sodium salt of polygalacturonic acid (Sigma-Aldrich) from citrus fruit (0.5%) and agarized with low calcium Oxoid No. 1 agar (2%, Oxoid, Basingstoke, UK). The medium was sterilized following the procedure of Ayers *et al.* (1966), then the pH was adjusted to 5 and 8, respectively, by adding NaOH 1 M. Polymethylgalacturonase activity was determined through the above-cited culture medium (Eriksson & Pettersson, 1975) enriched with Sigma citrus pectin (0.5%) and agarized with calcium-rich Oxoid No. 3 agar (2%).

The growth media was sterilized following the procedure of Durrands & Cooper (1988) and the pH was adjusted to 8 by adding HCl 1 M.

Forty microlitres of each treated pathogen suspension ( $10^5$  conidia  $\text{mL}^{-1}$ ) were placed in holes (three equidistant holes per dish) punched in the medium with a 5 mm-sterile cork borer. Control plates were inoculated with the same volumes of conidial suspension exposed to distilled water. After incubation at 20°C for 48 h, enzyme activities were determined with specific colorimetric methods visualizing a clear zone of substrate degradation around the holes. For xylanase, the clearing halo was visualized by staining with Congo red, then destaining with NaCl and measured as described above. Instead, the halo produced by pectic hydrolase activities was visualized by 1% cetyl methyl ammonium bromide (CTAB) dissolved in distilled water, heated at 30°C. The diameter of enzymes halos (mm) was measured with a ruler. For each treatment, five plates (replicates) were considered. Each assay was repeated twice.

## 2.7 | Spectroscopic analysis of yeasts secondary metabolites

Flasks containing 100 mL of NYDB medium (NYDA without agar) were inoculated with 100  $\mu\text{L}$  of antagonists cell suspensions ( $10^8$  cells  $\text{mL}^{-1}$ ) and then incubated at 25°C in a rotary shaker (250 rpm) for 1, 24 and 48 h. After each incubation time, the liquid cultures were centrifuged at  $5000 \times g$  for 20 min at room temperature and the supernatant collected in sterile tubes (2 mL), stored at -80°C and lyophilized. The control consisted of a liquid medium (NYDB) depleted of yeast cells. The supernatant was analyzed by FT-IR spectroscopy to obtain a rapid and non-destructive characterization of their main molecular components. Infrared spectra were recorded with a Bruker ALPHA series FT-IR spectrophotometer (Bruker, Ettlingen, Germany) equipped with an apparatus for attenuated total reflectance (ATR-Diamond crystal). The spectra were collected from 4000 to 400  $\text{cm}^{-1}$  and averaged over 100 scans (resolution = 4  $\text{cm}^{-1}$ ): four spectra were measured for each sample for each sampling time.



## 2.8 | *In vivo* assay

Tomatoes were surface disinfected by 1 min immersion in sodium hypochlorite solution (1%), rinsed with tap water, air-dried at room temperature and inoculated through artificial wounds by a sterile nail (3 mm × 3 mm × 3 mm) on the equator (one wound) with 15 µL of each antagonist suspension ( $10^8$  cells mL<sup>-1</sup>) (15 strains). After 1 h, 15 µL of *B. cinerea* suspension ( $10^5$  conidia mL<sup>-1</sup>) were inoculated by placement in the treated tomato wound. The control consisted of fruit inoculated with SDW. Tomatoes were kept at 20°C and 90% of relative humidity (RH). The disease incidence was detected after 6 days from the inoculation. Sixty tomatoes for each antagonist and control were used. The experiment was conducted three times.

## 2.9 | Statistical analysis

Data were analysed by one-way analysis of variance (ANOVA). Statistical comparison of means was carried out by using Fisher's LSD test ( $P < 0.05$ ). Data were reported as mean values ± standard deviation (SD) of two and three independent experiments. All analyses were performed with the software Statgraphic Plus Version 2.1 (Statistical Graphics Corp., USA 1996).

# 3 | Results

## 3.1 | *In vitro* assays

The effect of the three-tested *Aureobasidium* species on *B. cinerea* mycelial growth in the co-culturing condition and to the exposure of non-volatile metabolites produced by the strains was determined.

All the strains significantly reduced the pathogen colony growth. *Aureobasidium melanogenum* showed a lower efficacy compared to *A. pullulans* and *A. subglaciale* (17.8%, 22% and 27.8% respectively on average). Within each species, the strains AP3 (*A. pullulans*), AM9 (*A.*

*melanogenum*), and AS15 (*A. subglaciale*) resulted the most active against *B. cinerea* compared to the other tested strains (Fig. 1).

Concerning the pathogen colony growth, the non-volatile metabolites of all the tested strains significantly inhibited *B. cinerea* mycelial growth ranging from 95% to 100% (data not shown).

### 3.2 | Antagonists metabolites influence on *Botrytis cinerea* pathogenic enzymes

The effect of secondary metabolites produced by *A. pullulans*, *A. melanogenum*, *A. subglaciale* strains was evaluated on the activity of virulence key factors of *B. cinerea*. Three lytic enzymes: xylanase, pectinase, and polygalacturonase, were considered. Among these, xylanase activity produced by *B. cinerea* was the prevalent as shown by control samples (320 mm<sup>2</sup> halo area) compared to pectinase and polygalacturonase (180 and 90 mm<sup>2</sup>, respectively) (Fig. 2). Almost all the strains inhibited these enzymes. Conversely, the metabolites produced by the strain *A. melanogenum* AM10 and almost all the strains of *A. subglaciale* seemed to stimulate the fungal xylanase activity. An increase of 51.9% and 12.6% (on average, AS11, AS12, AS14 strains enzyme activity) was recorded, respectively for *A. melanogenum* AM10 strain and *A. subglaciale* strains. Conversely, *A. pullulans* showed the highest percentage of reduction (54.8%) (on average, AP1 and AP3 strains) of the above-mentioned enzyme.

Pectinase activity was mainly inhibited by *A. pullulans* strains on average by 47.6%, followed by *A. subglaciale* (40.1%), and *A. melanogenum* (30.5%). The most efficient strains were AP5 and AS15. Polygalacturonase enzyme resulted in the lowest detectable activity; however, *A. pullulans* AP3 and AP5, *A. subglaciale* AS13, and AS15 strains were confirmed as the most active to inhibit the target enzyme (by 42.7%, 42.7% and 47.9%, 46.8%, respectively).

### 3.3 | Yeasts secondary metabolites characterization

The FT-IR spectroscopic measurements were carried out to determine the main non-volatile compounds produced by three randomly chosen yeast strains (AP5, AS15, and AM6), that previously showed a high inhibition of *B. cinerea* mycelial growth in the No-VOCs assays.

The spectra differences between the cultures of the various strains at 48 h and control are reported in Fig. 3: positive peaks are related to yeasts secondary metabolites produced during the 48 h of incubation, while negative peaks should be attributed to the consumption of the culture media. The peaks attributed to the nutrient medium were derived from the FT-IR spectrum of pure NYDB (data not shown), while the attribution of carbohydrates, enzymes, siderophores and antifungal compounds was based on previously reported protocols (Aghatabay *et al.*, 2014; Bae *et al.*, 2015; Cozar *et al.*, 2006; Dahiya *et al.*, 2016; Higgins *et al.*, 2006; Holmen *et al.*, 1997; Shingel, 2002; Tu, 1982; Um *et al.*, 2013; Yang *et al.*, 2006; Zhbankov *et al.*, 2000). Positive spectroscopic peaks were attributed to different classes of compounds already known to be produced by *Aureobasidium* strains (Freimoser *et al.*, 2019): complex carbohydrates (Bozoudi & Tsaltas, 2018), as pullulan and other glucans involved in the biofilm formation used as biocontrol mechanism by the yeasts; lytic enzymes (Bozoudi & Tsaltas, 2018), as amylases, cellulases, lipases, xylanases, proteases, laccase, chitinases and mannanases; antifungal compounds (Bozoudi & Tsaltas, 2018), as aureobasidins (derivatives of cyclic depsipeptides possessing, as typical amino acids, proline, leucine and phenylalanine) and liamocins (polyols containing mannitol headgroup acetylated with polyesters tails, Freimoser *et al.*, 2019); siderophores used as iron depriving agents (Bozoudi & Tsaltas, 2018). The three spectra differences showed in Fig. 4 present a quite similar pattern of negative and positive peaks (in particular AP5 and AS15), while AM6 showed some differences in several spectral regions (highlighted in blue): between 3000 and 2800  $\text{cm}^{-1}$  (C-H stretching vibrations), 1750–1500  $\text{cm}^{-1}$  (carbonyl derivatives, as amides, and aromatic compounds vibrations), 1450–1350  $\text{cm}^{-1}$  (C-H bending and carboxylate groups vibrations), 1200–900  $\text{cm}^{-1}$  (C-O and C-H vibrations); 850-650  $\text{cm}^{-1}$  (chain deformation and aromatic rings bending vibrations) and between 550 and 500  $\text{cm}^{-1}$  (chain deformation). More in detail, the intense positive peaks observed in AM6 at 1720, 1614,

1170, 1075, 1025, 792 and 724  $\text{cm}^{-1}$  can be all attributed to melanin-related vibrations (Pierce & Rast, 1995): melanin is a heterogeneous class of natural pigments produced by *A. melanogenum*. The other positive peaks present in both AP5 and AS15, but absent in AM6 are: 1630, 1589, 1520, 1396, 1088, 934, 664 and 535  $\text{cm}^{-1}$ : the attribution of most of them can be ambiguous, because of the chemical similarities among the secondary metabolites produced by the different *Aureobasidium* strains. Enzymes, aureobasidins, and siderophores are all characterized by the presence of the amide group ( $\text{HN-C=O}$ ), that can be attributed to 1630 (amide I), 1520 (amide II) and 664 (amide IV)  $\text{cm}^{-1}$  bands. Moreover, other common chemical groups to these three classes of metabolites are carboxylate ions ( $\text{COO}^-$ ) and aromatic rings to which the 1396 and 1589  $\text{cm}^{-1}$  bands can be attributed, respectively. Only the 535  $\text{cm}^{-1}$  band can be uniquely attributed to hydroxymate siderophores (Higgins *et al.*, 2006; Holmen *et al.*, 1997; Yang *et al.*, 2006) bound to ferric ions. Again, another source of ambiguity concerns carbohydrates-related vibrations (2854, 1088, 934 and 822  $\text{cm}^{-1}$ ): biofilm produced by *Aureobasidium* is composed of pullulan and a mixture of  $\alpha$  and  $\beta$ -glucans; most of the lytic enzymes (i.e. polygalacturonase) are heavily glycosylated proteins, while liamocins contain the mannitol moiety.

To sum up the previous findings, it can be stated that both AP5 and AS15 strains showed an increased concentration of complex carbohydrates, enzymes, siderophores and antifungal compounds that are not present in AM6 spectrum or are even negative (i.e. 2854, 1396, 934 and 822  $\text{cm}^{-1}$  bands).

### 3.4 | *In vivo* assay

Antagonists strains were applied as a preventative treatment against grey mould on tomato cv “Datterini”. After 6 days of incubation at 20°C, grey mould disease incidence was evaluated. AP5 and AS15 reduced *B. cinerea* incidence by 80.9% and 55.5% respectively, showing the best efficacy respect to all the other strains (Fig. 4). All the remaining strains inhibited rot of less than

50%. Interestingly, the strain AM10 increased disease incidence compared to the control, suggesting a stimulatory effect of this strain on the fungal pathogen.

#### 4 | Discussion

*Aureobasidium pullulans*, now separated as different species, is a ubiquitous oligotroph that can be found in various environments such as hypersaline waters in salterns (Cimerman *et al.*, 2000), on rocks and monuments (Urzi *et al.*, 1999), in the atmosphere as airborne spores (Punnapayak *et al.*, 2003), in food and feeds (Sasahara & Izumori, 2005), in deserts (Jiang *et al.*, 2017), in freshwater, and even in glaciers (Gostinčar *et al.*, 2014). This broad ecology can be attributed to its great adaptability to various environmental conditions, poly-extremo tolerance to cope with different stresses and to their nutritional versatility (Gostinčar *et al.*, 2011) being armoured with a plethora of extracellular enzymes (Gostinčar *et al.*, 2014). All these characteristics contribute importantly to the biocontrol potential of this microorganism.

In the present work, the antagonistic differences between the strains of three different *Aureobasidium* species, namely *A. pullulans*, *A. melanogenum*, and *A. subglaciale*, against the grey mould of tomato were evaluated for the first time. The selection of *Aureobasidium* strains for this study was based on previous experiments, where yeast strains from heterogeneous and extreme environments were tested as potential biocontrol agents (Zajc *et al.*, 2018). On the other hand, strains of *A. melanogenum* were used as a highly related outgroup of strains that could not be used in biocontrol due to their potential pathogenicity (Zajc *et al.*, 2019).

The phylogenetic diversity of these species could be linked also to their different antagonistic behaviour and efficacy to control plant pathogens. The present study confirmed *A. pullulans* strains together with *A. subglaciale* as being the most active to control *B. cinerea* in *in vivo* assay by reducing the mould incidence by 40.3% and 42.8% (on average) on tomatoes as well as in *in vitro*

co-culturing assay by showing inhibition of 22% and 27.8% (on average), respectively; also, almost a total inhibition by No-VOCs production of the pathogen mycelium growth was displayed.

Gostinčar *et al.*, (2014) demonstrated that, due to the abundant production of extracellular enzymes and the high number of sugar transporters, *A. pullulans* has ecological preference associated to plants compared to the other species. In fact, previous studies of post-harvest diseases control were all conducted by using strains of *A. pullulans* isolated from plant or fruit (Di Francesco *et al.*, 2018; Banani *et al.*, 2014; Parafati *et al.*, 2015; Schena *et al.*, 2003; Castoria *et al.*, 2001; Ippolito *et al.*, 2000). This was recently confirmed in a study of characterisation of various strains of antagonistic *A. pullulans* from leaves and fruits from different parts of the world (Zajc *et al.*, 2020).

With regards to *A. subglaciale*, its extreme tolerance to various environmental factors could confer it a major ability to compete with fungal pathogens in storage conditions. On the other hand, *A. subglaciale* has a much more restricted distribution and showed a great ability to resist to gamma irradiation, UV light, and heavy metal ions (Liu *et al.*, 2017). According to its physiology, *A. subglaciale* displayed two important characteristics: it is a typical epiphyte and it belongs to the group of psychrophilic species (Kachalkin, 2010). Nevertheless, some strains belonging to the two above-mentioned species can present different efficacy, probably because the efficacy may vary based on fungal pathogen and pathosystem (Whipps, 2001; Botha, 2011).

*Aureobasidium melanogenum* was previously recognized as an opportunistic human pathogen (Gostinčar *et al.*, 2014) and our results showed how it was the least active species against *B. cinerea*, confirming its inadequate use in biological control of post harvest fruit fungal diseases.

*Aureobasidium pullulans* is also well known for its biotechnological potential, among other things for the production of the polysaccharide pullulan (Cheng *et al.*, 2011) and its biocontrol use in agriculture (Sharma *et al.*, 2009). It also produces a large spectrum of extracellular enzymes (Molnarova *et al.*, 2013) and an antifungal peptide such as aureobasidin A (Takesako *et al.*, 1991). Yeast-like cells, mycelia and melanin-pigmented chlamydospores of genus *Aureobasidium* are particularly known for their biotechnological significance since this organism is a producer of the

extracellular polysaccharide (EPS) pullulan (poly- $\alpha$ -1,6-maltotriose) (de Hoog, 1993, Gostinčar *et al.*, 2014, Peterson *et al.*, 2013). The FT-IR characterization revealed that *A. pullulans* and *A. subglaciale* strains showed many similarities in the production of several classes of secondary metabolites compounds: in particular, peaks attributed to complex carbohydrates (such as pullulan or other  $\alpha$  and  $\beta$ -glucans), degrading enzymes, cyclic depsipeptides (aureobasidins), siderophores and other antifungal compounds were detected. Pullulan and other polysaccharides may improve the biofilm formation and therefore adhesion, thus explaining the better performance of these two species in both *in vitro* and *in vivo* assays, together with a higher production of degrading enzymes (i.e. amylases, proteases or lipases), aureobasidins (Bozoudi & Tsaltas, 2018; Freimoser *et al.*, 2019), and siderophores, confirmed by the diagnostic band at 530  $\text{cm}^{-1}$  (Cozar *et al.*, 2006). On the contrary, the spectrum of AM6 was characterized by both the absence of many of the bands attributed to the previous classes of non-volatile metabolites and the presence of several positive peaks attributed to melanin pigments that could explain its worst performance in the inhibition of *B. cinerea*.

Nigam (2013) asserted that filamentous microorganisms can produce and secrete a wide array of active compounds including hydrolytic enzymes and enzyme inhibitors. Indeed, yeast-like fungus *A. pullulans* produces different polymers, lipids, volatile compounds, enzymes, and secondary metabolites, that confer it a marked antagonistic activity against bacteria and fungi (Freimoser *et al.*, 2019; Prasongsuk *et al.*, 2018; Price *et al.*, 2013, 2017; Takesako *et al.*, 1991; Zain *et al.*, 2009) by inhibiting their virulence. Our results showed that almost all the tested yeast strains inhibited the enzymes pathogenic activity on *B. cinerea*. The species *A. pullulans* displayed the highest xylanase and pectinase inhibitory activity, enzymes directly related to the pathogenicity/virulence of a microorganism (Douaiher *et al.*, 2007; Kikot *et al.*, 2009) and responsible for the fruit spoilage (Al-Hindi *et al.*, 2011), and one of the major Cell Wall Degrading Enzyme (CWDE) of fruit fungal pathogens, respectively.

Nevertheless, in some cases, xylanase enzyme was enhanced by individual strains mainly belonging to *A. melanogenum* and *A. subglaciale* acting like pathogenic enzyme elicitors. It can be assumed that, thus, the efficacy of BCAs strongly depend on their ecological competence (Khol *et al.* 2019). Our data demonstrated that different species of *Aureobasidium* exerted a different efficacy against *B. cinerea*. This was especially obvious for *A. melanogenum*, that showed the lowest antagonistic activity compared to *A. pullulans* and *A. subglaciale*.

Moreover, all the tested varieties were able to produce No-VOCs, essential for biocontrol activity of *B. cinerea*. More investigations on the study of other mechanisms of action should be performed to understand the exact contribute of *A. pullulans*, and also of *A. subglaciale* mainly for the control of fungal pathogens in cold storage conditions, to improve their efficacy, whereas *A. melanogenum* cannot be considered a candidate for biocontrol as it was previously linked to human infections.

Interestingly, our study uncovers the wide potential of biocontrol strains of *Aureobasidium* spp. isolated from a range of different environments not linked to plants, such as car reservoir, clouds, and glaciers of Arctica. This could change the strategy of the search for new, more potent biocontrol strains of already known antagonists.

## Declaration of interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## References

- Aghatabay, N.M., Parali, O., Baydar, C., & Dulger, B. (2014). Synthesis, structural aspects, antimicrobial activity and ion transportation investigation of four new [2 + 2] condensed 24-membered cycloheterphane peptides. *Journal of Inclusion Phenomena and Macrocyclic Chemistry*, 79, 415–424.
- Al-Hindi, R.R., Al-Najada, A.R., & Mohamed, S.A. (2011). Isolation and identification of some fruit spoilage fungi: Screening of plant cell. *African Journal of Microbiology Research*, 5, 443–448.
- Ayers, W.A., Papavizas, G.C., & Diem, A.F. (1966). Polygalacturonate trans-eliminase and polygalacturonase production by *Rhizoctonia solani*. *Phytopathology*, 56, 1006–1011.



- Banani, H., Spadaro, D., Zhang, D., Matic, S., Garibaldi, A., & Gullino, M.L. (2014). Biocontrol activity of an alkaline serine protease from *Aureobasidium pullulans* expressed in *Pichia pastoris* against four postharvest pathogens on apple. *International Journal of Food Microbiology*, 182, 182–183.
- Bae, M., Chung, B., Oh, K. B., Shin, J., & Oh, D.C. (2015). Hormaomycins B and C: New antibiotic cyclic depsipeptides from a marine mudflat-derived *Streptomyces* sp. *Marine Drugs*, 13, 5187–5200.
- Botha, J., Pinard, D., Creux, N., Hussey, S., Maritz Olivier, C., Spokevicius, A., Bossinger, G., Mizrachi, E., & Myburg, A. (2011). *Characterising the role of the Eucalyptus grandis SND2 promoter in secondary cell wall biosynthesis*. Arraiald'Ajuda, Bahia, Brazil: BMC Proceedings BioMed Central Ltd. 105 pp.
- Bozoudi, T., & Tsaltas, T. (2018). The multiple and versatile roles of *Aureobasidium pullulans* in the vitivinicultural sector. *Fermentation*, 4, 85.
- Castoria, R., De Curtis, F., Lima, G., Caputo, L., Pacifico, S., & De Cicco, V. (2001). *Aureobasidium pullulans* (LS-30), an antagonist of postharvest pathogens of fruits: study on its modes of action. *Postharvest Biology and Technology*, 22, 7–17.
- Chan, K.G., Puthucheary, S.D., & Chan, X.Y. (2011). Quorum sensing in *Aeromonas* species isolated from patients in Malaysia. *Current Microbiology*, 62, 167–172.
- Cheng, K.C., Demirci, A., & Catchmark, J. (2011). Pullulan: biosynthesis, production, and applications. *Applied Microbiology and Biotechnology*, 92, 29–44.
- Cimerman, N.G., Cimerman, N., Zalar, P., Hoog, S., & Plemenitaš, A. (2000). Hypersaline waters in salterns-natural ecological niches for halophilic black yeasts. *FEMS Microbiology Ecology*, 32, 235–240.
- Cozar, O., Leopold, N., Jelic, C., Chis, V., David, L., Mocanu, A., & Tomoaia-Cotisel, M. (2006). IR, Raman and surface-enhanced Raman study of desferrioxamine B and its Fe(III) complex, ferrioxamine B. *Journal of Molecular Structure*, 788, 1–6.
- Dahiya, R., Singh, S., Sharma, A., Chennupati, S.V., & Maharaj, S. 2016. First total synthesis and biological screening of a Proline-rich cyclopeptide from a Caribbean marine sponge. *Marine Drugs*, 14, 228.
- de Hoog, G.S. (1993). Evolution of black yeasts: possible adaptation to the human host). *Antonie Van Leeuwenhoek International Journal of General and Molecular Microbiology*, 63, 105–109.
- Di Francesco, A., Di Foggia, M., & Baraldi, E. (2020). *Aureobasidium pullulans* volatile organic compounds as alternative postharvest method to control brown rot of stone fruits. *Food Microbiology*, 87, 103395.
- Di Francesco, A., Calassanzio, M., Ratti, C., Folchi, A., & Baraldi, E. (2018a). Molecular characterization of the two postharvest biological control agents *Aureobasidium pullulans* L1 and L8. *Biological Control*, 123, 53–59.
- Di Francesco, A., Mari, M., Ugolini, L., & Baraldi, E. (2018b). Effect of *Aureobasidium pullulans* strains against *Botrytis cinerea* on kiwifruit during storage and on fruit nutritional composition. *Food Microbiology*, 72, 67–72.
- Di Francesco, A., Roberti, R., & Mari, M. (2018c). Defense response against postharvest pathogens in hot water treated apples *Scientia Horticulturae*, 227, 181–186.
- Di Francesco, A., Milella, F., Roberti, R., & Mari, M. (2017). A preliminary investigation into *Aureobasidium pullulans* as a potential biocontrol agent against *Phytophthora infestans* of tomato. *Biological Control*, 114, 144–149.
- Di Francesco, A., Mari, M. (2014). Use of biocontrol agents in combination with physical and chemical treatments: Efficacy assessment. *Stewart Postharvest Review*, 10, 2.
- Douaiher, M.N., Nowak, E., Durand R., Halama, P., & Reignault, P. (2007). Correlative analysis of *Mycosphaerella graminicola* pathogenicity and cell wall-degrading enzymes produced *in vitro*: the importance of xylanase and polygalacturonase. *Plant Pathology*, 56, 79 – 86.


- Durrands, P.K., & Cooper, R.M. (1988). Selection and characterization of pectinase-deficient mutants of the vascular wilt pathogen *Verticillium albo-atrum*. *Physiological and Molecular Plant Pathology*, 32, 43–362.
- Eckert, J.W., & Ogawa, J.M. (1988). The chemical control of postharvest diseases: deciduous fruits, berries, vegetables and roots/tuber crops. *Annual Review of Phytopathology*, 26, 433–469.
- Eriksson, K.E., & Pettersson, B. (1975). Extracellular enzyme system utilized by the fungus *Sporotrichum pulverulentum* (*Chrysosporium lignorum*) for the breakdown of cellulose 1. Separation purification and physico-chemical characterization of five endo-1,4-p-glucanases. *European Journal of Biochemistry*, 51, 193–206.
- Freimoser, F., Rueda-Mejia, M.P., Tilocca, B., & Migheli, Q. (2019). Biocontrol yeasts: mechanisms and applications. *World Journal of Microbiology and Biotechnology*, 35, 154.
- Gostinčar, C., Turk, M., Zajc, J., & Gunde-Cimerman, N. (2019). Fifty *Aureobasidium pullulans* genomes reveal a recombining polyextremotolerant generalist. *Environmental Microbiology*, 21, 3638–3652.
- Gostinčar, C., Ohm, R.A., Kogej, T., Sonjak, S., Turk, M., Zajc, J., Zalar, P., Grube, M., Sun, H., Han, J., Sharma, A., Chiniquy, J., Ngan, C.Y., Lipzen, A., Barry, K., Grigoriev, I.V., & Gunde-Cimerman, N. (2014). Genome sequencing of four *Aureobasidium pullulans* varieties: biotechnological potential, stress tolerance, and description of new species. *BMC Genomics*, 15, 549.
- Gostinčar, C., Grube, M., & Gunde-Cimerman, N. (2011). Evolution of fungal pathogens in domestic environments? *Fungal Biology*, 115, 1008–1018.
- Higgins, F.S., Magliocco, L.G., & Colthup, N.B. (2006). Infrared and Raman spectroscopy study of alkyl hydroxamic acid and alkyl hydroxamate isomers. *Applied Spectroscopy*, 60, 279–287.
- Holmen, B.A., Tejedor-Tejedor, M.I., & Casey, W.H. (1997). Hydroxamate complexes in solution and at Goethite-water interface : a cylindrical internal reflection Fourier transform infrared spectroscopy study. *Langmuir*, 13, 2197–2206.
- Ippolito, A., El Ghaouth, C.L., Wilson, M., & Wisniewski, M. (2000). Control of postharvest decay of apple fruit by *Aureobasidium pullulans* and induction of defence responses. *Postharvest Biology and Technology*, 19, 265–272.
- Janisiewicz, W.J., Workoski, T.J., & Sharer, C. (2000). Characterizing the mechanism of biological control of postharvest diseases on fruits with a simple method to study competition for nutrients. *Phytopathology*, 90, 1196–1200.
- Kachalkin, V.A. (2010). New Data on the Distribution of Certain Psychrophilic Yeasts in Moscow Oblast. *Microbiology*, 79, 840–844.
- Kikot, G.E., Hours, R.A., & Alconada, T.M. (2009.) Contribution of cell wall degrading enzymes to pathogenesis of *Fusarium graminearum*: a review. *Journal of Basic Microbiology*, 49, 231–241.
- Klein, M.N., & Kupper, K.C. (2018). Biofilm production by *Aureobasidium pullulans* improves biocontrol against sour rot in citrus. *Food Microbiology*, 69, 1–10.
- Kohl, J., Kolnaar, R., & Ravensberg, W.J. (2019). Mode of action of microbial biological control agents against plant diseases: relevance beyond efficacy. *Frontiers in Plant Science*, 10, 845.
- Liu, T.T., Zhu, L.Y., Zhang, Z.P., Huang, H., Zhang, Z.& D., Jiang, L. (2017) Protective role of trehalose during radiation and heavy metal stress in *Aureobasidium subglaciale* F134. *Scientific Report*, 7, 1–9.
- Madhupani, D.S., & Adikaram, N.K.D. (2017). Delayed incidence of stem-end rot and enhanced defences in *Aureobasidium pullulans*-treated avocado (*Persea americana* Mill.) fruit. *Journal of Plant Diseases and Protection -New Series*, 12.

- Mari, M., Martini, M., Spadoni, A., Roussi, W., & Bertolini, P. (2012). Biocontrol of apple postharvest decay by *Aureobasidium pullulans*. *Postharvest Biology and Technology*, 73, 56–62.
- Mari, M., Guizzardi, M., Bruennli, M., & Folchi, A. (1996). Postharvest biological control of grey mould (*Botrytis cinerea* pers.: Fr.) on fresh-market tomatoes with *Bacillus amyloliquefaciens*. *Crop Protection*, 1, 699–705.
- Molnorova, J., Vadkertiová, R., & Stratilová, E. (2013). Extracellular enzymatic activities and physiological profiles of yeasts colonizing fruit trees. *Journal of Basic Microbiology*, 54, (S1).
- Nigam, S.P. (2013). Microbial Enzymes with special characteristics for biotechnological applications. *Biomolecules*, 3, 597–611.
- Parafati, L., Vitale, A., Restuccia, C., & Cirvilleri, G. (2015). Biocontrol ability and action mechanism of food-isolated yeast strains against *Botrytis cinerea* causing post-harvest bunch rot of table grape. *Food Microbiology*, 47, 85–92.
- Parafati, L., Vitale, A., Restuccia, C., & Cirvilleri, G. (2017). Performance evaluation of Volatile Organic Compounds by antagonistic yeasts immobilized on hydrogel spheres against grey, green and blue post-harvest decays. *Food Microbiology*, 63, 191–198.
- Peterson, S.W., Manitchotpisit, P., & Leathers, T.D. (2013). *Aureobasidium thailandense* sp. nov. isolated from leaves and wooden surfaces. *International Journal of Systematic and Evolutionary Microbiology*, 63, 790–795.
- Pierce, J.A., & Rast, D.M. (1995). A comparison of native and synthetic mushroom melanins by Fourier-transform infrared spectroscopy. *Phytochemistry*, 39, 49–55.
- Prasongsuk, S., Lotrakul, P., Ali, I., Bankeeree, W., & Punnapayak, H. (2018). The current status of *Aureobasidium pullulans* in biotechnology. *Folia Microbiologica*, 63, 129–140.
- Price, N.P.J., Manitchotpisit, P., Vermillion, K.E., Bowman, M.J., & Leathers, T.D. (2013). Structural characterization of novel extracellular liamocins (mannitol oils) produced by *Aureobasidium pullulans* strain NRRL 50380. *Carbohydrate Research*, 370, 24–32.
- Price, N.P.J., Bischoff, K.M., Leathers, T.D., Cossé, A.A., & Manitchotpisit, P. (2017). Polyols, not sugars, determine the structural diversity of anti-streptococcal liamocins produced by *Aureobasidium pullulans* strain NRRL 50380. *The Journal of Antibiotics*, 70, 136–141.
- Punnapayak, H., Sudhadham, M., Prasongsuk, S., & Pichayangkura, S. (2003). Characterization of *Aureobasidium pullulans* isolated from airborne spores in Thailand. *Journal of Indian Microbiology and Biotechnology*, 30, 89–94.
- Rekha, C.P., & Chandra, P.S. (2007). Pullulan as a promising biomaterial for biomedical applications: a perspective. *Trends in Biomaterial and Artificial Organs*, 20.
- Sasahara, H., & Izumori, K. (2005). Production of L-sorbitol from L-fructose by *Aureobasidium pullulans* LP23 isolated from soy sauce mash. *Journal of Bioscience and Bioengineering*, 100, 223–226.
- Schena, L., Nigro F., Pentimone, I., Ligorio, A., & Ippolito, A. (2003). Control of postharvest rots of sweet cherries and table grapes with endophytic isolates of *Aureobasidium pullulans*. *Postharvest Biology and Technology*, 30, 209–220.
- Sharma, R.R., Singh, D., & Singh, R. (2009). Biological control of postharvest diseases of fruits and vegetables by microbial antagonists: a review. *Biological Control*, 50, 205–221.
- Shingel, K.I. (2002). Determination of structural peculiarities of dextran, pullulan and g-irradiated pullulan by Fourier-transform IR spectroscopy. *Carbohydrate Research*, 337, 1445–1451.
- Singh, R.S., Saini, G.K., & Kennedy, J.F. (2008). Pullulan: microbial sources, production and applications. *Carbohydrate Polymer*, 73, 515–531.
- St Leger, R.J., St Leger, L., Joshi, D., & Roberts, W. (1997). Adaptation of proteases and carbohydrases of saprophytic: phytopathogenic and entomopathogenic fungi to the requirements of their ecological niches. *Microbiology*, 143, 1983–1992.




- Takesako, K., Ikai, K., Haruna, F., Endo, M., Shimanaka, K., Sono, E., Nakamura, T., Kato, I., & Yamaguchi, H. (1991). Aureobasidins, new antifungal antibiotics. Taxonomy, fermentation, isolation. *Journal of Antibiotics*, 44, 919–924.
- Tu, A.T. (1982). Raman spectroscopy in biology: principles and applications. In *Chapter 3: Peptides and Proteins; Chapter 4: Enzymes and Immunoglobulins; Chapter 8: Carbohydrates*. New York, USA: John Wiley & Sons.
- Um, S., Choi, T.J., Kim, H., Kim, B.Y., Kim, S.H., Lee, S.K., Oh, K.B., Shin, J., & Oh, D.C. (2013). Ohmyungsamycins A and B: cytotoxic and antimicrobial cyclic peptides produced by *Streptomyces* sp. from a volcanic island. *Journal of Organic Chemistry*, 78, 12321–12329.
- Urzi, C., Leo, F.D., Passo, C.L., Criseo, G. (1999). Intra-specific diversity of *Aureobasidium pullulans* strains isolated from rocks and other habitats assessed by physiological methods and by random amplified polymorphic DNA (RAPD). *Journal of Microbiological Methods*, 36, 95–105.
- Wang, W., Chi, Z., Liu, G., Buzdar, M.A., Chi, Z., & Gu, Q. (2009). Chemical and biological characterization of siderophore produced by the marine-derived *Aureobasidium pullulans* HN6.2 and its antibacterial activity. *Biometals*, 22, 965–972.
- Whipps, J.M. (2001). Microbial interactions and biocontrol in the rhizosphere. *Journal of Experimental Botany*, 52, 487–511.
- Jiang, H. (2017). Both a PKS and a PPTase are involved in melanin biosynthesis and regulation of *Aureobasidium melanogenum* XJ5-1 isolated from the Taklimakan desert. *Gene*, 602, 8–15.
- Yang, J., Bremer, P.J., Lamont, I.L., & McQuillan, A.J. (2006). Infrared spectroscopic studies of siderophore-related hydroxamic acid ligands adsorbed on titanium dioxide. *Langmuir*, 22, 10109–10117.
- Zain, M.E., Awaad, A.S., Razak, A.A., Maitland, D.J., Khamis, N.E., & Sakhawy, M.A. (2009). Secondary metabolites of *Aureobasidium pullulans* isolated from Egyptian soil and their biological activity. *Journal of Applied Sciences Research*, 5, 1582–1591.
- Zajc, J., Gostinčar, C., Ravnikar, M., & Gunde-Cimerman, N. (2018). Polyextremotolerant yeasts *Aureobasidium* spp.: antagonistic activity against phytopathogenic fungi. IOBC-WPRS working group "Biological and integrated control of plant pathogens - Proceedings of the xv meeting on Biocontrol products: from lab testing to product development" at Lleida (Spain) 133, 177–178.
- Zajc, J., Gostinčar, C., Cernoša, A., & Gunde-Cimerman, N. (2019). Stress-tolerant yeasts: Opportunistic pathogenicity versus biocontrol potential. *Genes*, 10, 42.
- Zajc, J., Černoša, A., Di Francesco, A., Castoria, A., De Curtis, F., Lima, G., Badri, H., Jijakli, H., Ippolito, A., Gostinčar, C., Zalar, C.P., Gunde-Cimerman, N., & Janisiewicz, W. J. (2020). Variability of characteristics of *Aureobasidium pullulans* isolates exhibiting biocontrol activity against fruit decay pathogens. *Fungal Genomics and Biology*, 10, 1–13.
- Zalar, P., Gostinčar, C., de Hoog, G.S., Uršič, V., Sudhadham, M., & Gunde-Cimerman, N. (2008). Redefinition of *Aureobasidium pullulans* and its varieties. *Studies in Mycology*, 61, 21–38.
- Zhang, D., Spadaro, D., Garibaldi, A., & Gullino, M.L. (2010). Efficacy of the antagonist *Aureobasidium pullulans* PL5 against postharvest pathogens of peach, apple and plum and its modes of action. *Biological Control*, 54, 172–180.
- Zhbankov, R.G., Firsov, S.P., Korolik, E.V., Petrov, P.T., Lapkovski, M.P., Tsarenkov, V.M., Marchewka, M.K., & Ratajczak, H. (2000). Vibrational spectra and the structure of medical biopolymers. *Journal of Molecular Structure*, 555, 85–96.

**Table 1** List of strains used in this study.


Species	Culture collection strain number	Present study number	Isolation habitat	Sampling site location
<i>Aureobasidium pullulans</i>	EXF-6519	AP1	felt of a metal roof tile	Slovenia (Mengeš)
	EXF-10507	AP2	marble block surface	Italy (Messina)
	EXF-10629	AP3	car petrol reservoir	Slovenia (Jezero)
	EXF-10650	AP4	acrylic painting	Slovenia (Solkan)
	EXF-10751	AP5	cloud sample	France
<i>Aureobasidium melanogenum</i>	EXF-3378 / CBS 110374	AM6	public fountain	Thailand
	EXF-3397	AM7	endoperitoneal fluid	Greece
	EXF-8016	AM8	bathroom, between faucet and sink	nd
	EXF-8429	AM9	tap water	Slovenia
	EXF-11028	AM10	aquarium water	Slovenia, Ljubljana
<i>Aureobasidium subglaciale</i>	EXF-2481 / CBS 123387	AS11	subglacial ice	Arctic; Svalbard, NyAlesund
	EXF-4632	AS12	leaves of Convallaria	Slovenia
	EXF-2425	AS13	subglacial ice	Arctic; Svalbard, NyAlesund
	EXF-2428	AS14	subglacial ice	Arctic; Svalbard, NyAlesund
	EXF-2450	AS15	glacial ice	Arctic; Svalbard, NyAlesund



Percentage of inhibition of *Botrytis cinerea* mycelial growth co-cultured with *Aureobasidium pullulans*, *A. melanogenum* and *A. subglaciale* for 5 days at 25°C.




Xylanase (XYL), pectinase (PEC) and polygalacturonase (POL) activity (measured as area of cleared halo in mm<sup>2</sup>) of *Botrytis cinerea* conidia suspension (10<sup>5</sup> conidia mL<sup>-1</sup>) after exposure to secondary metabolites (No-VOCs) of *A. pullulans*, *A. melanogenum* and *A. subglaciale* strains (6 hours of incubation at 25°C) or to distilled water (control). Data are the means of two independent experiments ± standard deviation, each consisting of five replicates per treatment. Data were analysed using one-way ANOVA; Fisher's Least Significant Differences (LSDPEC), and polygalacturonase (POL) activity (measured as area of cleared halo in mm<sup>2</sup>) Test at  $P < 0.05$  was used to separate differences among the means.



Difference FT-IR spectra of *A. pullulans* (AP5), *A. melanogenum* (AM6) and *A. subglaciale* (AS15) strains after 48 h of incubation and the beginning (1 h). Positive peaks are related to the production of non-volatile compounds discussed in the text. Negative peaks are usually related to the consumption of the culture medium and are highlighted by the # symbol. The blue rectangles indicate the spectral regions in which AM6 spectrum differs the most from both AP5 and AS15.





*In vivo* antagonistic effect by *Aureobasidium pullulans*, *A. melanogenum* and *A. subglaciale* against *Botrytis cinerea* of tomato. 'Datterini' were artificially inoculated with antagonists' cells mL<sup>-1</sup> suspensions (15 .L, cells and after 1 hour with conidial suspension (15 .L, conidia mL<sup>-1</sup>) of *B. cinerea* and incubated for 6 days at 20°C and 90% RH. Tomato inoculated with water and *B. cinerea* suspension was considered as a control treatment. Each value is the mean of three replicates (60 fruits) ± standard deviation. Different letters represent significant differences among the strains treatments according to Fisher's Least Significant Difference (LSD) Test at  $P < 0.05$ .