



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Human adaptation to changing coastal landscapes in the Eastern Adriatic: Evidence from Vela Spila cave, Croatia

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Dean, S., Pappalardo, M., Boschian, G., Spada, G., Forenbaher, S., Juračić, M., et al. (2020). Human adaptation to changing coastal landscapes in the Eastern Adriatic: Evidence from Vela Spila cave, Croatia. QUATERNARY SCIENCE REVIEWS, 244, 1-18 [10.1016/j.quascirev.2020.106503].

Availability:

This version is available at: <https://hdl.handle.net/11585/771653> since: 2022-02-14

Published:

DOI: <http://doi.org/10.1016/j.quascirev.2020.106503>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Silas Dean, Marta Pappalardo, Giovanni Boschian, Giorgio Spada, Stašo Forenbaher, Mladen Juračić, Igor Felja, Dinko Radić, Preston T. Miracle, *Human adaptation to changing coastal landscapes in the Eastern Adriatic: Evidence from Vela Spila cave, Croatia*, Quaternary Science Reviews, Volume 244, 2020, 106503.

The final published version is available online at:
<https://doi.org/10.1016/j.quascirev.2020.106503>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Human adaptation to changing coastal landscapes in the Eastern Adriatic: Evidence from Vela Spila cave, Croatia

Silas Dean¹, Marta Pappalardo^{1*}, Giovanni Boschian², Giorgio Spada³, Stašo Forenbaher⁴, Mladen Juračić⁵,
Igor Felja⁶, Dinko Radić⁷ Preston T. Miracle⁸

¹Department of Earth Sciences, University of Pisa, Italy

²Department of Biology, University of Pisa, Italy

³Dipartimento di Scienze Pure e Applicate, Università di Urbino ‘Carlo Bo’, Urbino, Italy

⁴Institute for Anthropological Research, Zagreb, Croatia

⁵Croatian Academy of Sciences and Arts, Croatia

⁶University of Zagreb, Faculty of Science, Dept. of Geology, Croatia

⁷Centar za kulturu, Vela Luka, Croatia

⁸Department of Archaeology, University of Cambridge, UK

*Corresponding author

KEYWORDS: Sea-level change; Environmental change; Human settlement patterns, Bathymetry; Prehistoric archaeology; Late Pleistocene-Holocene; Central Adriatic coast; Croatia

Abstract

In this paper new palaeogeographic and archaeological data from the prehistoric cave Vela Spila on the island of Korčula in Croatia are combined with new realizations of two glacial isostatic adjustment models in order to present relative sea-level change scenarios confronting the inhabitants of the cave at different time slices and to show how they experienced and adapted to

23 sea-level and climate change from the Late Pleistocene through the Holocene. Our results show that
24 from the Late Upper Palaeolithic until the Mesolithic, humans in the study area would have
25 experienced tens of metres of sea-level rise, at rates in some cases up to 12 mm per year, and, owing
26 to the relatively flat morphology of the now submerged plains, hundreds of meters of horizontal
27 coastline change in the plains to the north and south of the island. This evidence supports the
28 hypothesis that the rapid loss of these plains likely contributed to the human abandonment of the
29 cave after the Palaeolithic for about five thousand years, followed by significant changes in lifestyle
30 and diet in the Mesolithic. Our results have important implications for the study of how past human
31 groups, especially in vulnerable coastal areas, were affected by sea level, climate, and other
32 environmental changes. Vela Spila represents a case study of how changing environment and rising
33 seas can force significant alterations in human societies, even when there is no risk of inundation to
34 the settlement itself.

35 **1 Introduction**

36 Since prehistory, humans have adapted their settlement and mobility patterns in response to
37 environmental change (Blute, 2008; Bini et al., 2018; Butzer, 2005), using their decision-making
38 skills (Nunn, 2020), or even engineering skills (Galili et al., 2019) to overcome challenges created
39 by modifications to coastlines and their related ecosystems. The Eastern Adriatic region represents a
40 key area to investigate the relationship between environmental changes and cultural responses. The
41 ‘Great Adriatic Plain’ periodically exposed during the Pleistocene provided suitable conditions for
42 human settlement; its loss to rising sea level during the Late Pleistocene and Early Holocene
43 precipitated a significant restructuring of plant, animal, and human communities (Boschian and
44 Fusco, 2007; Miracle, 2007). The palaeoecological and archaeological evidence of the region is
45 reviewed by Pilaar Birch and Linden (2018).

46 Sea-level change has been a major cause of environmental change; it has modified coastal
47 landscapes by dramatically and cyclically altering coastal plains around shallow sea basins, and by
48 influencing the distribution of islands and their distance from the mainland. The outline and area of
49 many present-day islands were different during the Late Pleistocene owing to the last ice age and
50 the relative sea-level (RSL) changes associated with it; the islands' current isolation is unusual
51 during the last 70 ka (Weigelt et al., 2016). Consequently, the distribution of human and animal
52 communities in coastal areas (Carroll et al., 2012; Garg et al., 2018; Hewitt, 2000; McLaughlin et
53 al., 2018) was profoundly affected, resulting in major behavioural and cultural change. A robust
54 reconstruction of shoreline position at different time slices of the Late Pleistocene and Holocene is
55 essential for understanding human settlement and animal dispersal patterns.

56 Following the seminal work by van Andel (1989), reconstructing ancient coastlines in order to
57 elucidate their influence on hominin dispersal and cultural behaviour became an important research
58 goal. In archaeological literature, the reconstruction of ancient coastal landscapes has focused on
59 comparing present-day bathymetric surfaces with global eustatic curves (Dobson, 2014). However,
60 past RSL can be reconstructed more precisely by combining geophysical and geological techniques
61 (Pirazzoli, 2013; De Boer et al., 2014; Shennan et al., 2015).

62 Sea-level reconstructions must be combined with bathymetric data, which have also been widely
63 used to identify landforms, assess palaeoenvironments and palaeoshorelines, and address
64 archaeological questions in the central Mediterranean. Bathymetric mapping and archaeological
65 research have been combined in Sardinia (Palombo et al., 2017) for this purpose. In the Tyrrhenian
66 Sea west of Italy, seismic profiles have been used to detect terraces that may be associated with
67 ancient sea stands (Casalbore et al., 2017). Extensive single- and multibeam data were used
68 (Trincardi et al., 2014) to identify several drowned landforms in the western Adriatic. Bathymetric
69 data from Malta (Foglini et al., 2016) have shown drowned karstic features and river landforms, and
70 can answer questions about prehistoric fauna migration and human settlement. In the NW Adriatic,

71 prehistoric settlement surfaces and lagoon deposits have contributed to landform and RSL
72 reconstructions (Fontana et al., 2017). In the eastern coast of the Adriatic dozens of submerged
73 coastal prehistoric sites have been located, from Istria to southern Dalmatia, and excavations have
74 been initiated on several of those sites (Bekić, 2017; Bekić et al., 2015; Benjamin et al., 2011;
75 Benjamin and Bonsall, 2009; Čelhar et al., 2017; Koncani Uhač and Čuka, 2015; Radić Rossi,
76 2008). The majority of the sites belong to relatively recent prehistoric times (Bronze Age and Iron
77 Age), while only very few of them belong to Copper Age or Neolithic periods. Further excavation,
78 interdisciplinary investigation (collaboration of archaeologists and geologists), and analyses are
79 required before they can be used as accurate indicators of local sea-level change in prehistoric times
80 (Rossi, 2017)

81 This study integrates palaeoenvironmental and archaeological evidence from Vela Spila, a cave
82 situated on the island of Korčula, close to the central Dalmatian coast in Croatia (Fig 1) with a new
83 synthesis of bathymetric datasets. We Palaeogeographic reconstructions were created at time slices
84 relevant for key cultural events during cave occupation phases from the Late Upper Palaeolithic to
85 the Neolithic. The main goal of this work is to investigate the relationship between the shoreline
86 position and the cave. The results provide an understanding of how inhabitants of the cave
87 experienced and adapted to sea-level and climate changes since the Late Pleistocene.

88 This study incorporates both existing and new data and approaches that are discussed in light of
89 current research. In particular:

90 1) Novel archaeological evidence provided in this study compliments previous work on Vela Spila
91 (Branscombe et al., 2020; Cristiani et al., 2014; Farbstein et al., 2012; Radić, 2018; Rainsford et al.,
92 2014). Vela Spila (Fig 2) is one of the richest and most promising archaeological sites in the Eastern
93 Adriatic region, with a Late Pleistocene to Holocene stratigraphic sequence spanning the Late
94 Upper Palaeolithic (LUP) to the Bronze Age. The deposits include exceptionally well-preserved
95 cultural, human, animal, and vegetal remains that document important environmental and biological

96 changes during the Pleistocene-Holocene, and cultural changes during the Palaeolithic-Mesolithic
97 and Mesolithic-Neolithic transitions.

98 2) Data used in the current study to reconstruct bathymetry around Vela Spila represent a synthesis
99 of available bathymetric datasets for the area, and thus an improvement on the datasets used by
100 previous authors such as Sikora et al. (2014), which used satellite bathymetric data combined with
101 GIS analysis to reconstruct river drainage during the lowstand of the Last Glacial Maximum
102 (LGM). Bathymetric data, as a source of palaeolandscape data, are applicable to our study area
103 because of the karstic nature of the Eastern Adriatic - rainwater sinks into the endokarst instead of
104 creating high sediment flux onto the seabed (Pikelj and Juračić, 2013). Consequently, LGM
105 landforms and topography are still visible in bathymetric data, aside from limited fluvial
106 sedimentation.

107 3) Bathymetric data for the seabed around Korčula are combined with two updated geophysical
108 models of glacial-hydro-isostatic adjustment (GIA) realized specifically for the area under
109 investigation in order to reconstruct palaeogeographic phases; all previous authors used in the best
110 case older, less refined models, and Sikora (2014) used a global model. This implies differences of
111 tens of kilometres in the position of the shoreline at specific times. The use of GIA models is valid
112 for this subregion of the Eastern Adriatic because tidal notches documented by Faivre and Butorac
113 (2018) show relatively slower rates of subsidence here than in areas to the north, at least for the
114 Late Holocene.

115 Vela Spila represents a relevant case study on how changing environment and rising seas can force
116 significant changes in human societies, even when there is no direct risk of inundation for the
117 settlement itself.

118 **2 Regional setting**

119 **2.1 Adriatic sea**

120 The Adriatic Sea (Fig 1) is a NW-SE elongated basin of approximately 800 by 200 km that
121 separates the Italian peninsula from the Balkans (Trincardi et al., 2014). This basin can be divided
122 into three sub-basins: the northern, which is 35 m deep on average, the middle with an average
123 depth of 140 m, and the southern with maximum depths reaching 1200 m (Arteghiani et al., 1997).
124 This paper is focused on the island of Korčula, which lies on the eastern coast between the middle
125 and southern basins. The Eastern Adriatic bedrock is about 8,000 m of limestone and/or dolomite
126 (Vlahović et al., 2005, 2002) formed in shallow sedimentation basins between the Mid-Late
127 Triassic, Mesozoic and Palaeogene, followed by relatively thin Eocene flysch outcrops. The
128 tectonic deformation pattern of the Eastern Adriatic is characterised by a complex system of faults,
129 thrust faults and overthrusts formed since the Miocene and developed along the so-called
130 Dinaric Strike with NW-SE orientation (Vlahović et al., 2005, 2002). The typical relief pattern of
131 mountain and hill ridges parallel to the coastline (Fig 1) derives from this tectonic compressional
132 motif, followed by extensional processes and eventually repeated flooding during the Quaternary
133 sea-level fluctuations. The westernmost Dinarides were submerged by the postglacial RSL rise
134 (Kelleter, 2005) and now form the Croatian islands and the typically rocky Dalmatian Coast (Felja
135 et al., 2015; Pikelj and Juračić, 2013). Significantly for this study area, the Eastern Adriatic coast
136 and its hinterland are karstic and dominated by typical karst landforms, with few significant rivers
137 discharging sediments into the sea, so that the present-day bathymetry closely reflects LGM
138 landforms (Pikelj and Juračić, 2013). This is confirmed in the immediate study area by research
139 (Giglio et al., 2020) in the waters around the present-day Neretva delta, which shows river
140 sedimentation rates quickly approaching 0 mm/a (mm per year) within 10 km offshore. This aspect

141 of the coast contrasts with the wide and flat morphology of the north-eastern Adriatic, which was an
142 alluvial plain dominated by sandy river discharge when the LGM RSL was 120 m below present
143 (Trincardi et al., 2014). Though these submerged morphological features in the western Adriatic
144 have been identified by multibeam imagery (Trincardi et al., 2014), including drowned karst
145 landscapes (Taviani et al., 2012), public high-resolution data are less available for the rocky Eastern
146 Adriatic.

147 **2.2 Central-southern Dalmatia**

148 Korčula is an island in central-southern Dalmatia, separated from the Pelješac peninsula by a
149 narrow channel about 40-45 m deep. Like most Dalmatian islands (Kelletat, 2005), it represents the
150 top of a submerged ridge formed by tectonic overthrusting and uplift of reliefs which, in the case of
151 the southern islands, are rotated counterclockwise with reference to the dinaric strike. The island
152 landscape is rugged, controlled by the nature of the underlying Mesozoic limestones and dolomitic
153 limestones, and also characterised by several deep and elongated *polje* oriented along the structural
154 motif and by deep karstic depressions, whose bottoms are filled by *terra rossa/crvenica* soils
155 (Bonacci et al., 2012). The coasts are almost always steep and rocky. The channel between Korčula
156 and Hvar -the island immediately to the north – is at most about 70 m deep and represents the
157 prolongation of the Neretva river valley towards the Adriatic basin (Sikora et al., 2014). On the
158 westernmost tip of the island, a narrow, winding and shallow inlet protects the Vela Luka harbour.
159 In the now submerged zones between islands, areas at greater depth closer to the LGM lowstand of
160 -120 m (Peltier and Fairbanks, 2006) or -115 m (Sikora et al., 2014) had less exposure time to
161 erosion and other landscape forming processes than areas situated near highstand levels. As a result,
162 most morphology of these submerged areas consist of broad, flat areas; a fact often compounded by
163 the presence of alluvial sediments in areas near palaeo rivers.

164 **2.3 Vela Spila**

165 Vela Spila (Fig. 2) is a large cave situated 121 m above sea level on the western side of Spilski Rat,
166 a low hill overlooking Vela Luka harbour. The cave's single, semi-ellipsoidal chamber, about 50 m
167 by 30 m wide and 17 m high, opens in Cenomanian radiolaritic limestone whose layers dip
168 southwards (Cristiani et al., 2014). The whole inside surface of about 1000 m² is at present always
169 well-lit through a wide entrance facing west and two large holes in its ceiling.

170 The outside landscape has been strongly affected by human activities over long time scales,
171 probably since the cave was first inhabited during the Late Glacial. Most hillsides have extensive
172 19th century terraces for viticulture or growing olive trees, though many are now abandoned. Good
173 quality data about outside soils and their evolution through the end of the Late Glacial to Middle
174 Holocene are difficult to collect in these conditions: the typical features are thick stone walls
175 sustaining *terra rossa* backfills, alternating with patches of barren denuded limestone. Cave data are
176 in this case the best way to acquire information about climate and environment evolution in the area
177 since they are undisturbed by agricultural activity during the last two millennia.

178 **2.4 Regional relative sea level**

179 Global eustatic sea level has risen roughly 120 m since the LGM (Fairbanks, 1989; Peltier and
180 Fairbanks, 2006). GIA models (Lambeck and Purcell, 2005; Peltier, 2004; Peltier et al., 2015) have
181 been used to estimate timing and magnitude of these changes. Datable biological, geological and
182 archaeological proxies have been used to calibrate these models and provide more precise
183 chronological indications of past RSL (Rovere et al., 2016).

184 In the Eastern Adriatic area, fossil remains of *Lithophyllum sp.* and an ancient harbour (Faivre et al.,
185 2013; Vacchi et al., 2016) provide sea-level indicators in close proximity to the study area. Salt
186 marsh foraminifera have also been studied (Shaw et al., 2018) to reconstruct RSL in the last
187 millenium, and suggest a subsidence rate of 0.45 +/- 0.6 mm/a in Jadrtovac, ~ 100 km northwest of

188 Korčula. A dataset of speleothem dates from submerged caves exists (Juračić and Surić, 2010) but
189 most of these samples predate the LGM; a specimen from ca. 9.2 cal ka BP is relevant to the current
190 study, though these indicators constitute only a maximum possible elevation for RSL. Vacchi et al.
191 (2016) used the ICE-5G (VM2) GIA model (Peltier, 2004) of the last 15 ka to predict a RSL rise of
192 less than 1 m since 4 ka BP, preceded by a much faster increase. However, they also noted that the
193 sea-level points indicate an RSL approximately 1 m lower than the model during the last 3 ka, and
194 hypothesize a general subsidence trend in the wider area. The observational data in this study area
195 (Faivre et al., 2013; Vacchi et al., 2016), however, were mostly limited to the last 3 ka BP.
196 Unfortunately for central-southern Dalmatia no valuable index points from coastal depressions are
197 available to constrain relative sea level, like the ones obtained by Brunović et al. (2019) for the
198 north-eastern Adriatic. In this area Antonioli et al. (2007) applied a methodology similar to the one
199 used in this work using a GIA model by Lambeck & Purcell (2005) that predicts roughly similar
200 changes for the last 12 ka in Istria (north-eastern Adriatic), but with the stabilizing point occurring
201 at ca. 6 ka BP instead of 4 ka BP.

202 **3 Materials and methods**

203 **3.1 Bathymetric data**

204 This study utilizes a combination of publicly available bathymetric data sources. These include
205 digitized nautical charts, and SRTM15_PLUS data from NOAA ERDDAP (Sandwell et al., 2014a),
206 which is a 15'' version of the Shuttle Radar Topography Mission (SRTM) dataset containing
207 bathymetric data from a combination of sources, chiefly from satellite altimetry gravity elevations
208 (Smith and Sandwell, 1997). The bathymetric data have been combined with topographic digital
209 elevation data of 1'' resolution (~30 m) from the SRTM. Table 1 lists these datasets and their
210 characteristics.

DATASET	Projection	Horizontal Datum	Vertical Datum	Resolution	Type	Citation
NASA SRTM Shuttle radar	Geographic	WGS84	EGM96	1'' or ~22 m at latitude (emerged areas only)	Raster (with vector coastline)	(NASA, 2000)
SRTM15_PLUS	Geographic	WGS84	Unspecified	15'' or ~340 m at latitude	Raster	(Olson et al., 2014, 2016; Sandwell et al., 2014a)
Chart soundings	Geographic	Lat/Lon	Presumed to be mareograph derived lowest astronomical tide	Varies: 10^1 or 10^2 m distance between points in harbours or channels. 10^3 m distances in deep seas	Vector	(Croatian Geoportal, 2019; Državni Hidrografski Institut, 1998)

Non homogeneous

211 *Table 1. Overview of the bathymetric and topographic datasets.*

212 The bathymetric and nautical sounding datasets were combined into a single point cloud using
 213 QGIS so that the high resolution SRTM 1'' and nautical chart data supersedes the SRTM15_PLUS
 214 15'' data in emerged areas, and a raster interpolation was created using the GRASS GIS algorithm
 215 v.surf.rst. Satellite bathymetry, with its high vertical resolution but poor horizontal resolution, was
 216 also eliminated within ~500 m of all coastlines to favour the more accurate nautical chart soundings
 217 available in those areas.

218 **3.2 Uncertainties**

219 The vertical datum for the Croatian chart soundings used in this DEM are based on HRSDM71, i.e.
 220 Mean Lower Low Waters of Spring Tide from tide gauges at Dubrovnik, Split, Bakar, Rovinj and
 221 Koper (epoch 1971.5) (Domijan et al., 2005; HHI, 2019). A ± 0.5 m is a cautious uncertainty for
 222 these whole-number elevations. The satellite altimetry data is heavily post processed to remove
 223 wave noise (Sandwell et al., 2014b; Smith and Sandwell, 1997), and come from electromagnetic
 224 pulse return times between satellites and the ocean surface at time of sampling, but uncertainties

225 surrounding the geoid height, GPS technologies, and the ship soundings used for calibration make
226 calculating an overall vertical uncertainty difficult. See Tozen et al. (2019) for further discussion of
227 error distributions. In any case, the ~0.5 km spatial resolution of the satellite derived bathymetry in
228 the SRTM15 dataset and the heterogenous spatial resolution and unspecified XY precision of
229 nautical chart soundings mean that XY errors far outweigh the vertical Z uncertainties created by
230 harmonizing the datasets used in the DEM in this area where the largest recorded tidal ranges are
231 less than 0.5 m (tide-forecast.com, 2019). The present study excludes datapoints within 1 km of
232 coastlines, and gives precedence to chart soundings when they conflict with the satellite data. We
233 The SRTM15_PLUS dataset in the study area was also validated with the chart soundings and it
234 was noted that in most deep areas (~100 m) the predicted depths correspond well (less than 10 m
235 disparity) with available chart soundings.

236 **3.3 GIA modelling**

237 For this work, new realizations of GIA models calculated specifically for Korčula were used.
238 Previous regional studies based on an GIA models (Lambeck and Purcell, 2005), have used
239 calculations of GIA for Istria (Antonioli et al., 2007) and Vis (Vacchi et al., 2016) but these span
240 only the last 12-14 ka BP, while Vela Spila's archaeological record goes back to ca. 20 ka BP. The
241 GIA results were obtained by solving the sea-level equation with the program SELEN developed by
242 Spada and Stocchi (2007). In its current implementation, SELEN (version 4) accounts for the
243 migration of shorelines and rotational effects on sea level (Spada and Galassi, 2017). The
244 realizations derive from two different GIA models, based on different assumptions about the
245 melting chronologies of Late Pleistocene ice sheets, and the viscosity profile of the mantle. The first
246 model is the ICE-6G_C (VM5a) of Peltier et al. (2015) while the second has been progressively
247 developed by K. Lambeck and co-workers at the Australian National University (ANU) (for more
248 information, see e.g., Lambeck et al. (2003) and subsequent refinements). These two realizations

249 are henceforth referred to as I6G_S4 and ANU_S4, where S4 is an abbreviation for SELEN (version
250 4). Since calculating and illustrating an uncertainty for a GIA model is difficult owing to the large
251 number of variables, assumptions, and parameters that comprise the final prediction (Melini and
252 Spada, 2019), having two models allows us to use the difference between them as an envelope of
253 uncertainty. The RSL reconstructions, the bathymetric digital elevation map, and the archaeological
254 stratigraphy were analysed together in order to reconstruct the palaeogeographic and environmental
255 conditions during the relevant archaeological phases. A literature review for sea-level indicators in
256 the region was also undertaken in order to provide verification for the models.

257 **3.4 Archaeological data**

258 The most relevant palaeoenvironmental aspects inferred from the observation of the Vela Spila
259 sediments have been summarised from published work (Branscombe et al., 2020; Čečuk and Radić,
260 2005; Cristiani et al., 2014; Farbstein et al., 2012; Rainsford et al., 2014) and complimented with
261 unpublished observations made in the four cave key trenches (Fig 3) during the systematic
262 excavations carried out since 2010. Chronology used in this study is partly derived from published
263 radiocarbon dates collected (Table 2) from previously unpublished dates and published sources
264 (Čečuk and Radić, 2005; Farbstein et al., 2012; Rainsford et al., 2014; Srdoč et al., 1989), and
265 partly from the typology of the archaeological remains, the chronology of which is assessed
266 confidently in Dalmatia (Kaczanowska and Kozłowski, 2018; Komšo, 2008; Kozłowski and
267 Kaczanowska, 2004; Miracle and Forenbaher, 2005). In this study, dates referring to specific C¹⁴
268 results use cal a or cal ka BP, whereas dates specifying general periods, GIA models, archaeological
269 typology, or other methods use a or ka BP.

270 4 Results

271 4.1 Palaeoenvironmental evidence from archaeological stratigraphy

272 The most relevant palaeoenvironmental changes from Vela Spila sediments are outlined here from
 273 bottom to top in order to relate them to shoreline changes. The general stratigraphy of the Vela Spila
 274 archaeological deposit is presented in Fig 4. The cave bedrock was not found in any of the
 275 excavated trenches, despite maximum excavation depths of about 5 m below the present-day cave
 276 floor, so the question of the oldest occupation of the cave is still open. Table 2 lists the collected ¹⁴C
 277 for this site, with calibration results summarised in Fig. 5 and referenced below in the text as well.

Lab Code	14C Age (a BP)	2-sigma calibrated age (a BP)	Method	Material dated	Reference	Attribution	Context
Z-3987	16200±200	19,530- 19,190	Std	WC	This paper*, Rainsford <i>et. al.</i> 2014	Palaeolithic	LUP A
VERA- 2338	16140±60	19665- 19255	AMS	WC	Farbstein <i>et al.</i> 2012: TAB. 1	Palaeolithic	LUP A
VERA- 2339	15690±70	19130- 18775	AMS	AB	Farbstein <i>et al.</i> 2012: TAB. 1	Palaeolithic	LUP A
z-3993	14500±100	17950- 17410	Std	WC	Farbstein <i>et al.</i> 2012: TAB. 1	Palaeolithic	LUP C
z-3992	14100±100	17587- 17105	Std	WC	Farbstein <i>et al.</i> 2012: TAB. 1	Palaeolithic	LUP D
z-3991	13300±100	16280- 15700	Std	WC	Farbstein <i>et al.</i> 2012: TAB. 1	Palaeolithic	LUP E
z-3988	12800±100	15660- 14925	Std	WC	This paper*, Rainsford <i>et. al.</i> 2014	Palaeolithic	LUP G
z-3990	12700±100	15500- 14655	Std	WC	This paper	Palaeolithic	LUP G
z-3989	12700±100	15500- 14655	Std	WC	Farbstein <i>et al.</i> 2012: TAB. 1	Palaeolithic	LUP G
VERA- 2345	12290±40	14490- 14050	AMS	AB	Farbstein <i>et al.</i> 2012: TAB. 1	Palaeolithic	NYT
VERA- 2346	12269±40	14410- 14025	AMS	WC	Čečuk & Radić 2005: 34	Palaeolithic	NYT
Wk- 27370	12457±57	14995- 14230	AMS	AB	This paper*, Rainsford <i>et. al.</i> 2014	Palaeolithic	LUP I
VERA- 2343	8290±35	9425-9135	AMS	AB	This paper	Mesolithic	Meso A
VERA- 2344	8230±35	9395-9030	AMS	WC	Čečuk & Radić 2005: 62	Mesolithic	Meso A
VERA- 2341	8200±30	9270-9030	AMS	WC	Čečuk & Radić 2005: 62	Mesolithic	Meso A

z-3995	8200±70	9400-9005	Std	WC	This paper*, Rainsford <i>et. al.</i> 2014	Mesolithic	Meso B
z-3986	8200±70	9400-9005	Std	WC	This paper*, Rainsford <i>et. al.</i> 2014	Mesolithic	
OxA-18171	8110±37	9235-8985	AMS	HB	Forenbahe <i>et al.</i> 2013: 591	Mesolithic	Meso B
Wk-24217	8004±41	9010-8715	AMS	HB	This paper		Burial 2/1986
z-3994	7410±70	8375-8045	Std	WC	This paper*, Rainsford <i>et. al.</i> 2014	Mesolithic	Meso D
Wk-24216	7350±39	8305-8025	AMS	AB	This paper*, Rainsford <i>et. al.</i> 2014	Mesolithic	Meso D
Z-1967	7300±120	8375-7930	Std	WC	Čečuk & Radić 2005: 62	Early Neolithic?	Meso D?
VERA-2342	7175±35	8045-7935	AMS	AB	This paper	Mesolithic	Meso D?
VERA-2340	7200±30	8155-7950	AMS	WC	Čečuk & Radić 2005: 62	Mesolithic	Meso D?
Z-1968	7000±120	8030-7595	Std	WC	Čečuk & Radić 2005: 82	Early Neolithic?	Early Neo?
Wk-24219	6735±35	7665-7515	AMS	AB	This paper	Neolithic	Neo C
Wk-24218	6678 ± 34	7610-7485	AMS	AB	This paper*, Rainsford <i>et al.</i> 2014	Neolithic	Neo C
z-1742	5430±100	6410-5950	Std	WC	Srdoč <i>et al.</i> 1989: 86	Neolithic ?	Neo?

Table 2. Radiocarbon age dataset from Vela Spila. Std: standard method; AMS: accelerator mass spectrometry; WC: wood charcoal; AB: animal bone; HB: human bone. *Uncalibrated date published for the first time in this paper.

278 4.1.1 Palaeolithic

279 The lowermost unit of the sequence (Fig. 4) exposed so far is represented by a 3 m-thick talus
 280 made up of unsorted frost slabs with an open-work fabric, and with no traces of human occupation.
 281 Even if geochronometric dates are not available for this unit, it can be attributed to the LGM
 282 because of its sedimentological characteristics and of the radiocarbon age of the overlying units
 283 (Farbstein et al., 2012). This type of sediment is consistent with a cold climatic spell typically
 284 connected with an RSL lowstand.

285 The sediments testifying to the first human occupation of the cave directly overlie the LGM
 286 talus and include Upper Palaeolithic remains ¹⁴C constrained to approximately 19.7-14.0 cal ka BP
 287 (Fig. 5, Table 2). These strata are composed of frost slabs that decrease in quantity and become finer
 288 upwards to the top of the Late Upper Pleistocene units. Alternations of discontinuous fine and

289 coarse-grained layers and/or gravelly lenses, as well as some blocks, testify to warmer and colder
290 climate fluctuations, and to localised deposition of gravel from inhomogeneous gelifraction of the
291 ceiling. The silty loamy fine fraction of the sediments suggests colluvial inputs of windblown
292 sediments previously deposited outside the cave or, when more clayey and reddish, of material
293 deriving from the dismantling of soils. The top of the sequence is marked by a horizon of large
294 limestone slabs overlain by a 15 cm-thick layer of sub-primary tephra locally deformed by frost
295 action, ascribed to the Neapolitan Yellow Tuff (NYT) ejected from a volcano in the Campi Flegrei
296 near Naples (Radić et al., 2007), ^{14}C dated to 12.3 ± 0.04 ka BP (14.5–14.1 cal ka BP; Tab. 2; Fig. 5)
297 (Čečuk and Radić, 2005; Farbstein et al., 2012), which fits reasonably well the 14.9 ± 0.4 ka BP
298 $^{40}\text{Ar}/^{39}\text{Ar}$ age of the tephra (Deino et al., 2004) and much better its modelled age 14.3–13.9 cal ka
299 BP date obtained by Bayesian method (Blockley et al., 2008).

300 The anthropic component of these layers –abundant organic matter, charcoal, bone and stone
301 tools deriving from intense occupation of the cave – is dominant over non-anthropoc material and
302 gives the sediment a very dark, blackish colour value. Archaeological materials include Late Upper
303 Palaeolithic lithics of Epigravettian tradition (Farbstein et al., 2012; Rainsford et al., 2014), sharing
304 strict analogies with other technocomplexes of Dalmatia and its hinterland, e.g. Pupičina peć
305 (Miracle, 2007), Šandalja II (Janković et al., 2012; Karavanić, 2003; Richards et al., 2015), Vlakno
306 (Cvitkušić et al., 2018; Vukosavljević and Perhoč, 2017), Zemunica (Šošić Klindžić et al., 2015),
307 Kopačina (Vukosavljević et al., 2011; Vukosavljević and Perhoč, 2017), Badanj (Whallon, 1999),
308 Crvena Stijena (Basler, 1975; Kozłowski and Kaczanowska, 2004). The large mammal fauna is
309 dominated by *Cervus elaphus* (red deer), followed by the dry environment taxon *Equus hydruntinus*
310 (European ass). Other less frequent taxa are *Capreolus capreolus* (roe deer), *Bos/Bison* (wild
311 cattle/bison), *Sus scrofa* (wild boar) and *Lepus* sp. (hare). A few specimens of *Canis lupus* (wolf),
312 *Felis lynx* (lynx), *Felis silvestris* (wild cat), *Vulpes vulpes* (red fox) are also present. Smaller
313 vertebrate remains (e.g., rodents, birds, bats, reptiles, fish) are not abundant in the Pleistocene

314 layers. The rich and abundant faunal assemblage suggests a landscape characterised by wide-open
315 areas with wooded patches that was resource-rich, productive and able to sustain a large community
316 of hunters-gatherers.

317 4.1.2 *Abandonment phase*

318 A complex and not yet fully understood human occupation gap of approximately 4.6 ka
319 occurred between the latest Late Upper Palaeolithic and the earliest Mesolithic phases (Radić,
320 2018). At the foot of the southern wall of the cave, the Mesolithic layers overlie the NYT tephra
321 directly. Conversely, at the centre of the cave huge limestone blocks, up to 2.5 m large, rest on the
322 NYT and are partially covered by about 1.4 m of limestone rubble mixed with reddish silty loam –
323 with no cultural remains. The base of Mesolithic layers is here about 1 m lower than at the foot of
324 the wall and overlies the rubble, apparently without sedimentary gaps. This situation suggests that
325 throughout the Late glacial the topography of the inside of the cave had been pre-determined by the
326 shape of the LGM scree, which had partly filled the cave from the western entrance and along the
327 southern wall. The central area was strongly depressed, so that the accumulation of the Late Upper
328 Palaeolithic layers did not level it completely. Later on, the deposition of ceiling breakdown blocks
329 was triggered by frost in some still undated –posterior to NYT – cold phase of the Late Glacial and
330 was followed by rubble and colluvium of reddish silty loam during the subsequent climate
331 improvement. The fine component of these sediments include mica flakes and other minerals
332 typical of the mineral suite of loess or loess-like sediments of the Adriatic area, and their grain-size
333 supports this (Cremaschi, 1990; Wacha et al., 2011a, 2011b). These sediments can be found only in
334 the central part of the cave, for the following two reasons: i) large breakdown blocks and rubble
335 accumulated only under the ceiling opening, which is located above this area; ii) colluvium flowing
336 into the cave from the western entrance, and probably also from the ceiling opening, was
337 preferentially deposited in the topographic low.

338 4.1.3 Mesolithic

339 This topographic setting also influenced the deposition of the Mesolithic layers. Thickness and
340 composition of sediments including the Mesolithic cultural remains are variable throughout the
341 cave; the bottom areas of the cave are rich in ash and charcoal remains, whereas the central part of
342 the cave was still slightly depressed when people started occupying the site again around 9.4 cal ka
343 BP. In the centre these layers are somewhat thick (about 1.3 m) and largely composed of fine
344 sediment deriving from the runoff of reworked red soils and/or windblown sediments, whereas they
345 are much thinner (about 20-60 cm) and stony at the foot of the southern wall. These characteristics
346 suggest that the topography of the cave (Fig. 3) inside was different from the present-day one, with
347 a large depression at its centre, and that different areas of the cave were used for specific purposes
348 by humans.

349 The age of the Mesolithic occupation is ca. 9.4 to 8.0 cal ka BP (Farbstein et al., 2012;
350 Rainsford et al., 2014). Faunal remains from these layers indicate a reduction of the typical Late
351 Pleistocene open environment mammal taxa, with a shift towards associations preferring good
352 vegetative shrubby or woody cover, represented here mostly by *Vulpes vulpes* (fox), *Capreolus*
353 *capreolus* (roe deer), *Sus scrofa* (wild boar). Among the infrequent remains of other small-size taxa
354 are those of *Erinaceus roumanicus* (Northern white-breasted hedgehog) (Mauch Lenardić et al.,
355 2018).

356 The decrease in mammal taxa size and a smaller number of ungulate remains testify to a
357 generalised reduction of high-quality food resources, which was replaced by the exploitation of
358 marine taxa. The Mesolithic units include remains of *Delphinidae* sp. (dolphin) and *Monachus*
359 *monachus* (monk seal), as well as abundant *Scomber japonicus* (mackerel) and other fish taxa
360 (Rainsford et al., 2014). These marine vertebrates are associated with extremely abundant molluscs
361 like *Patella caerulea* and *Patella rustica* (limpet) and Trochidae, mostly *Phorcus turbinatus*

362 (turbinate monodont) (Branscombe et al. 2020). Several other marine mollusc taxa– mostly adapted
363 to rocky coasts – are also well represented, even if less abundant than the aforementioned ones;
364 continental molluscs are also common, as in most Adriatic area sites (Rizner et al., 2009), and
365 mostly represented by *Eobania vermiculata* (chocolate band snail). Other Mesolithic sites of
366 Dalmatia and neighbouring areas that share similar techno-typological and economic characteristics
367 are Pupićina peć (Miracle, 2007, 2002, 2001), Abri Šebrn (Miracle et al., 2000), Nugljanska peć
368 (Pilaar Birch and Miracle 2015), Vela Spilja Lošinj (Pilaar Birch and Miracle 2017), Vlakno cave
369 (Vukosavljević et al., 2014; Cvitkušić et al., 2018), Zemunica (Šošić Klindžić et al., 2015). The
370 remarkable amount of marine resource remains found in the cave, including large quantities of
371 mackerel bones, pierced *Columbella* (dove snail) shells, and a relatively few large fish bones
372 (possibly associated with deep-sea fishing), indicates an economic shift towards the sea.

373 4.1.4 Neolithic

374 The age of the Holocene sediments (including the earliest Neolithic remains of Vela Spila
375 ascribed to the Impresa Ware cultural horizon) is bracketed between 8.0 and 6.0 cal ka BP (Čečuk
376 and Radić, 2001). During the Neolithic there was virtually no deposition of clastic sediments
377 derived by soil erosion from the outside of the cave, which is interpreted as a sign of extended
378 vegetation on the slopes, favoured by mild climatic condition and evenly distributed rainfall.
379 Depositional rates were however high inside the cave, because penning of domesticated animals,
380 mostly sheep and goats, produced large quantities of dung that was intentionally burned. Typical
381 sediments of this phase are the so-called “layer-cake” or *fumier* sequences (Boschian and Miracle,
382 2007), characterised by long successions of cyclically alternating black and white layers including
383 ash, charcoal and other products of herbivore metabolism. From the Neolithic onwards, depositional
384 environments in caves change in this way all over the Mediterranean area (Angelucci et al., 2009),
385 demonstrating the new role of caves within a complex agropastoral system. Similar *fumier*

386 sequences were excavated in other caves on both sides of the Adriatic Sea, e.g. in several north
387 Adriatic Karst sites (Boschian and Montagnari-Kokelj, 2000), at Pupičina peć (Boschian, 2006;
388 Forenbaher and Miracle, 2006) in Istria, at Zemunica Peć (Šošić Klindžić et al., 2015) in central
389 Dalmatia, and at Grotta dei Piccioni and Grotta S. Angelo (Boschian and Iaconis, 2008) in central
390 Italy.

391 Neolithic deposits contain abundant bones of domestic animals (mostly sheep and goats), few
392 wild animal bones, some terrestrial and marine mollusc shells, many potsherds, chert from the
393 Gargano Peninsula (c. 125 km to the southwest) and obsidian from Lipari (c. 500 km to the
394 southwest). Evidence of soil erosion is again apparent in the sequence corresponding with the Late
395 Neolithic onwards. In this case erosion is interpreted as evidence of human activities (agriculture
396 and pastoralism) which caused deforestation.

397 A long-lasting and intensive Neolithic occupation of the site is represented by a complete
398 sequence of cultural facies and characteristic pottery of all major styles occurring in the eastern
399 Adriatic area during this period (Forenbaher et al., 2013), testifying to the sociocultural and
400 economic aspects of the Neolithic transition (Zvelebil and Lillie, 2000) and the further evolution of
401 the Neolithic towards Copper Age and Early Bronze Age.

402 Throughout the period of occupation subsistence depended heavily on herding. Faunal
403 assemblages are dominated by domestic animal remains; primarily sheep and goats, while cattle and
404 pigs are less common. Also present in relatively small quantities are terrestrial and marine molluscs,
405 while fish remains are quite rare.

406 Despite systematic attempts to recover seeds through flotation of soil samples, direct evidence
407 of agriculture remains scarce, but this should not be interpreted as indicating an absence of plant
408 cultivation. Productive arable land tends to be located some distance away from the heavily
409 karstified areas where caves abound, and a similar absence of cultivars was observed in other caves
410 in the region (Borojević et al., 2008; Forenbaher and Miracle, 2006). After the Neolithic, the

411 archaeological sequence continues through the Copper and Bronze Ages (Radić, 2018), with rich
 412 ceramic assemblages and sediments of mostly *terra rossa* colluvium and anthropic components
 413 represented by variable amounts of organic matter and stable sediments.

414 **4.2 Sea-level change and palaeogeography around Vela Spila**

415 According to the model predictions (Fig. 6), RSL was 101-107 m below present at the start of
 416 Vela Spila's Upper Palaeolithic occupational phase, and then climbed at a rate of about 5 to 6 mm/a.
 417 Combining the isostatic model with the bathymetric analysis, it is possible to infer that the
 418 landscape around the Island of Korčula changed remarkably over the site's prehistoric phases (Table
 419 3). Contemporaneous with the first detected Upper Palaeolithic occupation (starting at ~19.7 cal ka
 420 BP, Fig. 7A), the cave was ~8-9 km from the sea and 221-228 m above it, while the area south of
 421 western Korčula consisted of a broad plain where the present-day Lastovo island group appeared as
 422 a cluster of hills. 7-16% of the land within 20 km of the site (a day's round-trip travel at a
 423 comfortable walk) was inundated. Korčula was still also connected to the mainland in the east via
 424 the Pelješac Peninsula and Hvar Island to the north. To the SW of the present bay of Vela Luka, a 0-
 425 30 m deep sea existed, and to the NW Vis and Biševo were also not yet islands. South of Vela Luka
 426 bay, a natural depression indicated by the bathymetric data might have been filled by a small,
 427 shallow lake or marsh a few meters deep and a couple kilometres wide. To the N and NE of present
 428 day Korčula, the palaeo-Neretva river flowed in an alluvial plain between Hvar and Korčula, shown
 429 in GIS reconstructions carried out by Sikora et al. (2014).

430

Archaeological phase	Start (ka BP)	End (ka BP)	RSL at start (m)	RSL at end (M)	RSL change (m)	Average rate of change (mm/a)	Average Horizontal Coastline Change (m/a)	Vertical RSL Change in human lifetime (m)	Horizontal coastline change in human lifetime (m)
Palaeolithic	19.7	14.0	-107 to -100	-74 to -72	29 to 33	5 to 6	5	0.2	160

Abandonment	14.0	9.4	-74 to -72	-30 to -16	44 to 55	10 to 12	14	0.4	492
Mesolithic	9.4	8.0	-30 to -16	-13 to -7	9 to 17	7 to 12	0.5 in Vela Luka Bay	0.2 to 0.4	18 in Vela Luka Bay
Neolithic	8.0	6.0	-13 to -7	-3 to -1	6 to 10	3 to 5	0.1 in Vela Luka Bay	0.1 to 0.2	4 in Vela Luka Bay

Table 3. Palaeogeographic changes by archaeological period at Vela Spila. In the Mesolithic and Neolithic, coastal migration around the island becomes insignificant so the rates of incursion into the long, valley-like Vela Luka Bay are noted instead. N.b. the distinction between mm/a (mm per year) and m/a (meters per year).

431 By the end of the Palaeolithic occupation at 14.0 ka BP (Fig. 7B), ca. 5 ka later, RSL had risen
 432 29 to 33 m (Table 3), the coastline had advanced 4-10 km into Vela Luka Bay, and Vis and Biševo
 433 were likely already islands. At this point, about 40% of land within 20 km of the site was flooded.
 434 When the cave was abandoned, sometime after the NYT eruption between 14.9 and 13.9 cal ka BP
 435 (Blockley et al., 2008; Deino et al., 2004), any nearby lake or depression south of Vela Luka Bay
 436 had already become part of the Adriatic Sea, which had been advancing eastward up to 4 m/a
 437 (meters per year) in this area.

438 According to our models, the most dramatic RSL rise and landscape change around Korčula
 439 occurred during the roughly 5 ka (14.0-9.4 cal ka BP) period when the cave was abandoned. RSL
 440 rose ~44 to 55 m at an average of about 10–13 mm/a and the coastline advanced eastward by as
 441 much as 10 to 12 m/a or more to the north and south of the island (Table 3). Land connections to
 442 areas north and south of Korčula, such as present day Lastovo and Hvar, were probably severed
 443 during the first millennium of the abandonment phase.

444 Much of the coastlines of the islands advanced to within 1 km of their present position by ca.
 445 12 ka BP, after which horizontal changes in coastlines became much smaller: above the ~70 m
 446 isobath, slopes around the island tend to be much steeper (Fig. 8). A land bridge between the
 447 Pelješac Peninsula and Korčula may have persisted until between 12 and 10 ka BP, at which time
 448 the geography of the area became quite similar to the present-day. When the cave was finally
 449 reoccupied at the start of the Mesolithic at ~8.0 ka BP (Fig. 7C, Fig. 9), the surrounding landscape

450 had been totally transformed. About 80% of nearby lands were flooded. 30-50 m deep seas were 2
451 km away from the cave to the north and south, and a shallow (10-20 m) bay was only 1-2 km to the
452 west. No land connections with other present-day islands existed, though many of the smaller islets
453 around Korčula may still have been connected to it at times. By ca. 8.0 ka BP, at the beginning of
454 the Neolithic, most of the northern and southern coasts of the island were even closer to the present-
455 day shoreline, though some islets may still have been connected, and the inner part of Vela Luka
456 harbour had not been fully submerged.

457 **5 Discussion**

458 **5.1 Last Palaeolithic occupants facing rapidly inundating planes**

459 The Palaeolithic people at Vela Spila experienced average RSL change rates of 5–6 mm/a,
460 although the ANU_S4 realization does predict some phases of higher rates. The rate of RSL change
461 increased to 10–12 mm/a during the abandonment phase (Table 3). Hence the Late Upper
462 Palaeolithic coastline was relatively stable compared to that of the subsequent abandonment phase.
463 The cave was only 8–9 km distant from the sea, and the lack of shells in the archaeological
464 stratigraphy during this phase of inhabitation is not necessarily a consequence of distance from the
465 cave but, more likely, a preference for the terrestrial species which had been the traditional Upper
466 Palaeolithic food resource (Miracle, 2007; Mussi and Palombo, 2001; Stewart et al., 2003) and
467 would still have been widely available at this point, given that they occupy a large part of the diet.
468 The palaeo-Neretva's plain north of Korčula became submerged during the abandonment phase at
469 Vela Spila, and in those millennia the high rates of RSL change may not have left time for the
470 formation of submerged landforms such as palaeocoastlines or palaeodeltas (Sikora et al., 2014).

471 During the Palaeolithic phase, only a relatively small area of potential hunting grounds was
472 inundated in the plains north and south of Korčula. Still, the coastline moved east at a rate of nearly

473 5 m/a, so that in the average human lifespan of 35 years, a person would have witnessed ca. 160
474 meters of horizontal coastal change. The rate of change experienced by the Upper Palaeolithic
475 foragers in the Central Adriatic was 6-7 times greater than the present-day estimates of ongoing
476 RSL change (Church et al., 2013; Neumann et al., 2015). It is most likely that this landscape change
477 stressed littoral habitats and limited potential human exploitation of marine resources; however,
478 sediments and cultural remains show that human frequentation continued to be intensive in the
479 cave. Following on the Palaeolithic occupation at Vela Spila, the abandonment phase (from ca. 14.0
480 to 9.4 cal ka BP) appears to have seen even more intense changes in the surrounding
481 palaeogeography.

482 **5.2 Environmental reasons for abandoning the cave: rising sea or climate?**

483 The deposition into Vela Spila of in-washed NYT tephra (Fig. 4) coincided or preceded
484 approximately the end of the Late Upper Palaeolithic occupation and the beginning of a long gap in
485 the human record (14.0–9.4 cal ka BP), at the start of which the RSL was 74–72 m below present.
486 Since the eruption took place about 300 km SW of Vela Spila, it is worth speculating what role, if
487 any, this event may have played in inducing Upper Palaeolithic people to abandon the cave.
488 Cryoturbation features involving specifically the NYT sediment in the cave suggest that deposition
489 was associated or immediately followed by a cold spell, though it cannot be ascertained if it was
490 consequential to the eruption.

491 Rapid landscape changes caused by a high RSL rise rate may have had a greater influence on
492 human behaviour and settlement pattern change during this phase; predicted RSL changes were
493 much faster than during the Upper Palaeolithic cave occupation. Humans would have witnessed
494 nearly 0.40 m of RSL rise, corresponding to about 500 m of horizontal coastline displacement
495 during a 35-year average lifespan, i.e. about 14 m of horizontal eastward coastal change per year
496 immediately north and south of Korčula. This suggests a highly unstable or marshy littoral zone,

497 rapid flooding of the traditional foraging grounds of the Upper Palaeolithic groups, and stress on the
498 ecosystem at all trophic levels. Particularly rapid change (about 20 m rise) took place during the
499 centuries following the Meltwater Pulse 1A (MWP-1A), which has been dated to 14.7–14.3 ka BP
500 (Deschamps et al., 2012).

501 Volcanic eruptions such as the NYT on the other hand usually have an impact on climate for
502 only a few years, so that their consequences are not easily detected in $\delta^{18}\text{O}$ isotope or other climate
503 proxies (Aufgebauer et al., 2012). The NYT eruption also coincides with the end of a colder climate
504 phase highlighted at Lake Ohrid (Vogel et al., 2010), and with increased winter precipitation
505 (Bordon et al., 2009; Aufgebauer et al., 2012). If the settlement was already in a marginal position
506 due to climate changes and to the landscape changes highlighted by this study, a sequence of
507 particularly unfavourable years due to an eruption may have contributed to pushing the Late Upper
508 Palaeolithic people to a tipping point that eventually resulted in abandonment of the site.

509 The first two thousand years of the Vela Spila occupation gap overlap with, in addition to the
510 NYT and Meltwater Pulse 1A, the Bølling–Allerød warming, and were followed by the Younger
511 Dryas major dry/cold event (Gkinis et al., 2014), which is recorded throughout the Balkans in
512 various lacustrine sequences (Cvetkoska et al., 2014). At many Late Upper Palaeolithic sites in the
513 Eastern Adriatic region the Bølling–Allerød warming is associated an increase in of human
514 settlement (Miracle, 2007), and the same pattern is largely true on the Italian side of the Adriatic
515 (Mussi 2001). The occupational sequence at Vela Spila is in stark contrast to this wider pattern.

516 Younger Dryas effects in the Adriatic Plain are indicated at Vela Spila by the occurrence of
517 partially altered loess-like sediments colluviated into the cave. The precise timing of this
518 colluviation is currently unknown; it probably was partly during the Younger Dryas, but mostly
519 during the early Holocene. It is, however, unclear that the Bølling–Allerød or Younger Dryas
520 climate changes played a direct role in forcing the Late Upper Palaeolithic foragers to leave the site,
521 although the Younger Dryas is correlated with cultural change and adaptation throughout Europe

522 (Aura et al., 2011; Burdukiewicz, 2011; Mussi and Peresani, 2011; Weber et al., 2011). Instead, the
523 rapid RSL rise concurrent with the meltwater pulse 1A and Bølling–Allerød may have been the
524 environmental trigger for the abandonment of Vela Spila, since the cave had already been
525 abandoned by the time of the Younger Dryas at ca. 12 or 13 ka BP (Gkinis et al., 2014).

526

527 **5.3 Stabilising coastlines and the Mesolithic**

528 Benjamin et al. (2017) suggest that the Northern Hemisphere cooling during the Younger Dryas
529 slowed the RSL rise in the Mediterranean as in other areas in the world. The GIA models in this
530 work (Fig. 6) also predict a decrease in RSL rise rates following the Younger Dryas, but only few
531 local relative sea-level indicators are available for verification. By ca. 9.4 ka BP, when the model
532 predicts RSL rates were starting to level off, Mesolithic groups re-occupied the cave and continued
533 dwelling in it for the following ~ 1.5 ka . The corresponding units of the cave sequence show
534 evidence of the Preboreal and Boreal humid climate conditions indicated by colluvia. In this period
535 RSL continued rising quickly from between -30 to -16 below present, up to between -13 m and -7 m
536 – i.e. a rise of 9 m to 17 m in approximately 1.5 ka (Table 3), which gives a rate of 7 to 12 mm/a,
537 though this is a slower rate of change than during the abandonment phase. The sea was only about
538 1-2 km distant from the cave at the onset of the Mesolithic, compared to 8-9 km in the Upper
539 Palaeolithic.

540 The Mesolithic assemblage is dominated by the remains of shellfish, fish and other marine
541 creatures. The exploitation of marine resources by Mesolithic people is widespread in the
542 Mediterranean and the Atlantic, though the Mediterranean is supposed to be less productive
543 (Colonese et al., 2011). In general, resource depression and human population pressure have been
544 considered the main reason for shifting toward marine resources (Colonese et al., 2011), even if
545 there are major differences between regions. However, it does not appear that the exploitation of

546 fish and shellfish was ever the most important food resource in the Adriatic region; isotopic studies
547 on human remains show that “while the Mesolithic people of the Adriatic coast consumed some
548 marine protein, terrestrial protein constituted a significant part of the diet” (Lightfoot et al., 2011, p.
549 82). The balance between the need for resource diversification and expedience of marine resources
550 due to the approaching coastline is not always clear or consistent. As the Mesolithic progresses,
551 there is a decline in the use of fish and an increase in shellfish and snails at Vela Spila, along with a
552 decrease in terrestrial fauna remains. Fishing seems to shift from open-sea to inshore taxa
553 (Rainsford et al. 2014). This is partially in contrast with data from the northern Adriatic where the
554 coastline approached later: here, fishing shifts from freshwater to marine taxa and Early Mesolithic
555 intensive snail exploitation is substituted by consumption of marine species (Boschian and Pitti,
556 1984; Cremonesi et al., 1984).

557 These data suggest that, in the northern Adriatic, the need for resource diversification was the
558 driving force towards marine food well before landscape modification due to RSL rise. In contrast,
559 the same need started more or less contemporaneously at Vela Spila, but was immediately oriented
560 towards marine resources because of the earlier advance of the coastline. The decrease in other
561 resources is testified by the disappearance of red deer and other large ungulates from the Mesolithic
562 layers, presumably because of the early Holocene palaeoenvironmental turnover, and because
563 Korčula had become an island where faunal replacement was impossible as access to more
564 productive hunting grounds decreased.

565 The transition between Mesolithic and Neolithic started in the Eastern Adriatic at ca. 8.0 ka BP
566 (Forenbaher et al., 2013). Radiocarbon determinations and cultural assemblages from Vela Spila are
567 consistent with this date. The Mesolithic-Neolithic transition shortly follows the 8.2 ka event, which
568 was a disruption in thermohaline Atlantic circulation owing to meltwater input from North America
569 (Alley and Ágústsdóttir, 2005). The 8.2 ka BP event is associated with an estimated drop in winter
570 temperatures of more than 4° C, at least in Northeastern Greece (Pross et al., 2009) and the sea-

571 level impact of the event in the study area is approximately 1 m (Kendall et al., 2008), although this
572 perturbation is too rapid and small to appear in the GIA models. Furthermore, because the coastlines
573 were already near their present-day position abutting the high relief slopes of the island (Fig. 8), the
574 RSL rise itself would not have been significant for the inhabitants unless they had built immobile
575 coastal infrastructure, of which no evidence exists. Likewise, no clear fingerprint of the 8.2 event
576 exists in the archaeological record of the site. By the end of the Neolithic at ca. 6.0 ka BP, RSL had
577 risen to within 3 meters of present level, and sea-level changes from then on were much slower
578 (Fig. 6).

579 **6 Conclusions**

580 In this work we direct evidence from a detailed, extensively excavated archaeological
581 stratigraphy was used to understand human reactions to changing rates of RSL rise and horizontal
582 coastline shift in a coastal landmass that becomes an island.

583 In the Late Upper Palaeolithic the abandonment of Vela Spila cave was triggered by
584 environmental events that caused the area to be less attractive for humans. In particular, high rates
585 of sea-level rise likely stressed the least observant inhabitants in Vela Spila more sharply than other
586 environmental factors (e.g. climatic forcing) by submerging hunting grounds and reshaping marine
587 and terrestrial ecosystems in the rapidly disappearing Great Adriatic Plain, where the coastline
588 advanced tens of meters per year.

589 The coastline positions around Korčula island become stabilized only by ~12 ka BP as the
590 rising seas surpassed the 70 m isobath and encountered the steep morphology of the island's sides
591 (Fig. 8), though rapid RSL rise continued. This state, together with other environmental factors,
592 may have fostered the re-occupation of the cave at ca. 9.4 cal ka BP. In the Mesolithic, it was the
593 new insularity of the island that forced humans to exploit all resources available (e.g. seafood and

594 smaller fauna like foxes, rather than larger animals such as red deer). The disadvantages of this
595 insularity were initially overcome by skilled seafaring and later, more effectively, by the onset of
596 agriculture and pastoralism that accompanied, after 8 ka BP, a substantial slowing down of sea-level
597 rise.

598 This study also emphasizes the importance of potential future research projects aimed at
599 gathering underwater data, for example by more detailed bathymetric surveys or sediment coring in
600 relevant areas around Vela Luka bay such as the plains to the north and south, the palaeo Neretva,
601 and the strait between Korčula and the Pelješac, which could reveal additional drowned landforms
602 and improve resolution for sea-level change histories and the environmental transformation of
603 Korčula from mainland to island.

604 **7 Acknowledgements**

605 The authors of this paper would like to recognize the EU project no. 692249 “Smart Integration
606 of Genetics with Sciences of the Past in Croatia: Minding and Mending the Gap”
607 (<http://mendthegap.agr.hr/>) for providing most of the travel and workshop opportunities for many of
608 the authors to perform research for this paper. Archaeological excavations and analyses were also
609 supported by the British Academy (SG101073), the University of Cambridge, and St John’s
610 College, Cambridge.

611 **8 Works Cited**

612
613 Alley, R.B., Ágústsdóttir, A.M., 2005. The 8k event: cause and consequences of a major
614 Holocene abrupt climate change. *Quaternary Science Reviews* 24, 1123–1149.
615 <https://doi.org/10.1016/j.quascirev.2004.12.004>

- 616 Andel, T.H.V., 1989. Late Quaternary sea-level changes and archaeology. *Antiquity* 63, 733–
617 745. <https://doi.org/10.1017/S0003598X00076869>
- 618 Angelucci, D.E., Boschian, G., Fontanals, M., Pedrotti, A., Vergès, J.M., 2009. Shepherds and
619 karst: the use of caves and rock-shelters in the Mediterranean region during the Neolithic. *World*
620 *Archaeology* 41, 191–214. <https://doi.org/10.1080/00438240902843659>
- 621 Antonioli, F., Anzidei, M., Lambeck, K., Auriemma, R., Gaddi, D., Furlani, S., Orrù, P.,
622 Solinas, E., Gaspari, A., Karinja, S., Kovačić, V., Surace, L., 2007. Sea-level change during the
623 Holocene in Sardinia and in the northeastern Adriatic (central Mediterranean Sea) from
624 archaeological and geomorphological data. *Quaternary Science Reviews* 26, 2463–2486.
625 <https://doi.org/10.1016/j.quascirev.2007.06.022>
- 626 Artegiani, A., Paschini, E., Russo, A., Bregant, D., Raicich, F., Pinardi, N., 1997. The Adriatic
627 Sea General Circulation. Part II: Baroclinic Circulation Structure. *J. Phys. Oceanogr.* 27, 1515–
628 1532. [https://doi.org/10.1175/1520-0485\(1997\)027<1515:TASGCP>2.0.CO;2](https://doi.org/10.1175/1520-0485(1997)027<1515:TASGCP>2.0.CO;2)
- 629 Aufgebauer, A., Panagiotopoulos, K., Wagner, B., Schaebitz, F., Viehberg, F.A., Vogel, H.,
630 Zanchetta, G., Sulpizio, R., Leng, M.J., Damaschke, M., 2012. Climate and environmental change
631 in the Balkans over the last 17 ka recorded in sediments from Lake Prespa (Albania/F.Y.R. of
632 Macedonia/Greece). *Quaternary International*, Temporal and spatial corridors of *Homo sapiens*
633 *sapiens* population dynamics during the Late Pleistocene and Early Holocene 274, 122–135.
634 <https://doi.org/10.1016/j.quaint.2012.02.015>
- 635 Aura, J.E., Jordá, J.F., Montes, L., Utrilla, P., 2011. Human responses to Younger Dryas in the
636 Ebro valley and Mediterranean watershed (Eastern Spain). *Quaternary International* 242, 348–359.
- 637 Basler, Đ., 1975. Crvena stijena: zbornik radova. Zajednica kulturnih ustanova, Nikšić.
- 638 Bekić, L., 2017. Šimuni, novo podvodno brončanodobno nalazište u Zadarskoj županiji.
639 *Diadora: glasilo Arheološkog muzeja u Zadru* 31, 41–50.

- 640 Bekić, L., Pešić, M., Scholz, R., Meštrov, M., 2015. Podvodna arheološka istraživanja na
641 prapovijesnom nalazištu Pakoštane–Janice. *Diadora: glasilo Arheološkog muzeja u Zadru* 29, 7–22.
- 642 Benjamin, J., Bekić, L., Komšo, D., Uhač, I.K., Bonsall, C., 2011. Investigating the submerged
643 prehistory of the eastern Adriatic: Progress and prospects. *Submerged prehistory* 193–206.
- 644 Benjamin, J., Bonsall, C., 2009. An underwater archaeological survey in the northern Adriatic
645 Sea. *International Journal of Nautical Archaeology* 38, 163–172.
- 646 Benjamin, J., Rovere, A., Fontana, A., Furlani, S., Vacchi, M., Inglis, R.H., Galili, E., Antonioli,
647 F., Sivan, D., Miko, S., Mourtzas, N., Felja, I., Meredith-Williams, M., Goodman-Tchernov, B.,
648 Kolaiti, E., Anzidei, M., Gehrels, R., 2017. Late Quaternary sea-level changes and early human
649 societies in the central and eastern Mediterranean Basin: An interdisciplinary review. *Quaternary*
650 *International* 449, 29–57. <https://doi.org/10.1016/j.quaint.2017.06.025>
- 651 Bini, M., Pappalardo, M., Rossi, V., Noti, V., Amorosi, A., Sarti, G., 2018. Deciphering the
652 effects of human activity on urban areas through morphostratigraphic analysis: The case of Pisa,
653 Northwest Italy. *Geoarchaeology* 33, 43–51. <https://doi.org/10.1002/gea.21619>
- 654 Birch, S.E.P., Linden, M.V., 2018. A long hard road... Reviewing the evidence for
655 environmental change and population history in the eastern Adriatic and western Balkans during the
656 Late Pleistocene and Early Holocene. *Quaternary International, Impacts of gradual and abrupt*
657 *environmental changes on Late glacial to Middle Holocene cultural changes in Europe* 465, 177–
658 191. <https://doi.org/10.1016/j.quaint.2016.12.035>
- 659 Blockley, S.P., Ramsey, C.B., Pyle, D.M., 2008. Improved age modelling and high-precision
660 age estimates of late Quaternary tephras, for accurate palaeoclimate reconstruction. *Journal of*
661 *Volcanology and Geothermal Research* 177, 251–262.
- 662 Blute, M., 2008. CULTURAL ECOLOGY. *Encyclopedia of Archaeology*.
663 <https://doi.org/10.1016/B978-012373962-9.00098-4>

- 664 Bonacci, O., Ljubenković, I., Knežić, S., 2012. The water on a small karst island: the island of
665 Korčula (Croatia) as an example. *Environ Earth Sci* 66, 1345–1357. [https://doi.org/10.1007/s12665-](https://doi.org/10.1007/s12665-011-1345-9)
666 011-1345-9
- 667 Bordon, A., Peyron, O., Lézine, A.-M., Brewer, S., Fouache, E., 2009. Pollen-inferred Late-
668 Glacial and Holocene climate in southern Balkans (Lake Maliq). *Quaternary International, Rhythms*
669 *and Causalities of the Anthropisation Dynamic in Europe between 8500 and 2500 cal BP:*
670 *Sociocultural and/or Climatic Assumptions* 200, 19–30.
671 <https://doi.org/10.1016/j.quaint.2008.05.014>
- 672 Borojević, K., Forenbaher, S., Kaiser, T., Berna, F., 2008. Plant use at Grapčeva cave and in the
673 eastern Adriatic Neolithic. *Journal of Field Archaeology* 33, 279–303.
- 674 Boschian, G., 2006. Geoarchaeology of Pupićina cave, in: Miracle, P., Forenbaher, S. (Eds.),
675 *Prehistoric Herders of Northern Istria: The Archaeology of Pupićina Cave*. Archaeological Museum
676 of Istria, Pula, pp. 123–162.
- 677 Boschian, G., Fusco, F., 2007. Figuring Out No-One's Land Why Was the Karst Deserted in the
678 Late Glacial?, in: Whallon, R. (Ed.), *Late Paleolithic Environments and Cultural Relations Around*
679 *the Adriatic*. Archaeopress, Oxford, pp. 15-+.
- 680 Boschian, G., Iaconis, M., 2008. Geoarchaeology of the deposits of Grotta dei Piccioni and
681 Grotta Sant'Angelo (Abruzzo, Central Italy), in: Boschian, G. (Ed.), *Proceedings of the 2nd*
682 *International Conference on Soils and Archaeology, Atti Società Toscana Scienze Naturali, Mem.*
683 pp. 181–188.
- 684 Boschian, G., Miracle, P., 2007. Shepherds and caves in the karst of Istria (Croatia). *Atti della*
685 *Società Toscana di Scienze Naturali* 173–180.
- 686 Boschian, G., Montagnari-Kokelj, E., 2000. Prehistoric shepherds and caves in the Trieste Karst
687 (Northeastern Italy). *Geoarchaeology* 15, 331–371. [https://doi.org/10.1002/\(SICI\)1520-](https://doi.org/10.1002/(SICI)1520-6548(200004)15:4<331::AID-GEA3>3.0.CO;2-H)
688 6548(200004)15:4<331::AID-GEA3>3.0.CO;2-H

- 689 Boschian, G., Pitti, C., 1984. I livelli mesolitici della Grotta dell'Edera, in: *Il mesolitico sul*
690 *Carso Triestino*, Quaderni della Società per la Preistoria e Protostoria della Regione Friuli-Venezia
691 Giulia. Italo Svevo, Trieste, pp. 143–210.
- 692 Branscombe, T.L., Bosch, M.D., Miracle, P.T., 2020. Seasonal Shellfishing across the East
693 Adriatic Mesolithic-Neolithic Transition: Oxygen Isotope Analysis of *Phorcus turbinatus* from Vela
694 Spila (Croatia). *Environmental Archaeology* 1–14.
- 695 Burdukiewicz, J.M., 2011. Late Glacial hunter-gatherer reactions to the Younger Dryas cooling
696 event in the southern and eastern Baltic regions of Europe. *Quaternary International* 242, 302–312.
- 697 Butzer, K.W., 2005. Environmental history in the Mediterranean world: cross-disciplinary
698 investigation of cause-and-effect for degradation and soil erosion. *Journal of Archaeological*
699 *Science* 32, 1773–1800. <https://doi.org/10.1016/j.jas.2005.06.001>
- 700 Carroll, F.A., Hunt, C.O., Schembri, P.J., Bonanno, A., 2012. Holocene climate change,
701 vegetation history and human impact in the Central Mediterranean: evidence from the Maltese
702 Islands. *Quaternary Science Reviews* 52, 24–40. <https://doi.org/10.1016/j.quascirev.2012.07.010>
- 703 Casalbore, D., Falese, F., Martorelli, E., Romagnoli, C., Chiocci, F.L., 2017. Submarine
704 depositional terraces in the Tyrrhenian Sea as a proxy for paleo-sea level reconstruction: Problems
705 and perspective. *Quaternary International*, *Quaternary coastal and marine studies in Central*
706 *Mediterranean* 439, 169–180. <https://doi.org/10.1016/j.quaint.2016.02.027>
- 707 Čečuk, B., Radić, D., 2005. Višeslojno pretpovijesno nalazište - Vela Luka, Otok Korčula.
708 Višeslojno pretpovijesno nalazište - Vela Luka, Otok Korčula, Korčula.
- 709 Čečuk, B., Radić, D., 2001. Vela Špilja - preliminarni rezultati dosadašnjih istraživanja, in:
710 Čečuk, B. (Ed.), *Arheološka istraživanja na području otoka Korčule i Lastova*. Hrvatsko arheološko
711 društvo, Zagreb, pp. 75–118.

- 712 Čelhar, M., Parica, M., Ilkić, M., Vujević, D., 2017. A Bronze Age underwater site near the islet
713 of Ričul in northern Dalmatia (Croatia). *SKYLLIS-Zeitschrift für maritime und limnische*
714 *Archäologie und Kulturgeschichte* 17, 21.
- 715 Church, J.A., Clark, P.U., Cazenave, A., Gregory, J.M., Jevrejeva, S., Levermann, A.,
716 Merrifield, M.A., Milne, G.A., Nerem, R.S., Nunn, P.D., Payne, A.J., Pfeffer, W.T., Stammer, D.,
717 Unnikrishnan, A.S., 2013. Sea Level Change, in: *The Physical Science Basis: Contribution of*
718 *Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*
719 *Change*. pp. 1137–1216.
- 720 Colonese, A.C., Mannino, M.A., Mayer, D.B.-Y., Fa, D.A., Finlayson, J.C., Lubell, D., Stiner,
721 M.C., 2011. Marine mollusc exploitation in Mediterranean prehistory: an overview. *Quaternary*
722 *international* 239, 86–103.
- 723 Cremaschi, M., 1990. Stratigraphy and palaeoenvironmental significance of the loess deposits
724 on Susak Island (Dalmatian archipelago). *Quaternary International* 5, 97–106.
- 725 Cremonesi, G., Meluzzi, C., Pitti, C., Wilkens, B., 1984. Grotta Azzurra: scavi 1982 (nota
726 preliminare), in: *Il mesolitico sul Carso Triestino, Quaderni della Società per la Preistoria e*
727 *Protostoria della Regione Friuli-Venezia Giulia*. Italo Svevo, Trieste, pp. 21–37.
- 728 Cristiani, E., Farbstein, R., Miracle, P., 2014. Ornamental traditions in the Eastern Adriatic: The
729 Upper Palaeolithic and Mesolithic personal adornments from Vela Spila (Croatia). *Journal of*
730 *Anthropological Archaeology* 36, 21–31. <https://doi.org/10.1016/j.jaa.2014.06.009>
- 731 Croatian Geoportal, 2019. Topografska karta u mjerilu 1:25000.
- 732 Cvetkoska, A., Levkov, Z., Reed, J.M., Wagner, B., 2014. Late Glacial to Holocene climate
733 change and human impact in the Mediterranean: The last ca. 17ka diatom record of Lake Prespa
734 (Macedonia/Albania/Greece). *Palaeogeography, Palaeoclimatology, Palaeoecology* 406, 22–32.
735 <https://doi.org/10.1016/j.palaeo.2014.04.010>

- 736 Cvitkušić, B., Radović, S., Vujević, D., 2018. Changes in ornamental traditions and subsistence
737 strategies during the Palaeolithic-Mesolithic transition in Vlakno cave. *Quaternary International*,
738 *Quaternary of Croatia* 494, 180–192. <https://doi.org/10.1016/j.quaint.2017.08.053>
- 739 De Boer, B., Stocchi, P., Van De Wal, R., 2014. A fully coupled 3-D ice-sheet-sea-level model:
740 algorithm and applications. *Geoscientific Model Development* 7, 2141–2156.
- 741 Deino, A.L., Orsi, G., de Vita, S., Piochi, M., 2004. The age of the Neapolitan Yellow Tuff
742 caldera-forming eruption (Campi Flegrei caldera – Italy) assessed by $^{40}\text{Ar}/^{39}\text{Ar}$ dating method.
743 *Journal of Volcanology and Geothermal Research*, *The Neapolitan Volcanoes: Vesuvius, Campi*
744 *Flegrei and Ischia* 133, 157–170. [https://doi.org/10.1016/S0377-0273\(03\)00396-2](https://doi.org/10.1016/S0377-0273(03)00396-2)
- 745 Deschamps, P., Durand, N., Bard, E., Hamelin, B., Camoin, G., Thomas, A.L., Henderson,
746 G.M., Okuno, J., Yokoyama, Y., 2012. Ice-sheet collapse and sea-level rise at the Bølling warming
747 14,600 years ago. *Nature* 483, 559–564. <https://doi.org/10.1038/nature10902>
- 748 Dobson, J.E., 2014. *Aquaterra Incognita: Lost Land Beneath The Sea*. *Geographical Review*
749 104, 123–138. <https://doi.org/10.1111/j.1931-0846.2014.12013.x>
- 750 Domijan, N., Leder, N., Čupić, S., 2005. Visinski datumi Republike Hrvatske (Vertical datums
751 of the Republic of Croatia), in: *Zbornik radova trećeg hrvatskog kongresa o katastru s*
752 *međunarodnim sudjelovanjem*. Croatia.
- 753 Državni Hidrografski Institut, 1998. *Jadransko More*, charts MK17, MK-19, MK-20, MK-21,
754 MK-22, MK-23, MK-24, 535.
- 755 Fairbanks, R.G., 1989. A 17,000-year glacio-eustatic sea level record: influence of glacial
756 melting rates on the Younger Dryas event and deep-ocean circulation. *Nature* 342, 637–642.
757 <https://doi.org/10.1038/342637a0>
- 758 Faivre, S., Bakran-Petricioli, T., Horvatinčić, N., Sironić, A., 2013. Distinct phases of relative
759 sea level changes in the central Adriatic during the last 1500 years — influence of climatic

- 760 variations? *Palaeogeography, Palaeoclimatology, Palaeoecology* 369, 163–174.
761 <https://doi.org/10.1016/j.palaeo.2012.10.016>
- 762 Faivre, S., Butorac, V., 2018. Recently submerged tidal notches in the wider Makarska area
763 (Central Adriatic, Croatia). *Quaternary International* 494, 225–235.
764 <https://doi.org/10.1016/j.quaint.2017.07.020>
- 765 Farbstein, R., Radić, D., Brajković, D., Miracle, P.T., 2012. First Epigravettian Ceramic
766 Figurines from Europe (Vela Spila, Croatia). *PLOS ONE* 7, e41437.
767 <https://doi.org/10.1371/journal.pone.0041437>
- 768 Felja, I., Fontana, A., Furlani, S., Bajraktarević, Z., Paradžik, A., Topalović, E., Rossato, S.,
769 Čosović, V., Juračić, M., 2015. Environmental changes in the lower Mirna River valley (Istria,
770 Croatia) during Upper Holocene. *Geologia Croatica* 68, 209–224.
771 <https://doi.org/10.4154/GC.2015.16>
- 772 Foglini, F., Prampolini, M., Micallef, A., Angeletti, L., Vandelli, V., Deidun, A., Soldati, M.,
773 Taviani, M., 2016. Late Quaternary coastal landscape morphology and evolution of the Maltese
774 Islands (Mediterranean Sea) reconstructed from high-resolution seafloor data. *Geological Society,*
775 *London, Special Publications* 411, 77–95.
- 776 Fontana, A., Vinci, G., Tasca, G., Mozzi, P., Vacchi, M., Bivi, G., Salvador, S., Rossato, S.,
777 Antonioli, F., Asioli, A., Bresolin, M., Di Mario, F., Hajdas, I., 2017. Lagoonal settlements and
778 relative sea level during Bronze Age in Northern Adriatic: Geoarchaeological evidence and
779 paleogeographic constraints. *Quaternary International* 439, 17–36.
780 <https://doi.org/10.1016/j.quaint.2016.12.038>
- 781 Forenbaier, S., Kaiser, T., Miracle, P.T., 2013. Dating the East Adriatic Neolithic. *European*
782 *Journal of Archaeology* 16, 589–609. <https://doi.org/10.1179/1461957113Y.0000000038>

- 783 Forenbaher, S., Miracle, P.T., 2006. Pupićina Cave and the spread of farming in the eastern
784 Adriatic, in: Miracle, P., Forenbaher, S. (Eds.), Prehistoric Herders of Northern Istria: The
785 Archaeology of Pupićina Cave. Archaeological Museum of Istria, Pula, pp. 483–523.
- 786 Galili, E., Benjamin, J., Eshed, V., Rosen, B., McCarthy, J., Horwitz, L.K., 2019. A submerged
787 7000-year-old village and seawall demonstrate earliest known coastal defence against sea-level rise.
788 PLoS ONE 14. <https://doi.org/10.1371/journal.pone.0222560>
- 789 Garg, K.M., Chattopadhyay, B., Wilton, P.R., Malia Prawiradilaga, D., Rheindt, F.E., 2018.
790 Pleistocene land bridges act as semipermeable agents of avian gene flow in Wallacea. *Molecular*
791 *Phylogenetics and Evolution* 125, 196–203. <https://doi.org/10.1016/j.ympev.2018.03.032>
- 792 Giglio, F., Romano, S., Albertazzi, S., Chiarini, F., Ravaioli, M., Ligi, M., Capotondi, L., 2020.
793 Sediment Dynamics of the Neretva Channel (Croatia Coast) Inferred by Chemical and Physical
794 Proxies. *Applied Sciences* 10, 807. <https://doi.org/10.3390/app10030807>
- 795 Gkinis, V., Simonsen, S.B., Buchardt, S.L., White, J.W.C., Vinther, B.M., 2014. Water isotope
796 diffusion rates from the NorthGRIP ice core for the last 16,000 years – Glaciological and
797 paleoclimatic implications. *Earth and Planetary Science Letters* 405, 132–141.
798 <https://doi.org/10.1016/j.epsl.2014.08.022>
- 799 Govorčin, D.P., Juračić, M., Horvatinčić, N., Onofri, V., 2001. Holocene sedimentation in the
800 Soline Channel (Mljet Lakes, Adriatic Sea). *Natura Croatica* 10, 247–258.
- 801 Hewitt, G., 2000. The genetic legacy of the Quaternary ice ages. *Nature* 405, 907–913.
802 <https://doi.org/10.1038/35016000>
- 803 HHI, 2019. Hrvatski Hidrografski Institut [WWW Document]. Hrvatski Hidrografski Institut.
804 URL <http://www.hhi.hr/en/mareo> (accessed 12.28.19).
- 805 Janković, I., Ahern, J.C.M., Karavanić, I., Stockton, T., Smith, F.H., 2012. Epigravettian
806 Human Remains and Artifacts from Šandalja II, Istria, Croatia. *PaleoAnthropology*. Official
807 Publication of the Paleoanthropology Society. 2012, 87–122.

- 808 Juračić, M., Surić, M., 2010. Late Pleistocene–Holocene environmental changes–records from
809 submerged speleothems along the Eastern Adriatic coast (Croatia). *Geologia Croatica* 63, 155–169.
- 810 Kaczanowska, M., Kozłowski, J.K., 2018. The Mesolithic lithic industries of the eastern
811 Adriatic zone. *Folia Quaternaria* 191–215.
- 812 Karavanić, I., 2003. L'industrie aurignacienne de la grotte de Šandalja II (Istrie, Croatie) dans
813 le contexte de la région de l'Est de l'Adriatique. *L'Anthropologie* 107, 577–602.
814 <https://doi.org/10.1016/j.anthro.2003.10.003>
- 815 Kelletat, D.H., 2005. Dalmatian Coasts. *Encyclopedia of coastal science, Encyclopedia of earth*
816 *sciences series*.
- 817 Kendall, R.A., Mitrovica, J.X., Milne, G.A., Törnqvist, T.E., Li, Y., 2008. The sea-level
818 fingerprint of the 8.2 ka climate event. *Geology* 36, 423–426. <https://doi.org/10.1130/G24550A.1>
- 819 Komšo, D., 2008. Mezolitik u Hrvatskoj. *Opusc. Archaeol. Rad. Arheol. zavoda* 30, 55–92.
- 820 Koncani Uhač, I., Čuka, I., 2015. M., 2015–Doprinos poznavanju podmorskog eneolitičkog
821 nalazišta u uvali Zambratija, *Histria archaeologica* 46. Pula 25–73.
- 822 Kozłowski, J.K., Kaczanowska, M., 2004. Gravettian/Epigravettian sequences in the Balkans
823 and Anatolia. *Mediterranean Archaeology and Archaeometry* 4, 5–18.
- 824 Lambeck, K., Purcell, A., 2005. Sea-level change in the Mediterranean Sea since the LGM:
825 model predictions for tectonically stable areas. *Quaternary Science Reviews* 24, 1969–1988.
826 <https://doi.org/10.1016/j.quascirev.2004.06.025>
- 827 Lambeck, K., Purcell, A., Johnston, P., Nakada, M., Yokoyama, Y., 2003. Water-load definition
828 in the glacio-hydro-isostatic sea-level equation. *Quaternary Science Reviews* 22, 309–318.
829 [https://doi.org/10.1016/S0277-3791\(02\)00142-7](https://doi.org/10.1016/S0277-3791(02)00142-7)
- 830 Lightfoot, E., Boneva, B., Miracle, P.T., Šlaus, M., O'Connell, T.C., 2011. Exploring the
831 Mesolithic and Neolithic transition in Croatia through isotopic investigations. *Antiquity* 85, 73–86.
832 <https://doi.org/10.1017/S0003598X00067442>

- 833 Mauch Lenardić, J., Oros Sršen, A., Radović, S., 2018. Quaternary fauna of the Eastern
834 Adriatic (Croatia) with the special review on the Late Pleistocene sites. *Quaternary International*
835 494, 130–151. <https://doi.org/10.1016/j.quaint.2017.11.028>
- 836 McLaughlin, T.R., Stoddart, S., Malone, C., 2018. Island risks and the resilience of a
837 prehistoric civilization. *World Archaeology* 50, 570–600.
838 <https://doi.org/10.1080/00438243.2018.1515656>
- 839 Melini, D., Spada, G., 2019. Some remarks on Glacial Isostatic Adjustment modelling
840 uncertainties. *Geophys J Int* 218, 401–413. <https://doi.org/10.1093/gji/ggz158>
- 841 Miracle, P., 2007. The Late Glacial "Great Adriatic Plain": "Garden of Eden" or "No Man's
842 Land" during the Epipalaeolithic? A view from Istria (Croatia), in: *Late Paleolithic Environments*
843 and Cultural Relations around the Adriatic, BAR International Series. Archaeopress, pp. 41–51.
- 844 Miracle, P., 2002. Mesolithic meals from Mesolithic middens, in: Miracle, P., Milner, N. (Eds.),
845 *Consuming Passions and Patterns of Consumption*. McDonald Inst of Archeological, pp. 65–88.
- 846 Miracle, P., 2001. Feast or famine? Epipalaeolithic subsistence in the northern Adriatic basin. 1
847 28, 177–197. <https://doi.org/10.4312/dp.28.10>
- 848 Miracle, P., Galanidou, N., Forenbahe, S., 2000. Pioneers in the hills: Early Mesolithic
849 foragers at Šebrn Abri (Istria, Croatia). *European Journal of Archaeology* 3, 293–329.
850 <https://doi.org/10.1179/146195700807860837>
- 851 Miracle, P.T., Forenbahe, S., 2005. Neolithic and Bronze-Age Herders of Pupičina Cave,
852 Croatia. *Journal of Field Archaeology* 30, 255–281.
- 853 Mussi, M., Palombo, M.R., 2001. Les carnivores du Pliocène supérieur et du Pléistocène de la
854 péninsule italienne: quelles implications pour le peuplement humain. Brugal, J. Ph., Fosse, P.(eds),
855 *Actes du XIVe Congrès UISPP «Hommes & Carnivores au Paléolithique»*, Liège 597–609.
- 856 Mussi, M., Peresani, M., 2011. Human settlement of Italy during the Younger Dryas.
857 *Quaternary international* 242, 360–370.

858 NASA, 2000. Shuttle Radar Topography Mission (SRTM) 1 Arc-Second Global Elevation
859 Data.

860 Neumann, B., Vafeidis, A.T., Zimmermann, J., Nicholls, R.J., 2015. Future Coastal Population
861 Growth and Exposure to Sea-Level Rise and Coastal Flooding - A Global Assessment. PLOS ONE
862 10, e0118571. <https://doi.org/10.1371/journal.pone.0118571>

863 Nunn, P.D., 2020. In Anticipation of Extirpation How Ancient Peoples Rationalized and
864 Responded to Postglacial Sea Level Rise. *Environmental Humanities* 12, 113–131.
865 <https://doi.org/10.1215/22011919-8142231>

866 Olson, C.J., Becker, J.J., Sandwell, D.T., 2016. SRTM15_PLUS: Data Fusion of Shuttle Radar
867 Topography Mission (SRTM) Land Topography With Measured and Estimated Seafloor
868 Topography (NCEI Accession 0150537).

869 Olson, C.J., Becker, J.J., Sandwell, D.T., 2014. A new global bathymetry map at 15 arcsecond
870 resolution for resolving seafloor fabric: SRTM15_PLUS. AGU Fall Meeting Abstracts 2014,
871 OS34A-03.

872 Palombo, M.R., Antonioli, F., Lo Presti, V., Mannino, M.A., Melis, R.T., Orru, P., Stocchi, P.,
873 Talamo, S., Quarta, G., Calcagnile, L., Deiana, G., Altamura, S., 2017. The late Pleistocene to
874 Holocene palaeogeographic evolution of the Porto Conte area: Clues for a better understanding of
875 human colonization of Sardinia and faunal dynamics during the last 30 ka. *Quaternary International*
876 439, 117–140. <https://doi.org/10.1016/j.quaint.2016.06.014>

877 Peltier, W.R., 2004. Global Glacial Isostasy and the Surface of the Ice-Age Earth: The Ice-5g
878 (vm2) Model and Grace. *Annual Review of Earth and Planetary Sciences* 32, 111–149.
879 <https://doi.org/10.1146/annurev.earth.32.082503.144359>

880 Peltier, W.R., Argus, D.F., Drummond, R., 2015. Space geodesy constrains ice age terminal
881 deglaciation: The global ICE-6G_C (VM5a) model. *Journal of Geophysical Research: Solid Earth*
882 120, 450–487. <https://doi.org/10.1002/2014JB011176>

- 883 Peltier, W.R., Fairbanks, R.G., 2006. Global glacial ice volume and Last Glacial Maximum
884 duration from an extended Barbados sea level record. *Quaternary Science Reviews, Critical*
885 *Quaternary Stratigraphy* 25, 3322–3337. <https://doi.org/10.1016/j.quascirev.2006.04.010>
- 886 Pikelj, K., Juračić, M., 2013. Eastern Adriatic Coast (EAC): Geomorphology and Coastal
887 Vulnerability of a Karstic Coast. *Journal of Coastal Research* 289, 944–957.
888 <https://doi.org/10.2112/JCOASTRES-D-12-00136.1>
- 889 Pirazzoli, P.A., 2013. Geomorphological Indicators, in: *Encyclopedia of Quaternary Science*.
890 Elsevier, pp. 377–384. <https://doi.org/10.1016/B978-0-444-53643-3.00133-3>
- 891 Pross, J., Kotthoff, U., Müller, U.C., Peyron, O., Dormoy, I., Schmiedl, G., Kalaitzidis, S.,
892 Smith, A.M., 2009. Massive perturbation in terrestrial ecosystems of the Eastern Mediterranean
893 region associated with the 8.2 kyr B.P. climatic event. *Geology* 37, 887–890.
894 <https://doi.org/10.1130/G25739A.1>
- 895 Radić, D. (Ed.), 2018. *Island of Korčula: From the earliest times to the end of prehistory*.
896 Centar za kulturu “Vela Luka,” Vela Luka, Croatia.
- 897 Radić, D., Lugović, B., Marjanac, L., 2007. Neapolitan Yellow Tuff (NYT) from theleistocene
898 sediments in Vela Spila on the island of Korčula: a valuable chronostratigraphic marker of the
899 transition from the Palaeolithic to the Mesolithic. *Opvscvla archaeologica* 31, 20.
- 900 Radić Rossi, I., 2008. Arheološka baština u podmorju Kaštelanskog zaljeva. *Archaeologia*
901 *Adriatica* 2, 489–506.
- 902 Rainsford, C., O’Connor, T., Miracle, P., 2014. Fishing in the Adriatic at the Mesolithic–
903 Neolithic transition: Evidence from Vela Spila, Croatia. *Environmental Archaeology* 19, 311–320.
904 <https://doi.org/10.1179/1749631414Y.0000000018>
- 905 Richards, M.P., Karavanić, I., Pettitt, P., Miracle, P., 2015. Isotope and faunal evidence for high
906 levels of freshwater fish consumption by Late Glacial humans at the Late Upper Palaeolithic site of

- 907 Šandalja II, Istria, Croatia. *Journal of Archaeological Science* 61, 204–212.
908 <https://doi.org/10.1016/j.jas.2015.06.008>
- 909 Rizner, M., Vukosavljević, N., Miracle, P., 2009. The paleoecological and paleodietary
910 significance of edible land snails (*Helix* sp.) across the Pleistocene-Holocene transition on the
911 eastern Adriatic coast, in: *MESO 2005: The 7th International Conference on the Mesolithic in*
912 *Europe*.
- 913 Rossi, I.R., 2017. Proučavanje promjena morske razine na arheološkim nalazištima u priobalju.
914 *Vjesnik za Arheologiju i Povijest Dalmatinsku* 110.
- 915 Rovere, A., Stocchi, P., Vacchi, M., 2016. Eustatic and Relative Sea Level Changes. *Curr Clim*
916 *Change Rep* 2, 221–231. <https://doi.org/10.1007/s40641-016-0045-7>
- 917 Sandwell, D.T., Becker, J.J., Smith, W.H.F., 2014a. SRTM15_PLUS Estimated Topography, 15
918 seconds, Global, v1.
- 919 Sandwell, D.T., Müller, R.D., Smith, W.H.F., Garcia, E., Francis, R., 2014b. New global marine
920 gravity model from CryoSat-2 and Jason-1 reveals buried tectonic structure. *Science* 346, 65–67.
921 <https://doi.org/10.1126/science.1258213>
- 922 Shaw, T.A., Plater, A.J., Kirby, J.R., Roy, K., Holgate, S., Tutman, P., Cahill, N., Horton, B.P.,
923 2018. Tectonic influences on late Holocene relative sea levels from the central-eastern Adriatic
924 coast of Croatia. *Quaternary Science Reviews* 200, 262–275.
925 <https://doi.org/10.1016/j.quascirev.2018.09.015>
- 926 Shennan, I., Long, A.J., Horton, B.P. (Eds.), 2015. *Handbook of Sea-Level Research*. John
927 Wiley & Sons, Ltd, Chichester, UK.
- 928 Sikora, M., Mihanović, H., Vilibić, I., 2014. Paleo-coastline of the Central Eastern Adriatic Sea,
929 and Paleo-Channels of the Cetina and Neretva rivers during the last glacial maximum. *Acta*
930 *Adriatica* 55, 3–18.

- 931 Smith, W.H.F., Sandwell, D.T., 1997. Global Sea Floor Topography from Satellite Altimetry
932 and Ship Depth Soundings. *Science* 277, 1956–1962. <https://doi.org/10.1126/science.277.5334.1956>
- 933 Šošić Klindžić, R., Radović, S., Težak Gregl, T., Šlaus, M., Perhoć, Z., Altherr, R., Hulina, M.,
934 Gerometta, K., Boschian, G., Vukosavljević, N., Ahern, J.C.M., Janković, I., Richards, M.,
935 Karavanić, I., 2015. Late Upper Paleolithic, Early Mesolithic and Early Neolithic from the cave site
936 Zemunica near Bisko (Dalmatia, Croatia) 12, 3–46.
- 937 Spada, G., Galassi, G., 2017. Extent and dynamic evolution of the lost land aquaterra since the
938 Last Glacial Maximum. *Comptes Rendus Geoscience* 349, 151–158.
- 939 Spada, G., Stocchi, P., 2007. SELEN: A Fortran 90 program for solving the “sea-level
940 equation.” *Computers & Geosciences* 33, 538–562. <https://doi.org/10.1016/j.cageo.2006.08.006>
- 941 Srdoč, D., Obelić, B., Horvatinčić, N., Bronić, I.K., Sliepčević, A., 1989. Rudjer Bošković
942 Institute radiocarbon measurements XI. *Radiocarbon* 31, 85–98.
- 943 Stewart, J.R., Van Kolfschoten, M., Markova, A., Musil, R., 2003. The mammalian faunas of
944 Europe during oxygen isotope stage three. Neanderthals and modern humans in the European
945 landscape during the last glaciation 60, 103–130.
- 946 Taviani, M., Angeletti, L., Campiani, E., Ceregato, A., Foglini, F., Maselli, V., Morsilli, M.,
947 Parise, M., Trincardi, F., 2012. Drowned karst landscape offshore the Apulian margin (Southern
948 Adriatic Sea, Italy). *Journal of Cave and Karst Studies* 74, 197–212.
949 <https://doi.org/10.4311/2011JCKS0204>
- 950 tide-forecast.com, 2019. Tide Times and Tide Charts Worldwide [WWW Document].
951 [Meteo365.com](https://www.tide-forecast.com). URL <https://www.tide-forecast.com> (accessed 7.27.19).
- 952 Tozer, B., Sandwell, D.T., Smith, W.H.F., Olson, C., Beale, J.R., Wessel, P., 2019. Global
953 Bathymetry and Topography at 15 Arc Sec: SRTM15+. *Earth and Space Science* 6, 1847–1864.
954 <https://doi.org/10.1029/2019EA000658>

- 955 Trincardi, F., Campiani, E., Correggiari, A., Foglini, F., Maselli, V., Remia, A., 2014.
- 956 Bathymetry of the Adriatic Sea: The legacy of the last eustatic cycle and the impact of modern
957 sediment dispersal. *Journal of Maps* 10, 151–158. <https://doi.org/10.1080/17445647.2013.864844>
- 958 Vacchi, M., Marriner, N., Morhange, C., Spada, G., Fontana, A., Rovere, A., 2016. Multiproxy
959 assessment of Holocene relative sea-level changes in the western Mediterranean: Sea-level
960 variability and improvements in the definition of the isostatic signal. *Earth-Science Reviews* 155,
961 172–197. <https://doi.org/10.1016/j.earscirev.2016.02.002>
- 962 Vlahović, I., Tišljarić, J., Velić, I., Matičec, D., 2005. Evolution of the Adriatic Carbonate
963 Platform: Palaeogeography, main events and depositional dynamics. *Palaeogeography,*
964 *Palaeoclimatology, Palaeoecology* 220, 333–360. <https://doi.org/10.1016/j.palaeo.2005.01.011>
- 965 Vlahović, I., Tišljarić, J., Velić, I., Matičec, D., 2002. The Karst Dinarides are Composed of
966 Relics of a Single Mesozoic Platform: Facts and Consequences. *Geologia Croatica* 55, 171–183.
- 967 Vogel, H., Wagner, B., Zanchetta, G., Sulpizio, R., Rosén, P., 2010. A paleoclimate record with
968 tephrochronological age control for the last glacial-interglacial cycle from Lake Ohrid, Albania and
969 Macedonia. *J Paleolimnol* 44, 295–310. <https://doi.org/10.1007/s10933-009-9404-x>
- 970 Vukosavljević, N., Perhoč, Z., 2017. Lithic raw material procurement of the Late Epigravettian
971 hunter-gatherers from Kopačina Cave (island of Brač, Dalmatia, Croatia). *Quaternary International,*
972 *Prehistoric hunter-gatherers and farmers in the Adriatic and neighboring regions* 450, 164–185.
973 <https://doi.org/10.1016/j.quaint.2016.09.017>
- 974 Vukosavljević, N., Perhoč, Z., Altherr, R., 2014. Pleistocene-Holocene transition in the Vlakno
975 Cave on the island of Dugi otok (Dalmatia, Croatia)—lithic perspective. *Prilozi Instituta za*
976 *arheologiju u Zagrebu* 31, 5–72.
- 977 Vukosavljević, N., Perhoč, Z., Čečuk, B., Karavanić, I., 2011. Late Glacial knapped stone
978 industry of Kopačina Cave. *Vjesnik za arheologiju i povijest dalmatinsku* 104, 7–54.

979 Wacha, L., Frechen, M., Crnjaković, M., 2011a. The loess chronology of the Island of Susak,
980 Croatia. *Quaternary Science International* 60, 153–169.

981 Wacha, L., Pavlaković, S.M., Novothny, Á., Crnjaković, M., Frechen, M., 2011b.
982 Luminescence dating of Upper Pleistocene loess from the Island of Susak in Croatia. *Quaternary*
983 *International* 234, 50–61.

984 Weber, M.-J., Grimm, S.B., Baales, M., 2011. Between warm and cold: Impact of the Younger
985 Dryas on human behavior in Central Europe. *Quaternary International* 242, 277–301.

986 Weigelt, P., Steinbauer, M.J., Cabral, J.S., Kreft, H., 2016. Late Quaternary climate change
987 shapes island biodiversity. *Nature* 532, 99.

988 Whallon, R., 1999. The lithic tool assemblages at Badanj within their regional context, in:
989 Bailey, G.N., Adam, E., Panagopoulou, E., Perles, C., Zachos, K. (Eds.), *The Palaeolithic*
990 *Archaeology of Greece and Adjacent Areas*, Monograph of the British School at Athens. Presented
991 at the Proceedings of the ICOPAG Conference, British School at Athens, Athens.

992 Wunsam, S., Schmidt, R., Müller, J., 1999. Holocene lake development of two Dalmatian
993 lagoons (Malo and Veliko Jezero, Isle of Mljet) in respect to changes in Adriatic sea level and
994 climate. *Palaeogeography, Palaeoclimatology, Palaeoecology* 146, 251–281.

995 Zvelebil, M., Lillie, M.C., 2000. Transition to agriculture in Eastern Europe, in: Price, T.D.
996 (Ed.), *Europe's First Farmers*. Cambridge University Press, New York.

997

998

999 **Figures captions**

1000 Fig 1. Map of the Adriatic Sea and study area. A: general map of the Adriatic Sea, divided into
 1001 three sub-basins, and surrounding areas. B: Korčula and surrounding Islands. Tectonic information:
 1002 Korbar, 2009; Surić et al., 2014. Map and relief data: GEBCO and GADM.

1003 Fig 2. Vela Spila and surroundings. Top: View from the cave towards the SW, showing Vela
 1004 Luka town and harbour (photo from P.T. Miracle). Middle left: View of Vela Luka bay looking
 1005 northeast. Hvar and the mainland in background. Vela Spila is in the centre hill of the image (photo
 1006 from G. Boschian) Middle right: Another view from Vela Spila, but looking west out Vela Luka bay.
 1007 The innermost harbour is visible (photo from F. Rinaldi). Bottom: Interior photomosaic of Vela
 1008 Spila. The most recently excavated trenches are on the left (photo from G. Boschian).

1009 Fig 3. Plan of Vela Spila cave, with locations of archaeological excavations. 1: dripline of
 1010 entrance and of windows in ceiling; 2: excavations pre-2000; 3: excavations 2000-2018; A-A'
 1011 location of the profile in fig. 4.

1012 Fig 4. General stratigraphy of the Vela Spila archaeological deposit. From left to right: cultural
 1013 horizons and informal Quaternary stratigraphy, photo and main lithostratigraphic subdivisions of
 1014 vertical profile (A-A' in fig. 3). NYT: Neapolitan Yellow Tuff; red lines: erosive boundaries. Note
 1015 the wavy limits of NYT and overlying units, due to frost action in sediment.

1016 Fig 5. Summary of calibrated dates from Vela Spila. Note the absence of archaeological
 1017 material for dating during the abandonment phase.

1018 Fig 6. Sea level reconstructions for Vela Luka. Sea-level realizations for the Vela Luka area of
 1019 Korčula. Dashed line: I6G_S4 realization of the ICE-6G_C (VM5a) (Peltier et al., 2015). Dotted
 1020 line: ANU_S4 realization of Lambeck/ANU (Lambeck et al., 2003; Lambeck and Purcell, 2005).
 1021 Sea-level indicators from Faivre et al. (2013), Juračić and Surić (2010), and Mljet lakes (Govorčin
 1022 et al., 2001; Wunsam et al., 1999) shown in green. Archaeological periods for the site reconstructed
 1023 with 14C dates from Table 2. Exact period delineations are an abstraction, in reality date
 1024 uncertainties and gradual cultural shifts mean the transitions were nuanced.

1025 Fig 7. Palaeogeography and sea-level change at Vela Spila. Palaeogeographic reconstruction of
 1026 the area around Korčula at different phases of Vela Spila's settlement history. A) Start of the
 1027 Palaeolithic ca. 19.7 cal ka BP. B) Start of the abandonment phase ca. 14.0 cal ka BP. C) Start of the
 1028 Mesolithic ca. 9.4 cal ka BP. Vela Spila location shown by red dot. Light blue shows ICE6G
 1029 predicted RSL; dark blue shows Lambeck/ANU predicted RSL. Dotted blue line indicates probable
 1030 location of palaeo-Neretva river according to Sikora et al.(2014). For each time-slice, a red line or
 1031 lines indicates distance to sea from Vela Spila. The crosshatched circle indicates the percent of land
 1032 within a 20 km radius from the cave that is inundated. Map elevations interpolated from datasets
 1033 listed in Table 1.

1034 Fig 8. Slope vs. depth in Western Korčula. Profile of 15 km transects AB and CD from Fig 5C,
 1035 showing the higher slopes that predominate above the 70 m isobath (note, Y axis in left panels is
 1036 exaggerated). Right panel shows slope in degrees vs. depth below sea level along transect CD.
 1037 Slopes increase above -70 m. This morphology results in negligible horizontal coastline change
 1038 after ca. 12 ka BP.

1039 Fig 9. A zoomed in view of the palaeo shorelines at the start of the Mesolithic (~9.4 ka BP) at
 1040 Vela Spila, overlaid on modern orthophotos (Croatian Geoportal, 2019). ANU and ICE models for
 1041 sea-level are shown. The situation of the site on a hill at ~121 m ASL above the harbour valley is
 1042 evident. To the south and southeast, where the town of Vela Luka is now situated the western end of
 1043 the flat karstic depression known as Blatsko Polje can be seen.

1044

Figure 1

[Click here to access/download;Figure;Fig1_regional_map_double_column.tif](#)








Figure 3





OxCal v4.3.2 Bronk Ramsey (2017); r5 IntCal13 atmospheric curve (Reimer et al 2013)





Figure 6









Figure 8





Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: