

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

How siderophore production can influence the biocontrol activity of *Aureobasidium pullulans* against *Monilinia laxa* on peaches

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

alessandra di francesco, elena baraldi (2021). How siderophore production can influence the biocontrol activity of *Aureobasidium pullulans* against *Monilinia laxa* on peaches. *BIOLOGICAL CONTROL*, 152(January 2021), 1-6 [10.1016/j.biocontrol.2020.104456].

Availability:

This version is available at: <https://hdl.handle.net/11585/775215> since: 2020-10-20

Published:

DOI: <http://doi.org/10.1016/j.biocontrol.2020.104456>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

How siderophore production can influence the biocontrol activity of *Aureobasidium pullulans* against *Monilinia laxa* on peaches

Alessandra Di Francesco, Elena Baraldi

PII: S1049-9644(20)30683-6
DOI: <https://doi.org/10.1016/j.biocontrol.2020.104456>
Reference: YBCON 104456

To appear in: *Biological Control*

Received Date: 16 October 2019
Revised Date: 30 June 2020
Accepted Date: 8 October 2020

Please cite this article as: Di Francesco, A., Baraldi, E., How siderophore production can influence the biocontrol activity of *Aureobasidium pullulans* against *Monilinia laxa* on peaches, *Biological Control* (2020), doi: <https://doi.org/10.1016/j.biocontrol.2020.104456>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

How siderophore production can influence the biocontrol activity of *Aureobasidium pullulans* against *Monilinia laxa* on peaches

Alessandra Di Francesco*, Elena Baraldi

Department of Agricultural Sciences, University of Bologna, Viale G. Fanin, 46, 40127 Bologna, Italy

*corresponding author: alessand.difrancesc3@unibo.it

Abstract

In the present study *Aureobasidium pullulans* strains L1 and L8 were shown both *in vitro* and *in vivo* to compete for iron with *Monilinia laxa* through the secretion of siderophores (1.2 and 1.4 mg ml⁻¹, respectively) and to prevent postharvest fruit decay of peaches. The two strains reduced mycelial growth and conidial germination of *M. laxa* specially in presence of lower iron concentrations (5 and 10 µg l⁻¹ FeCl₃), confirming a better efficacy when nutrients are scarce. In *in vivo* assay, *A. pullulans* L1 and L8 strains inhibited pathogen virulence, reducing by 83.5% and 84.4% on average respectively the peach lesion diameter for each tested iron solutions (5, 10, 20 µg l⁻¹ FeCl₃). The highest iron solution slowed down the antagonists' action and conversely increase the pathogen aggressiveness. Results shows that *A. pullulans* L1 and L8 strains compete with *M. laxa* for iron, so revealing new biocontrol aspects. Both strains showed the capability to decrease the accumulation of iron competing with some fungal pathogens and reducing their virulence. These results provide new perspectives for the use of biocontrol agents in agriculture.

Keywords: Yeast - Iron – Stone fruit – *Monilinia laxa* - Siderophore

1. Introduction

Monilinia laxa is the most common species in European stone fruit orchards, and able to cause postharvest losses that reach high values (59%) (Larena et al., 2005). *Monilinia* control depends on an integrated strategy based on fungicide use and cultural practices. However, to overcome the issues related to the use of fungicides, alternative pathogen control strategies have been investigated, such as the use of biocontrol agents (BCAs) (Di Francesco et al., 2016). *Aureobasidium pullulans* L1 and L8 strains showed a high capability to control *Monilinia* spp. on stone fruit as previously reported by Mari et al. (2012).

Among the various modes of action involved in the control of postharvest diseases by antagonistic yeasts, competition for nutrients and space is considered one of the most important, particularly to control wound pathogens for which the availability of exogenous nutrients is crucial during the early stage of infection (Elad and Chet 1987, Mekbib et al., 2011, Bautista-Rosales et al., 2013, Di Francesco et al., 2016, 2017, 2018a).

In fact, previous studies reported how the antifungal ability of *A. pullulans* (Janisiewicz et al., 2000; Bencheqroun et al., 2006), *Pichia caribbica* (Xu et al., 2013), and *Pichia guilliermondii* (Chanchaichaovivat et al., 2008) against *Penicillium expansum*, *Rhizopus stolonifer*, and *Colletotrichum capsici* respectively, can be influenced by the presence or the absence of exogenous nutrients (Druvefors et al., 2005, Bencheqroun et al., 2006, Liu et al., 2010).

Besides the role of competing for nutrients, iron may also play a role in the Biocontrol Agents (BCAs) modes of action (Raaska and Mattila-Sandholm 1995) through the production of chelating molecules (siderophores).

These are molecules able to chelate low weight ferric compounds enhancing the effectiveness of BCAs by subtracting iron to pathogen inhibiting its growth and metabolic activity (Riquelme, 1996). Microorganisms can produce a variety of chelating agents, which solubilize ferric iron and transport it into the cell (Liu et al., 2013, Calvente et al., 2001a, Sanz Ferramola et al., 2013). Under competitive conditions when nutrient availability is a limiting factor, the production of siderophores can represent a distinctive feature for a microorganism. It is known that yeasts produce only a type

of siderophore: hydroxamate (Riquelme, 1996). The siderophores production could play an important role in the biocontrol of pathogens because by iron sequestering their growth and metabolic activity were inhibited or slowed down (Riquelme 1996). Calvente et al. (2001a) demonstrated that rhodotorulic acid, a siderophore produced by *Rhodotorula glutinis* improved the biological control of blue rot of apples caused by *P. expansum*. Also, *Bacillus subtilis* produced siderophores playing an important role in the control of *Fusarium oxysporum* (Yu et al., 2011). Other studies reported the siderophores ability to inhibit the growth of certain pathogenic fungi, like *Pythium ultimum* and *Sclerotinia sclerotiorum* (Hamdan et al., 1991, McLoughlin et al., 1992), establishing that these molecules could be considered as a potential mechanism of action against several fungal pathogens. Parafati et al. (2015) hypothesized that competition for iron was as one of the main mechanisms of action for *Metschnikowia pulcherrima* against *Botrytis cinerea*. In a previous study, *A. pullulans* L1 and L8 strains yet showed competition for nutrients like amino acids and space (Di Francesco et al., 2017, 2018a) against *Monilinia laxa* and *B. cinerea*. In the present work, we investigated for the first time the involvement of iron in the biocontrol activity of both strains against the brown rot agent of peaches. Specifically, the aims of this study were to determine the ability of L1 and L8 to produce siderophores through i) the chrome azurol S (CAS) assay, ii) the spectrophotometer quantification, and iii) the *in vitro* and *in vivo* competition for iron with *M. laxa* by testing three different concentrations of FeCl_3 (5, 10 and 20 $\mu\text{g ml}^{-1}$).

2. Materials and methods

2.1 Fruit

Peaches (*Prunus persica* (L.) Batsch) cv 'Redhaven' were harvested at commercial maturity in orchards located in Cadriano (Bologna, Italy). Fruit were stored at 0 °C and used within 5 days from harvest and wounded by a sterile nail (3x3x3 mm) on opposite sides of the equatorial area.

2.2 Pathogen and antagonists

Monilinia laxa (ML4 strain from CRIOF-DipSA collection) (Di Francesco et al., 2017) was chosen for the present study for its aggressiveness and grown as by Martini et al. (2016). Conidia from pathogen colonies 10 days-old, grown on tomato agar (20 g of Agar Technical, Oxoid Basingstoke, Hampshire, UK; 750 ml distilled water to which 250 ml tomato sauce was added after sterilization) at 25 °C, were collected and suspended in sterile distilled water containing 0.05 %-(v/v) Tween 80. The concentration of each conidia suspension was quantified with haematocytometer and adjusted to a concentration of 10^5 conidia ml⁻¹ with sterile distilled water. *Aureobasidium pullulans* L1 and L8 strains (Di Francesco et al., 2018b), were maintained on nutrient yeast dextrose agar (NYDA: 8 g of nutrient broth, 5 g of yeast extract, 10 g of dextrose and 15 g of agar in 1 L of distilled water) at 4 °C until use. Two days before trials, each antagonist was inoculated on NYDA and incubated at 25 °C for 2 days. The yeast cells were collected in sterile distilled water and adjusted with haematocytometer to a concentration of 10^8 cells ml⁻¹.

2.3 CAS assay

CAS-blue agar was prepared according to Schwyn and Neilands (1987). Dishes were prepared with 30 ml of culturing medium for each antagonist and fungus: NYDA for L1 and L8 and PDA for *M. laxa*. Then the solidified growth media was cut in two halves, one of which was replaced by CAS-blue agar. The dishes were inoculated placing in the centre of the borderline, between the two media, a 6-mm-plug of L1, L8 or *M. laxa* derived from a colony in active growth. The plates were incubated in the dark at 25 °C until the agar colour-change. The CAS reaction was determined by measuring the advance of the colour-change in the CAS-blue agar from blue to different shades of red. The control plates were not inoculated but incubated under the same conditions described above. The sample unit was represented by 10 dishes and the experiment was repeated twice.

2.4 Spectrophotometer siderophore assay

For the siderophore production by yeasts and *M. laxa*, one loop of L1 and L8 cells and one of *M. laxa* conidia were transferred in a siderophore solution as described in Wang et al. (2009) and incubated at 25 °C at 200 rpm for 24 h. In addition, L1 or L8 cells and *M. laxa* were co-cultured in the siderophore culture solution and incubated in the same conditions above mentioned. Aliquots of 5 ml of each sample's suspension were inoculated into 45 ml of new siderophore solution and suddenly incubated for 120 h at 25 °C at 200 rpm. The supernatant of the centrifuged (5.000 rpm for 20 min) cultures (500 µl) was added to 2.5 ml of a solution containing 5 mM FeCl₃ and used for quantitative determination of siderophore by a spectrophotometer at 440 nm (Infinite® 200 PRO-Tecan) after 30 min of incubation at 25 °C. A standard curve (0, 0.05, 0.1, 0.25, 0.5, 0.75 and 1.0 mg ml⁻¹) was prepared with deferoxamine mesylate (Sigma-Aldrich) (one type of hydroxamates) as chelating agent standard (Calvente et al., 2001). The siderophore amount was extrapolated from the deferoxamine mesylate standard curve. The sample unit was represented by three flasks (replicates) for each condition. The experiment was repeated twice.

2.5 *In vitro* competition for iron

The competition for nutrient exerted by L1 and L8 *A. pullulans* strains on *M. laxa* mycelial growth was investigated through the dual interaction between the pathogen and the yeast strains. The experiment conducted with the two strains and the pathogen was assessed on Minimal Medium Agar (MMA: Na₂HPO₄ 6 g, KH₂PO₄ 3 g, NaCl 0.5 g, NH₄Cl 1 g, MgSO₄ 1M, CaCl₂ 0.1M, Sucrose 30 g, Agar technical 20 g) amended with 0, 5, 10 and 20 µg ml⁻¹ of FeCl₃ (Saravanakumar et al., 2008). *Monilinia laxa* mycelial plugs (6 mm of diameter) from 7-day-old colony were inoculated at 30 mm of distance from the plate edge. *Aureobasidium pullulans* L1 and L8 cells were taken from 48 h culture on NYDA by sterile loop and were streaked at 30 mm from the edge of the other side of the plate. Plates were incubated at 25 °C for 5 days and the radial growth of pathogen mycelium was measured. Plates inoculated only with *M. laxa* plug represented the control. The sample unit was represented by five plates (replicates) for each condition. The experiment was performed twice.

130

131 2.6 Influence of iron concentrations on antagonists' biocontrol activity against *Monilinia laxa*
 132 germination and germ tube elongation

133 The competition activity for iron of L1 and L8 was assayed on the germination and germ tube
 134 elongation of *M. laxa* conidia. Tissue culture plates (Costar, Corning Inc., Corning, NY) and culture
 135 plate inserts Millicell-CM (Millipore Corp., Bedford, MA) were used as reported by Janisiewicz et
 136 al., (2000) with some modifications (Di Francesco et al., 2017). Aliquots (120 μ l) of minimal
 137 medium broth (without agar) amended with 0, 5, 10 and 20 μ g ml⁻¹ of FeCl₃ were dispensed in the
 138 wells of culture plates with 40 μ l of the pathogen conidia suspension (10⁵ conidia ml⁻¹), while the
 139 same aliquots of L1 or L8 cell suspensions (10⁸ cells ml⁻¹) were dispensed inside the cylinder
 140 inserts, without physical contact between antagonist and pathogen. The plates were placed at 25 °C
 141 on a rotary shaker at 50 rpm; after 6 h of incubation, cylinders were removed from the wells and 20
 142 μ l of the medium were transferred to a glass slide for microscope (Nikon Eclipse TE2000–E)
 143 observations. The percentage of conidia germination and the germ tube elongation (μ m) were
 144 determined (90 conidia per treatment, 3 microscopic fields with 30 conidia each). Cells without iron
 145 solutions addition were considered as a control. The experiment was performed twice.

146

147 2.7 *In vivo* competition for iron

148 Peach fruits (20) cv 'Redhaven' were wounded and inoculated with 20 μ l of each yeast suspension
 149 10⁸ cell ml⁻¹; after 1 h at room temperature fruit were inoculated with 20 μ l of the pathogen conidia
 150 suspension (10⁵ cells ml⁻¹) and right after its drying, 20 μ l of iron solutions, at concentrations above
 151 mentioned, were added. Fruit treated with sterile distilled water instead of yeast cell suspensions
 152 represented the control. Fruit were stored at 20 °C and 90% of relative humidity (RH) for 7 days
 153 and the lesion diameters determined by the pathogen were recorded. The sample unit was
 154 represented by 45 peaches (15 for replicate) for each antagonist, iron concentration, and control.
 155 The experiment was repeated twice.

2.8 Statistical analysis

Data were statistically handled by one-way analysis of variance (ANOVA). Statistical comparison of means was carried out to reveal the differences between treatments using Tukey's HSD Test ($\alpha = 0.05$). Data were reported as mean values \pm standard error (SE) of two experiments. Before analysis of data, homogeneity of variance was tested by the Kruskal–Wallis test. All analyses were performed with Statgraphics software (version centurion 15.0).

3. Results

3.1 *Aureobasidium pullulans* L1, L8 strains siderophore production

In order to verify the siderophore production by the antagonists and *M. laxa*, CAS-blue agar was used as screening assay for the target microorganisms. Our results showed that both L1 and L8 induced a colour change of CAS-blue agar from blue to dark orange indicating that both *A. pullulans* strains are siderophore producers. However, despite the two strains required the same time (twelve days) to change the CAS-blue-agar from blue to dark orange, the colour change halo produced by both yeasts was different in size, being 20 mm for L1, and 27 mm for L8 (Table 1) (Fig. 1). In fact, L1 strain after 12 days of incubation did not determined a total colour change of CAS-blue agar half, as instead L8. Conversely, *M. laxa* produced a barrage between the two media without any colour change of the indicative agar. Control plates, without plug inoculation, not changed colour after incubation conditions.

3.2 Spectrophotometer quantification of siderophores

To quantify the siderophore production by *A. pullulans* L1 and L8 strains and the influence of *M. laxa* on this activity, a spectrophotometer assay with deferoxamine mesylate standard curve was conducted. Both strains showed a considerable activity, producing 1.2 and 1.4 mg ml⁻¹ of

siderophore by L1 and L8 respectively, while *M. laxa* produced only 0.64 mg ml⁻¹ of siderophore, a quantity not detectable by CAS-blue-agar.

Siderophore quantity was not influenced by *M. laxa* showing the same amount with respect to the yeasts grown alone (Fig. 2).

3.3 *In vitro* competition for iron

Competition for iron was tested co-culturing in MM plates amended with FeCl₃ (0, 5, 10, or 20 µg ml⁻¹) L1 and L8 strains with *M. laxa* isolate. As shown by Table 2, both *A. pullulans* L1 and L8 strains reduced *M. laxa* mycelium diameter by 41.3% and 43.3%, respectively when iron concentration was not present in MM.

L1 and L8 competition was mainly stimulated by the presence of 10 µg ml⁻¹ and 20 µg ml⁻¹ of FeCl₃, both leading to an inhibition of fungal pathogen mycelial growth corresponding to 45% and 47.6% and 43% and 45%, respectively.

Aureobasidium pullulans strain L8 was slightly more effective with respect to L1 in competing for iron, determining a higher inhibition of *M. laxa* mycelial growth on average by 44.7% and 42.9%, respectively.

3.4 Influence of iron concentrations on *M. laxa* germination and germ tube elongation

After 6 h of incubation at 25 °C in MM broth without iron and in MM broth with 5 µg ml⁻¹ of FeCl₃, *M. laxa* conidial germination was reduced by both L1 and L8 strain of 50%, as showed in Table 3. The presence of 10 µg ml⁻¹ of FeCl₃ in the culture medium induced a higher antagonistic activity by both strains against *M. laxa* conidial germination with respect to the other iron concentrations (5 and 20 µg ml⁻¹ of FeCl₃). In effect, conidial germination of *M. laxa* was reduced by 62.5% and 68.7%, respectively by L1 and L8. Conversely, 20 µg ml⁻¹ of FeCl₃ inhibited the strains antagonistic activity, that showed a reduction of *M. laxa* conidia germination of ~35%.

As regards to *M. laxa* germ tube length, the highest inhibition values exerted by L1 and L8 were mainly showed with 5 and 10 $\mu\text{g ml}^{-1}$ of FeCl_3 solution by 59.5%, 63.1% and by 65.9%, 67.6%, respectively. Nevertheless, 10 $\mu\text{g ml}^{-1}$ of FeCl_3 stimulated mostly pathogen germ tube elongation (10.4%) with respect to the other concentrations (Table 4). Anyway, L8 strain was always slightly more effective with respect to L1 in determining a higher inhibition of *M. laxa* conidial germination and germ tube elongation on average by 52.1%, 50.2%, and by 57.7%, 57.1%, respectively.

3.5 *In vivo* competition for iron

The influence of the addition of aliquots of iron solutions at different concentrations on fungal virulence and yeasts antagonistic activity on peach fruit was measured. *Monilinia laxa* growth on peach fruit in absence of antagonists was slightly enhanced by 6.8% only at FeCl_3 concentrations of 10 or 20 $\mu\text{g ml}^{-1}$ whereas no influence was observed on the pathogen aggressiveness at 5 $\mu\text{g ml}^{-1}$, the lowest FeCl_3 concentrations (Fig. 3).

However, L1 and L8 *A. pullulans* strains showed a great capability to inhibit *M. laxa* aggressiveness, reducing the peach lesion diameters by 83.5% and 84.4% on average respectively, at each iron concentration. Furthermore, the highest concentration of iron (20 $\mu\text{g ml}^{-1}$) seemed to slow down the antagonists action and conversely increase the pathogen aggressiveness. In fact, both L1 and L8 controlled *M. laxa* incidence in presence of 20 $\mu\text{g ml}^{-1}$ FeCl_3 on average by 88% with respect to 92%, in presence of the other iron concentrations (data not reported).

4. Discussion

Aureobasidium pullulans (L1 and L8) are considered promising alternatives to fungicides in the reduction of the stone fruits postharvest diseases' incidence caused by *Monilinia* spp. (Mari et al., 2012; Di Francesco et al., 2017). In our work, competition for nutrients was studied and in particular the competition for iron, a nutrient that played a significant role in biocontrol interactions (Raaska and Mattila-Sandholm 1995).

As known, iron is an essential nutrient for all organisms (Wang et al., 2009) acting as a cofactor for enzymes and regulatory proteins involved in many cellular processes (Miethke and Marahiel 2007; Li et al., 2018) but not promptly available due to the low solubility in alkaline environments. One of the main strategy used by microorganisms (such as yeasts) and by plants to obtain iron is the secretion of siderophores (Li et al., 2018), which are low-molecular-weight compounds that chelate iron in the extracellular phase re-entering it in the cells by specific membrane transporters (Hider and Kong 2010). More than 500 siderophores have been discovered from microorganisms and plants and their chemical diversity often renders a variety of biological functions beyond capture iron (Ho et al., 2019). Yeasts produce only hydroxamate-type siderophore (Riquelme 1996) derived from the amino acid ornithine and classified into four structural families: fusarines, coprogens, ferrichromes and rhodotorulic acid (Johnson 2008). Wang et al., (2009) showed that *A. pullulans* could produce 1.1 mg ml⁻¹ hydroxamate type siderophore. In our experiments, both L1 and L8 were evaluated for their capability to produce siderophore by CAS screening assay (Schwyn and Neilands 1987). This assay was used for the screening of siderophore producing microorganisms by the induction of a colour change of the growth medium: from blue to orange, red, or purple. Our results showed that L1 and L8 strains produced siderophore, displaying after twelve days of incubation a colour change of the CAS agar, from blue to red. Moreover, L8 strain induced a more intense agar colour change and extended halo with respect to L1. Conversely, *M. laxa* was not able to produce siderophores in solid medium but produced a clear barrage between the common fungal culture medium (PDA) and the siderophore screening agar (CAS). In fact, CAS agar assay is mainly useful for the identification of microorganisms capable to produce siderophores in high concentrations and characterized by a high affinity for iron (III). Most likely, the assay was not so sensitive to detect the low siderophores concentration produced by *M. laxa*. In effect, these results were supported by the quantitative analysis conducted with the spectrophotometer. Both yeasts were found to produce high levels of chelating molecules (1.2 and 1.4 mg ml⁻¹, respectively) whit respect to the pathogen that seemed instead to be a poor siderophores

259 producer (0.64 mg ml⁻¹). Probably siderophores role vary among different fungal pathosystems
 260 (Chen et al., 2013) and according to the strain intrinsic characteristics. In fact, the importance of
 261 siderophores for fungal pathogenesis was firstly demonstrated for the pathogens such as
 262 *Cochliobulus heterostrophus* (Lee et al., 2005; Oide et al., 2006), *Alternaria* spp. (Chen et al.,
 263 2013), *Ustilago maydis* (Mei et al., 1993) and for the human pathogen *Aspergillus fumigatus*
 264 (Hissen et al., 2005; Schrettl et al., 2007; Chen et al., 2013).

265 In our study, the siderophore production by the yeasts was not affected by the presence of *M. laxa*,
 266 probably for the nutritional and biochemical preferences of the fungus. According to these results,
 267 we can suppose that *M. laxa* probably mainly compete for other nutrients rather than for iron; on the
 268 other side, *A. pullulans* strains L1 and L8 showed a high effective antagonistic action against *M.*
 269 *laxa* in presence of iron (10 µg l⁻¹ FeCl₃), displaying a great capability to create critical conditions
 270 for the pathogen growth.

271 As other microorganisms (Droby et al., 1989, Bencheqroun et al., 2006) both L1 and L8, reduced
 272 their antagonistic efficacy with the addition of high concentrations of exogenous nutrients, e.g. with
 273 20 µg l⁻¹ of FeCl₃ both in *in vitro* and in *in vivo* assays.

274 In fact, the highest reduction of conidial germination, brown rot severity and incidence (data not
 275 shown) in fruit was revealed in presence of lower iron concentrations, confirming that both strains
 276 displayed their antagonistic efficacy when exogenous nutrients were scarce (Calvente et al., 1999,
 277 Calvo et al., 2008, Sanz Ferramola et al., 2013). Plante and Labbè (2019) showed how in
 278 *Schizosaccaromyces pombe* the siderophore synthetase expression remains upregulated under low
 279 iron conditions.

280 According with previous observations on nutritional requirements of *Monilinia* spp. germination
 281 (Byrde and Willetts 1977), in the present research, the germination rate of the pathogen conidia
 282 resulted low when grown in minimal medium. This confirmed that the target pathogen is nutrient-
 283 dependent and needs an adequate amount of supplements for its growth: both conidial germination
 284 and hyphal development (Bencheqroun et al., 2006, Di Francesco et al., 2017). Moreover, the two

BCAs notably reduced *M. laxa* germ tube elongation: a fungal characteristic directly related to the pathogen aggressiveness (Fujii 2004). In fact, one critical step of germination is the germ tube elongation that can be critically blocked by iron deficiency (Plante and Labbè 2019), in this case caused by the antagonistic action of both L1 and L8 to compete for nutrients.

In conclusion, *A. pullulans* L1 and L8 strains can compete with *M. laxa* for nutrients such as amino acids as previously reported by Di Francesco et al. (2017, 2018a) and in the present study for iron, a vital element for cell proliferation and survival for almost all organisms. Therefore, the ability of *A. pullulans* strains L1 and L8 to acquire iron can counteract the expression of fungal pathogen genes required for the production of siderophores and virulence in fruits (Chen et al., 2013). This information can be very important to exploit L1 and L8 activity in a commercial formulation characterized by a right and efficient composition. More investigations are necessary to isolate and characterize siderophores produced by both yeast strains and possibly verify their environmental involvement in absorption and metabolization of metals. Nevertheless, in our study the strong efficacy exerted by L1 and L8 strains can be attributed most likely to a synergic activity of different mechanisms of action.

Conflict of Interest

The authors declare that they have no conflict of interests.

Authors contribution

Dr. Alessandra Di Francesco conducted all the experiments, interpreted data, and drafted the manuscript. Prof. Elena Baraldi coordinated the experimental plan, has corrected, and approved the final draft of the manuscript.

References

- 311 Bautista-Rosales, P. U., Calderon-Santoya, M., Servin-Villegas, R., Ochoa-Alvarez, N. A.,
 312 Ragazzo-Sanchez, J. A., 2013. Action mechanisms of the yeast *Meyerozyma caribbica* for the
 313 control of the phytopathogen *Colletotrichum gloesporioides* in mangoes. Biol. Control 65, 293-301
 314
- 315 Bencheqroun, S. K., Bajji, M., Massart, S., EI-Jaafari, S., Jijakli, M. H., 2006. *In vitro* and *in situ*
 316 study of postharvest apple blue mould biocontrol by *Aureobasidium pullulans*: evidence for the
 317 involvement of competition for nutrients. Postharvest Biol. Technol. 46, 28–135
 318
- 319 Byrde, R. J. W., Willetts, H. J., 1977. The Brown Rot Fungi of Fruit: Their Biology and Control.
 320 Pergamon Press, Oxford, United Kingdom
 321
- 322 Calvente, V., Benuzzi, D., Sanz de Tosetti, M. I., 1999. Antagonistic action of siderophores from
 323 *Rhodotorula glutinis* upon the postharvest pathogen *Penicillium expansum*. Int. Biodet. Biodeg. J.
 324 43,167-172
 325
- 326 Calvente, V., de Orellano, M. E., Sansone, G., Benuzzi, D., Sanz de Tosetti, M. I., 2001a. Effect of
 327 nitrogen source and pH on siderophore production by *Rhodotorula* strains and their application to
 328 biocontrol of phytopathogenic moulds. J. Ind. Microbiol. Biotechnol. 26, 226-229
 329
- 330 Calvente, V., Orellano, M. E., Sansone, G., Sanz de Tosetti, M. I., 2001b. A simple agar plate assay
 331 for screening siderophore producer yeasts. J. Microbiol. Meth. 47, 273-279
 332
- 333 Calvo, J., Calvente, V., Orellano, M. E., Benuzzi, D., Sanz de Tosetti, M. I., 2008. Control of
 334 *Penicillium expansum* and *Botrytis cinerea* on apple fruit by mixture of bacteria and yeast. Food
 335 Bioprocess Tech. 3, 644-650
 336

- 337 Chanchaichaovivat, A., Panijpan, B., Ruenwongsa, P., 2008. Putative modes of action of *Pichia*
338 *guilliermondii* strain R13 in controlling chilli anthracnose after harvest. *Biol. Control* 47, 207-215
339
- 340 Chen, L. H., Lin, C. H., Chung, K. R., 2013. A nonribosomal peptide synthetase mediates
341 siderophore production and virulence in the citrus fungal pathogen *Alternaria alternata*. *Mol. Plant*
342 *Pathol.* 14, 497–505
343
- 344 Di Francesco A., Martini C., Mari M., 2016. Biological control of postharvest diseases by microbial
345 antagonists: how many mechanisms of action? *Eur. J. Plant Pathol.* 145, 711-717
346
- 347 Di Francesco, A., Ugolini, L., D'aquino, S., Pagnotta, E., Mari, M., 2017. Biocontrol of *Monilinia*
348 *laxa* by *Aureobasidium pullulans* strains: Insights on competition for nutrients and space. *Int. J.*
349 *Food Microbiol.* 248, 32-38
350
- 351 Di Francesco, A., Mari, M., Ugolini, L., Baraldi, E., 2018a. Effect of *Aureobasidium pullulans*
352 strains against *Botrytis cinerea* on kiwifruit during storage and on fruit nutritional composition.
353 *Food Microbiol.* 72, 67-72
354
- 355 Di Francesco, A., Calassanzio, M., Ratti, C., Folchi, A., Baraldi, E., 2018b. Molecular
356 characterization of the two postharvest biological control agents *Aureobasidium pullulans* L1 and
357 L8. *Biol. Control* 123, 53-59
358
- 359 Droby, S., Chalutz, E., Wilson, C. L., Wisniewski, M., 1989. Characterization of the biocontrol
360 activity of *Debaryomyces hansenii* in the control of *Penicillium digitatum* on grape fruit. *Can. J.*
361 *Microbiol.* 35, 794
362

- 363 Druvefors, U. A., Passoth, V., Schnurer, J., 2005. Nutrient effects on biocontrol of *Penicillium*
364 *roqueforti* by *Pichia anomala* J121 during airtight storage of wheat. J. Appl. Environ. Microbiol.
365 71, 1865-1869
366
- 367 Elad, Y., Chet, I., 1987. The role of competition for nutrients in biocontrol of *Pythium* damping off
368 by bacteria. Phytopathology 77, 190-195
369
- 370 Fujii, T. 2004. Germination of spore and decomposition of apple fruit tissue by hypha in
371 *Penicillium expansum*. Mycotoxins 54, 2
372
- 373 Hamdan, H., Weller, D. M., Thomashow, L. S., 1991. Relative importance of fluorescent
374 siderophores and other factors in biological control of *Gaeumannomyces graminis* var. *Tritici* by
375 *Pseudomonas fluorescens* 2-79 and M4-80R. J. Appl. Environ. Microbiol. 57, 3270-3277
376
- 377 Hider RC, Kong X, 2010. Chemistry and biology of siderophores. Natural product reports 27, 637-
378 657
379
- 380 Hissen, A. H. T., Wan, A. N. C., Warwas, M. L., Pinto, L. J., Moore, M. M., 2005. The *Aspergillus*
381 *fumigatus* siderophore biosynthetic gene *sidA*, encoding L-ornithine N5-oxygenase, is required for
382 virulence. Infect. Immun. 73, 5493-5503
383
- 384 Ho, Y. N., Lee, H. J., Hsieh, C. T., Peng, C. C., Yang, Y. L., 2019. Chemistry and biology of
385 salicyl-capped siderophores. Nat. Prod. Chem. 59, 431-49
386

- 387 Janisiewicz, W. J., Tworowski, T. J., Sharer, C., 2000. Characterizing the mechanism of biological
 388 control of postharvest diseases on fruits with a simple method to study competition for nutrients.
 389 Phytopathology 90, 1196–1200
 390
- 391 Johnson, L., 2008. Iron and siderophores in fungal-host interactions. Mycol. Res. 112, 170–183
 392
- 393 Larena, I., Torres, R., De Cal, A., Linan, M., Melgarejo, P., Domenichini, P., Bellini, A., Mandrin,
 394 J.F., Lichou, J., Ochoa de Eribe, X., Usall J., 2005. Biological control of postharvest brown rot
 395 (*Monilinia* spp.) of peaches by field applications of *Epicoccum nigrum*. Biol. Control 32, 305-310
 396
- 397 Lee, B. N., Kroken, S., Chou, D. Y. T., Robbertse, B., Yoder, O. C., Turgeon, B. G., 2005.
 398 Functional analysis of all nonribosomal peptide synthetases in *Cochliobolus heterostrophus* reveals
 399 a factor, NPS6, involved in virulence and resistance to oxidative stress. Eukaryotic Cell 4, 545–555
 400
- 401 Li, C., Zhu, L., Pan, D., Li, S., Xiao, H., Zhang, Z., Shen, X., Wang, Y., Long, M., 2018.
 402 Siderophore-mediated iron acquisition enhances resistance to oxidative and aromatic compound
 403 stress in JMP134. Appl. Environ. Microbiol. 85
 404
- 405 Liu, X., Fang, W., Liu, L., Yu, T., Lou, B., Zheng, X., 2010. Biological control of postharvest sour
 406 rot of citrus by two antagonistic yeasts. Lett. Appl. Microbiol. 51, 30–35
 407
- 408 Liu, P., Luo, L., Long, C., 2013. Characterization of competition for nutrients in the biocontrol of
 409 *Penicillium italicum* by *Kloeckera apiculata*. Biol. Control 67, 157-162
 410

- 411 Mari, M., Martini, C., Guidarelli, M., Neri, F., 2012. Postharvest biocontrol of *Monilinia laxa*,
412 *Monilinia fructicola* and *Monilinia fructigena* on stone fruit by two *Aureobasidium pullulans*
413 strains. Biol. Control 60, 132-140
- 414
- 415 Martini, C., Guidarelli, M., Di Francesco, A., Ceredi, G., Mari, M., 2016. Characterization of
416 thiophanate methyl resistance in Italian *Monilinia fructicola* isolates. J. Plant Pathol. 98 (3)
- 417
- 418 McLoughlin, T. J., Quinn, J. P., Bettermann, A., Bookland, R., 1992. *Pseudomonas cepacia*
419 suppression of sunflower wilt fungus and role of antifungal compounds in controlling the disease.
420 Appl. Environ. Microbiol. 58, 1760–1763
- 421
- 422 Mei, B. G., Budde, A. D., Leong, S. A., 1993. Sid1, a gene initiating siderophore biosynthesis in
423 *Ustilago maydis*: molecular characterization, regulation by iron, and role in phytopathogenicity.
424 Proc. Natl. Acad. Sci. USA 90, 903–90
- 425
- 426 Mekbib, S. B., Regnier, T. J. C., Korsten, L., 2011. Efficacy and mode of action of yeast
427 antagonists for control of *Penicillium digitatum* in oranges. Trop. Plant Pathol. 36, 233–240
- 428
- 429 Miethke, M., Marahiel, M. A., 2007. Siderophore-based iron acquisition and pathogen control.
430 Microb. Mol. Biology Rev. 71, 413-51
- 431
- 432 Oide, S., Moeder, W., Krasnoff, S., Gibson, D., Haas, H., Yoshioka, K., Turgeon, B. G., 2006.
433 NPS6, encoding a nonribosomal peptide synthetase involved in siderophore-mediated iron
434 metabolism, is a conserved virulence determinant of plant pathogenic ascomycetes. Plant Cell 18,
435 2836–2853

436

437 Parafati, L., Vitale, A., Restuccia, C., Cirvilleri, G., 2015. Biocontrol ability and action mechanism
 438 of food-isolated yeast strains against *Botrytis cinerea* causing post-harvest bunch rot of table grape.
 439 Food Microbiol. 47, 85-92

440

441 Plante, S., Labbè, S., 2019. Spore germination requires ferrichrome biosynthesis and the
 442 siderophore transporter Str1 in *Schizosaccharomyces pombe*. Genetics 211, 893-911

443

444 Raaska, L., Mattila-Sandholm, T., 1995. Effects of iron level on the antagonistic action of
 445 siderophores from non-pathogenic *Staphylococcus* spp. J. Microbiol. Biotechnol. 15, 480-485

446

447 Riquelme, M., 1996. Fungal siderophores in plant-microbe interactions. Microbiology 12, 537–546

448

449 Sanz Ferramola, M. I., Benuzzi, D., Calvente, V., Calvo, J., Sansone, G., Cerutti, S., Raba, J., 2013.
 450 The use of siderophores for improving the control of postharvest diseases in stored fruits and
 451 vegetables. Microbial pathogens and strategies for combating them: science, technology and
 452 education (A. Méndez-Vilas, Ed.), pp. 1385-1394

453

454 Saravanakumar, D., Ciavarella, A., Spadaro, D., Garibaldi, A., Gullino, M. L., 2008.
 455 *Metschnikowia pulcherrima* strain MACH1 outcompetes *Botrytis cinerea*, *Alternaria alternata* and
 456 *Penicillium expansum* in apple through iron depletion. Postharvest Biol. Technol. 49, 121-128

457

458 Schrettl, M., Bignell, E., Kragl, C., Sabiha, Y., Loss, O., Eisendle, M., Wallner, A., Arst, H. N.,
 459 Haynes, K., Haas, H., 2007. Distinct roles for intra- and extra-cellular siderophores during
 460 *Aspergillus fumigatus* infection. PLoS Pathogen 28, 1195-207

461

- 462 Schwyn, B., Neilands, J. B., 1987. Universal chemical assay for the detection and determination of
463 siderophores. *Anal. Biochem.* 160, 47-56
464
- 465 Wang, W., Chi, Z., Liu, G., Buzdar, M. A., Chi, Z., Gu, Q., 2009. Chemical and biological
466 characterization of siderophore produced by the marine-derived *Aureobasidium pullulans* HN6.2
467 and its antibacterial activity. *Biometals* 22, 965–972
468
- 469 Xu, B., Zhang, H., Chen, K., Xu, Q., Yao, Y., Gao, H., 2013. Biocontrol of postharvest *Rhizopus*
470 decay of peaches with *Pichia caribbica*. *Curr. Microbiol.* 67, 255-261
471
- 472 Yu, X., Ai, C., Xin, L., Zhou, G., 2011. The siderophore producing bacterium, *Bacillus subtilis*
473 CAS15, has a biocontrol effect on *Fusarium* wilt and promotes the growth of pepper. *Eur. J. Soil*
474 *Biol.* 47, 138


475

476

477 Figure 1

478 CAS assay performed with *Aureobasidium pullulans* L1, L8 strains and *Monilinia laxa* (isolate

479 ML4).



480

481

482

483

484

485

486 Figure 2

487 Quantitative (mg ml^{-1}) determination of siderophore at 440 nm by spectrophotometer, on488 deferoxamine mesylate standard curve, of *Aureobasidium pullulans* L1 and L8 strains and489 *Monilinia laxa* (M), alone and co-cultured. Data are the means of two independent490 experiments \pm standard error, each consisting of three flasks per treatment. Different letters indicate491 significant differences at $\alpha=0.05$ according to Tukey's HSD Test.




Figure 3

In vivo antagonistic effect of L1 and L8 *Aureobasidium pullulans* strains on *Monilinia laxa* in peach fruits. Fruits were artificially inoculated with 20 μ l of yeasts cell suspensions (10^8 cells ml⁻¹); after air drying they were inoculated with 20 μ l of *M. laxa* conidia suspension (10^5 conidia ml⁻¹) and finally with an iron suspension (0, 5, 10 and 20 μ g ml⁻¹ FeCl₃) (0, 5, 10, 20). Control consisted of peaches inoculated with water instead of yeasts suspensions. Data are the means of two independent experiments \pm standard error, each consisting of 45 fruit per treatment. For each treatment group (Control, L1, and L8) different letters indicate significant differences at $\alpha = 0.05$ according to Tukey's HSD Test.




Table 1

CAS assay for analysis of siderophore production by *Aureobasidium pullulans* (L1 and L8) and *Monilinia laxa* (isolate ML4). Data are the means of two independent experiments \pm standard error, each consisting of ten replicates per treatment. Different letters within the same column indicate significant differences at $\alpha = 0.05$ according to Tukey's HSD Test.

Microorganism	Growth time (days)*	Color change	CAS reaction **
<i>Monilinia laxa</i>	10 \pm 0.4b	-	0 \pm 0.0a
L1	12 \pm 0.0a	dark orange	20 \pm 1.9b
L8	12 \pm 0.0a	dark orange	27 \pm 1.5c

518

519 *Days required for the fungal mycelium to cover the non-CAS half of the plate.

520 ** mm of advance of colour change front in the CAS blue agar after three weeks of incubation.

521

522

523 Table 2

524 Mycelium diameter (mm) of *Monilinia laxa* growth in dual culture with *Aureobasidium pullulans*525 L1 and L8 strains in minimal medium agar plates amended with 0, 5, 10 and 20 $\mu\text{g ml}^{-1}$ FeCl_3 and

526 incubated at 25 °C for 5 days. Control plates were inoculated only with pathogen plug (6 mm).

527 Data are the means of two independent experiments \pm standard error, each consisting of five528 replicates per treatment. Within the same FeCl_3 treatment (column), the same lower-case letters529 represent no significant differences at $\alpha = 0.05$ according to Tukey's HSD Test. Within the same

530 yeast strains or pathogen isolate (control) (row), the same upper-case letters represent no significant

531 differences at $\alpha = 0.05$ according to Tukey's HSD Test.

532

533

	Treatment (FeCl_3)			
	0 $\mu\text{g ml}^{-1}$	5 $\mu\text{g ml}^{-1}$	10 $\mu\text{g ml}^{-1}$	20 $\mu\text{g ml}^{-1}$
Control	60.0 \pm 0.6cA	60.0 \pm 1.2bA	63.0 \pm 1.0cB	61.0 \pm 1.5cAB
L1	35.2 \pm 1.9bB	35.0 \pm 0.7aB	34.7 \pm 1.3bA	34.9 \pm 1.1bA
L8	34.0 \pm 1.1aB	34.4 \pm 1.0aB	33.0 \pm 0.8aA	33.6 \pm 0.8aAB

534

535

536

537

538 Table 3

539 Effect of *Aureobasidium pullulans* (L1 and L8) on conidia germination (%) of *Monilinia laxa*540 grown in culture medium (minimal medium) amended with 5, 10 or 20 $\mu\text{g ml}^{-1}$ FeCl_3 and incubated

at 25°C for 6 hours. Data are the means of two independent experiments \pm standard error, each consisting of 90 conidia per treatment. Within the same FeCl₃ treatment (column), the same lower-case letters represent no significant differences at $\alpha = 0.05$ according to Tukey's HSD Test. Within the pathogen isolate (control) and yeast strains (row), the same upper-case letters represent no significant differences at $\alpha = 0.05$ according to Tukey's HSD Test.

	Treatment (FeCl ₃)			
	0 $\mu\text{g ml}^{-1}$	5 $\mu\text{g ml}^{-1}$	10 $\mu\text{g ml}^{-1}$	20 $\mu\text{g ml}^{-1}$
Control	40 \pm 1.3cA	41 \pm 1.8bA	48 \pm 0.8cC	43 \pm 1.5bB
L1	20 \pm 2.1bB	19 \pm 1.7aB	18 \pm 1.4bA	28 \pm 2.1aC
L8	19 \pm 0.9aB	19 \pm 1.5aB	15 \pm 1.2aA	29 \pm 1.8aC

549

550

551

552 Table 4

553 Effect of *Aureobasidium pullulans* (L1 and L8) on *Monilinia laxa* conidia germ tube elongation
 554 (μm) grown in liquid culture medium (minimal medium) amended with 0, 5, 10 and 20 $\mu\text{g ml}^{-1}$
 555 FeCl₃ at 25 °C for 6 hours. Data are the means of two independent experiments \pm standard error,
 556 each consisting of 90 conidia per treatment. Within the same FeCl₃ treatment (column), the same
 557 lower-case letters represent no significant differences significant differences at $\alpha = 0.05$ according
 558 to Tukey's HSD Test. Within the pathogen isolate (control) and yeast strains (row), the same upper-
 559 case letters represent no significant differences significant differences at $\alpha = 0.05$ according to
 560 Tukey's HSD Test.

561

	Treatment (FeCl ₃)			
	0µg ml ⁻¹	5 µg ml ⁻¹	10 µg ml ⁻¹	20 µg ml ⁻¹
Control	16.2±1.3cA	16.3±1.6cA	18.2±2.1cB	16.5±1.8bA
L1	7.7±2.1aB	6.6±1.8bA	6.2±2.3bA	8.1±2.1aC
L8	8.3±1.5bC	6.0±1.6aB	5.7±3.5aA	8.2±1.8aC

562

563

564 **Conflict of Interest**

565 The authors declare that they have no conflict of interests.

566

567 **Authors contribution**

568 Dr. Alessandra Di Francesco conducted all the experiments, interpreted data, and drafted the
 569 manuscript. Prof. Elena Baraldi coordinated the experimental plan, has corrected, and approved the
 570 final draft of the manuscript.

571

572

573 **Highlights**

574

575 - Competition for nutrients is one of the most important mechanism of action for BCAs

576 - Iron plays a role in BCAs modes of action through the production of siderophores

577 - Siderophores can inhibit the growth of certain pathogenic fungi

578 - *Aureobasidium pullulans* L1 and L8 strains compete for iron against *Monilinia laxa*

579

580