



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Please, don't do it! Fifteen years of progress of non-invasive brain stimulation in action inhibition

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Borgomaneri S., Serio G., Battaglia S. (2020). Please, don't do it! Fifteen years of progress of non-invasive brain stimulation in action inhibition. CORTEX, 132, 404-422 [10.1016/j.cortex.2020.09.002].

Availability:

This version is available at: <https://hdl.handle.net/11585/780994> since: 2022-01-24

Published:

DOI: <http://doi.org/10.1016/j.cortex.2020.09.002>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Borgomaneri, S., Serio, G., & Battaglia, S. (2020). Please, don't do it! Fifteen years of progress of non-invasive brain stimulation in action inhibition. *Cortex*, 132, 404–422.
<https://doi.org/10.1016/j.cortex.2020.09.002>

The final published version is available online at:
<https://doi.org/10.1016/j.cortex.2020.09.002>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

Please, don't do it!

Fifteen years of progress of non-invasive brain stimulation in action inhibition

Sara Borgomaneri^{1,2*}, Gianluigi Serio¹ & Simone Battaglia¹

¹ Centro studi e ricerche in Neuroscienze Cognitive, Dipartimento di Psicologia, Università di Bologna, Campus di Cesena, 47521, Cesena, Italy; ² IRCCS Fondazione Santa Lucia, 00179 Rome, Italy.

*Correspondence should be addressed to: Sara Borgomaneri. Department of Psychology, University of Bologna. Viale Berti Pichat 5, Bologna, 40127, Italy. e-mail: sara.borgomaneri@unibo.it

Abstract

The ability to inhibit prepotent responses is critical for survival. Action inhibition can be investigated using a stop-signal task (SST), designed to provide a reliable measure of the time taken by the brain to suppress motor responses. Here we review the major research advances using the combination of this paradigm with the use of non-invasive brain stimulation techniques in the last **fifteen** years. We highlight new methodological approaches to understanding and exploiting several processes underlying action control, which is critically impaired in several psychiatric disorders. In this review we present and discuss existing literature demonstrating i) the importance of the use of non-invasive brain stimulation in studying human action inhibition, unveiling the neural network involved ii) the critical role of prefrontal areas, including the pre-supplementary motor area (pre-SMA) and the inferior frontal gyrus (IFG), in inhibitory control iii) the neural and behavioural evidence of proactive and reactive action inhibition. As the main result of this review, the specific literature demonstrated the crucial role of pre-SMA and IFG as evidenced from the field of noninvasive brain stimulation studies. Finally, we discuss the critical questions that remain unanswered about how such non-invasive brain stimulation protocols can be translated to therapeutic treatments.

Keywords: Action inhibition; stop-signal task; transcranial magnetic stimulation; transcranial direct current stimulation; Action Inhibition Network.

1. Introduction

1 The control of voluntary action involves a series of different processes that need to be harmoniously
2 orchestrated, such as choosing from a range of possible actions as well as inhibiting responses as
3
4 circumstances demand. The ability to inhibit prepotent ongoing actions is crucial to prevent
5
6 potentially harmful behavioral outcomes. The ability to inhibit prepotent responses can be
7
8 investigated experimentally using a stop-signal task (SST), designed to provide a sensitive measure
9
10 of the time taken by the brain to inhibit or suppress inappropriate motor responses (Lappin &
11
12 Eriksen, 1966; Logan, Cowan, & Davis, 1984; Vince, 1948). It requires participants to respond to a
13
14 go stimulus and to abort the ongoing response when a stop signal is presented. To measure the
15
16 participant's performance on the SST, the stop signal reaction time (SSRT), an index of inhibition,
17
18 is computed based on Logan and Cowan's notion (Logan et al., 1984). Logan and Cowan proposed
19
20 the existence of a race between a go process, triggered by the presentation of the go stimulus, and a
21
22 stop process, triggered by the presentation of the stop signal.

23
24
25
26
27
28
29
30
31 When the stop process terminates before the go process, the response is inhibited, but when the go
32
33 process terminates before the stop process, the response is activated. Thus, unpredictable stop
34
35 signals require withholding the response when a go response has already been initiated. The
36
37 stopping process needs to be estimated from a stochastic model, the so-called 'race model' (Logan
38
39 et al., 1984), which estimates the SSRT from the distribution of 'go' reaction times and the
40
41 observed probability of responding on 'stop' trials for a given stop-signal delay. This measure is an
42
43 indicator of inhibitory control, with a lower value indicating a more rapid ability to respond to a
44
45 stop signal (Cai et al., 2015). The time between presentation of the go stimulus and presentation of
46
47 the stop signal is termed the 'stop signal delay' (SSD). Critically, the motor inhibition will be
48
49 successful only when the SSD is short enough to allow inhibitory processes to stop the ongoing
50
51 motor program, while an increased SSD will result in an increased likelihood of failure to inhibit
52
53 the response to the go stimulus. SSD can be fixed or dynamically modified trial-by-trial based on
54
55 the participant's performance (Verbruggen & Logan, 2009). However, estimated SSRT gives the
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

measure of the duration of the inhibitory process by revealing the time necessary for successful motor inhibition (Logan et al., 1984).

Recent theories (Aron, 2011; Braver, Gray, & Burgess, 2007) propose that inhibitory control can be divided into proactive mechanisms (response slowing in anticipation of a stop-signal) and reactive mechanisms (outright stopping in response to a stop-signal, measured by means of the SSRT). An example of these mechanisms includes the possibility to brake roughly if a person gets in your way while you are driving your car, as a reaction to the rapid and unexpected change in the context (reactive inhibition). However, one could also pre-emptively act so as not to drive too fast, as the probability of encountering a bypassing pedestrian in the street is often high, and therefore this approach would almost guarantee a successful outcome (proactive inhibition and working memory). Typically, proactive inhibitory control is measured by the increase in reaction times in the go trial when the probability of a stop-signal is high (Cai et al., 2015; Meyer & Bucci, 2016; Zandbelt, Van Buuren, Kahn, & Vink, 2011) or by an index of the proactive inhibitory control, named ‘preparatory cost’ (PC) (Cai et al., 2015). Crucially, a low preparatory cost indicates a weaker anticipation of a stop-signal, while a higher preparatory cost indicates better proactive inhibitory control.

Several brain areas have been associated with the mechanisms underlying inhibitory control, with a network including left and right inferior frontal gyrus (IFG) (Aron, Fletcher, Bullmore, Sahakian, & Robbins, 2003; Chevrier, Noseworthy, & Schachar, 2007; Leung & Cai, 2007; Zhang, Geng, & Lee, 2017), dorsolateral prefrontal cortex (dlPFC) (Bari & Robbins, 2013; Leung & Cai, 2007), anterior cingulate (ACC) (Ito, Stuphorn, Brown, & Schall, 2003; Zhang et al., 2017), pre-supplementary motor area (pre-SMA) (Bari & Robbins, 2013; Leung & Cai, 2007), supplementary motor area (SMA) (Bari & Robbins, 2013; Li, Huang, Constable, & Sinha, 2006; Zhang et al., 2017), bilateral superior temporal gyri (Zhang et al., 2017), parietal cortex (Bari & Robbins, 2013; Leung & Cai, 2007; Zhang et al., 2017), insula (Bari & Robbins, 2013; Leung & Cai, 2007; Zhang et al., 2017), basal ganglia (Bari & Robbins, 2013; Chevrier et al., 2007; Zhang et al., 2017), cerebellum (Clark, King, & Turner, 2020), frontal eye fields (FEF) and supplementary eye field

(SEF) (Dillon & Pizzagalli, 2007; Hanes & Schall, 1996; Leung & Cai, 2007; Stuphorn & Schall, 2006; Stuphorn, Taylor, & Schall, 2000), all implicated by a range of studies (see Zhang et al., 2017 for meta-analysis). The critical and specific role of each of these areas in action inhibition is still debated. Recent studies, for instance, have demonstrated that lesions to IFG cause a deficit of response inhibition, as measured using tasks that require the suppression of an initiated manual response (Aron et al., 2003) or the suppression of a reflexive saccade (Hodgson et al., 2007). In addition, human patient studies revealed that damage to another section of the prefrontal cortex, the right superior frontal regions (including pre-SMA and SMA), raised SSRT in the SST (Floden & Stuss, 2006 for a review see Mostofsky & Simmonds, 2008). In support of the critical role of pre-SMA in action control, several studies with macroelectrode stimulation in epilepsy patients show that pre-SMA stimulation leads to the arrest of ongoing vocal or manual movements (Lüders et al., 1988; Mikuni et al., 2006; Swann et al., 2012) and a single case report of a patient with a lesion of the pre-SMA extending to cingulate and superior frontal gyri also showed a behavioral stopping deficit for a stop-change task (Nachev, Wydell, O'Neill, Husain, & Kennard, 2007). From all this evidence, it seems clear that dorsomedial damage impairs stopping; however, lesions were not restricted to the pre-SMA, preventing the possibility to draw any conclusions. Indeed, macrostimulation of both the pre-SMA and the right IFG can induce motor arrest (Fried et al., 1991; Lüders et al., 1988), and stimulation of the pre-SMA evoked responses in the right IFG, corresponding well to a white-matter connection, and also the stopping-related task evoked responses (Swann et al., 2012). As suggested by the multiple domain hypothesis of response inhibition (Rubia et al., 2001), distinct parts of the frontal lobes may be responsible for different aspects of inhibitory control, for instance the recruitment of the dlPFC seems to be relevant under conditions in which greater cognitive demand (i.e., more working memory load) is necessary to guide response inhibition (Criaud & Boulinguez, 2013; Jahfari, Stinear, Claffey, Verbruggen, & Aron, 2010; Mostofsky et al., 2003). Indeed, although the modulation of behavior when expecting a stopping stimulus (proactive inhibition) is a proposed function linked to the pre-SMA (Aron, 2011;

1 Boulinguez, Ballanger, Granjon, & Benraiss, 2009; Chikazoe et al., 2009; Forstmann et al., 2008;
2 Jahfari et al., 2010; Zandbelt & Vink, 2010), some hypotheses posit the dlPFC as a candidate for
3 proactive inhibition due to the working memory component in such behavior (Criaud &
4 Boulinguez, 2013; Jahfari et al., 2010; Mostofsky et al., 2003). Importantly, monkey evidence
5 recorded firing in the pre-SMA during both stop trials and non-stop trials (Stuphorn & Emeric,
6 2012). Firing peaks during non-stop trials suggested that the pre-SMA could induce proactive
7 inhibition throughout phasic firing due to its reactive function. In addition, switching from
8 repetitive to new movements became worse by disrupting pre-SMA activity (Rushworth, Hadland,
9 Paus, & Sipila, 2002). Thus, the pre-SMA appears to be recruited during both response inhibition
10 and action switching.
11

12 Critically, all the aforementioned studies employed functional magnetic resonance (Aron et al.,
13 2003; Bari & Robbins, 2013; Chevrier et al., 2007; Clark et al., 2020; Dillon & Pizzagalli, 2007;
14 Hanes & Schall, 1996; Ito et al., 2003; Leung & Cai, 2007; Li, Huang, et al., 2006; Stuphorn &
15 Schall, 2006; Stuphorn et al., 2000; Zhang et al., 2017) or patients with brain lesions (Aron et al.,
16 2003; Floden & Stuss, 2006; Fried et al., 1991; Hodgson et al., 2007; Lüders et al., 1988; Mikuni et
17 al., 2006; Nachev et al., 2007; Swann et al., 2012). Although neuroimaging studies provide
18 evidence that links areas in the brain to task performance, this approach is unable to provide a
19 causal link showing which areas play an essential role. Similarly, human patient studies cannot
20 avoid associated limitations as a result of neural plasticity or reorganization. A way to solve this
21 issue, is to use non-invasive brain stimulation (NIBS) techniques to selectively manipulate in
22 healthy participants single cortical components of the action inhibition network (AIN) to investigate
23 their - specific - contribution in the several processes underlying action control (i.e., inhibition,
24 selection, competition and switching of actions). Recently, there has been a growing interest in the
25 application of different non-invasive brain stimulation techniques to induce neuroplasticity and to
26 modulate cognition and behavior (Huang et al., 2017). In this review, we report and compare,
27 whenever possible, the main results of action inhibition studies that undertook to investigate the
28

1 critical role of the key nodes of the AIN by means of NIBS using an SST paradigm. In particular,
2 only peer-reviewed published studies in English language were included. Studies were required to
3
4 report experimental designs on healthy volunteers, using TMS (we excluded studies testing motor
5
6 evoked potentials) or tDCS in human participants with a control condition (sham stimulation, active
7
8 control stimulation or baseline performance). Only studies implementing the stop-signal task and
9
10 reporting stop-signal reaction time (SSRT) as outcome were included. Our aim is to understand the
11
12 crucial role of cortical brain regions of the AIN in motor action suppression and to disclose which
13
14 are the most efficient NIBS protocols.
15
16
17

18
19 TMS is a non-invasive neural modulation technique with valuable potential in both neuroscience
20
21 (Bergmann, Karabanov, Hartwigsen, Thielscher, & Siebner, 2016) and clinical studies (Lefaucheur
22
23 et al., 2014). In particular, it has been shown that repetitive TMS (rTMS) can temporarily modify
24
25 brain function for minutes to hours (Hamada et al., 2007; Huang, Edwards, Rounis, Bhatia, &
26
27 Rothwell, 2005; Iyer, Schleper, & Wassermann, 2003; Jung, Shin, Jeong, & Shin, 2008).
28
29 Stimulations at low (≤ 1 Hz) and high (≥ 5 Hz) frequency can decrease or increase neuronal
30
31 excitability, respectively. Among rTMS protocols, theta-burst stimulation (TBS) can produce long
32
33 after-effects (>20 min) using relatively short-term stimulation (typically, 40– 190 s) at a higher
34
35 frequency (50 Hz) (Huang et al., 2005). In particular, continuous TBS (cTBS; 40 s train of
36
37 uninterrupted TBS is given for a total 600 pulses) is supposed to suppress cortical excitability,
38
39 while intermittent TBS (iTBS; 2 s train of TBS is repeated every 10 s for a total of 190 s and 600
40
41 pulses) is thought to facilitate it (Huang et al., 2005). While using tDCS, direct electric current,
42
43 passing through two saline soaked sponge electrodes placed over the participant's skull, is applied
44
45 to modulate human brain excitability (George & Aston-Jones, 2010). Through this type of
46
47 stimulation, feeble electric currents (1-2 mA) are conducted through two electrodes (anode and
48
49 cathode), which increase or decrease neuron activity by changing the membrane potential (Nitsche
50
51 & Paulus, 2000). Although the exact functioning of tDCS is not entirely clear, it is well established
52
53 that several minutes of stimulation with the anode placed over a target area increases its cortical
54
55
56
57
58
59
60
61
62
63
64
65

1 activity, whereas placing the cathode sustainably reduces it. Thus, tDCS could increase or decrease
2 neuron activity and also modulate behavior, and the effects depend on the duration and the current
3 density of the stimulation (Nitsche et al., 2008).
4
5

6
7 The endeavor to understand the neural circuits of action inhibition in humans originated
8 approximately 15 years ago; here we review the major advances, highlighting new methodological
9 approaches to understanding and exploiting several processes underlying action control. In
10 particular, we critically examine different non-invasive brain stimulation protocols in SST and the
11 specific results obtained, aiming at trying to delineate and suggest important factors that can
12 contribute to modulate inhibition performance. Hence, our goal is to provide fundamental
13 implications of clinical relevance for the recent enhancing in the understanding of psychiatric
14 disorders and thus improving relative treatments in this area of research. A summary of the
15 experimental results, methods and stimulation protocols adopted in each NIBS study is reported in
16 Table 1 and Table 2, while Figure 1 and 2 show the stimulation site in each NIBS-SST experiment.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33 34 **2. Transcranial Magnetic Stimulation and action inhibition**

35
36 Although the cognitive mechanisms underlying response inhibition have been studied for several
37 years (Logan, 1981, 1994), the neural mechanism subtending this process is still a matter of debate.
38
39 The most accredited theory proposes that the human prefrontal cortex is responsible for executive
40 control, but it is contentious whether discrete prefrontal regions are specialized to carry out domain-
41 specific functions (Aron, Robbins, & Poldrack, 2004; Duncan & Owen, 2000; Goldman-Rakic,
42 1987; Rowe, Toni, Josephs, Frackowiak, & Passingham, 2000). Results from several studies
43 suggested that prefrontal regions share control over a range of cognitive processes, including those
44 involved in the inhibition and selection of responses (Duncan & Owen, 2000). Others, however,
45 have argued that mechanisms of response inhibition are governed by a discrete network of brain
46 regions in the parietal and prefrontal cortex (Aron et al., 2003; Garavan, Ross, & Stein, 1999;
47 Morita, Nakahara, & Hayashi, 2004; Rubia, Smith, Brammer, & Taylor, 2003). A way to solve this
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

issue is to use TMS in healthy volunteers to temporarily interfere with the activity of a selective region of the prefrontal cortex to assess its causal role in a discrete function. In 2006, Chambers and colleagues undertook to test the hypothesis that discrete regions of the right prefrontal and parietal cortex (e.g., IFG and middle frontal gyrus- MFG or angular gyrus -AG) selectively govern response inhibition in the healthy brain. To investigate this issue, the authors asked participants to undergo an SST after receiving different sessions of 15 min (1 Hz) of offline rTMS at an average of 92% resting **motor threshold (rMT)** to the right IFG, MFG, or AG. In the SST, participants had to identify a target go stimulus as rapidly as possible. In 25% of the trials, a stop signal was presented, instructing participants to withhold their response. To manipulate the difficulty of successfully inhibiting, the stop signal was presented randomly at various delays after the go signal. Moreover, to maximize the sensitivity of their TMS protocol, stop signals were presented at SSDs that that yielded 25–75% correct inhibitions. Finally, to measure possible effects of cortical reorganization over time, participants received two consecutive blocks of rTMS per session, each followed by the SST, and pupil diameter was recorded throughout the experiment as an index of physiological arousal. Results showed that rTMS over rIFG impact the **response** inhibition performance (i.e., lower percentage of correct inhibitions and longer SSRT) compared to sham, in both the left and right hands selectively in the first block after rTMS, while in the second block after rTMS no significant effects of rTMS on behavior were found. Moreover, no significant effects of rTMS were observed on go-RT or on error rates nor in the change of pupil diameter over time, excluding that diminished arousal over time may explain the lack of significant effect in the second block. To conclude, these data successfully demonstrated the critical role of the rIFG in inhibiting or overriding prepotent responses of both hands. However, such data do not imply that IFG is the sole mediator of response inhibition, since this brain region is richly interconnected with a range of cortical and subcortical structures (Durstson et al., 2003; Vink et al., 2005), which can be affected by the deactivation of the IFG and may contribute to action control. Moreover, the rTMS effect over IFG was selectively observed in the first block and it tended to disappear during the second block.

1 These results suggest that the AIN is able to functionally reorganize, directing elsewhere the critical
2 process ascribed to IFG (Siebner & Rothwell, 2003), such as to the right MFG, parietal cortex, or
3
4 homologous structures in the left hemisphere (Hester, Murphy, & Garavan, 2004). The following
5
6 year, the same research group (Chambers et al., 2007) investigated whether a range of inhibitory
7
8 behaviors including response inhibition and response selection may be subtended by a common
9
10 neural substrate. To test this hypothesis, participants were tested in their ability to overcome
11
12 competing response tendencies (i.e., flanker task), and in the ability to withhold an initiated action
13
14 (i.e., SST) after two separate sessions (20 min each) of offline 1 Hz rTMS at an average of 92%
15
16 rMT over left and right IFG and dorsal premotor cortex (dPMC). The combined flanker/SST
17
18 required participants to identify the direction of the central arrow as rapidly and accurately as
19
20 possible with their left or right hand, ignoring the distractors and withholding their response when
21
22 the stop signal occurred. Results from this behavioral data revealed that the ability to inhibit a
23
24 prepotent response (in the SST) is closely related to the degree of competition between responses
25
26 (in the flanker task SSRT reduced significantly when incongruent distractors flanked the central
27
28 arrow target, relative to conditions in which the flankers were congruent or neutral). The rTMS data
29
30 showed that the rTMS over rIFG slowed SSRT on incongruent trials relative to both sham and
31
32 dPMC conditions (but see Yang, Khalifa, & Völlm, 2018 for null results on SSRT after the
33
34 excitatory stimulation of the rIFG; 45 trains of 10-Hz rTMS stimulation session at 100% of the rMT
35
36 consisted of 900 pulses in total with a 2-s duration of each train and a 10-s interval between each
37
38 train). No effects were found in congruent trials, demonstrating that rTMS over the rIFG selectively
39
40 increased SSRT in trials in which a suppression of a competing response is required. Moreover,
41
42 analysis on go trials showed that performance tended to improve after stimulation of right dPMC
43
44 compared with the sham and rIFG stimulation conditions. Data from the experiment in which IFG
45
46 and dPMC were stimulated in the left hemisphere showed no significant effect of the rTMS either in
47
48 the SSRT or in the go trials analysis. To sum up, these data confirm previous findings (Aron et al.,
49
50 2003; Chambers et al., 2006; Hodgson et al., 2007) on the causal role of the right (and not the left)
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 IFG in stop-signal inhibition, selectively when response selection was placed under competition
2 (i.e., incongruent flanker trials). Therefore, the rIFG appears to be especially crucial for erasing
3 responses under conditions of increased response competition, even though rIFG stimulation did not
4 alter the effect of competition on response execution. The rTMS over rIFG may have delayed the
5 release of an inhibitory trigger or reduced the speed of the stopping process once triggered, possibly
6 due to the existence of direct connections between the IFG and the subthalamic nucleus of the basal
7 ganglia (Aron, Behrens, Smith, Frank, & Poldrack, 2007; Aron & Poldrack, 2006; Nambu, Tokuno,
8 & Takada, 2002). Moreover, stimulation of the right dPMC tended to enhance execution
9 performance in all conditions, facilitating RT of both hand responses regardless of the degree of
10 competitive response selection. This result is in contrast with results obtained in previous rTMS
11 studies of response selection (Koski, Molnar-Szakacs, & Iacoboni, 2005; Praamstra, Kleine, &
12 Schnitzler, 1999; Schluter, Rushworth, Passingham, & Mills, 1998), possibly due to several
13 methodological differences such as the behavioral paradigms, rTMS protocols, and site localization.
14 However, the observed double dissociation of the rTMS effects between the rIFG and right dPMC
15 leads authors to the conclusion that response inhibition and execution rely on distinct neural
16 processes despite activating a common cortical network. From this evidence, it seems clear that the
17 rIFG has a predominant inhibitory role in updating behavior; however, other theorists posit that this
18 area may have a role in attention, mediating detection of stimulus change (Sharp et al., 2010),
19 whereas other brain regions, such as the pre-SMA, are effectively involved in action inhibition or in
20 both processes (Hampshire, Chamberlain, Monti, Duncan, & Owen, 2010). Alternatively, the rIFG
21 could have a general role in updating action plans (Mars, Piekema, Coles, Hulstijn, & Toni, 2007).
22 The key difference between the inhibition and updating accounts is that the inhibition account
23 assumes that rIFG is selectively crucial for stopping, whereas the updating account assumes that
24 rIFG is crucial for different forms of updating (including stopping). To investigate whether the rIFG
25 effectively has a role in updating behavior, Verbruggen and colleagues (2010) employed cTBS over
26 rIFG, over the more dorsal right inferior frontal junction (rIFJ) and over pre-SMA or sham in
27

1 separate sessions, and tested participants in a context cueing paradigm. In this paradigm, which is
2 able to dissociate several control processes involved in updating behavior, the identity of a cue
3 indicated different task contexts (ignore, dual, and stop). When the cue changed color, participants
4 initiated a manual response. In a minority of trials, the cue turned bold after a variable delay: in the
5 signal-ignore trials, participants had to ignore the change and execute the response; in the dual-
6 signal trials, participants had to execute an additional response by pressing an alternative key; and
7 in stop-signal trials, participants had to try to withhold the response. Importantly, to disentangle
8 visual-detection account and the action-updating account, the colored cue turned bold for 250 ms
9 after a variable delay (Stimulus Onset Asynchrony - SOA). In signal-ignore and dual-signal trials,
10 SOAs were fixed (100, 250, or 400 ms); in stop-signal trials, SOA was dynamically adjusted.
11 Results showed that cTBS over the rIFG and rIFJ impaired performance in both stop and dual tasks
12 compared to sham, suggesting that both areas are not selectively crucial for inhibitory control. In
13 contrast to the results from cTBS over rIFG in which the dual-task effect did not interact with SOA,
14 a strong interaction with SOA was found for the rIFJ stimulation. These results indicate that rIFG
15 has a crucial role in action control, updating action plans, while rIFJ has a role in detecting visual
16 changes of stimulus features. No effects were found when cTBS was applied over pre-SMA. These
17 data demonstrated that rIFG is critical for stopping a response, but also that the cognitive role of
18 rIFG is not purely one of inhibitory control (see data from dual-task) but seems to be not attentional
19 (see data on the lack of interactions with SOA).

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46 In 2015, Dippel & Beste tested the effect of an iTBS over the rIFG (together with cTBS over the
47 rIFG and sham) and investigated the possible role of IFG in multi-component behavior. To test this
48 issue, participants performed an SST in which the ongoing action was occasionally interrupted
49 following delivery of a stop stimulus, while an alternative response had to be executed upon the
50 presentation of an auditory stimulus. The auditory stimulus was presented either simultaneously or
51 following the stop stimulus, so that when presented simultaneously, participants could process the
52 two actions either serially or in parallel, while when the auditory signal followed the stop, it was

1 only possible to process the two actions in a serial manner. Results showed that iTBS over the rIFG
2 induced a more efficient strategy of task goal activation, while cTBS of the rIFG induced a less
3 efficient strategy of task goal activation compared with sham TBS (i.e., cTBS slowed the RT to the
4 change stimulus, whereas the iTBS protocol increased it), while no effect on SSRT were found. It is
5 important to mention that the present study is not completely comparable to the classical SST and
6 therefore results may be ascribed to the use of a modified version of SST. However, these results
7 are consistent with studies suggesting that the rIFG is unlikely to reflect a specific module for
8 inhibitory demands but is instead also part of other cognitive domains (Erika-Florence, Leech, &
9 Hampshire, 2014).

10 The first study that employed rTMS for investigating the critical role of pre-SMA in action
11 inhibition was carried out by Chen, Muggleton, Tzeng, Hung, & Juan in 2009. Pre-SMA represents
12 an important hub in the action inhibition network since it has direct connections to both the rIFG
13 and the subthalamic nucleus, which are considered to be the main neural substrates for inhibitory
14 control (Aron et al., 2007); however, its critical role in this process had never been tested. Chen and
15 coworkers therefore decided to test the critical role of pre-SMA by using an online 10 Hz rTMS
16 protocol (i.e., while participants are doing the SST). The SST required participants to make a key-
17 press response when a dot was presented on the left with the left finger, or with the right index
18 finger when the dot was presented on the right. In stop trials, a central fixation dot reappeared and
19 acted as an instruction to withhold responses to the peripheral target. In the rTMS blocks, at the
20 onset of the go signal, two TMS pulses with an inter-pulse interval of 100 ms at 60% of the
21 **maximum stimulator output (MSO)** were delivered over the pre-SMA or vertex (i.e., the active
22 control site) in separate sessions. rTMS over pre-SMA elongated the SSRT compared to vertex
23 stimulation and without TMS. Since Chambers and colleagues in 2006 found differences in SSRT
24 selectively in the first block, an additional analysis took into consideration data from separate sub-
25 sessions. Results showed that the pre-SMA TMS effect seems more substantial in the first sub-
26 session (i.e., longer SSRT in the first compared to the second sub-session). Indeed, no main effect

1 for stimulation sites was found in the second sub-session. These data demonstrate the critical role of
2 pre-SMA in the inhibitory control for both hands, in line with previous fMRI data (Li, Huang, et al.,
3
4 2006), which found that activation of the pre-SMA was correlated with shorter SSRTs, suggesting
5
6 that greater activity in pre-SMA is indicative of more efficient inhibitory control. The fact that the
7
8 effects are visible only during the first session may suggest that the role of pre-SMA is dependent
9
10 on familiarity with the task or that its functional weight in inhibitory control may change over time.
11
12 Although informative, the effect of the elongated SSRT when the TMS was applied before the stop
13
14 signal left open two alternative hypotheses: the TMS may have disrupted the stopping preparation
15
16 or the stopping process itself. To investigate the issue further, another study (Cai, George,
17
18 Verbruggen, Chambers, & Aron, 2012) tested the role of the right pre-SMA in modulation of
19
20 response tendencies versus implementation of the stopping process by using a conditional SST. In
21
22 this task, participants were asked to cancel their response selectively whenever the arrow go signal
23
24 turns red and points in the critical direction. During the task, dual TMS pulses were delivered over
25
26 right pre-SMA or vertex at 125/150 ms or at 175/200 ms after the go signal, while no pulses were
27
28 delivered on the remaining half of the trials. Results showed that SSRT on TMS trials in the pre-
29
30 SMA session was significantly longer than SSRT in the vertex TMS condition or in trials without
31
32 TMS, in line with previous findings (Chen et al., 2009). However, such effect of SSRT increase
33
34 could relate to a disruption in setting up the stopping rule or to disruption in implementing stopping,
35
36 and therefore, in a second experiment, the authors employed a selective SST, which involves a trial-
37
38 by-trial changing stopping rule and allows to dissociate setting up the stopping rule from
39
40 implementing stopping (Aron & Verbruggen, 2008). Specifically, the selective SST has a
41
42 foreknowledge period (maybe stop right hand), a delay, a go stimulus (move both hands together)
43
44 and, then, in some trials a stop signal requiring the participant to stop the cued hand and continue
45
46 with the other is presented. The stop signal is not informative of which hand to stop; therefore, the
47
48 participant needs to encode the cue presented in the foreknowledge period. To interfere with the
49
50 period corresponding to setting up the selective stopping rule, dual pulses were delivered at 100/125
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

ms or at 150/175 ms after the offset of the task cue, while at 125/150 or 175/200 ms after the onset of the stop signal or after the average SSD in go trials to interfere with the period corresponding to implementing the stopping process. No pulses were administered in the remaining trials. In trials in which TMS was delivered over pre-SMA, SSRT was significantly longer than SSRT in trials without TMS or when the vertex was stimulated. In both experiments, no significant effect of the TMS over pre-SMA was found in other behavioral indices, ruling out possible distraction or discomfort effects. These data suitably demonstrated that TMS over the right pre-SMA specifically disrupted the implementation of stopping process, since the stimulation after the stop signal was too late to affect setting up the stopping rule. A later study (Obeso, Cho, et al., 2013) combined the cTBS (20 s block consisted of 3 pulses at a rate of 50 Hz repeated every 200 ms at 80% of the active MT; AMT) over pre-SMA or sham with the H2 15O positron emission tomography (PET) scans during the SST and provided evidences on the causal role of the pre-SMA in reactive inhibition (reduced SSRT relative to sham, as found in a subsequent study which found this effect to be specific for cTBS over pre-SMA and not IFG; Obeso et al., 2017). Moreover, cTBS had an impact on the regional cerebral blood flow (rCBF) in several frontal areas, such as in the left pre-SMA, left IFG, right premotor and right inferior parietal cortex, suggesting that the stimulation may have a major impact not only locally in the underlying cortex but also distally in interconnected areas, which may also be critical in the observed results or may represent a compensatory activation against the TBS-induced “disruptive” effect. In the same year, the same research group undertook to further investigate whether pre-SMA has a role in action inhibition (both reactive and proactive) and/or in the switching of actions, and whether this area acts in concert with the rIFG (Obeso, Robles, Muñoz-Marrón, & Redolar-Ripoll, 2013). Participants were involved in a modified version of the SST in which they have to respond to the arrow’s directions (go trials), and to stop any movement when the arrow was followed by an infrequent cross. In the switch condition, the arrow turns blue, requiring switching to a new movement. Before the task, cTBS over the rIFG or sham were applied. To see the influence of rIFG in pre-SMA functioning, following cTBS, participants

1 performed the stop switching task while receiving single TMS pulses (100 ms after stimulus; 60%
2 of the MSO) over the right pre-SMA or vertex. Results on stopping processes showed that online
3
4 TMS pulses over the right pre-SMA (after sham cTBS but not after cTBS over IFG) altered the
5
6 stopping process (i.e., prolonged SSRT), and the preparation to inhibit an action (i.e., reduced
7
8 response times needed to adapt between different task conditions during proactive inhibition), but
9
10 no such worsening was observed in switch trials, when participants received online TMS pulses
11
12 over both IFG and pre-SMA, suggesting the lack of a critical role of these two regions at least in the
13
14 tested temporal window (i.e., 100 ms). To conclude, these data confirmed the crucial role of pre-
15
16 SMA in reactive and proactive inhibition. Moreover, it is obvious that the successful inhibition of
17
18 an action requires the suppression of activity in the primary motor cortex; however, the way in
19
20 which action inhibition commands reach M1 is still a matter of debate. In 2013 Zandbelt,
21
22 Bloemendaal, Hoogendam, Kahn, & Vink undertook to investigate how the regions of the AIN
23
24 interact to exert inhibitory control over M1 and whether these regions have a causal involvement in
25
26 proactive and reactive inhibition. Each participant underwent three separate rTMS-fMRI sessions,
27
28 in which they received rTMS stimulation over the rIFC, over the pre-SMA (i.e., supplementary
29
30 motor complex; SMC) and sham. rTMS consists of 20 trains of 30 6-Hz pulses at 90% of rMT with
31
32 an intertrain interval of 25 sec, followed by 600 1-Hz pulses at 110% of rMT. Immediately
33
34 afterwards, participants were asked to perform the stop-signal anticipation task in the scanner.
35
36 Importantly, to test both proactive and reactive inhibition, the stop-signal probability was
37
38 manipulated across trials. Typically, as stop-signal probability increases, participants proactively
39
40 slow down responding to increase inhibition accuracy when a stop-signal occurs. Results showed
41
42 that rIFC and pre-SMA stimulation did not influence proactive inhibition (i.e., no changes in RT
43
44 were detected) but only reactive inhibition. Indeed, participants became faster in reactive stopping
45
46 (i.e., shorter SSRT) after both rIFC and pre-SMA stimulation. Interestingly, this effect was
47
48 associated with increased M1 deactivation. Furthermore, rIFC and pre-SMA stimulation increased
49
50 right striatal activation, implicating an involvement of fronto-striatal pathways in reactive
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 inhibition. Finally, rIFC stimulation altered pre-SMA activation, while pre-SMA stimulation did not
2 alter rIFC activation, indicating that rIFC lies upstream from pre-SMA.
3

4 Although converging evidence of these NIBS studies suggests a causal role of rIFC in successful
5 response inhibition, most of the above-mentioned studies focused exclusively on the SST as a
6 measure of action cancellation, without assessing the critical role of the rIFC in action restraint.
7 Thus, to fill this gap, Dambacher and colleagues in 2014 measured action restraint with a go/no-go
8 task and action cancellation in an SST, after separate sessions of cTBS (at 100% of the AMT) or
9 sham stimulation of the right anterior insula (rAI), right superior frontal gyrus, and pre-SMA.
10 Results showed higher false alarms rates in inhibition trials in the go/no-go task and SST after
11 rTMS over rAI, demonstrating that the stimulation reduced both the ability to restrain and cancel
12 responses, while disruption of the right superior frontal gyrus specifically impaired the ability to
13 restrain from responding (higher false alarms rates in inhibition trials selectively in the go/no-go
14 task), while leaving the ability for action cancellation largely intact. Inhibitory processes were not
15 affected by the stimulation applied to the right middle frontal gyrus and pre-SMA. These findings
16 emphasize the role of IFG for global inhibition, whereas superior frontal regions seem to be
17 specifically relevant for successful action restraint.
18

19 To examine the causal links between several regions of the action inhibition network, such as the
20 frontal regions and basal ganglia, in 2015 Watanabe and coworkers measured human fMRI activity
21 during SST before and after excitatory/inhibitory rTMS of the pre-SMA. To induce long-lasting
22 changes in brain activity, the authors used quadripulse rTMS (QPS), the effects of which are
23 supposed to continue for a relatively long time after the stimulation (from 30 min to 2 h; Hamada et
24 al., 2007). The QPS consisted of a train of four monophasic magnetic pulses, delivered with four
25 magnetic stimulators connected to a specially designed combining module at 90% of AMT. The
26 inter-burst interval was set at 5 s. One rTMS block had 360 consecutive bursts, each of which
27 comprised four magnetic pulses separated by interstimulus intervals of 5 or 50 ms, for excitatory or
28 inhibitory rTMS respectively. Each participant underwent an SST performed in an fMRI scanner
29

1 before and after a session of rTMS (excitatory, inhibitory of the pre-SMA or sham). Excitatory
2 rTMS significantly decreased SSRT compared to the sham condition, while inhibitory rTMS
3
4 significantly increased SSRT. Indeed, such behavioral changes were correlated with neural changes
5
6 in striatum (STR) and globus pallidus pars interna (GPi), which were affected by the stimulation of
7
8 the pre-SMA. In contrast, such causal effects were not observed in the activity and functional
9
10 interactions involving the right IFC and the subthalamic nucleus (SN), suggesting that the rTMS
11
12 over the pre-SMA affected the indirect pre-SMA–STR–GPi pathway by possible top-down
13
14 regulation from the pre-SMA to the GPi via STR during successful response inhibition. In an
15
16 attempt to further investigate this point and to exclude the possible role of attentional capture of the
17
18 stop signal presentation, Lee and colleagues (2016) investigated the causal relations of the pre-SMA
19
20 and the rIFG in a conditional SST by combining cTBS and a continue condition task, which
21
22 requires the same motor response as in a go trial but captures attention as in a stop trial, since
23
24 participants were requested to respond to a continue signal with a go response rather than a stop
25
26 response. The analysis took into account individual differences in the degree of slowing in continue
27
28 trials (high versus low) and showed that the selective disruption of right pre-SMA activity
29
30 significantly influenced the continue process only in the low-slowness group, whereas disruption of
31
32 the rIFG did not lead to any significant changes in performance irrespective of the degree of
33
34 slowing. The authors concluded that the pre-SMA plays a critical role in response slowing in
35
36 continue trials during conditional stopping, and it is likely that its efficiency in updating motor
37
38 planning and reinitiating an inhibited response was associated with the amount of slowing, as
39
40 previously suggested by fMRI data (Sharp et al., 2010). In the same year, Xu and colleagues
41
42 investigated the effect of online single pulses of TMS at different intensities (i.e., high =120%,
43
44 medium = 80%, and low = 40% of the individual rMT) over the pre-SMA during fMRI within the
45
46 fronto-basal-ganglia network and the relation of these changes to the response inhibition
47
48 performance. The authors found that TMS induced a BOLD signal increase within the fronto-basal-
49
50 ganglia inhibitory network. In particular, the high-intensity TMS changed the BOLD signal in the
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 right pallidum and left caudate, which also correlated specifically with the SSRT performance (i.e.,
2 increased in connectivity and decreased in SSRT). These results suggest that the widespread effect
3
4 of pre-SMA TMS, at least at the suprathreshold level, was immediate and related to the task-free
5
6
7 neural activity associated with response inhibition.
8

9 To go a step further in establishing the temporal and causal interplay between the pre-SMA and the
10 rIFG, Allen, Singh, Verbruggen, & Chambers (2018) combined the high temporal and reasonable
11 spatial resolution of magnetoencephalography (MEG) with TMS to elucidate the temporal
12 dynamics of the interplay between the IFC and the pre-SMA in an SST. Indeed, very few studies
13 have addressed this issue, and only one study on a single subject, has used intracranial recording to
14 demonstrate that activity related to stopping in the pre-SMA can precede IFC activation (Swann et
15 al., 2012). Allen and colleagues employed the same SST in both MEG and TMS sessions. As TMS
16 was functionally guided according to the MEG results, participants underwent the TMS session on a
17 separate day, after their MEG session. Single pulses of TMS were delivered at 110% of each
18 participant's rMT and the timing of the TMS pulse was dependent upon an individual participant's
19 tROI, which represents the limited period during which either the pre-SMA or the IFC can exert an
20 influence, being situated between the onset of the stop signal and the action. Data showed larger
21 SSRT during active TMS relative to sham, but no differences were found across temporal
22 stimulation conditions (i.e., the temporal order in which TMS was applied to the regions, initial pre-
23 SMA TMS or initial IFC TMS), demonstrating that the application of TMS to both regions
24 disrupted stopping, consistent with the critical role of both the pre-SMA and IFC in inhibitory
25 control. Importantly, however, the temporal order in which TMS was applied to the regions made
26 no difference to the disruption of stopping, in contrast with the idea that information is transmitted
27 from the pre-SMA to the right IFC, and that the pre-SMA may therefore be the cortical source of
28 the stop signal. These results were additionally supported by MEG data, which suggested that the
29 time course of activity originating from the two regions during response inhibition is roughly
30 simultaneous, possibly mirroring high levels of mutually interdependent activity. Granger causality
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 tests suggested that while neither region has been shown to initiate or precipitate a cascade of events
2 in the response inhibition process, it is possible that there is an ongoing drive from pre-SMA to IFC
3 during the interplay between the regions during stopping. Hence, the authors concluded that a pre-
4 SMA to IFC drive may correspond to a proactive state of preparation, as opposed to a specific
5 direction of influence in the actual stopping of an action. The effectiveness of different TBS
6 protocols was tested in a recent study (Ji et al., 2019) which tested their efficiency in affecting the
7 response inhibition ability by interfering with the pre-SMA activity. Changes in performance in an
8 SST were assessed comparing inhibition performance in two sessions performed pre and post
9 different TMS protocols (i.e., 5-Hz, 25-Hz at 110% of the rMT, intermittent (i) TBS at 70% of the
10 rMT, and sham stimulation). Results showed performance improvement (i.e., shorter SSRT) in the
11 iTBS session compared to sham, while no effects were observed in the 5-Hz and 25-Hz rTMS
12 protocols. This effect, however, only occurred during the second session of the SST after iTBS.
13 However, the effect of iTBS was not reproduced if participants only received sham or iTBS.
14 Therefore, further investigation is needed to account for the high intersubjective variability.

15 So far, NIBS studies have mainly tested the crucial role of the pre-SMA and rIFG in action control,
16 while, to the best of our knowledge, only one study undertook to investigate the critical role of the
17 FEF in action control (Muggleton, Chen, Tzeng, Hung, & Juan, 2010). Interestingly, a single unit
18 recordings study in monkeys (Thompson, Biscoe, & Sato, 2005) found that manual responses are
19 associated with inhibition of FEF movement neurons when the task does not require a saccade to be
20 made, suggesting that the FEF might serve a general function related to inhibitory control in the
21 absence of the involvement of eye movements. By applying TMS over the FEF during an SST,
22 Muggleton and colleagues (2010) found that TMS over the FEF had no effect on response
23 generation (indexed by go-RT) but disrupted inhibitory control (i.e., longer SSRT compared to
24 control vertex stimulation and no TMS). This data suggested that the FEF may play a role in
25 inhibiting actions that do not require eye movements. Finally, Osada and colleagues in 2019
26 investigated the crucial role of the posterior parietal cortex, which has reciprocal projections with

1 the prefrontal cortex (Cavada & Goldman- Rakic, 1989), in action inhibition. The authors found
2 that the stimulation over the intraparietal sulcus (IPS) (single pulses of TMS delivered at 120%
3 rMT), disrupted performance in the SST (i.e., longer SSRT) when stimulation was applied 30–0 ms
4 before stopping.
5
6

7
8 In summary, the FEF and IPS, like the IFG and pre-SMA, are involved in successful inhibitory
9 performance with TMS delivered over this area, resulting in elevated SSRT and indicative of
10 altered inhibitory control.
11
12
13
14
15

16
17
18
19 -----
20
21 Please_insert_Coloured_Figure_1_near_here
22
23
24

25
26 -----
27
28 -----
29
30
31
32 Please_insert_Table_1_near_here
33
34
35
36 -----
37
38
39
40

41 **3. Transcranial direct current stimulation and action inhibition**

42 A way to gain further evidence about the neural network implied in the action inhibition is the use
43 of tDCS. Thus, the application of tDCS to the AIN would be a useful adjuvant intervention
44 modality for modulation of response inhibition and its related dynamic behavioral changes. One of
45 the first hypotheses of using this protocol in the SST framework was that excitatory tDCS, with the
46 anode placed over the pre-SMA, would facilitate inhibitory control and that inhibitory tDCS, with
47 the cathode placed over the pre-SMA, would impair performance. Such effects would not only
48 confirm the critical role of the pre-SMA in the inhibitory control network, but, importantly, may
49 demonstrate that a behavioral improvement in inhibitory control is possible by means of the
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

excitatory stimulation of this area. Such tDCS-induced improvement may be fruitful as an initial step towards developing clinical treatments aiming at enhancing inhibitory control (Schroeder, Schwippel, Wolz, & Svaldi, 2020).

In the seminal study of Hsu and coworkers (2011), participants performed an SST before and after tDCS administration (1.5 mA, 10 min). Two groups of participants were tested: one group underwent tDCS **with the anode and cathode placed over the left pre-SMA** (tDCS-pre-SMA) or as sham in separate sessions, while another group was stimulated **with the anode and cathode placed over the left primary motor cortex (tDCS-M1)** or as sham. The results showed that **excitatory** tDCS selectively over the pre-SMA improved the efficiency of inhibitory control (i.e., reduced errors). Conversely, **inhibitory** tDCS **of the pre-SMA activity** showed a tendency towards impaired inhibitory control. Thus, the main finding of the study was that selective error rates in the stop signal trials were significantly modulated by the tDCS polarity: they were reduced by **excitatory** stimulation and increased by **inhibitory** stimulation, confirming the critical role of the pre-SMA in inhibitory control (see also Liang et al., 2014 for similar findings after **excitatory** stimulation of the pre-SMA activity and Yu, Tseng, Hung, Wu, & Juan, 2015, who also demonstrated that **excitatory tDCS-induced BOLD increases in the ventromedial prefrontal cortex (vmPFC), which was positively correlated with participants' improvement in stopping efficiency**). No effects on RT or SSRT were observed either in the tDCS-pre-SMA or in the tDCS-M1 group **(for similar results see Bender, Filmer, & Dux, 2017; Reinhart & Woodman, 2014)**.

The lack of significant results in the SSRT may be due to the fact that the authors set the SSD based on the participants' responses, and thus SSD made the task too simple, preventing SSRT estimation in some participants.

To further investigate the role of the pre-SMA, in a later study (Kwon & Kwon, 2013) the authors examined whether tDCS over the pre-SMA and M1 alters the SSRT using the SST. Participants were tested in two separate sessions, before and after tDCS administration (1 mA, 10 min) **with the anode placed** over the pre-SMA, or over the primary motor cortex (M1) as active control condition,

1 or sham stimulation. Findings demonstrated significant reductions in the SSRT and longer RT after
2 pre-SMA stimulation compared to both M1 and sham stimulations (see Kwon et al., 2013 for a
3 decrease of the SSRT after and during tDCS with the anode placed over M1). No significant effect
4 after administration of the tDCS was found in the go trials. These data suggested that excitatory
5 tDCS administered over the pre-SMA resulted in enhancement of the inhibitory control.
6

7 Within the AIN, response inhibition has also been localized in the IFG, based on both functional
8 brain imaging and lesion-based approaches (Aron et al., 2003; Chevrier et al., 2007; Leung & Cai,
9 2007; Zhang et al., 2017). In the seminal study by Jacobson, Javitt, & Lavidor (2011), the authors
10 used the SST to investigate whether the rIFG is crucial for action inhibition, by applying excitatory
11 and inhibitory tDCS (1 mA, 10 min). Participants underwent different tDCS conditions: anode
12 placed unilaterally over the rIFG, cathode placed unilaterally over the rIFG, bilateral anode/cathode
13 placed over the rIFG and sham. Furthermore, in a control task, participants were required to
14 discriminate visual shapes similar to those presented in the SST; however, no stop signal was given
15 and, in a control experiment, the right angular gyrus was stimulated as control site (rAG). The
16 authors found that the activation of the rIFG by unilateral excitatory stimulation significantly
17 improved response inhibition relative to a sham condition (i.e., reduced SSRT; which was also
18 recently confirmed in a study demonstrating strengthened behavioral response inhibition to external
19 food images compared to sham after excitatory rIFG stimulation; Chen, Jackson, Dong, Zhang, &
20 Chen, 2019), whereas excitatory stimulation of both the rIFC and rAG did not affect RT in the go
21 trials of the SST and control task. Furthermore, the SST was not affected by tDCS delivered over
22 the control site (rAG). The authors only found a near-significant effect for bilateral stimulation and
23 suggested that the left IFG (lIFG) may also be involved in response inhibition. Indeed, only when
24 the activity of the rIFG was facilitated (by unilateral excitatory stimulation) did the participants
25 show reduced SSRT compared to the bilateral stimulation in which the rIFG was facilitated and
26 lIFG was inhibited.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 In addition, when only the rIFG was inhibited (by placing the cathode unilaterally), participants
2 showed longer SSRT compared with the bilateral stimulation condition in which the rIFG was
3 inhibited and the lIFG was facilitated. These findings suggest a critical role of both the right and left
4 IFG in action control.
5
6

7
8
9 In the following year, Ditye, Jacobson, Walsh, & Lavidor (2012) investigated the effects of tDCS
10 placing the anode over the rIFG in combination with a behavioral training using an SST. This study
11 aimed to investigate the additional role of repetitive behavioral training combined with tDCS over
12 the rIFG. The authors investigated the effect of the training integrated with rIFG-tDCS on
13 behavioral inhibition and specifically tested whether i) multiple-session training is effective in
14 improving the ability to inhibit responses and ii) the stimulation of the rIFG would further
15 induce/facilitate training-induced improvements. Participants were randomly assigned to an
16 excitatory tDCS group or to a control group without tDCS. Prior to performing the task, the
17 experimental group received excitatory tDCS (1.5 mA, 15 min) over the rIFG. Importantly, tDCS
18 was delivered every day (over a total of 5 days), except on the last day, because the fifth day was
19 aimed to investigate the sustainability of the effects of tDCS, 24 h after stimulation had stopped.
20 Results suggested that behavioral training effectively improved the ability to inhibit responses and
21 that the combination of SST training with tDCS generated a greater effect than multiple-session
22 training alone. The only significant difference across groups was found at day 3 and 4 (i.e., reduced
23 SSRT in the group with the anode placed over the rIFG) but not at day 5, suggesting that the
24 beneficial effect of tDCS is rather short-lived. An important limitation of this study is the lack of a
25 placebo stimulation group, preventing the possibility to rule out unspecific effects of brain
26 stimulation.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52
53 To sum up, the rIFG and pre-SMA have been the main focus of these attempts to modulate response
54 stopping by means of NIBS. However, other cortical areas such as the dorsolateral prefrontal cortex
55 (dlPFC) have been found to be implicated in inhibitory control (Bari & Robbins, 2013; Leung &
56 Cai, 2007). Therefore, to investigate the crucial role of the right dlPFC (rdlPFC), Stramaccia and
57
58
59
60
61
62
63
64
65

1 coworkers (2015) delivered tDCS (1.5 mA, 20 min) over two target areas: the rIFG and rdIPFC.
2 The participants, divided into five experimental groups based on whether their tDCS stimulation
3 was with the anode placed over the rIFG or the rdIPFC, with the cathode placed over the rIFG or the
4 rdIPFC, or sham. The SST was performed 15 min after the tDCS stimulation (delayed task). Results
5 revealed reduced SSRT in the group with the anode placed over the rIFG relative to sham (but see
6 Stramaccia, Penolazzi, Altoè, & Galfano, 2017). This finding suggests that the rdIPFC does not
7 have a crucial role in response stopping. It is worth noting that this response stopping improvement
8 is unlikely to result from a general cognitive enhancement, since no significant effects were found
9 in the go trials. Indeed, such improvement is more likely to reflect increased inhibitory
10 performance, due to the selective results on SSRT. However, this null result would not necessarily
11 imply that the rdIPFC plays no role in inhibitory processing, but it is possible that the rdIPFC
12 stimulation is short-lasting and hence not evident in the delayed protocol used in this study. Null
13 effects after tDCS in SSRT were also found when placing the anode over the orbitofrontal cortex
14 (OFC) (Ouellet et al., 2015), demonstrating that inhibitory motor control is primarily mediated by
15 non-OFC regions (e.g., pre-SMA, IFG and FEF). In the study by Mansouri and colleagues (2017),
16 participants performed an SST while they listened to high-tempo music, low-tempo music, or
17 background-noise. SST was performed before and after tDCS (1.5 mA, 10 min) with the anode over
18 the rdIPFC or sham. Practice led to a decrease in SSRT and an increase in RT, which was found to
19 be significant only in the low-tempo music condition and for the background-noise condition (sham
20 stimulation). The authors suggested that practice-related changes in response inhibition may be
21 attenuated by high-tempo music. Indeed, such attenuation was restored by the excitatory stimulation
22 of the rdIPFC, demonstrating that the tDCS modulatory effect was dependent on the background
23 music (i.e., high-tempo music), which may improve the inhibitory performance.

24 To further investigate the role of rdIPFC in the action inhibition Friehs and Frings (2018) used an
25 SST and stimulated the rdIPFC in a pre-post experimental design employing excitatory tDCS (0.5
26 mA, 19 min). The authors contrasted the effect of an excitatory stimulation with sham and they

1 expected to observe an increase in inhibitory functions after tDCS with the anode placed over the
2 rdIPFC, as evidenced by a reduction in SSRT. At the beginning participants performed a pre-test
3 SST block; subsequently half the sample underwent the excitatory tDCS stimulation, whereas the
4 other half received a sham stimulation. Afterward, participants were tested in a post-test SST. The
5 results showed that applying tDCS with the anode placed over the rdIPFC relative to sham reduced
6 SSRT and omission errors. Importantly, since RT were not affected by the stimulation, possible
7 speed-up of responses due to the stimulation can be excluded. Although Stramaccia et al., 2015
8 were not able to point out the involvement of dlPFC, this study suggests a possible role of the
9 rdIPFC in cognitive inhibition processes and further emphasizes that inhibitory control is a
10 malleable mechanism. A similar and subsequent study (Friebs & Frings, 2019) used the same
11 experimental design placing the cathode over the rdIPFC (0.5 mA, 19 min) to potentially hamper
12 the response inhibition process as measured by the SST. Results showed that the inhibitory tDCS
13 decreased performance by increasing SSRT, without affecting omission errors in stop trials and go-
14 RT. Taken together, these results support the idea of a critical role of the dlPFC in response
15 inhibition and suggest that polarity-dependent modulation of the response-inhibition process is
16 possible with excitatory stimulation boosts inhibition, whereas inhibitory stimulation diminishes it,
17 although to a lesser degree, in line with previous findings over pre-SMA (Hsu et al., 2011).
18 Finally, a recent work (Fehring et al., 2019) investigated whether excitatory tDCS effects in action
19 inhibition depend on the level of learning in a cognitive task. To this aim, participants completed
20 the SST either before and after tDCS stimulation (1.5 mA, 10 min) with the anode placed over the
21 rdIPFC or sham tDCS stimulation was delivered in two distinct sessions, separated by a one-week
22 washout period. This experimental design was implemented to test within-session learning (changes
23 in performance on the same day) and across-sessions learning (changes in performance after one
24 week). Results showed that excitatory tDCS had an across-week effect on behavioral measures (i.e.,
25 longer go-RT and shorter SSRT), suggesting that the impact of the tDCS was dependent on the
26 level of experience learning. When excitatory stimulation was delivered in the first week (before the

1 occurrence of across session learning), shorter SSRT was observed in the post-tDCS testing
2 (within-session learning), however when it was delivered in the second week, the magnitude of
3 within-session learning was significantly attenuated (i.e., no effect on SSRT was observed). Taken
4 together, these results suggested that plastic changes induced by learning may have altered the
5 susceptibility of the prefrontal cortex to be further modulated by tDCS. However, such practice-
6 related learning seems to saturate or attenuate the plasticity of the neurocircuitry; and therefore,
7 attenuates tDCS differences across sections delivered in separate weeks.
8
9

10
11
12
13
14
15
16
17 In summary, according to several studies, tDCS administered with the anode placed over the IFG,
18 pre-SMA, dlPFC and M1 reduced SSRT, improving behavioural performance. However, it is
19 important to consider that all these studies investigated the critical role of each of these components
20 independently, therefore, information about the neural interplay among these areas remains obscure.
21
22

23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

As mentioned in the introduction, inhibitory control can be divided into proactive mechanisms (response slowing in anticipation of a stop-signal) and reactive mechanisms (outright stopping in response to a stop-signal). Therefore, an important issue concerns the relationship between proactive and reactive inhibitory control, and whether the same neural substrate subserves both mechanisms.

To assess the causal role of the rIFG and IPL, which was found to be more strongly activated in the stop than go trials (Congdon et al., 2010; Hughes et al., 2014), in proactive and reactive inhibitory control, Cai and colleagues in 2015 asked participants to complete an SST after the administration of tDCS (1.5 mA, 15 min) with the anode placed over the rIFG, rIPL or the visual primary cortex (V1) as active control condition in a within-subjects design. To index proactive inhibitory control, the authors employed the preparatory cost (PC) index, which was calculated including only correct go trials by subtracting the mean RT of the first two post-stop go trials from that of the later post-stop go trials (i.e., higher PC indicates better proactive inhibitory control). Instead, reactive inhibitory control was indexed by SSRT. Compared to the V1 stimulation, excitatory stimulation of the rIFG facilitated both proactive and reactive control (i.e., after rIFG stimulation larger PC and

1 greater reduction of SSRT was found, see Li et al., 2019 for similar results on SSRT). No
2 significant results were obtained by stimulating the IPL. Moreover, since there was a significant
3 negative correlation between the PC index and the SSRT, the authors conducted a mediation
4 analysis to examine whether the tDCS effect on reactive control was mediated by proactive control.
5 The results showed no significant mediation effect, and the effect of stimulation condition on the
6 SSRT change was still significant after adding the mediator. Thus, the results suggest that the tDCS
7 effect on reactive inhibitory control was not mediated by facilitation of proactive inhibitory control.
8 Another work investigated the role of the rIFC in the proactive and reactive inhibitory process
9 (Cunillera, Fuentemilla, Brignani, Cucurell, & Miniussi, 2014) using a combination of go/no-go and
10 SST. TDCS (1.5 mA, 18 min) with the anode placed over the rIFC or sham were delivered in a
11 within-subjects design. The task consisted in a choice-reaction go/no-go task in which there were
12 two variant perceptual complexity conditions, based on the visual discriminability of the stimuli.
13 Results showed shorter SSRT in the excitatory rIFC condition relative to sham, reflecting a
14 behavioral inhibitory improvement. Moreover, excitatory tDCS slowed down the RT, and reduced
15 error rates when the stimuli discriminability was easier. These results indicate a possible increase in
16 task control, which was interpreted as a proactive inhibitory process. A subsequent work (Cunillera,
17 Brignani, Cucurell, Fuentemilla, & Miniussi, 2016) used the same task in a within-subject design in
18 which online tDCS (1.5 mA, 20 min) with the anode placed over the rIFC or sham stimulation were
19 delivered during the SST. The authors found a significant increase in RT for excitatory tDCS of the
20 rIFC activity relative to sham, which was interpreted as increased proactive inhibition. However, in
21 contrast to the previous result of the same authors (Cunillera et al., 2014), excitatory stimulation
22 was not observed to be effective in modulating reactive inhibition, as indicated by the lack of
23 significant effect in the SSRT. A possible explanation for the null effect of tDCS may be related to
24 the different protocol of tDCS used between these studies (i.e., online vs offline stimulation). An
25 additional study (Castro-Meneses, Johnson, & Sowman, 2016) compared reactive inhibition across
26 manual and vocal effector systems (i.e., key pressing and vocal responses, respectively), examined

1 the effects of tDCS (1.5 mA, 15 min) with the anode placed over the right prefrontal cortex (rPFC)
2 and looked at the relationship between reactive and proactive inhibition. Results showed that both
3
4 vocal and manual reactive inhibition was faster (i.e., shorter SSRT) during and immediately after
5
6
7 excitatory tDCS relative to sham. Moreover, a greater level of proactive inhibition (i.e., calculated
8
9 by subtracting the mean of the go-RTs of the relevant stop condition from the mean of the go-RTs
10
11 of the irrelevant stop condition; see Chikazoe et al., 2009) enhanced the reactive inhibition (i.e.,
12
13 when more proactive inhibition is implemented, SSRT gets shorter), supporting the hypothesis that
14
15 the rPFC is part of a core network for reactive inhibition and supports the previous contention that
16
17 proactive inhibition is possibly modulated via pre-activating the reactive inhibition network.
18
19

20
21 In an attempt to improve the spatial focality of tDCS, a recent study (Hogeveen et al., 2016) used
22
23 high-definition tDCS (HD-tDCS) or conventional tDCS in two separate groups of participants to
24
25 stimulate the rIFC during an SST or during a control task (choice reaction task; CRT). A third
26
27 group of participants completed the same behavioral protocol but received conventional tDCS to a
28
29 control site (mid-occipital cortex). The study employed a pretest–posttest design, repeated over two
30
31 sessions. Thus, in the first session, participants completed the SST, then they underwent one of the
32
33 three tDCS conditions while doing the same task, followed by another SST. The second session was
34
35 equivalent to the first, with the exception that participants completed a choice reaction time task
36
37 (CRT). SSRT analysis revealed significant performance improvements (i.e., shorter SSRT) in
38
39 inhibitory control in both the conventional and the HD-tDCS group compared to the control group
40
41 (mid-occipital cortex stimulation). The results suggest that HD-tDCS and conventional tDCS
42
43 (delivered over rIFC) have statistically similar effects on response inhibition.
44
45
46
47
48
49

50
51 Finally, in a recent study by Sandrini and colleagues (2020), participants performed an SST before
52
53 (on a first day) and after (day 2) tDCS with the anode placed over the rIFC (1.5 mA, 20 min) or
54
55 sham in separate groups of participants. The study design included a resting-state functional
56
57 magnetic resonance imaging (rsfMRI) before and after tDCS, and an event-related fMRI (efMRI)
58
59 during the SST immediately following the post-tDCS rsfMRI on day 2.
60
61
62
63
64
65

1 Consistent with previous findings (Jacobson et al., 2011; Stramaccia et al., 2015), results (calculated
2 on the difference in SSRT between session 2 and session 1) showed a significant difference
3 between the **excitatory** and sham tDCS groups, with shorter SSRT found in the group **with the**
4 **anode placed over the rIFC** relative to the sham group. In addition, no significant differences were
5 observed between groups in RT. **Regarding the fMRI results, rsfMRI revealed a significant increase**
6 **in the functional connectivity between the rIFC and caudate, and between the rIFC, right pre-SMA,**
7 **right inferior parietal cortex (rIPC), and rdIPFC after excitatory tDCS. Moreover, efMRI results**
8 **showed that the excitatory tDCS strengthened the functional connectivity between the right pre-**
9 **SMA and subthalamic nuclei during stop responses, suggesting that the stimulation of the rIFC**
10 **induces neural modulation in several brain regions, such as the rIFC, rdIPFC, right pre-SMA, basal**
11 **ganglia, and their interconnected brain regions (e.g., rIPC).**

12 To conclude, the aforementioned tDCS studies suggested that proactive and reactive action
13 inhibition mechanisms are partially independent and the existing studies converge in suggesting the
14 crucial role of the rIFG in both reactive and proactive action control while, to the best of our
15 knowledge, to date no studies have undertaken to specifically test the role of the pre-SMA **in the**
16 **proactive and reactive mechanisms** by using tDCS.

17 -----
18
19 Please_insert_Coloured_Figure_2_near_here
20
21 -----
22
23 -----

24
25
26 Please_insert_Table_2_near_here
27
28 -----

29 **Conclusions**

1 Action control represents a complex mechanism that subserves several different processes such as
2 inhibition, selection, competition and switching of actions with a composite network of prefrontal
3 areas (AIN) that have a role in orchestrating such processes. Indeed, deficits in performance in
4 action control have been seen in various psychiatric disorders, including attention-deficit
5 hyperactivity disorder and conduct disorder (Schachar & Logan, 1990; Schachar, Tannock, &
6 Logan, 1993; Schachar, Tannock, Marriott, & Logan, 1995). It has also been shown to be altered in
7 cocaine dependent men (Li, Milivojevic, Kemp, Hong, & Sinha, 2006), autism (Kana, Keller,
8 Minshew, & Just, 2007) and obsessive compulsive disorder (Chamberlain, Blackwell, Fineberg,
9 Robbins, & Sahakian, 2005), while impulsive-violent offenders show elevated SSRT (Chen,
10 Muggleton, Juan, Tzeng, & Hung, 2008). Therefore, disclosing the specific and critical role of the
11 different components of the AIN represents a crucial challenge to pave the way for designing novel
12 NIBS therapeutic interventions aimed at enhancing the ability to improve cognitive control and
13 inhibit potentially dangerous actions. Here we have reviewed the major research advances using the
14 combination of SST with the use of NIBS over the last fifteen years (Juan & Muggleton, 2012).
15 Although sometimes controversial, results suggest that both the pre-SMA and IFG (especially the
16 right) may play a crucial role in (reactive and proactive) control of actions. The most controversial
17 findings derived from studies assessing the proactive inhibitory control. Such inconsistency may be
18 ascribed to the different paradigms used to investigate this process, relative to the consistency in the
19 paradigms used to measure the reactive inhibitory (usually by means of classical SST). FEF
20 represents an additional critical area, at least in reactive action control that does not require eye
21 movements, while dlPFC was found to be critical in the proactive (but not reactive) control of
22 action. An unresolved issue is whether the left IFG plays a crucial role in action inhibition since, to
23 the best of our knowledge, only few studies employed NIBS to test its critical role (Chambers et al.,
24 2007; Jacobson et al., 2011) and they reported partially contradicting results. Therefore we believe
25 that the most parsimonious account is to suggest that both the right and left IFG are probably
26 involved in response inhibition (in line with fMRI data of bilateral activations); however, future

1 studies should directly investigate the critical role of the left IFG involvement in response
2 inhibition.

3
4 To conclude, it seems that pre-SMA and IFG do not play a selective role in inhibitory control but in
5 a range of inhibitory behaviors including response inhibition, selection and competition. One
6 potential limitation of the NIBS techniques is that the magnetic/electrical field falls off very rapidly
7 with distance from the TMS coil/tDCS electrodes, meaning that superficial areas of cortex are easy
8 to stimulate, but those deep in a sulcus or far from the scalp surface are more complicated to
9 stimulate (Siebner, Hartwigsen, Kassuba, & Rothwell, 2009). Consequently, some of the AIN
10 components cannot be investigated by NIBS, such as the subcortical structures (e.g., basal ganglia),
11 thus their crucial role remains obscure. Moreover, it needs to be taken into account that tDCS have
12 been found to produce spatially widespread electric fields (EF) on the cerebral cortex, hence,
13 individual modeling is required to plan EF doses when focal montages are used (Mikkonen, Laakso,
14 Tanaka, & Hirata, 2020). Beside this limitation, several progresses have been achieved in the last
15 fifteen years in healthy participants, however future studies should assess not only the critical role
16 of the single nodes of the AIN but also their interactions by studying their effective connectivity
17 (Zanon, Borgomaneri, & Avenanti, 2018) and/or their causal interactions (Chiappini et al., 2020;
18 Fiori et al., 2017, 2016). All this information is fundamental to carefully assess the clinical and
19 applied potentialities of NIBS in clinical populations (Plewnia, Schroeder, & Wolkenstein, 2015).

20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 **Acknowledgments**

47 This work was supported by grants from Ministero della Salute, Italy [GR-2018-12365733]
48 awarded to Sa.B.

49 50 51 52 53 54 55 56 **Declaration of Interests**

57 The authors declare no competing interests
58
59
60
61
62
63
64
65

References

- 1
2 Allen, C., Singh, K. D., Verbruggen, F., & Chambers, C. D. (2018). Evidence for parallel activation
3
4 of the pre-supplementary motor area and inferior frontal cortex during response inhibition: A
5
6 combined MEG and TMS study. *Royal Society Open Science*, *5*(2), 171369.
7
8
9 <https://doi.org/10.1098/rsos.171369>
10
11
12 Aron, A. R. (2011). From reactive to proactive and selective control: Developing a richer model for
13
14 stopping inappropriate responses. *Biological Psychiatry*, *69*(12), 55–68.
15
16
17 <https://doi.org/10.1016/j.biopsych.2010.07.024>
18
19
20 Aron, A. R., Behrens, T. E., Smith, S., Frank, M. J., & Poldrack, R. A. (2007). Triangulating a
21
22 cognitive control network using diffusion-weighted magnetic resonance imaging (MRI) and
23
24 functional MRI. *The Journal of Neuroscience*, *27*(14), 3743–3752.
25
26
27 <https://doi.org/10.1523/JNEUROSCI.0519-07.2007>
28
29
30 Aron, A. R., Fletcher, P. C., Bullmore, E. T., Sahakian, B. J., & Robbins, T. W. (2003). Stop-signal
31
32 inhibition disrupted by damage to right inferior frontal gyrus in humans. *Nature Neuroscience*,
33
34 *6*(2), 115–116. <https://doi.org/10.1038/nn1003>
35
36
37 Aron, A. R., & Poldrack, R. A. (2006). Cortical and subcortical contributions to stop signal
38
39 response inhibition: Role of the subthalamic nucleus. *Journal of Neuroscience*, *26*(9), 2424–
40
41 2433. <https://doi.org/10.1523/JNEUROSCI.4682-05.2006>
42
43
44 Aron, A. R., Robbins, T. W., & Poldrack, R. A. (2004). Inhibition and the right inferior frontal
45
46 cortex. *Trends in Cognitive Sciences*, *8*(4), 170–177. <https://doi.org/10.1016/j.tics.2004.02.010>
47
48
49 Aron, A. R., & Verbruggen, F. (2008). Stop the Presses. Dissociating a selective from a global
50
51 mechanism for stopping. *Psychological Science*, *19*(11), 1146–1153.
52
53
54 <https://doi.org/10.1111/j.1467-9280.2008.02216.x>
55
56
57 Bari, A., & Robbins, T. W. (2013). Inhibition and impulsivity: Behavioral and neural basis of
58
59 response control. *Progress in Neurobiology*, *108*, 44–79.
60
61 <https://doi.org/10.1016/j.pneurobio.2013.06.005>
62
63
64
65

1 Bender, A. D., Filmer, H. L., & Dux, P. E. (2017). Transcranial direct current stimulation of
2 superior medial frontal cortex disrupts response selection during proactive response inhibition.
3
4 *NeuroImage*, 158, 455–465. <https://doi.org/10.1016/j.neuroimage.2016.10.035>
5
6

7 Bergmann, T. O., Karabanov, A., Hartwigsen, G., Thielscher, A., & Siebner, H. R. (2016).
8
9 Combining non-invasive transcranial brain stimulation with neuroimaging and
10
11 electrophysiology: Current approaches and future perspectives. *NeuroImage*, 140, 4–19.
12
13 <https://doi.org/10.1016/j.neuroimage.2016.02.012>
14
15

16
17 Boulinguez, P., Ballanger, B., Granjon, L., & Benraiss, A. (2009). The paradoxical effect of
18
19 warning on reaction time: Demonstrating proactive response inhibition with event-related
20
21 potentials. *Clinical Neurophysiology*, 120(4), 730–737.
22
23 <https://doi.org/10.1016/j.clinph.2009.02.167>
24
25

26
27 Braver, T. S., Gray, J. R., & Burgess, G. C. (2007). Explaining the many varieties of working
28
29 memory variation: Dual mechanisms of cognitive control. In A. R. A. Conway, C. Jarrold, & J.
30
31 Kane (Eds.), & A. Miyake & J. N. Towse (Ed.), *Variation in working memory* (pp. 76–106).
32
33 Oxford University Press.
34
35

36
37 Cai, W., George, J. S., Verbruggen, F., Chambers, C. D., & Aron, A. R. (2012). The role of the
38
39 right presupplementary motor area in stopping action: Two studies with event-related
40
41 transcranial magnetic stimulation. *Journal of Neurophysiology*, 108(2), 380–389.
42
43 <https://doi.org/10.1152/jn.00132.2012>
44
45

46
47 Cai, Y., Li, S., Liu, J., Li, D., Feng, Z., Wang, Q., ... Xue, G. (2015). The role of the frontal and
48
49 parietal cortex in proactive and reactive inhibitory control: A transcranial direct current
50
51 stimulation study. *Journal of Cognitive Neuroscience*, 28(1), 177–186.
52
53 https://doi.org/10.1162/jocn_a_00888
54
55

56
57 Castro-Meneses, L. J., Johnson, B. W., & Sowman, P. F. (2016). Vocal response inhibition is
58
59 enhanced by anodal tDCS over the right prefrontal cortex. *Experimental Brain Research*,
60
61 234(1), 185–195. <https://doi.org/10.1007/s00221-015-4452-0>
62
63
64
65

1 Cavada, C., & Goldman-Rakic, P. S. (1989). Posterior parietal cortex in rhesus monkey: I.

2 Parcellation of areas based on distinctive limbic and sensory corticocortical connections.

3
4
5 *Journal of Comparative Neurology*, 287(4), 393–421. <https://doi.org/10.1002/cne.902870402>

6
7 Chamberlain, S. R., Blackwell, A. D., Fineberg, N. A., Robbins, T. W., & Sahakian, B. J. (2005).

8
9 The neuropsychology of obsessive compulsive disorder: The importance of failures in

10
11 cognitive and behavioural inhibition as candidate endophenotypic markers. *Neuroscience and*

12
13 *Biobehavioral Reviews*, 29(3), 399–419. <https://doi.org/10.1016/j.neubiorev.2004.11.006>

14
15
16
17 Chambers, C. D., Bellgrove, M. A., Gould, I. C., English, T., Garavan, H., McNaught, E., ...

18
19 Mattingley, J. B. (2007). Dissociable mechanisms of cognitive control in prefrontal and

20
21 premotor cortex. *Journal of Neurophysiology*, 98(6), 3638–3647.

22
23
24 <https://doi.org/10.1152/jn.00685.2007>

25
26
27 Chambers, C. D., Bellgrove, M. A., Stokes, M. G., Henderson, T. R., Garavan, H., Robertson, I. H.,

28
29 ... Mattingley, J. B. (2006). Executive “brake failure” following deactivation of human frontal

30
31 lobe. *Journal of Cognitive Neuroscience*, 18(3), 444–455.

32
33
34 <https://doi.org/10.1162/089892906775990606>

35
36
37 Chen, C. Y., Muggleton, N. G., Juan, C. H., Tzeng, O. J. L., & Hung, D. L. (2008). Time pressure

38
39 leads to inhibitory control deficits in impulsive violent offenders. *Behavioural Brain Research*,

40
41 187(2), 483–488. <https://doi.org/10.1016/j.bbr.2007.10.011>

42
43
44 Chen, C. Y., Muggleton, N. G., Tzeng, O. J. L., Hung, D. L., & Juan, C. H. (2009). Control of

45
46 prepotent responses by the superior medial frontal cortex. *NeuroImage*, 44(2), 537–545.

47
48
49 <https://doi.org/10.1016/j.neuroimage.2008.09.005>

50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
Chen, S., Jackson, T., Dong, D., Zhang, X., & Chen, H. (2019). Exploring effects of single-session

anodal tDCS over the inferior frontal gyrus on responses to food cues and food cravings

among highly disinhibited restrained eaters: A preliminary study. *Neuroscience Letters*, 706,

211–216. <https://doi.org/10.1016/j.neulet.2019.05.035>

Chevrier, A. D., Noseworthy, M. D., & Schachar, R. (2007). Dissociation of response inhibition and

1 performance monitoring in the stop signal task using event-related fMRI. *Human Brain*
2 *Mapping*, 28(12), 1347–1358. <https://doi.org/10.1002/hbm.20355>

3
4 Chiappini, E., Borgomaneri, S., Marangon, M., Turrini, S., Romei, V., & Avenanti, A. (2020).

5
6
7 Driving associative plasticity in premotor-motor connections through a novel paired

8
9 associative stimulation based on long-latency cortico-cortical interactions. *Brain Stimulation*,

10
11
12 In press. <https://doi.org/10.1016/j.brs.2020.08.003>

13
14 Chikazoe, J., Jimura, K., Hirose, S., Yamashita, K. I., Miyashita, Y., & Konishi, S. (2009).

15
16 Preparation to inhibit a response complements response inhibition during performance of a
17 stop-signal task. *Journal of Neuroscience*, 29(50), 15870–15877.

18
19
20
21 <https://doi.org/10.1523/JNEUROSCI.3645-09.2009>

22
23
24 Clark, S. V., King, T. Z., & Turner, J. A. (2020). Cerebellar contributions to proactive and reactive

25
26 control in the stop signal task: A systematic review and meta-analysis of functional magnetic
27 resonance imaging studies. *Neuropsychology Review*, Online ahead of print.

28
29
30
31 <https://doi.org/10.1007/s11065-020-09432-w>

32
33
34 Congdon, E., Mumford, J. A., Cohen, J. R., Galvan, A., Aron, A. R., Xue, G., ... Poldrack, R. A.

35
36 (2010). Engagement of large-scale networks is related to individual differences in inhibitory
37 control. *NeuroImage*, 53(2), 653–663. <https://doi.org/10.1016/j.neuroimage.2010.06.062>

38
39
40
41 Criaud, M., & Boulinguez, P. (2013). Have we been asking the right questions when assessing

42
43 response inhibition in go/no-go tasks with fMRI? A meta-analysis and critical review.

44
45
46 *Neuroscience and Biobehavioral Reviews*, 37(1), 11–23.

47
48
49 <https://doi.org/10.1016/j.neubiorev.2012.11.003>

50
51 Cunillera, T., Brignani, D., Cucurell, D., Fuentemilla, L., & Miniussi, C. (2016). The right inferior

52
53 frontal cortex in response inhibition: A tDCS–ERP co-registration study. *NeuroImage*, 140,
54 66–75. <https://doi.org/10.1016/j.neuroimage.2015.11.044>

55
56
57
58 Cunillera, T., Fuentemilla, L., Brignani, D., Cucurell, D., & Miniussi, C. (2014). A simultaneous

59
60 modulation of reactive and proactive inhibition processes by anodal tDCS on the right inferior

frontal cortex. *PLoS ONE*, 9(11), e113537. <https://doi.org/10.1371/journal.pone.0113537>

1
2 Dambacher, F., Sack, A., Lobbstaël, J., Arntz, A., Brugmann, S., & Schuhmann, T. (2014). The
3
4 role of right prefrontal and medial cortex in response inhibition: Interfering with action
5
6 restraint and action cancellation using transcranial magnetic brain stimulation. *Journal of*
7
8 *Cognitive Neuroscience*, 26(8), 1775–1784. https://doi.org/10.1162/jocn_a_00595
9

10
11 Dillon, D. G., & Pizzagalli, D. A. (2007). Inhibition of action, thought, and emotion: A selective
12
13 neurobiological review. *Applied and Preventive Psychology*, 12(3), 99–114.
14
15 <https://doi.org/10.1016/j.appsy.2007.09.004>
16
17

18
19 Dippel, G., & Beste, C. (2015). A causal role of the right inferior frontal cortex in implementing
20
21 strategies for multi-component behaviour. *Nature Communications*, 6, 6587.
22
23 <https://doi.org/10.1038/ncomms7587>
24
25

26
27 Ditye, T., Jacobson, L., Walsh, V., & Lavidor, M. (2012). Modulating behavioral inhibition by
28
29 tDCS combined with cognitive training. *Experimental Brain Research*, 219(3), 363–368.
30
31 <https://doi.org/10.1007/s00221-012-3098-4>
32
33

34
35 Duncan, J., & Owen, A. M. (2000). Common regions of the human frontal lobe recruited by diverse
36
37 cognitive demands. *Trends in Neurosciences*, 23(10), 475–483. [https://doi.org/10.1016/S0166-](https://doi.org/10.1016/S0166-2236(00)01633-7)
38
39 [2236\(00\)01633-7](https://doi.org/10.1016/S0166-2236(00)01633-7)
40

41
42 Durston, S., Tottenham, N. T., Thomas, K. M., Davidson, M. C., Eigsti, I. M., Yang, Y., ... Casey,
43
44 B. J. (2003). Differential patterns of striatal activation in young children with and without
45
46 ADHD. *Biological Psychiatry*, 53(10), 871–878. [https://doi.org/10.1016/S0006-](https://doi.org/10.1016/S0006-3223(02)01904-2)
47
48 [3223\(02\)01904-2](https://doi.org/10.1016/S0006-3223(02)01904-2)
49
50

51
52 Erika-Florence, M., Leech, R., & Hampshire, A. (2014). A functional network perspective on
53
54 response inhibition and attentional control. *Nature Communications*, 5, 4073.
55
56 <https://doi.org/10.1038/ncomms5073>
57

58
59 Fehring, D. J., Illipparampil, R., Acevedo, N., Jaberzadeh, S., Fitzgerald, P. B., & Mansouri, F. A.
60
61 (2019). Interaction of task-related learning and transcranial direct current stimulation of the
62
63
64
65

prefrontal cortex in modulating executive functions. *Neuropsychologia*, 131, 148–159.

<https://doi.org/10.1016/j.neuropsychologia.2019.05.011>

Fiori, F., Chiappini, E., Candidi, M., Romei, V., Borgomaneri, S., & Avenanti, A. (2017). Long-latency interhemispheric interactions between motor-related areas and the primary motor cortex: A dual site TMS study. *Scientific Reports*, 7(1), 14936. <https://doi.org/10.1038/s41598-017-13708-2>

Fiori, F., Chiappini, E., Soriano, M., Paracampo, R., Romei, V., Borgomaneri, S., & Avenanti, A. (2016). Long-latency modulation of motor cortex excitability by ipsilateral posterior inferior frontal gyrus and pre-supplementary motor area. *Scientific Reports*, 6, 38396. <https://doi.org/10.1038/srep38396>

Floden, D., & Stuss, D. T. (2006). Inhibitory control is slowed in patients with right superior medial frontal damage. *Journal of Cognitive Neuroscience*, 18(11), 1843–1849. <https://doi.org/10.1162/jocn.2006.18.11.1843>

Forstmann, B. U., Dutilh, G., Brown, S., Neumann, J., Von Cramon, D. Y., Ridderinkhof, K. R., & Wagenmakers, E. J. (2008). Striatum and pre-SMA facilitate decision-making under time pressure. *Proceedings of the National Academy of Sciences of the United States of America*, 105(45), 17538–17542. <https://doi.org/10.1073/pnas.0805903105>

Fried, I., Katz, A., McCarthy, G., Sass, K. J., Williamson, P., Spencer, S. S., & Spencer, D. D. (1991). Functional organization of human supplementary motor cortex studies by electrical stimulation. *Journal of Neuroscience*, 11(11), 3656–3666. <https://doi.org/10.1523/jneurosci.11-11-03656.1991>

Friebs, M. A., & Frings, C. (2018). Pimping inhibition: Anodal tDCS enhances stop-signal reaction time. *Journal of Experimental Psychology: Human Perception and Performance*, 44(12), 1933–1945. <https://doi.org/10.1037/xhp0000579>

Friebs, M. A., & Frings, C. (2019). Cathodal tDCS increases stop-signal reaction time. *Cognitive, Affective and Behavioral Neuroscience*, 19(5), 1129–1142. <https://doi.org/10.3758/s13415->

- 1
2 Garavan, H., Ross, T. J., & Stein, E. A. (1999). Right hemispheric dominance of inhibitory control:
3
4 An event-related functional MRI study. *Proceedings of the National Academy of Sciences of*
5
6 *the United States of America*, 96(14), 8301–8306. <https://doi.org/10.1073/pnas.96.14.8301>
7
8
- 9 George, M. S., & Aston-Jones, G. (2010). Noninvasive techniques for probing neurocircuitry and
10
11 treating illness: Vagus nerve stimulation (VNS), transcranial magnetic stimulation (TMS) and
12
13 transcranial direct current stimulation (tDCS). *Neuropsychopharmacology*, 35(1), 301–316.
14
15 <https://doi.org/10.1038/npp.2009.87>
16
17
- 18 Goldman-Rakic, P. S. (1987). Circuitry of primate prefrontal cortex and regulation of behaviour by
19
20 representational memory. In *Handbook of physiology: The nervous system* (Mountcastl, pp.
21
22 373–417). Bethesda: American Physiological Society.
23
24
- 25 Hamada, M., Hanajima, R., Terao, Y., Arai, N., Furubayashi, T., Inomata-Terada, S., ... Ugawa, Y.
26
27 (2007). Quadro-pulse stimulation is more effective than paired-pulse stimulation for plasticity
28
29 induction of the human motor cortex. *Clinical Neurophysiology*, 118(12), 2672–2682.
30
31 <https://doi.org/10.1016/j.clinph.2007.09.062>
32
33
- 34 Hampshire, A., Chamberlain, S. R., Monti, M. M., Duncan, J., & Owen, A. M. (2010). The role of
35
36 the right inferior frontal gyrus: Inhibition and attentional control. *NeuroImage*, 50(3), 1313–
37
38 1319. <https://doi.org/10.1016/j.neuroimage.2009.12.109>
39
40
- 41 Hanes, D. P., & Schall, J. D. (1996). Neural control of voluntary movement initiation. *Science*,
42
43 274(5286), 427–430. <https://doi.org/10.1126/science.274.5286.427>
44
45
- 46 Hester, R., Murphy, K., & Garavan, H. (2004). Beyond common resources: The cortical basis for
47
48 resolving task interference. *NeuroImage*, 23(1), 202–212.
49
50 <https://doi.org/10.1016/j.neuroimage.2004.05.024>
51
52
- 53 Hodgson, T., Chamberlain, M., Parris, B., James, M., Gutowski, N., Husain, M., & Kennard, C.
54
55 (2007). The role of the ventrolateral frontal cortex in inhibitory oculomotor control. *Brain*,
56
57 130(6), 1525–1537. <https://doi.org/10.1093/brain/awm064>
58
59
60
61
62
63
64
65

- 1 Hogeveen, J., Grafman, J., Abozeria, M., David, A., Bikson, M., & Hauner, K. K. (2016). Effects of
2 high-definition and conventional tDCS on response inhibition. *Brain Stimulation*, *9*(5), 720–
3 729. <https://doi.org/10.1016/j.brs.2016.04.015>
4
5
6
7 Hsu, T. Y., Tseng, L. Y., Yu, J. X., Kuo, W. J., Hung, D. L., Tzeng, O. J. L., ... Juan, C. H. (2011).
8 Modulating inhibitory control with direct current stimulation of the superior medial frontal
9 cortex. *NeuroImage*, *56*(4), 2249–2257. <https://doi.org/10.1016/j.neuroimage.2011.03.059>
10
11
12
13
14 Huang, Y. Z., Edwards, M. J., Rounis, E., Bhatia, K. P., & Rothwell, J. C. (2005). Theta burst
15 stimulation of the human motor cortex. *Neuron*, *45*(2), 201–206.
16
17
18
19 <https://doi.org/10.1016/j.neuron.2004.12.033>
20
21
22 Huang, Y. Z., Lu, M. K., Antal, A., Classen, J., Nitsche, M., Ziemann, U., ... Rothwell, J. (2017).
23 Plasticity induced by non-invasive transcranial brain stimulation: A position paper. *Clinical*
24 *Neurophysiology*, *128*(11), 2318–2329. <https://doi.org/10.1016/j.clinph.2017.09.007>
25
26
27
28
29 Hughes, M. E., Budd, T. W., Fulham, W. R., Lancaster, S., Woods, W., Rossell, S. L., & Michie, P.
30 T. (2014). Sustained brain activation supporting stop-signal task performance. *European*
31 *Journal of Neuroscience*, *39*(8), 1363–1369. <https://doi.org/10.1111/ejn.12497>
32
33
34
35
36 Ito, S., Stuphorn, V., Brown, J. W., & Schall, J. D. (2003). Performance monitoring by the anterior
37 cingulate cortex during saccade countermanding. *Science*, *302*(5642), 120–122.
38
39
40
41 <https://doi.org/10.1126/science.1087847>
42
43
44 Iyer, M. B., Schleper, N., & Wassermann, E. M. (2003). Priming stimulation enhances the
45 depressant effect of low-frequency repetitive transcranial magnetic stimulation. *Journal of*
46 *Neuroscience*, *23*(34), 10867–10872. <https://doi.org/10.1523/jneurosci.23-34-10867.2003>
47
48
49
50
51 Jacobson, L., Javitt, D. C., & Lavidor, M. (2011). Activation of inhibition: Diminishing impulsive
52 behavior by direct current stimulation over the inferior frontal gyrus. *Journal of Cognitive*
53 *Neuroscience*, *23*(11), 3380–3387. https://doi.org/10.1162/jocn_a_00020
54
55
56
57
58 Jahfari, S., Stinear, C. M., Claffey, M., Verbruggen, F., & Aron, A. R. (2010). Responding with
59 restraint: What are the neurocognitive mechanisms? *Journal of Cognitive Neuroscience*, *22*(7),
60
61
62
63
64
65

1479–1492. <https://doi.org/10.1162/jocn.2009.21307>

1
2 Ji, G. J., Wei, J. J., Liu, T., Li, D., Zhu, C., Yu, F., ... Hu, P. (2019). Aftereffect and reproducibility
3
4 of three excitatory repetitive TMS protocols for a response inhibition task. *Frontiers in*
5
6 *Neuroscience*, 13, 1155. <https://doi.org/10.3389/fnins.2019.01155>

7
8
9 Juan, C. H., & Muggleton, N. G. (2012). Brain stimulation and inhibitory control. *Brain*
10
11 *Stimulation*, 5(2), 63–69. <https://doi.org/10.1016/j.brs.2012.03.012>

12
13
14 Jung, S. H., Shin, J. E., Jeong, Y. S., & Shin, H. I. (2008). Changes in motor cortical excitability
15
16 induced by high-frequency repetitive transcranial magnetic stimulation of different stimulation
17
18 durations. *Clinical Neurophysiology*, 119(1), 71–79.
19
20
21 <https://doi.org/10.1016/j.clinph.2007.09.124>

22
23
24 Kana, R. K., Keller, T. A., Minshew, N. J., & Just, M. A. (2007). Inhibitory control in high-
25
26 functioning autism: Decreased activation and underconnectivity in inhibition networks.
27
28 *Biological Psychiatry*, 62(3), 198–206. <https://doi.org/10.1016/j.biopsych.2006.08.004>

29
30
31 Koessler, L., Maillard, L., Benhadid, A., Vignal, J. P., Felblinger, J., Vespignani, H., & Braun, M.
32
33 (2009). Automated cortical projection of EEG sensors: Anatomical correlation via the
34
35 international 10-10 system. *NeuroImage*, 46(1), 64–72.
36
37
38 <https://doi.org/10.1016/j.neuroimage.2009.02.006>

39
40
41 Koski, L., Molnar-Szakacs, I., & Iacoboni, M. (2005). Exploring the contributions of premotor and
42
43 parietal cortex to spatial compatibility using image-guided TMS. *NeuroImage*, 24(2), 296–305.
44
45
46 <https://doi.org/10.1016/j.neuroimage.2004.09.027>

47
48 Kwon, J. W., Nam, S. H., Lee, N. K., Son, S. M., Choi, Y. W., & Kim, C. S. (2013). The effect of
49
50 transcranial direct current stimulation on the motor suppression in stop-signal task.
51
52 *NeuroRehabilitation*, 32(1), 191–196. <https://doi.org/10.3233/NRE-130836>

53
54
55
56 Kwon, Y. H., & Kwon, J. W. (2013). Response inhibition induced in the stop-signal task by
57
58 transcranial direct current stimulation of the pre-supplementary motor area and primary
59
60 sensorimotor cortex. *Journal of Physical Therapy Science*, 25(9), 1083–1086.
61
62

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

<https://doi.org/10.1589/jpts.25.1083>

Lacadie, C. M., Fulbright, R. K., Rajeevan, N., Constable, R. T., & Papademetris, X. (2008). More accurate Talairach coordinates for neuroImaging using nonlinear registration. *NeuroImage*, 42(2), 717–725. <https://doi.org/10.1016/j.neuroimage.2008.04.240>

Lappin, J. S., & Eriksen, C. W. (1966). Use of a delayed signal to stop a visual reaction-time response. *Journal of Experimental Psychology*, 72(6), 805–811. <https://doi.org/10.1037/h0021266>

Lee, H. W., Lu, M. S., Chen, C. Y., Muggleton, N. G., Hsu, T. Y., & Juan, C. H. (2016). Roles of the pre-SMA and rIFG in conditional stopping revealed by transcranial magnetic stimulation. *Behavioural Brain Research*, 296, 459–467. <https://doi.org/10.1016/j.bbr.2015.08.024>

Lefaucheur, J.-P. P., André-Obadia, N., Antal, A., Ayache, S. S., Baeken, C., Benninger, D. H., ... Garcia-Larrea, L. (2014). Evidence-based guidelines on the therapeutic use of repetitive transcranial magnetic stimulation (rTMS). *Clinical Neurophysiology*, 125(11), 2150–2206. <https://doi.org/10.1016/j.clinph.2014.05.021>

Leung, H. C., & Cai, W. (2007). Common and differential ventrolateral prefrontal activity during inhibition of hand and eye movements. *Journal of Neuroscience*, 27(37), 9893–9900. <https://doi.org/10.1523/JNEUROSCI.2837-07.2007>

Li, C. S., Huang, C., Constable, R. T., & Sinha, R. (2006). Imaging response inhibition in a stop-signal task: Neural correlates independent of signal monitoring and post-response processing. *Journal of Neuroscience*, 26(1), 186–192. <https://doi.org/10.1523/JNEUROSCI.3741-05.2006>

Li, C. S., Milivojevic, V., Kemp, K., Hong, K., & Sinha, R. (2006). Performance monitoring and stop signal inhibition in abstinent patients with cocaine dependence. *Drug and Alcohol Dependence*, 85(3), 205–212. <https://doi.org/10.1016/j.drugalcdep.2006.04.008>

Li, L. M., Violante, I. R., Leech, R., Hampshire, A., Opitz, A., McArthur, D., ... Sharp, D. J. (2019). Cognitive enhancement with Salience Network electrical stimulation is influenced by network structural connectivity. *NeuroImage*, 185, 425–433.

<https://doi.org/10.1016/j.neuroimage.2018.10.069>

1
2 Liang, W. K., Lo, M. T., Yang, A. C., Peng, C. K., Cheng, S. K., Tseng, P., & Juan, C. H. (2014).

3
4 Revealing the brain's adaptability and the transcranial direct current stimulation facilitating
5 effect in inhibitory control by multiscale entropy. *NeuroImage*, *90*, 218–234.

6
7
8
9 <https://doi.org/10.1016/j.neuroimage.2013.12.048>

10
11 Logan, G. D. (1981). Attention, automaticity, and the ability to stop a speeded choice response. In J.

12
13 Long & A. D. Baddeley (Eds.), *Attention and performance IX*. Hillsdale, NJ: Erlbaum.

14
15 Logan, G. D. (1994). On the ability to inhibit thought and action: A users' guide to the stop signal

16
17 paradigm. In D. Dagenbach & T. H. Carr (Eds.), *Inhibitory processes in attention, memory,*
18
19 *and language* (pp. 189–239). Academic Press.

20
21
22 Logan, G. D., Cowan, W. B., & Davis, K. A. (1984). On the ability to inhibit simple and choice

23
24 reaction time responses: A model and a method. *Journal of Experimental Psychology: Human*

25
26
27
28 *Perception and Performance*, *10*(2), 276–291. <https://doi.org/10.1037/0096-1523.10.2.276>

29
30
31 Lüders, H., Lesser, R. P., Dinner, D. S., Morris, H. H., Wyllie, E., & Godoy, J. (1988). Localization

32
33 of cortical function: New information from extraoperative monitoring of patients with

34
35 epilepsy. *Epilepsia*, *29*(s2), S56–S65. <https://doi.org/10.1111/j.1528-1157.1988.tb05799.x>

36
37
38 Mansouri, F. A., Acevedo, N., Illipparampil, R., Fehring, D. J., Fitzgerald, P. B., & Jaberzadeh, S.

39
40 (2017). Interactive effects of music and prefrontal cortex stimulation in modulating response

41
42 inhibition. *Scientific Reports*, *7*(1), 18096. <https://doi.org/10.1038/s41598-017-18119-x>

43
44
45 Mars, R., Piekema, C., Coles, G., Hulstijn, W., & Toni, I. (2007). On the Programming and

46
47 Reprogramming of Actions. *Cerebral Cortex*, *17*(12), 2972–2979.

48
49
50 <https://doi.org/10.1093/cercor/bhm022>.

51
52 Meyer, H. C., & Bucci, D. J. (2016). Neural and behavioral mechanisms of proactive and reactive

53
54 inhibition. *Learning and Memory*, *23*(10), 504–514. <https://doi.org/10.1101/lm.040501.115>

55
56
57 Mikkonen, M., Laakso, I., Tanaka, S., & Hirata, A. (2020). Cost of focality in TDCS:

58
59 Interindividual variability in electric fields. *Brain Stimulation*, *13*(1), 117–124.

<https://doi.org/10.1016/j.brs.2019.09.017>

1
2 Mikuni, N., Ohara, S., Ikeda, A., Hayashi, N., Nishida, N., Taki, J., ... Hashimoto, N. (2006).

3
4 Evidence for a wide distribution of negative motor areas in the perirolandic cortex. *Clinical*
5
6
7 *Neurophysiology*, 117(1), 33–40. <https://doi.org/10.1016/j.clinph.2005.08.021>

8
9 Morita, M., Nakahara, K., & Hayashi, T. (2004). A rapid presentation event-related functional
10
11 magnetic resonance imaging study of response inhibition in macaque monkeys. *Neuroscience*
12
13 *Letters*, 356(3), 203–206. <https://doi.org/10.1016/j.neulet.2003.11.066>

14
15 Mostofsky, S. H., Schafer, J. G. B., Abrams, M. T., Goldberg, M. C., Flower, A. A., Boyce, A., ...
16
17 Pekar, J. J. (2003). fMRI evidence that the neural basis of response inhibition is task-
18
19 dependent. *Cognitive Brain Research*, 17(2), 419–430. <https://doi.org/10.1016/S0926->
20
21
22
23
24
25 6410(03)00144-7

26
27 Mostofsky, S. H., & Simmonds, D. J. (2008). Response inhibition and response selection: Two
28
29 sides of the same coin. *Journal of Cognitive Neuroscience*, 20(5), 751–761.
30
31
32 <https://doi.org/10.1162/jocn.2008.20500>

33
34 Muggleton, N. G., Chen, C. Y., Tzeng, O. J. L., Hung, D. L., & Juan, C. H. (2010). Inhibitory
35
36 control and the frontal eye fields. *Journal of Cognitive Neuroscience*, 22(12), 2804–2812.
37
38
39
40 <https://doi.org/10.1162/jocn.2010.21416>

41
42 Nachev, P., Wydell, H., O'Neill, K., Husain, M., & Kennard, C. (2007). The role of the pre-
43
44 supplementary motor area in the control of action. *NeuroImage*, 36(3–3), T155–T163.
45
46
47 <https://doi.org/10.1016/j.neuroimage.2007.03.034>

48
49 Nambu, A., Tokuno, H., & Takada, M. (2002). Functional significance of the cortico-subthalamo-
50
51 pallidal “hyperdirect” pathway. *Neuroscience Research*, 43(2), 111–117.
52
53
54
55 [https://doi.org/10.1016/S0168-0102\(02\)00027-5](https://doi.org/10.1016/S0168-0102(02)00027-5)

56
57 Nitsche, M. A., Cohen, L. G., Wassermann, E. M., Priori, A., Lang, N., Antal, A., ... Pascual-
58
59 Leone, A. (2008). Transcranial direct current stimulation: State of the art 2008. *Brain*
60
61 *Stimulation*, 1(3), 206–223. <https://doi.org/10.1016/j.brs.2008.06.004>

1 Nitsche, M. A., & Paulus, W. (2000). Excitability changes induced in the human motor cortex by
2 weak transcranial direct current stimulation. *Journal of Physiology*, 527(3), 633–639.

3
4
5 <https://doi.org/10.1111/j.1469-7793.2000.t01-1-00633.x>

6
7 Obeso, I., Cho, S. S., Antonelli, F., Houle, S., Jahanshahi, M., Ko, J. H., & Strafella, A. P. (2013).
8
9 Stimulation of the pre-SMA influences cerebral blood flow in frontal areas involved with
10
11 inhibitory control of action. *Brain Stimulation*, 6(5), 769–776.

12
13
14 <https://doi.org/10.1016/j.brs.2013.02.002>

15
16
17 Obeso, I., Robles, N., Muñoz-Marrón, E., & Redolar-Ripoll, D. (2013). Dissociating the role of the
18
19 pre-SMA in response inhibition and switching: A combined online and offline TMS approach.
20
21 *Frontiers in Human Neuroscience*, 7, 150. <https://doi.org/10.3389/fnhum.2013.00150>

22
23
24 Obeso, I., Wilkinson, L., Teo, J. T., Talelli, P., Rothwell, J. C., & Jahanshahi, M. (2017). Theta
25
26 burst magnetic stimulation over the pre-supplementary motor area improves motor inhibition.
27
28 *Brain Stimulation*, 10(5), 944–951. <https://doi.org/10.1016/j.brs.2017.05.008>

29
30
31 Osada, T., Ohta, S., Ogawa, A., Tanaka, M., Suda, A., Kamagata, K., ... Konishi, S. (2019). An
32
33 essential role of the intraparietal sulcus in response inhibition predicted by parcellation-based
34
35 network. *Journal of Neuroscience*, 39(13), 2509–2521.

36
37
38 <https://doi.org/10.1523/JNEUROSCI.2244-18.2019>

39
40
41 Ouellet, J., McGirr, A., Van den Eynde, F., Jollant, F., Lepage, M., & Berlim, M. T. (2015).
42
43 Enhancing decision-making and cognitive impulse control with transcranial direct current
44
45 stimulation (tDCS) applied over the orbitofrontal cortex (OFC): A randomized and sham-
46
47 controlled exploratory study. *Journal of Psychiatric Research*, 69, 27–34.

48
49
50 <https://doi.org/10.1016/j.jpsychires.2015.07.018>

51
52
53 Plewnia, C., Schroeder, P. A., & Wolkenstein, L. (2015). Targeting the biased brain: Non-invasive
54
55 brain stimulation to ameliorate cognitive control. *The Lancet Psychiatry*, 2(4), 351–356.

56
57
58 [https://doi.org/10.1016/S2215-0366\(15\)00056-5](https://doi.org/10.1016/S2215-0366(15)00056-5)

59
60
61 Praamstra, P., Kleine, B. U., & Schnitzler, A. (1999). Magnetic stimulation of the dorsal premotor

cortex modulates the Simon effect. *NeuroReport*, 10(17), 3671–3674.

<https://doi.org/10.1097/00001756-199911260-00038>

Reinhart, R. M. G., & Woodman, G. F. (2014). Causal control of medial-frontal cortex governs electrophysiological and behavioral indices of performance monitoring and learning. *Journal of Neuroscience*, 34(12), 4214–4227. <https://doi.org/10.1523/JNEUROSCI.5421-13.2014>

Rowe, J. B., Toni, I., Josephs, O., Frackowiak, R. S. J., & Passingham, R. E. (2000). The prefrontal cortex: Response selection or maintenance within working memory? *Science*, 288(5471), 1656–1660. <https://doi.org/10.1126/science.288.5471.1656>

Rubia, K., Russell, T., Overmeyer, S., Brammer, M. J., Bullmore, E. T., Sharma, T., ... Taylor, E. (2001). Mapping motor inhibition: Conjunctive brain activations across different versions of go/no-go and stop tasks. *NeuroImage*, 13(2), 250–261. <https://doi.org/10.1006/nimg.2000.0685>

Rubia, K., Smith, A. B., Brammer, M. J., & Taylor, E. (2003). Right inferior prefrontal cortex mediates response inhibition while mesial prefrontal cortex is responsible for error detection. *NeuroImage*, 20(1), 351–358. [https://doi.org/10.1016/S1053-8119\(03\)00275-1](https://doi.org/10.1016/S1053-8119(03)00275-1)

Rushworth, M. F. S., Hadland, K. A., Paus, T., & Sipila, P. K. (2002). Role of the human medial frontal cortex in task switching: A combined fMRI and TMS study. *Journal of Neurophysiology*, 87(5), 2577–2592. <https://doi.org/10.1152/jn.2002.87.5.2577>

Sandrini, M., Xu, B., Volochayev, R., Awosika, O., Wang, W. T., Butman, J. A., & Cohen, L. G. (2020). Transcranial direct current stimulation facilitates response inhibition through dynamic modulation of the fronto-basal ganglia network. *Brain Stimulation*, 13(1), 96–104. <https://doi.org/10.1016/j.brs.2019.08.004>

Schachar, R., & Logan, G. D. (1990). Impulsivity and inhibitory control in normal development and childhood psychopathology. *Developmental Psychology*, 26(5), 710–720. <https://doi.org/10.1037/0012-1649.26.5.710>

Schachar, R., Tannock, R., & Logan, G. (1993). Inhibitory control, impulsiveness, and attention deficit hyperactivity disorder. *Clinical Psychology Review*, 13(8), 721–739.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

[https://doi.org/10.1016/S0272-7358\(05\)80003-0](https://doi.org/10.1016/S0272-7358(05)80003-0)

Schachar, R., Tannock, R., Marriott, M., & Logan, G. (1995). Deficient inhibitory control in attention deficit hyperactivity disorder. *Journal of Abnormal Child Psychology*, 23(4), 411–437. <https://doi.org/10.1007/BF01447206>

Schluter, N. D., Rushworth, M. F., Passingham, R. E., & Mills, K. R. (1998). Temporary interference in human lateral premotor cortex suggests dominance for the selection of movements. A study using transcranial magnetic stimulation. *Brain*, 121(Pt 5), 785–799. <https://doi.org/10.1093/brain/121.5.785>.

Schroeder, P. A., Schwippel, T., Wolz, I., & Svaldi, J. (2020). Meta-analysis of the effects of transcranial direct current stimulation on inhibitory control. *Brain Stimulation*, 13(5), 1159–1167. <https://doi.org/10.1016/j.brs.2020.05.006>

Sharp, D. J., Bonnelle, V., De Boissezon, X., Beckmann, C. F., James, S. G., Patel, M. C., & Mehta, M. A. (2010). Distinct frontal systems for response inhibition, attentional capture, and error processing. *Proceedings of the National Academy of Sciences of the United States of America*, 107(13), 6106–6111. <https://doi.org/10.1073/pnas.1000175107>

Siebner, H R, & Rothwell, J. (2003). Transcranial magnetic stimulation: New insights into representational cortical plasticity. *Experimental Brain Research*, 148(1), 1–16. <https://doi.org/10.1007/s00221-002-1234-2>

Siebner, Hartwig R., Hartwigsen, G., Kassuba, T., & Rothwell, J. C. (2009). How does transcranial magnetic stimulation modify neuronal activity in the brain? Implications for studies of cognition. *Cortex*, 45(9), 1035–1042. <https://doi.org/10.1016/j.cortex.2009.02.007>

Stramaccia, D. F., Penolazzi, B., Altoè, G., & Galfano, G. (2017). TDCS over the right inferior frontal gyrus disrupts control of interference in memory: A retrieval-induced forgetting study. *Neurobiology of Learning and Memory*, 144, 114–130. <https://doi.org/10.1016/j.nlm.2017.07.005>

Stramaccia, D. F., Penolazzi, B., Sartori, G., Braga, M., Mondini, S., & Galfano, G. (2015).

1 Assessing the effects of tDCS over a delayed response inhibition task by targeting the right
2 inferior frontal gyrus and right dorsolateral prefrontal cortex. *Experimental Brain Research*,
3
4 233(8), 2283–2290. <https://doi.org/10.1007/s00221-015-4297-6>
5
6

7 Stuphorn, V., & Emeric, E. E. (2012). Proactive and reactive control by the medial frontal cortex.
8
9 *Frontiers in Neuroengineering*, 5, 9. <https://doi.org/10.3389/fneng.2012.00009>
10

11 Stuphorn, V., & Schall, J. D. (2006). Executive control of countermanding saccades by the
12
13 supplementary eye field. *Nature Neuroscience*, 9, 925–931. <https://doi.org/10.1038/nn1714>
14
15

16 Stuphorn, V., Taylor, T. L., & Schall, J. D. (2000). Performance monitoring by the supplementary
17
18 eye field. *Nature*, 408, 857–860. <https://doi.org/10.1038/35048576>
19
20

21 Swann, N. C., Cai, W., Conner, C. R., Pieters, T. A., Claffey, M. P., George, J. S., ... Tandon, N.
22
23 (2012). Roles for the pre-supplementary motor area and the right inferior frontal gyrus in
24
25 stopping action: Electrophysiological responses and functional and structural connectivity.
26
27 *NeuroImage*, 59(3), 2860–2870. <https://doi.org/10.1016/j.neuroimage.2011.09.049>
28
29
30

31 Thompson, K. G., Biscoe, K. L., & Sato, T. R. (2005). Neuronal basis of covert spatial attention in
32
33 the frontal eye field. *Journal of Neuroscience*, 25(41), 9479–9487.
34
35 <https://doi.org/10.1523/JNEUROSCI.0741-05.2005>
36
37

38 Verbruggen, F., Aron, A. R., Stevens, M. A., & Chambers, C. D. (2010). Theta burst stimulation
39
40 dissociates attention and action updating in human inferior frontal cortex. *Proceedings of the*
41
42 *National Academy of Sciences of the United States of America*, 107(31), 13966–13971.
43
44 <https://doi.org/10.1073/pnas.1001957107>
45
46
47

48 Verbruggen, F., & Logan, G. D. (2009). Models of response inhibition in the stop-signal and stop-
49
50 change paradigms. *Neuroscience and Biobehavioral Reviews*, 33(5), 647–661.
51
52 <https://doi.org/10.1016/j.neubiorev.2008.08.014>
53
54

55 Vince, M. A. (1948). The intermittency of control movements and the psychological refractory
56
57 period. *British Journal of Psychology. General Section*, 38, 149–157.
58
59 <https://doi.org/10.1111/j.2044-8295.1948.tb01150.x>
60
61
62
63
64
65

1 Vink, M., Kahn, R. S., Raemaekers, M., van den Heuvel, M., Boersma, M., & Ramsey, N. F.

2 (2005). Function of striatum beyond inhibition and execution of motor responses. *Human*
3 *Brain Mapping*, 25(3), 336–344. <https://doi.org/10.1002/hbm.20111>
4
5

6
7 Watanabe, T., Hanajima, R., Shirota, Y., Tsutsumi, R., Shimizu, T., Hayashi, T., ... Konishi, S.

8
9 (2015). Effects of rTMS of pre-supplementary motor area on fronto basal ganglia network
10 activity during stop-signal task. *Journal of Neuroscience*, 35(12), 4813–4823.
11

12
13 <https://doi.org/10.1523/JNEUROSCI.3761-14.2015>
14

15
16
17 Xu, B., Sandrini, M., Wang, W. T., Smith, J. F., Sarlls, J. E., Awosika, O., ... Cohen, L. G. (2016).

18
19 PreSMA stimulation changes task-free functional connectivity in the fronto-basal-ganglia that
20
21 correlates with response inhibition efficiency. *Human Brain Mapping*, 37(9), 3236–3249.
22

23
24 <https://doi.org/10.1002/hbm.23236>
25

26
27 Yang, C. C., Khalifa, N., & Völlm, B. (2018). Excitatory repetitive transcranial magnetic

28
29 stimulation applied to the right inferior frontal gyrus has no effect on motor or cognitive
30
31 impulsivity in healthy adults. *Behavioural Brain Research*, 347, 1–7.
32

33
34 <https://doi.org/10.1016/j.bbr.2018.02.047>
35

36
37 Yu, J., Tseng, P., Hung, D. L., Wu, S. W., & Juan, C. H. (2015). Brain stimulation improves

38
39 cognitive control by modulating medial-frontal activity and preSMA-vmPFC functional
40

41
42 connectivity. *Human Brain Mapping*, 36(10), 4004–4015. <https://doi.org/10.1002/hbm.22893>
43

44
45 Zandbelt, B. B., Bloemendaal, M., Hoogendam, J. M., Kahn, R. S., & Vink, M. (2013).

46
47 Transcranial magnetic stimulation and functional MRI reveal cortical and subcortical
48
49 interactions during stop-signal response inhibition. *Journal of Cognitive Neuroscience*, 25(2),
50

51
52 157–174. https://doi.org/10.1162/jocn_a_00309
53

54
55 Zandbelt, B. B., van Buuren, M., Kahn, R. S., & Vink, M. (2011). Reduced proactive inhibition in

56
57 schizophrenia is related to corticostriatal dysfunction and poor working memory. *Biological*

58
59 *Psychiatry*, 70(12), 1151–1158. <https://doi.org/10.1016/j.biopsych.2011.07.028>
60

61
62 Zandbelt, B. B., & Vink, M. (2010). On the role of the striatum in response inhibition. *PLoS ONE*,

5(11), e13848. <https://doi.org/10.1371/journal.pone.0013848>

1
2 Zanon, M., Borgomaneri, S., & Avenanti, A. (2018). Action-related dynamic changes in inferior
3
4 frontal cortex effective connectivity: a TMS/EEG coregistration study. *Cortex*, 108, 193–209.

5
6
7 <https://doi.org/10.1016/j.cortex.2018.08.004>

8
9 Zhang, R., Geng, X., & Lee, T. M. C. (2017). Large-scale functional neural network correlates of
10
11 response inhibition: An fMRI meta-analysis. *Brain Structure and Function*, 222(9), 3973–

12
13
14 3990. <https://doi.org/10.1007/s00429-017-1443-x>
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure legend

Please use color for both:




Figure 1. Talairach coordinates of the targeted cortical sites in the TMS studies employing SST reconstructed using Surf Ice (<https://www.nitrc.org/projects/surfice>).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 2. Electrodes placement for stimulation in the tDCS studies employing SST were converted into Talairach coordinates (Koessler et al., 2009) and the targeted cortical sites were reconstructed using Surf Ice (<https://www.nitrc.org/projects/surfice>).

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

<i>Study</i>	<i>Stimulation</i>	<i>Talairach Coordinates (x, y, z)</i>	<i>n</i>	<i>Frequency</i>	<i>Duration</i>	<i>Intensity</i>	<i>Online/Offline</i>	<i>Task</i>	<i>Main findings on inhibitory control</i>
Chambers et al. (2006)	rTMS	rIFG: 58, 20, 15	16	1 Hz	15 min	92% rMT	offline	SST	Increased SSRT
Chambers et al. (2007)	rTMS	rIFG: 60, 20, 15 lIFG: -64, 18, 14	16	1 Hz	20 min	92% rMT	offline	Flanker-SST	Increased SSRT after rIFG stimulation
Chen et al. (2009)	rTMS	pre-SMA: -4, 32, 51	9	10 Hz	2 pulses	60% MSO	online	SST	Increased SSRT
Muggleton et al. (2010)	rTMS	FEF: 33, 9, 59	9	10 Hz	2 pulses	65% MSO	online	SST	Increased SSRT
Verbruggen et al. (2010)	cTBS	rIFG: 55, 16, 7 pre-SMA: 4, 36, 55	18	50 Hz	40s	70% rMT	offline	Dual SST	Increased SSRT after rIFG stimulation
Cai et al. (2012)	dual-pulse TMS	pre-SMA: 10, 11, 64	16	/	2 pulses	110% rMT	online	conditional SST	Increased SSRT
Obeso, Cho et al. (2013)	cTBS	pre-SMA: 6, 22, 46	16	50 Hz	20s	80% AMT	offline	SST	Reduced SSRT
Obeso, Robles et al. (2013)	cTBS or single-pulse TMS	pre-SMA: 6, 20, 44 rIFG: 53, 24, 44	16	50 Hz /	40s	80% AMT 60% MSO	offline online	modified SST	Increased SSRT with single-pulse TMS over pre-SMA

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Zandbelt et al. (2013)	rTMS	rIFC: 53, 13, 1 rSMC: 8, 4, 61	21	6 Hz + 1 Hz	20 min	90% rMT + 110% rMT	offline	modified SST	Reduced SSRT after rIFC and rSMC rTMS
Dambacher et al. (2014)	cTBS	pre-SMA: 3, 5, 49	11	50 Hz	40 s	100% AMT	offline	SST and Go/no-Go	No difference in SSRT
Dippel & Bestie (2015)	cTBS and iTBS	rIFG: 54, 19, 11	18	50 Hz	40 s	70% rMT	offline	Stop-change task	No difference in SSRT
Watanabe et al. (2015)	Quadri-pulse rTMS	pre-SMA: 6, 10, 56	9	200 Hz 20 Hz	30 min	90% AMT	offline	SST	Reduced SSRT (excitatory rTMS) Increased SSRT (inhibitory rTMS)
Lee et al. (2016)	cTBS	pre-SMA: 20, 10, 56 rIFG: 42, 17, 18	24	50 Hz	40s	40% MSO	offline	conditional SST	No difference in SSRT
Xu et al. (2016)	single-pulse TMS	pre-SMA: 10, 12, 46	17	/	1 pulse	40% rMT + 80% rMT + 120% rMT	offline	SST	Increased connectivity within the inhibitory network which correlated with reduced SSRT
Obeso et al. (2017)	cTBS	pre-SMA: 6, 27, 40 rIFG: 41, 28, 16	14	50 Hz	40s	80% AMT	offline	conditional SST	Reduced SSRT after cTBS over pre- SMA
Allen et al. (2018)	single-pulse TMS	pre-SMA: 6, 20, 50 rIFC: 53, 20, 6	17	6 – 130 Hz	1 pulse	110% rMT	online	SST	Increased SSRT
Yang et al. (2018)	rTMS	rIFG: 50, 31, -1	20	10 Hz	9 min	100% rMT	offline	SST	No difference in SSRT

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Ji et al. (2019)	rTMS rTMS iTBS	pre-SMA: 6, 10, 56	38	5 Hz 25 Hz 50 Hz	18 min 16 min 39 min	110% rMT 110% rMT 70% rMT	offline	SST	Reduced SSRT after iTBS
Osada et al. (2019)	single-pulse TMS	IPS: 43, -43, 47	22	/	1 pulse	120% rMT	online	SST	Increased SSRT

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. Summary of TMS findings in inhibitory control investigations. Studies that reported coordinates based on the Montreal Neurological Institute were converted into Talairach coordinates through the application of the Yale BioImage Suite Package (Lacadie, Fulbright, Rajeevan, Constable, & Papademetris, 2008).

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

<i>Study</i>	<i>Stimulation</i>	<i>Stimulation Reference</i>	<i>Localization</i>	<i>n</i>	<i>Intensity</i>	<i>Duration</i>	<i>Electrodes' size</i>	<i>Online/Offline</i>	<i>Task</i>	<i>Main findings on inhibitory control</i>
Hsu et al. (2011)	Anodal / Cathodal	Left cheek	pre-SMA: -4, 32, 51	28	1.5 mA	10 min	16 cm ²	offline	SST	No difference in SSRT after anodal or cathodal
Jacobson et al. (2011)	Anodal / Cathodal	Left orbito-frontal cortex	rIFG: crossing point between T4-Fz and F8-Cz lIFG: the crossing point between T3-Fz and F7-Cz	22	1 mA	10 min	25 cm ²	offline	SST	Reduced SSRT after anodal Increased SSRT after cathodal
Ditye et al. (2012)	Anodal	Left orbito-frontal cortex	rIFG: crossing point between T4-Fz and F8-Cz	22	1.5 mA	15 min	35 cm ²	offline	SST	Reduced SSRT
Kwon & Kwon (2013)	Anodal	Left supraorbital area	pre-SMA: 4 cm anterior to Cz	40	1 mA	10 min	35 cm ²	offline	SST	Reduced SSRT
Kwon et al. (2013)	Anodal	Left supraorbital area	M1: C4	40	1 mA	10 min	35 cm ²	online / offline	SST	Reduced SSRT
Cunillera et al. (2014)	Anodal	Left IFC	rIFC: crossing point between T4-Fz and F8-Cz	22	1.5 mA	18 min	9 cm ²	offline	SST and Go/no-Go	Reduced SSRT
Liang et al., (2014)	Anodal	Left cheek	pre-SMA: Fz	18	1.5 mA	10 min	16 cm ²	offline	SST	Reduced SSRT

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Reinhart & Woodman (2014)	Anodal / Cathodal	Right cheek	medial-frontal cortex: FCz	18	1.5 mA	20 min	19 cm ²	offline	SST	No difference in SSRT
Stramaccia et al. (2015)	Anodal / Cathodal	Left supraorbital area	rIFG: crossing point between T4-Fz and F8-Cz	115	1.5 mA	20 min	16 cm ²	offline	SST	Reduced SSRT after anodal
Cai et al. (2015)	Anodal	Left cheek	rIFG: crossing point between F4 and F8	22	1.5 mA	15 min	25 cm ²	offline	SST	Reduced SSRT
Ouellet et al. (2015)	Anodal	Right and left OFC	OFC: Fp1 and Fp2	45	1.5 mA	30 min	35 cm ²	offline	SST	No difference in SSRT
Yu et al. (2015)	Anodal	Left cheek	pre-SMA: Fz	23	2 mA	20 min	16 cm ²	offline	SST	Reduced SSRT
Cunillera et al. (2016)	Anodal	Left IFC	rIFC: crossing point between T4-Fz and F8-Cz	13	1.5 mA	20 min	9 cm ²	online	SST and Go/no-Go	No difference in SSRT
Castro-Meneses et al. (2016)	Anodal	Left cheek	rPFC: crossing point between T4-Fz and F8-Cz	14	1.5 mA	15 min	25 cm ²	online / offline	SST	Reduced SSRT
Hogeveen et al. (2016)	HD-tDCS or Anodal	Vertex: Cz	rIFC: FC6	46	1 mA	20 min	25 cm ²	offline	SST	Reduced SSRT
Bender et al. (2017)	Anodal / Cathodal	Right mastoid	pre-SMA: 2, 15, 51	18 36	0.7 mA	9 min 13 min	25 cm ²	offline	modified SST	No difference in SSRT

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Mansouri et al. (2017)	Anodal	Right supraorbital area	ldIPFC: F3	73	1.5 mA	10 min	10 cm ²	offline	modified SST	Reduced SSRT
Stramaccia et al. (2017)	Anodal / Cathodal	Left supraorbital area	rIFG: FC4	72	1.5 mA	20 min	16 cm ²	online	SST	No difference in SSRT
Chen et al. (2019)	Anodal	Left cheek	rIFG: crossing point between F4 and F8	57	1.5 mA	20 min	25 cm ²	offline	modified SST	Reduced SSRT
Friehs & Frings (2018)	Anodal	Left deltoid muscle	rdIPFC: F4	59	0.5 mA	19 min	9 cm ²	offline	SST	Reduced SSRT
Friehs & Frings (2019)	Cathodal	Left deltoid muscle	rdIPFC: F4	45	0.5 mA	19 min	9 cm ²	offline	SST	Increased SSRT
Fehring et al. (2019)	Anodal	Right supraorbital area	ldIPFC: F3	73	1.5 mA	10 min	10 cm ²	offline	SST	Reduced SSRT
Li et al. (2019)	Anodal / Cathodal	Right shoulder	rIFG: F8	24	2 mA	12 min	12 cm ²	online	SST	Reduced SSRT after anodal
Sandrini et al. (2020)	Anodal	Left supraorbital area	rIFC: 53, 28, 3	30	1.5 mA	20 min	25 cm ²	offline	SST	Reduced SSRT

Table 2. Summary of tDCS findings in inhibitory control investigations.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65