

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Application of different techniques on stone fruit (*Prunus* spp.) drying and assessment of physical, chemical and biological properties: Characterization of dried fruit properties

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Vakula A., Tepic Horecki A., Pavlic B., Jokanovic M., Ognjanov V., Milovic M., et al. (2021). Application of different techniques on stone fruit (*Prunus* spp.) drying and assessment of physical, chemical and biological properties: Characterization of dried fruit properties. JOURNAL OF FOOD PROCESSING AND PRESERVATION, 45(2), 1-18 [10.1111/jfpp.15158].

Availability:

This version is available at: <https://hdl.handle.net/11585/800884> since: 2021-02-27

Published:

DOI: <http://doi.org/10.1111/jfpp.15158>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

Application of different techniques on stone fruit (*Prunus* spp.) drying and assessment of physical, chemical and biological properties: Characterization of dried fruit properties

Anita Vakula^a, Aleksandra Tepić Horecki^a, Branimir Pavlić^a, Marija Jokanović^a, Vladislav Ognjanov^b, Maja Milović^b, Nemanja Teslić^c, Giuseppina Parpinello^d, Marlies Decler^{e,f}, Zdravko Šumić^{a*}

^aUniversity of Novi Sad, Faculty of Technology Novi Sad, Bulevar cara Lazara 1, 21000 Novi Sad, Serbia

^bUniversity of Novi Sad, Faculty of Agriculture, Trg Dositeja Obradovića 8, 21000 Novi Sad, Serbia

^cUniversity of Novi Sad, Institute of Food Technology, Bulevar cara Lazara 1, 21000 Novi Sad, Serbia

^dUniversity of Bologna, Department of Agricultural and Food Sciences, Piazza Goidanich 60, 47521 Cesena (FC), Italy

^eGhent University, Faculty of Bioscience Engineering, Laboratory of Food Microbiology and Food Preservation, Belgium

^fGhent University, Faculty of Pharmaceutical Sciences, Laboratory of Food Analysis, Belgium

*Corresponding author:

E-mail: sumic@uns.ac.rs, Telephone: +381 21 485 3718, <https://orcid.org/0000-0002-9770-0139>

This is the peer reviewed version of the following article: *Application of different techniques on stone fruit (Prunus spp.) drying and assessment of physical, chemical and biological properties: Characterization of dried fruit properties*, by Anita Vakula, Aleksandra Tepić Horecki, Branimir Pavlić, Marija Jokanović, Vladislav Ognjanov, Maja Milović, Nemanja Teslić, Giuseppina Parpinello, Marlies Decler, Zdravko Šumić, which has been published in final form in *Journal of Food Processing and Preservation* Volume 45, Issue 2, February 2021, e15158, DOI <https://doi.org/10.1111/jfpp.15158>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

Abstract

Drying of stone fruit with different techniques and characterization of physical, chemical and biological properties of convective dried, vacuum dried and lyophilised stone fruit, as well as analysing and structuring the data sets by principal component analysis (PCA) were obtained in this paper. Drying technique significantly influenced the shear force, hardness, springiness and cohesiveness of dried apricot NS4 (Novi Sad 4) samples ($p < 0.05$); the total phenolic, flavonoid and monomeric anthocyanin content of dried sour cherry Feketicka, sweet cherry Lapins, sweet cherry Sweet Heart and plum Toptase samples and also the antioxidant capacity of dried sour cherry Feketicka, plum Anna Spath and peach Lela samples. The most noticeable differentiations of the stone fruit groups of samples dried with convective and vacuum drying and lyophilisation were observed in raw and dried sour cherry Feketicka and Erdi Botermo samples.

Keywords: stone fruit; physical, chemical and biological properties; drying techniques; principal component analysis.

Practical applications

The results and conclusions obtained in this research have various application in food industry in many aspects. First, part of the fruit varieties investigated in this study were developed at the Faculty of Agriculture and their application in the fruit drying industry has been thoroughly investigated. In addition, application of different drying techniques on different stone fruit species were applied in this research. Finally, the possibilities of preservation the most important quality indicators of dried fruit was observed. The impact of obtained conclusions and results in the field of agricultural and food industry is significant, since they could be applied in the industrial processes.

DR. ALEKSANDRA N. TEPIĆ HORECKI (Orcid ID : 0000-0002-2479-0313)

DR. BRANIMIR PAVLIC (Orcid ID : 0000-0002-3551-7478)

DR. ZDRAVKO M. ŠUMIĆ (Orcid ID : 0000-0002-9770-0139)

Article type : Original Article

1. Introduction

Sour cherry (*P. cerasus*), sweet cherry (*P. avium*), apricot (*P. armeniaca*), plum (*P. domestica*) and peach (*P. persica*) are the most prevalent stone fruit in Serbia. First matured are sour and sweet cherries which are followed by apricots, plums and peaches (Wills, Scriven, & Greenfield, 1983). Due to their widespreadness, good processing potential and good sensory and nutritional characteristics, certain varieties of stone fruit have been investigated by many authors (Esti, Cinquanta, Sinesio, Moneta, & Di Matteo, 2002; Toğrul & Pehlivan, 2003; Vargas, Jablonski, Flôres, & Rios, 2017; Wojdyło, Figiel, Lech, Nowicka, & Oszmiański, 2014). Stone fruit has been investigated also in the field of application of different extraction techniques, such as for example paper by Garofulić, Dragović-Uzelac, Jambrak and Jukić (2013) the effect of microwave assisted extraction on the isolation of anthocyanins and phenolic acids from sour cherry Marasca was observed. Furthermore, in the paper by Zaghdoudi et al. (2015) accelerated solvent extraction of carotenoids from peach, apricot, among other fruit, were investigated while conventional and ultrasound-assisted extraction of anthocyanins from blackberry and sweet cherry cultivars were described in the paper by Oancea, Grosu, Ketney and Stoia (2013).

The consumption of stone fruit can bring many benefits for human health since fruit in general is rich in bioactive compounds such as phenolic compounds (Tomás-Barberán et al., 2001), anthocyanins, carotenoids, vitamin C and organic acids (Wills et al., 1983). However,

the amount of bioactive compounds in fresh fruit, as well as in certain fruit product, is influenced by their stability which is related to their oxidation and environmental sensitivity (Leong & Oey, 2012). Valuable components of stone fruit made it prevalent raw material of processing industry.

In order to extend the shelf life and its usage throughout the year, fruit is usually preserved with drying, freezing or made into jams, compotes or juices. One of the ways of food preservation is the reduction of the water availability (Mulet, Cárcel, Sanjuan, & Bon, 2003). During the drying process, water from the raw material, which is necessary for microorganism's growth and enzymatic activity, has been removed. This prevents the growth of microorganisms since the minimum water activity at which microorganisms can grow is 0.60 (Beuchat et al., 2013). Many enzymes will function minimally in the 0.65-0.70 range (Lopez et al. 1997) which indicates that lower water activity results in reduced enzyme activity. Drying process has also been used prior to extraction of secondary metabolites from different plants such as for example applied and described in the paper by Rangkadilok et al. (2007) where evaluation of free radical scavenging and antityrosinase activities of standardized longan fruit extract and thus one part of the fruit was dried before extraction process. Also in the paper by Assefa and Keum (2017) different drying methods (microwave-, oven-, freeze-, and air-drying) were applied for yuzu (*Citrus junos* Sieb ex Tanaka) drying prior to extraction of polyphenols and antioxidants.

The commonly used type of drying in the food industry is convective drying process with hot air (Mujumdar & Beke, 2003). However, the presence of oxygen and high temperatures involved in drying process negatively influence physical, chemical and biological properties of dried products (Larrosa, Cadaval, & Pinto, 2015). Also, drying usually precedes extraction processes in order to reduce water content and to preserve polyphenol content and so undesirable drying conditions may influence degradation of bioactive compounds which could further lead to lower extraction yields. On the other hand, application of high temperatures during drying may influence release of bounded polyphenols (Papoutsis et al., 2018). But, also when high temperatures are applied part of the polyphenols might be oxidized and converted to other compounds (Abhay, Hii, Law, Suzannah, & Djaeni, 2016). Disadvantages of convective drying, in terms of negative influence of oxygen presence and high applied temperatures could be successfully solved by using vacuum drying technique. During vacuum drying, applied vacuum decreases the pressure in the chamber at the given temperature and the water vapour has constantly being removed from vacuum chamber

(Bourdoux, Li, Rajkovic, Devlieghere, & Uyttendaele, 2016). Combinations of vacuum drying with infrared, ultrasound or microwave drying were investigated by Chen, Guo and Wu, 2016; Pu and Sun 2016; Xie et al. 2017. The type of vacuum drying which includes sublimation process is freeze drying or lyophilisation. Beside all advantages relative to vacuum and convective drying, lyophilisation is not commonly used for fruit drying due to higher drying cost and energy consumption and low efficiency (Pei et al., 2014).

Even though, stone fruit and their products were thoroughly investigated (Aghbashlo, Kianmehr, & Hassan-Beygi, 2010; Celik, Demirkol, Durmus, & Tarakci, 2020; Doymaz, 2014; Doymaz & İsmail, 2011; Goyal, Kingsly, Manikantan, & Ilyas, 2007; Ihns, Diamante, Savage, & Vanhanen, 2011; Ouaabou et al., 2020), in known and accessible databases there are no papers investigating different types of fruit, drying techniques and characterization of the most important quality of dried products in the framework of a single research. Encouraged with these facts, authors' main goal was drying of two varieties of each stone fruit type and characterization of their physical, chemical and biological properties. Furthermore, another goal was to compare physical, chemical and biological properties in terms of applied drying techniques on the one hand and in terms of certain variety of each type of stone fruit on the other, by PCA utilization.

2. Materials and Methods

Samples

Fresh stone fruit samples were purchased at the Faculty of Agriculture, University of Novi Sad. Two varieties of each type: sour cherry (Feketicka (SCF) and Erdi Botermo (SCEB)); sweet cherry (Lapins (SCL) and Sweet Heart (SCSH)); apricot (Buda (AB) and NS4 (ANS4)); plum (Anna Spath (PAS) and Toptaste (PTT)) and peach (Lela (PL) and Fairtime (PFT)) were collected from the experimental field of Faculty of Agriculture, Novi Sad, at Rimski Šančevi (Serbia).

After the measuring of flesh/stone ratio and mass per unit area, samples were prepared for each type of drying. The stones were carefully moved from each sample. Sour and sweet cherries were dried directly while apricots and peaches were first cut on two halves and four quarters, respectively. Samples intended for convective and vacuum drying were not frozen

previously, while for lyophilisation samples were frozen and stored at -20 °C until drying. For each drying, 300 g of the sample was measured.

Investigated properties of fresh stone fruit samples are given in supplementary material (Table S1).

Chemicals

The following reagents were purchased from Sigma-Aldrich Chem (Steinheim, Germany): Folin-Ciocalteu reagent, (±)-catechin, gallic acid, 2,2-diphenyl-1-picrylhydrazyl (DPPH), ABTS (2,2'-azino-bis-(-3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt), TPZT (2,4,6-tris(2-pyridyl)-s-triazine), iron (III)-chloride and Iron (II)-sulfate heptahydrate and potassium persulfate. Trolox (6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid) was purchased from Sigma-Aldrich (Milano, Italy). Sodium acetate and hydrochloric acid were obtained from Merck (Darmstadt, Germany). Formic acid was obtained from VWR International (Milan, Italy). Acetonitrile (HPLC grade and LC-MS grade), methanol (absolute, LC-MS grade) and glacial formic acid (99%, ULC-MS) were purchased from BioSolve BV (Valkenswaard, the Netherlands). Ammonium acetate was obtained from Merck (Darmstadt, Germany). Methanol (HiPerSolv Chromanorm HPLC grade) was supplied by VWR International (Zaventem, Belgium). Ultrafree[®]-MC centrifugal filter devices (0.22 µm) were supplied by Millipore (Bredford, MA, USA). Water was purified on a Milli-Q[®] SP Reagent water system from Millipore Corp (Brussels, Belgium). Valinomycin (VAL) (10 mg, solid standard) and beauvericin (BEA), enniatins (ENN) (A, A1, B, B1) (1 mg, solid standard) were purchased from Sigma-Aldrich (Diegem, Belgium).

All other chemicals were of analytical and HPLC grade. Experimental design

Experimental design and used are presented in Figure 1 and Table 1, respectively. A total of forty stone fruit samples, each weighing 300 g, were used in experiments. ten fresh (F) samples (SCF-F, SCEB-F, SCL-F, SCSH-F, AB-F, ANS4-F, PAS-F, PTT-F, PL-F, PFT-F), ten samples dried by convective drying (C) (SCF-C, SCEB-C, SCL-C, SCSH-C, AB-C, ANS4-C, PAS-C, PTT-C, PL-C, PFT-C), ten samples dried by vacuum drying (V) (SCF-V, SCEB-V, SCL-V, SCSH-V, AB-V, ANS4-V, PAS-V, PTT-V, PL-V, PFT-V) and ten samples dried by lyophilisation (L) (SCF-L, SCEB-L, SCL-L, SCSH-L, AB-L, ANS4-L, PAS-L, PTT-L, PL-L, PFT-L).

Independent variables were temperature, pressure and time during drying which are presented in Table 2 for each sample. Dependent variables were four physical parameters: moisture content (MC), water activity (a_w), total colour change (ΔE), texture (shear force, penetration force, hardness, springiness, cohesiveness and chewiness); three chemical parameters: total phenolic (TPC), flavonoid (TFC) and monomeric anthocyanin content (TMAC) and one biological parameter (antioxidant capacity (FRAP, DPPH and ABTS test)). Also, toxin analysis (BEA and ENNs (A, A1, B, B1)) were performed on fresh and dried samples. The number of repetition has been mentioned for each analyse separately.

Figure 1. Experimental design-flow chart

Table 1. Fresh and dried stone fruit-abbreviations

Drying techniques

Vacuum drying process was described in detail by Šumić, Tepić, Vidović, Jokić, & Malbaša (2013), while convective drying and lyophilisation procedures have been described in detail by Šumić et al. (2016). Vacuum and convective drying was continued until no mass change was detected (final moisture content in equilibrium). The conditions of applied drying methods are presented in Table 2.

Table 2. Conditions of convective drying, vacuum drying and lyophilisation.

Analyses

-Texture analysis

Instrumental texture measurements were performed using a Texture Analyser (TE32, Stable Micro Systems, UK). Preparation of fresh samples for texture analysis was the same as preparation for drying process, explained in detail in section Samples (Material and Methods). For dried sour and sweet cherry samples there were also no special preparations, while dried apricot, plum and peach samples were cut in pieces dimensions 2 cm x 2 cm x 1

cm. All texture analyses were performed at room temperature. The texture analyses were performed twelve times for statistical purpose.

Shearing Test

The shear force was measured using Craft Knife Adapter. Instrumental settings for shear force analyses were the following: test speed – 1.0 mm/s; load cell – 5 kg. Shear force has been expressed as force (g) required for cutting of the samples.

Penetration Test

The penetration test was carried using 2 mm stainless Cylinder probes. Instrumental settings: test speed – 2.0 mm/s; load cell – 5 kg were set for penetration force analysis. The sample was positioned centrally relative to the Cylinder probe. Penetration force has been expressed as force (g) required for penetration through the samples.

Texture Profile Analysis (TPA)

Texture profile analysis (TPA) was performed as described by Bourne (1978), using Texture Analyser (TE32, Stable Micro Systems, UK) equipped with a cylindrical plate of 36 mm in diameter. Dried samples were compressed twice to 40% of their original thickness at a constant speed of 1 mm/s. Hardness, springiness, cohesiveness and chewiness were determined by using TPA test. Hardness and chewiness were expressed as force (g) necessary for sample compression unit, while springiness and cohesiveness are dimensionless values. Hardness was defined by peak force during the first compression cycle; springiness as the rate at which a deformed sample goes back to its undeformed condition after the deforming force is removed; cohesiveness was calculated as the ratio of the area under the second curve to the area under the first curve and chewiness was obtained by multiplying hardness, cohesiveness, and springiness (Bianchi et al., 2016).

-Antioxidant capacity assays

Approximately 10 g of fresh and 5 g of dried stone fruit samples were ground in a blender before the extraction. Grounded samples were transferred to a volumetric flask and 50 mL of methanol, as extraction solvent, was added. Extraction was carried out for 24 h at the room temperature. The obtained extracts were filtered, placed into a glass bottles and stored to prevent oxidative damage until analysis.

FRAP assay

The sample ability to reduce Fe^{3+} was measured using slightly modified method firstly presented by Benzie and Strain (1996). The FRAP reagent was freshly prepared from 300 mM acetate buffer (pH=3.6), 10 mM 2,4,6-tris(2-pyridyl)-s-triazine (TPZT) in 40 mM HCl solution and 20 mM FeCl_3 aqueous solution. Solutions were mixed in ratio 10:1:1 (v/v/v). Previously diluted extracts and FRAP reagent were mixed (0.1 mL + 1.9 mL) and stored to incubate in the dark at 37 °C for 10 min. The measurements were performed at 593 nm, in duplicates, with UV–VIS spectrophotometer (Cary 60, Agilent Technologies, Santa Clara, USA). The results were finally reported as mg of Fe^{2+} equivalents per g of dry weight.

DPPH assay

The sample ability to scavenge 2,2-diphenyl-1-picrylhydrazyl free radicals (DPPH \cdot) was measured using a modified method originally presented by Brand-Williams, Cuvelier and Berset, et al. (1995). Briefly, methanolic solution of the DPPH reagent (65 μM) was freshly prepared and adjusted with methanol to reach absorbance of 0.70 (± 0.02). DPPH reagent and previously diluted extracts were mixed (2.9 mL + 0.1 mL) in the 10 mm plastic cuvettes and incubated at room temperature for 60 min. Free radical scavenging measurements were performed at 517 nm, in duplicates with UV–VIS spectrophotometer (Cary 60, Agilent Technologies, Santa Clara, USA). The obtained results were reported as mg of Trolox equivalents per g of dry weight.

ABTS assay

The ABTS free radical scavenging ability of samples was measured using a modified method originally described by Re et al. (1999). ABTS stock solution was freshly prepared from mixture (1:1, v/v) of 2.45 mM potassium persulfate aqueous solution and 7mM ABTS (2,2'-azino-bis-(-3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt) aqueous solution and left in the dark at room temperature for 16 h. A stock solution was diluted using 300 mM acetate buffer (pH=3.6) to an absorbance of 0.70 (± 0.02). Previously diluted extracts and ABTS reagent were mixed (0.1 mL + 2.9 mL) and stored in the dark at room temperature for 300 minutes. The measurements were performed at 734 nm, in duplicates with UV–VIS spectrophotometer (Cary 60, Agilent Technologies, Santa Clara, USA). The results were finally reported as mg of Trolox equivalents per g of dry weight.

-Total phenolics content

The content of total phenolic content in methanolic extracts was determined by Folin-Ciocalteu procedure (Singleton and Rossi, 1965), using gallic acid as a standard. Content of

total phenolic content has been expressed as mg of gallic acid equivalent per 100 g of dry weight of samples (mg GAE/100 g DW). Experiments were replicated three times and results are expressed as mean values. Absorbance was measured at 765 nm.

The UV/Vis spectrophotometer (model 6300 Spectrophotometer, Jenway, UK) was used for all spectrophotometric methods.

-Toxin Analysis

Primary stock solutions were prepared by dissolving the solid standard in acetonitrile (1 mg/mL). All stock solutions, except VAL, were stored at -20 °C. VAL was stored at 4 °C. Working solutions of 10 µg/mL were prepared in acetonitrile and then stored at 4 °C and renewed monthly. Mixture solutions (BEA and ENNs) were prepared prior to each experiment by diluting the working solution in acetonitrile. Initially, 2 g of fresh and dried stone fruit samples were homogenized and transferred into 50 mL extraction tubes. The method for toxin analysis is explained in detail in the paper by Decleer, Rajkovic, Sas, Madder and De Saeger (2016).

Description of analysis (moisture content, water activity, total colour change, total flavonoid content and monomeric anthocyanin content) is shown in detail by Tepić-Horecki et al. (2018).

Statistical analysis

All the data were analysed by univariate analysis of variance (ANOVA, $p < 0.05$) in order to differentiate the samples using an $\alpha = 0.05$ criterion and Tukey's Multiple Comparison Test. Principal Component Analysis (PCA) was applied in order to analyse and structurize the obtained results. Statistica 10.0 (StatSoft Inc., Tulsa, OK, USA) was used both for the PCA and ANOVA.

3. Results and Discussion

Physical, chemical and biological characterization has been performed in stone fruit dried with convective drying, vacuum drying and lyophilisation. Dried stone fruit characterization was observed by measuring MC, aw, ΔE , texture analysis (shear force, penetration force,

hardness, springiness, cohesiveness and chewiness), TPC, TFC, TMC, antioxidant capacity (FRAP, DPPH and ABTS test) and toxin analysis (BEA and ENNs (A, A1, B, B1)).

Deviations are a normal occurrence in the case of technological processing of different plant material with usage of completely different techniques. For this reason, several iterations were performed and the results were statistically processed in order to make conclusions with the statistically defined security that are discussed for each individual parameter in this part of the paper.

3.1. Physical properties

3.1.1. Moisture content (MC) (Table 3)

In dried SCEB and SCL samples drying techniques significantly influenced ($p < 0.05$) the MC. It could be seen that, both for sour and sweet cherry samples, the lowest and the highest MC were obtained in convective and vacuum dried samples, respectively. In apricot and peach samples, ANS4 and PFT, respectively, it was not noticed significant difference of MC ($p < 0.05$) between convective and vacuum dried samples. It was also noticed that in both plum varieties (PTT and PAS) and in peach PL samples, drying techniques significantly influenced the MC.

3.1.2. Water activity (a_w) (Table 3)

It could be seen that drying techniques influenced a_w values in all types and varieties of stone fruit. The lowest a_w values in all types of dried fruit were obtained in lyophilised samples (SCEB-L, SCSH-L, ANS4-L, PTT-L, PL-L) and the highest ones were obtained in vacuum dried samples (SCEB-V, SCL-V, AB-V, PAS-V, PL-V).

3.1.3. Total colour change (ΔE) (Table 3)

In dried sour cherry samples, the lowest ΔE was obtained in convective dried samples (SCF-C and SCEB-C), while the highest ΔE was observed in lyophilised samples (SCF-L and SCEB-L). Concerning these results, it was assumed that in case of drying sour cherry samples (SCF and SCEB), pre-freezing of fruit and long drying time, significantly influenced the increase of ($p < 0.05$) ΔE , beside the absence of oxygen and low temperatures. Significantly different ($p < 0.05$) values of ΔE were obtained between SCL-V and SCSH-V sweet cherry varieties. Namely, in SCL and SCSH varieties, the lowest ΔE values were obtained in vacuum dried and lyophilised samples, respectively, while the highest ΔE values were noticed in lyophilised and convective dried samples, respectively. In dried apricot samples AB and also in peach samples PL it was noticed that applied drying techniques significantly influence ($p < 0.05$) ΔE . The value of ΔE of vacuum dried peaches, obtained by Kwon, Kim and Youn (2013), was 14.31 which is higher than ΔE obtained for PL-V sample (7.39) and lower than ΔE observed for PFT-V sample (36.92). Both in dried AB and ANS4 samples, the lowest ΔE values were obtained in vacuum dried samples (AB-V and ANS4-V), while the highest values of ΔE were observed in lyophilised samples (AB-L and ANS4-L).

Table 3. Experimentally obtained values of MC, a_w and ΔE of dried samples.

3.1.4. Texture analysis

Shear force (Table 4)

The lowest shear force obtained in all dried samples was 147.4 g (ANS4-L), while the highest one was noticed in PTT-L sample (3856.7 g). It could be seen that both the highest and the lowest values of shear force of all dried samples were observed in lyophilised samples. Applied drying techniques significantly influenced ($p < 0.05$) the shear force of ANS4 samples.

Penetration force (Table 4)

It could be seen that in all convective dried samples 5000 g penetration force was observed. However, all these samples were too hard for penetration by sphere of instrument. Since the weight of 5000 g was used during measurement, all values for penetration force were noticed

as > 5000 g. In different varieties of the same type of plum fruit, the lowest and the highest penetration force were observed, PTT-V (509.3 g) and PAS-L (2852.9 g), respectively. Based on the obtained results for dried SCEB, SCL, SCSH, PTT, PAS, PL and PFT samples, drying techniques significantly influenced ($p < 0.05$) the penetration force.

Hardness (Table 4)

Hardness, as primary texture property, is defined as force necessary to attain a given deformation (Szczesniak, 2002). PAS-V and PTT-C were the samples with the lowest (252.4 g) and the highest (6377.8 g) hardness of all dried samples. Drying techniques significantly influenced ($p < 0.05$) the hardness of SCSH and ANS4 samples.

Springiness (Table 4)

Springiness is defined as rate at which a deformed material goes back to its undeformed condition after the deforming force is removed (Szczesniak, 2002). Springiness of fresh samples varied between 0.29 (ANS4-F) and 0.94 (SCF-F), while in dried samples this range was between 0.28 (PFT-L) and 0.80 (SCF-C). Based on the observed results for dried SCF, SCL and ANS4 samples, drying techniques significantly influenced ($p < 0.05$) the springiness.

Cohesiveness (Table 4)

The extent to which a material can be deformed before it ruptures defines the cohesiveness parameter (Szczesniak, 2002). The lowest cohesiveness of fresh samples was obtained in PL-F sample (0.13), while cohesiveness obtained for PAS-F sample was the highest (0.71 g). The lowest cohesiveness of all dried samples was 0.24 g obtained in ANS4-L sample, while the highest one (0.95 g) was noticed in PTT-L sample. It could be noticed that for dried samples ANS4 and PFT, drying techniques significantly influenced ($p < 0.05$) the cohesiveness.

Chewiness (Table 4)

Chewiness, a product of hardness, cohesiveness and springiness, is defined as energy required to masticate a solid food to a state ready for swallowing (Szczesniak, 2002). Chewiness of fresh samples varied between 26.5 g (ANS4-F) and 806.8 g (PTT-F), while in dried samples this range was between 79.4 g (AB-L) and 1637.5 g (SCEB-C).

Table 4. Experimentally obtained values of dried samples texture.

3.2. Chemical properties

3.2.1. Total phenolic content (TPC) (Figure 2.a)

The lowest TPC of all investigated fresh samples was obtained in AB-F sample (194.94 mg GAE/100 g DW), while SCF-F was sample with the highest TPC (1713.23 mg GAE/100 g DW). TPC in sweet cherry, obtained by Serradilla et al. (2011) varied between 59.05 and 117 mg gallic acid/100 g raw weight. TPC of all dried samples varied between 199.27 mg GAE/100 g DW obtained in AB-L sample and 1605.34 mg GAE/100 g DW noticed in SCF-V sample. Drying techniques significantly influenced ($p < 0.05$) the TPC of dried SCF, SCL, SCSH, ANS4, PTT, PAS and PFT samples and also did not significantly influence ($p < 0.05$) TPC of AB samples. Kwon et al. (2013) applied cold vacuum drying on peaches and observed TPC of dried peach was 4.03 mg GAE (gallic acid equivalents)/g. The behaviour of the polyphenols compounds in cases when they are exposed to different drying conditions are presented in the paper by Papoutsis et al. (2018) and also described in the introduction part of this paper. Also, influence of different drying techniques on different types of plant material is described in papers by Alfaro, Mutis, Quiroz, Seguel, and Scheuermann (2014) and Heredia, Barrera and Andrés (2007) where effects of drying techniques on murtilla fruit polyphenols and antioxidant capacity drying of cherry tomato by a combination of different dehydration techniques were investigated, respectively.

Figure 2. Total phenolic (a), flavonoid (b) and monomeric anthocyanin (c) content observed in fresh, convective dried, vacuum dried and lyophilised samples.

3.2.2. Total flavonoid content (TFC) (Figure 2.b)

As a subgroup of phenolic compounds, flavonoids contribute to antioxidant profile of fresh and dried fruit. Thus, these two parameters are often investigated together such as in research

by Hooshmand and Arjmandi (2009) where phenolic and flavonoid content in dried plum powder was 22.4 mg/100 g. Different drying parameters influence TFC preservation to a greater or lesser extent. TFC of fresh samples varied between 68.44 mg CE/100 g DW (PAS-F) and 780.26 mg CE/100 g DW (SCF-F). It was noticed that both the highest values of TPC and TFC were observed in the same SCF-F sample. In dried samples AB-C and SCF-V, the lowest (92.42 mg CE/100 g DW) and the highest (824.32 mg CE/100 g DW) TFC contents, were obtained respectively. Based on the observed results for dried SCF, SCL, SCSH, AB, ANS4, PTT, PAS, PL and PFT samples, drying techniques significantly influenced ($p < 0.05$) the TFC.

3.2.3. Total monomeric anthocyanin content (TMAC) (Figure 2.c)

Content of anthocyanins, widely appeared red colorants of fruits and vegetables, were already investigated by Serradilla et al. (2011) where observed content of total content of these compounds varied between 4.06 and 39.44 mg-cyanidin-3-O-rutinoside/100 g raw weight. The lowest TMAC of fresh samples was obtained in PAS-F sample (1.07 mg CGE/100 g DW), while SCL-F was sample with the highest TMAC (46.31 mg CGE/100 g DW). TMAC of all dried samples varied between 0.29 mg CGE/100 g DW obtained in PAS-C sample and 262.15 mg CGE/100 g DW noticed in PTT-V sample. In dried SCF, SCEB, SCL, SCSH, ANS4 and PTT samples applied drying techniques significantly influenced ($p < 0.05$) the obtained TMAC.

3.3. Biological properties

3.3.1. Antioxidant properties (Figure 3)

Antioxidant capacity of fresh sweet cherries, obtained by Serradilla et al. (2011) was in the range from 317.92 to 439.10 mg Trolox/100 g raw weight. Also, Wojdyło, Figiel, Lech, Nowicka and Oszmiański (2014) observed antioxidant capacity using FRAP and ABTS tests in fresh sour cherries (61.31 and 159.052 mg Trolox/g DW respectively); convective dried sour cherries (from 31.807 to 41.591 mg Trolox/g DW and from 80.348 and 119.589 mg

Trolox/g DW, respectively), vacuum-microwave dried sour cherries (from 36.860 to 47.458 mg Trolox/g DW and from 117.036 to 171.178 mg Trolox/g DW, respectively) and lyophilised sour cherries (60.227 and 151.293 mg Trolox/g DW). Three antioxidant tests, FRAP, DPPH and ABTS, were analysed in order to obtain antioxidant capacity of fresh and dried stone fruit.

FRAP test (Figure 3.a)

The lowest antioxidant capacity, obtained by FRAP test, of all investigated fresh samples was obtained in PAS-F sample (0.1521 mg Fe²⁺/g DW), while SCL-F was sample with the highest antioxidant capacity (0.7197 mg Fe²⁺/g DW). Antioxidant capacity of all dried samples was in the range from 0.1953 mg Fe²⁺/g DW obtained in PAS-V sample to 2.7481 mg Fe²⁺/g DW noticed in the vacuum dried SCF-V sample. Drying techniques significantly influenced ($p < 0.05$) the antioxidant capacity of dried SCF, AB, PTT, PAS and PL samples.

DPPH test (Figure 3.b)

Antioxidant capacity obtained by DPPH test in all fresh samples varied between 1.5947 mg Trolox/g DW observed in PAS-F sample and 17.3361 mg Trolox/g DW noticed in SCF-F sample. Antioxidant capacity of all dried samples was in the range from 1.6505 mg Trolox/g DW obtained in AB-C sample to 26.7122 mg Trolox/g DW observed in PTT-C sample. Based on the obtained results of dried SCF, ANS4, PAS, PL and PFT samples, it was noticed that drying techniques significantly influenced ($p < 0.05$) the antioxidant capacity.

ABTS test (Figure 3.c)

Antioxidant capacity obtained by ABTS test of investigated fresh fruit samples was in the range from 2.6720 mg Trolox/g DW obtained in PAS-F sample to 11.7142 mg Trolox/g DW noticed in SCL-F sample. Antioxidant capacity of all dried samples varied between 3.3675 mg Trolox/g DW obtained in SCF-V sample and 58.3583 mg Trolox/g observed in PTT-C sample. It could be seen that drying techniques significantly influenced ($p < 0.05$) the antioxidant capacity of dried SCF, SCSH, ANS4, PTT, PAS and PL samples.

Figure 3. Antioxidant capacity of fresh, convective dried, vacuum dried and lyophilised samples obtained by FRAP (a), DPPH (b) and ABTS (c) test.

Since FRAP, ABTS and DPPH assays present three different assays of the antioxidant capacity in terms of the different compounds that sample has ability to scavenge or reduce, it was chosen to analyse all three assays in order observe complete picture of the antioxidant profile of fresh and dried stone fruit samples. Compatibility between these assays are in accordance with the results obtained in papers by Maria do Socorro et al (2010); Popović, Štajner, Kevrešan, and Bijelić (2012) and Vakula, Šumić, Zeković, Tepić Horecki, Pavlić (2019).

Based on the Figure 2.a and Figures 2.a, b and c, it could be seen that the TPC content of most of the fresh and dried stone fruit samples are in accordance with all three investigated antioxidative tests (FRAP, DPPH and ABTS). There are also certain disagreements that could be explained with the fact that the phenolic compounds are just a part of the compounds which make the total antioxidant capacity in stone fruit samples. Other important compounds such as flavonoid and anthocyanins compounds, which are also investigated in this research also have important role in term of antioxidant profile of fresh and dried stone fruit. For this reason, all these compounds are chosen to be investigated in this paper, to make as complete picture as it is possible in terms of the quality properties of fresh and dried stone fruit samples. Accordingly, similar results in terms of the accordance between total phenolic content and antioxidant capacity are presented in the papers by Igual, García-Martínez, Martín-Esparza, and Martínez-Navarrete (2012); Madrau et al. (2009) and Sultana, Anwar, Ashraf, and Saari (2012).

3.4. Toxin analysis

Different types of toxins in various fruits were investigated by many authors (Drusch & Ragab, 2003; MacDonald et al., 1999). The presence of emerging *Fusarium* mycotoxins

(ENNs and BEA) was determined in samples of nuts and dried fruits commercialized in Valencia, by Tolosa, Font, Mañes, and Ferrer (2013). Based on the research by Tolosa et. al. (2013), the average levels of BEA and ENNs A, A1, B, B1 in dried fruit were 0.007, 0.242, 0.011, 0.058, 0.022 mg/kg, respectively.

In this study, the presence of BEA and ENNs A, A1, B, B1 was investigated in fresh and dried stone fruit. Based on the obtained results, it was noticed that all fresh and dried samples were free of all investigated toxins.

3.5. Chemometric analysis

Principal Component Analysis (PCA) was applied on obtained data in order to get a better overview of the similarities between convective dried, vacuum dried and lyophilised stone fruit based on their physical, chemical and biological properties. Multivariate statistical approach i.e. PCA, was applied in order to differentiate samples dried with different drying techniques and also to find correlations between investigated physical, chemical and biological parameters. Therefore, physical parameters used for PCA were MC, a_w , ΔE , texture, chemical parameters were TPC, TFC, TMAC, while biological parameter was antioxidant capacity.

Figure 4. Bi-plot distribution of PC1 and PC2 (a) and PC1 and PC3 (b) for grouping investigated physical, chemical and biological parameters.

Figure 5. Score plot of PC1 and PC2 for grouping fresh, convective dried, vacuum dried and lyophilised samples with prominent sour cherry (a), sweet cherry (b), apricot (c), plum (d) and peach (e) samples.

PCA was carried out in order to reduce the number of dimensions of the complex system with 14 grouping variables. It could be seen that the PC1 and PC2 accounted for 65.02% (Figure 4.a), while PC2 and PC3 accounted 57.45% (Figure 4.b) of the total variance of the model. These first three Principal Components (PC) accounted for 76.6% of the total variance of the model. PC1 was negatively correlated with all grouping parameters except MC, a_w and springiness. On the other hand, PC2 was negatively correlated with all investigated physical parameters and positively correlated with all texture analysis except springiness and ΔE . Accordingly, PC3 was negatively correlated with shear force, cohesiveness, ΔE and TMAC

while it positively correlated with penetration force, hardness, chewiness, TFC and all antioxidative tests.

The distribution of the samples was significantly influenced by the type of both fruit and drying techniques, since the following three groups: fresh, convective dried, vacuum dried and lyophilised samples of the each type of fruit could be grouped together (Figure 5). Numbering of dried samples is presented in Table 2, while fresh samples were numbered as SCF, SCEB, SCL, SCSH, AB, ANS4, PTT, PAS, PL and PFT from 1 to 10, respectively. The most clear differences of these three groups could be seen in dried sour cherry, sweet cherry and apricot samples (Figure 5.a,b,c), while group of fresh samples clearly differentiated from other groups in all types of investigated stone fruit. Accordingly, groups of fresh samples were characterized by the high values of both MC and a_w , which could be expected since these two parameters are in direct correlation. Based on these grouping of fresh samples, it was also expected that groups of convective dried fruit were in opposite upper left quadrant for sour cherry (Figure 5.a), sweet cherry (Figure 5.b) and apricot (Figure 5.c). However, groups of plum (Figure 5.d) and peach (Figure 5.e) convective dried samples were both in left and right opposite upper quadrant. It could be seen that groups of convective dried sour and sweet cherries as well as apricot samples were characterized by the highest values of ΔE and all texture analyses, except springiness. The clearest group of vacuum dried and lyophilised samples was obtained in sour cherry samples (Figure 5.a) in down left quadrant, opposite of groups of fresh and convective samples. Group of vacuum dried and lyophilised samples in other fruit samples was randomised and it differentiated less clearly compared to sour cherry vacuum dried and lyophilised samples group (Figure 5.b,c,d,e).

4. Conclusions

Characterization of physical, chemical and biological properties of convective dried, vacuum dried and lyophilised stone fruit, as well as analysing and structuring the data sets by principal component analysis (PCA) were obtained in this paper.

The lowest moisture content (6.54%), water activity (0.255) and total colour change (3.15) of all investigated dried samples were observed in lyophilised plum Toptaste; lyophilised sour

cherry Erdi Botermo and lyophilised apricot NS4; convective dried sour cherry Feketicka, respectively.

Based on the results obtained for texture analyses of all dried samples, it could be concluded that in dried apricot NS4 samples applied drying techniques significantly influenced ($p < 0.05$) the obtained shear force, hardness, springiness and cohesiveness. On the other hand, properties of variety of sweet cherry did not significantly influence ($p < 0.05$) shear force, penetration force, hardness, springiness and chewiness during vacuum drying.

The highest total phenolic and flavonoid content of all dried samples were obtained in the vacuum dried sour cherry Feketicka sample, while the highest total monomeric anthocyanin content was observed in vacuum dried plum Toptase sample. Drying techniques significantly influenced ($p < 0.05$) the total phenolic, flavonoid and monomeric anthocyanin content in dried sour cherry Feketicka, sweet cherry Lapins, sweet cherry Sweet Heart and plum Toptase samples.

Based on the results obtained with DPPH and ABTS antioxidant tests, the convective dried plum Toptase sample possesses the highest antioxidant capacity, while FRAP test showed that the sample with the highest antioxidant capacity was vacuum dried sour cherry Feketicka. Drying techniques significantly influenced ($p < 0.05$) the antioxidant capacity of dried sour cherry Feketicka, plum Anna Spath and peach Lela samples, based on results obtained from all three antioxidative tests.

The first three Principal Components (PC) accounted for 76.6% of the total variance of the model. The most apparent differentiations of the groups of fresh; convective dried; and vacuum dried and lyophilised samples of the each type of investigated stone fruit were obtained in fresh and dried sour cherry Feketicka and Erdi Botermo samples.

5. Acknowledgments

This work was supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia [Project 451-03-68/2020-14/ 200134].

The authors would like to thank Professor Andreja Rajković, Faculty of Bioscience engineering, Ghent University, for cooperation in the framework of Erasmus+ programme for academic staff mobility, Key Action 1.

6. References

Abhay, S. M., Hii, C. L., Law, C. L., Suzannah, S., & Djaeni, M. (2016). Effect of hot-air drying temperature on the polyphenol content and the sensory properties of cocoa beans. *International Food Research Journal*, 23.

Aghbashlo, M., Kianmehr, M. H., & Hassan-Beygi, S. R. (2010). Drying and rehydration characteristics of sour cherry (*Prunus cerasus* L.). *Journal of Food Processing and Preservation*, 34, 351–365. doi: <https://doi.org/10.1111/j.1745-4549.2008.00310.x>

Assefa, A. D., & Keum, Y. S. (2017). Effect of extraction solvent and various drying methods on polyphenol content and antioxidant activities of yuzu (*Citrus junos* Sieb ex Tanaka). *Journal of Food Measurement and Characterization*, 11, 576–585.

Benzie, I. F., & Strain, J. J. (1996). The ferric reducing ability of plasma (FRAP) as a measure of “antioxidant power”: the FRAP assay. *Analytical Biochemistry*, 239, 70–76. doi: <https://doi.org/10.1006/abio.1996.0292>

Beuchat, L. R., Komitopoulou, E., Beckers, H., Betts, R. P., Bourdichon, F., Fanning, S., & Ter Kuile, B. H. (2013). Low–water activity foods: increased concern as vehicles of foodborne pathogens. *Journal of Food Protection*, 76, 150–172. doi: <https://doi.org/10.4315/0362-028X.JFP-12-211>

Bianchi, T., Guerrero, L., Gratacós-Cubarsí, M., Claret, A., Argyris, J., Garcia-Mas, J., & Hortós M. (2016). Textural properties of different melon (*Cucumis melo* L.) fruit types: Sensory and physical-chemical evaluation. *Scientia Horticulturae*, 201, 46–56. doi: <https://doi.org/10.1016/j.scienta.2016.01.028>

Bourdoux, S., Li, D., Rajkovic, A., Devlieghere, F., & Uyttendaele, M. (2016). Performance of Drying Technologies to Ensure Microbial Safety of Dried Fruits and Vegetables. *Comprehensive Reviews in Food Science and Food Safety*, 15, 1056–1066. doi: <https://doi.org/10.1111/1541-4337.12224>

Bourne, M. (1978). Texture profile analysis. *Food Technology*, 32, 62–66.

Brand-Williams, W., Cuvelier, M. E., & Berset, C. (1995). Use of a free radical method to evaluate antioxidant activity. *LWT - Food Science and Technology*, 28, 25–30. doi: [https://doi.org/10.1016/S0023-6438\(95\)80008-5](https://doi.org/10.1016/S0023-6438(95)80008-5)

Celik, O. F., Demirkol, M., Durmus, Y., & Tarakci, Z. (2020). Effects of drying method on the phenolics content and antioxidant activities of cherry laurel (*Prunus laurocerasus* L.). *Journal of Food Measurement and Characterization*, 14, 48–54.

Chen, Z. G., Guo, X. Y., & Wu, T. (2016). A novel dehydration technique for carrot slices implementing ultrasound and vacuum drying methods. *Ultrasonics Sonochemistry*, 30, 28–34. doi: <https://doi.org/10.1016/j.ultsonch.2015.11.026>

Decleer, M., Rajkovic, A., Sas, B., Madder, A., & De Saeger, S. (2016). Development and validation of ultra-high-performance liquid chromatography–tandem mass spectrometry methods for the simultaneous determination of beauvericin, enniatins (A, A1, B, B1) and cereulide in maize, wheat, pasta and rice. *Journal of Chromatography A*, 1472, 35–43. doi: <https://doi.org/10.1016/j.chroma.2016.10.003>

Doymaz, İ., & İsmail, O. (2011). Drying characteristics of sweet cherry. *Food and Bioproducts Processing*, 89, 31–38. doi: <https://doi.org/10.1016/j.fbp.2010.03.006>

Doymaz, I. (2014). Suitability of thin-layer drying models for infrared drying of peach slices. *Journal of food processing and preservation*, 38, 2232–2239. doi: <https://doi.org/10.1111/jfpp.12277>

Drusch, S., & Ragab, W. (2003). Mycotoxins in fruits, fruit juices, and dried fruits. *Journal of Food Protection*, 66, 1514–1527. doi: <https://doi.org/10.4315/0362-028X-66.8.1514>

Esti, M., Cinquanta, L., Sinesio, F., Moneta, E., & Di Matteo, M. (2002). Physicochemical and sensory fruit characteristics of two sweet cherry cultivars after cool storage. *Food Chemistry*, 76, 399–405. doi: [https://doi.org/10.1016/S0308-8146\(01\)00231-X](https://doi.org/10.1016/S0308-8146(01)00231-X)

Garofulić, I. E., Dragović-Uzelac, V., Jambrak, A. R., & Jukić, M. (2013). The effect of microwave assisted extraction on the isolation of anthocyanins and phenolic acids from sour cherry Marasca (*Prunus cerasus* var. Marasca). *Journal of Food Engineering*, 117, 437–442. doi: <https://doi.org/10.1016/j.jfoodeng.2012.12.043>

- Goyal, R. K., Kingsly, A. R. P., Manikantan, M. R., & Ilyas, S. M. (2007). Mathematical modelling of thin layer drying kinetics of plum in a tunnel dryer. *Journal of Food Engineering*, 79, 176-180. doi: <https://doi.org/10.1016/j.jfoodeng.2006.01.041>
- Hooshmand, S., & Arjmandi, B. H. (2009). Dried plum, an emerging functional food that may effectively improve bone health. *Ageing research Reviews*, 8, 122-127. doi: <https://doi.org/10.1016/j.arr.2009.01.002>
- Igual, M., García-Martínez, E., Martín-Esparza, M. E., & Martínez-Navarrete, N. (2012). Effect of processing on the drying kinetics and functional value of dried apricot. *Food Research International*, 47, 284-290. doi: <https://doi.org/10.1016/j.foodres.2011.07.019>
- Ihns, R., Diamante, L. M., Savage, G. P., & Vanhanen, L. (2011). Effect of temperature on the drying characteristics, colour, antioxidant and beta-carotene contents of two apricot varieties. *International Journal of Food Science & Technology*, 46, 275-283. doi: <https://doi.org/10.1111/j.1365-2621.2010.02506.x>
- Kwon, G. M., Kim, J. W., & Youn, K. (2013). Effect of different pre-treatments on the physicochemical and antioxidant activities of cold-vacuum dried peaches. *Korean Journal of Food Science and Technology*, 45, 466-472. doi: <https://doi.org/10.9721/KJFST.2013.45.4.466>
- Larrosa, A. P., Cadaval, T. R., & Pinto, L. A. (2015). Influence of drying methods on the characteristics of a vegetable paste formulated by linear programming maximizing antioxidant activity. *LWT-Food Science and Technology*, 60, 178-185. doi: <https://doi.org/10.1016/j.lwt.2014.08.003>
- Leong, S. Y., & Oey, I. (2012). Effects of processing on anthocyanins, carotenoids and vitamin C in summer fruits and vegetables. *Food Chemistry*, 133, 1577-1587. doi: <https://doi.org/10.1016/j.foodchem.2012.02.052>
- Lopez, A., Pique, M. T., Boatella, J., Parcerisa, J., Romero, A., Ferrá, A., & Garcí, J. (1997). Influence of drying conditions on the hazelnut quality. II. Enzymatic activity. *Drying Technology*, 15, 979-988. doi: <https://doi.org/10.1080/07373939708917272>
- MacDonald, S., Wilson, P., Barnes, K., Damant, A., Massey, R., Mortby, E., Shepherd, M. J. (1999). Ochratoxin A in dried vine fruit: method development and survey. *Food Additives & Contaminants*, 16, 253-260. doi: <https://doi.org/10.1080/026520399284019>

Madrau, M. A., Piscopo, A., Sanguinetti, A. M., Del Caro, A., Poiana, M., Romeo, F. V., & Piga, A. (2009). Effect of drying temperature on polyphenolic content and antioxidant activity of apricots. *European Food Research and Technology*, 228, 441. doi: 10.1007/s00217-008-0951-6

Maria do Socorro, M. R., Alves, R. E., de Brito, E. S., Pérez-Jiménez, J., Saura-Calixto, F., & Mancini-Filho, J. (2010). Bioactive compounds and antioxidant capacities of 18 non-traditional tropical fruits from Brazil. *Food Chemistry*, 121, 996–1002. doi: <https://doi.org/10.1016/j.foodchem.2010.01.037>

Mujumdar, A. S., & Beke, J. (2003). Grain drying: Basic principles, in *Handbook of postharvest technology: Cereals, fruits, vegetables, tea, and spices* (eds Chakraverty, A., Mujumdar, A. S., Raghavan, G. S. V., Ramaswamy, H. S.). Marcel Dekker, New York, USA, pp. 119–139.

Mulet, A., Cárcel, J. A., Sanjuan, N., Bon, J. (2003). New food drying technologies-Use of ultrasound. *Food Science and Technology International*, 9, 215–221. doi: <https://doi.org/10.1177/1082013203034641>

Oancea, S., Grosu, C., Ketney, O., & Stoia, M. (2013). Conventional and ultrasound-assisted extraction of anthocyanins from blackberry and sweet cherry cultivars. *Acta Chimica Slovenica*, 60, 383–389.

Ouaabou, R., Nabil, B., Ouhammou, M., Idlimam, A., Lamharrar, A., Ennahli, S., Hanine, H., & Mahrouz, M. (2020). Impact of solar drying process on drying kinetics, and on bioactive profile of Moroccan sweet cherry. *Renewable Energy*, 151, 908–918.

Papoutsis, K., Vuong, Q. V., Golding, J. B., Hasperué, J. H., Pristijono, P., Bowyer, M. C., Scarlett, M. C., & Stathopoulos, C. E. (2018). Pretreatment of citrus by-products affects polyphenol recovery: A review. *Food Reviews International*, 34, 770–795.

Pei, F., Yang, W. J., Shi, Y., Sun, Y., Mariga, A. M., Zhao, L. Y. & Hu, Q. H. (2014). Comparison of freeze-drying with three different combinations of drying methods and their influence on colour, texture, microstructure and nutrient retention of button mushroom (*Agaricus bisporus*) slices. *Food and Bioprocess Technology*, 7, 702–710. doi: <https://doi.org/10.1007/s11947-013-1058-z>

Popović, B. M., Štajner, D., Kevrešan, S., & Bijelić, S. (2012). Antioxidant capacity of cornelian cherry (*Cornus mas* L.)—Comparison between permanganate reducing antioxidant capacity and other antioxidant methods. *Food Chemistry*, 134, 734–741. doi: <https://doi.org/10.1016/j.foodchem.2012.02.170>

Pu, Y. Y., & Sun, D. W. (2016). Prediction of moisture content uniformity of microwave-vacuum dried mangoes as affected by different shapes using NIR hyperspectral imaging. *Innovative Food Science & Emerging Technologies*, 33, 348–356. doi: <https://doi.org/10.1016/j.ifset.2015.11.003>

Rangkadilok, N., Sitthimonchai, S., Worasuttayangkurn, L., Mahidol, C., Ruchirawat, M., & Satayavivad, J. (2007). Evaluation of free radical scavenging and antityrosinase activities of standardized longan fruit extract. *Food and Chemical Toxicology*, 45, 328–336. doi: <https://doi.org/10.1016/j.fct.2006.08.022>

Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., & Rice-Evans, C. (1999). Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine*, 26, 1231–1237. doi: [https://doi.org/10.1016/S0891-5849\(98\)00315-3](https://doi.org/10.1016/S0891-5849(98)00315-3)

Serradilla, M. J., Lozano, M., Bernalte, M. J., Ayuso, M. C., López-Corrales, M., & González-Gómez D. (2011). Physicochemical and bioactive properties evolution during ripening of ‘Ambrunés’ sweet cherry cultivar. *LWT-Food Science and Technology*, 44, 199–205. doi: <https://doi.org/10.1016/j.lwt.2010.05.036>

Sultana, B., Anwar, F., Ashraf, M., & Saari, N. (2012). Effect of drying techniques on the total phenolic contents and antioxidant activity of selected fruits. *Journal of Medicinal Plants Research*, 6, 161–167. doi: <https://doi.org/10.5897/JMPR11.916>

Šumić, Z., Tepić, A., Vidović, S., Jokić, S., & Malbaša, R. (2013). Optimization of frozen sour cherries vacuum drying process. *Food Chemistry*, 136, 55–63. doi: <https://doi.org/10.1016/j.foodchem.2012.07.102>

Šumić, Z., Vakula, A., Tepić, A., Čakarević, J., Vitas, J., & Pavlić, B. (2016). Modelling and optimization of red currants vacuum drying process by response surface methodology (RSM). *Food Chemistry* 203, 465–475. doi: <https://doi.org/10.1016/j.foodchem.2016.02.109>

Szczesniak, A. S. (2002). Texture is a sensory property. *Food Quality and Preference*, 13, 215–225. doi: [https://doi.org/10.1016/S0950-3293\(01\)00039-8](https://doi.org/10.1016/S0950-3293(01)00039-8)

Tepić Horecki, A., Vakula, A., Pavlić, B., Jokanović, M., Malbaša, R., Vitas, J., Jaćimović, V., & Šumić, Z. (2018). Comparative drying of cornelian cherries: Kinetics modeling and physico- chemical properties. *Journal of Food Processing and Preservation*, 42, e13562. doi: <https://doi.org/10.1111/jfpp.13562>

Toğrul, İ. T., & Pehlivan, D. (2003). Modelling of drying kinetics of single apricot. *Journal of Food Engineering*, 58, 23–32. doi: [https://doi.org/10.1016/S0260-8774\(02\)00329-1](https://doi.org/10.1016/S0260-8774(02)00329-1)

Tolosa, J., Font, G., Mañes, J., & Ferrer, E. (2013). Nuts and dried fruits: natural occurrence of emerging *Fusarium* mycotoxins. *Food Control*, 33, 215–220. doi: <https://doi.org/10.1016/j.foodcont.2013.02.023>

Tomás-Barberán, F. A., Gil, M. I., Cremin, P., Waterhouse, A. L., Hess-Pierce, B., & Kader, A. A. (2001). HPLC– DAD– ESIMS analysis of phenolic compounds in nectarines, peaches, and plums. *Journal of Agricultural and Food Chemistry*, 49, 4748–4760. doi: <https://doi.org/10.1021/jf0104681>

Vakula, A., Šumić, Z., Zeković, Z., Tepić Horecki, A., & Pavlić, B. (2019). Screening, influence analysis and optimization of ultrasound- assisted extraction parameters of cornelian cherries (*Cornus mas* L.). *Journal of Food Processing and Preservation*, 43, e14226. doi: <https://doi.org/10.1111/jfpp.14226>

Vargas, E. F. D., Jablonski, A., Flôres, S. H., & Rios, A. D. O. (2017). Waste from peach (*Prunus persica*) processing used for optimisation of carotenoids ethanolic extraction. *International Journal of Food Science & Technology*, 52, 757–762. doi: <https://doi.org/10.1111/ijfs.13332>

Wang, M., Jiang, N., Wang, Y., Jiang, D., & Feng, X. (2017). Characterization of phenolic compounds from early and late ripening sweet cherries and their antioxidant and antifungal activities. *Journal of Agricultural and Food Chemistry*, 65, 5413–5420. doi: <https://doi.org/10.1021/acs.jafc.7b01409>

Wills, R. B., Scriven, F. M., & Greenfield, H. (1983) Nutrient composition of stone fruit (*Prunus* spp.) cultivars: apricot, cherry, nectarine, peach and plum. *Journal of the Science of Food and Agriculture*, 34, 1383–1389. doi: <https://doi.org/10.1002/jsfa.2740341211>

Wojdyło, A., Figiel, A., Lech, K., Nowicka, P., & Oszmiański, J. (2014). Effect of convective and vacuum–microwave drying on the bioactive compounds, color, and antioxidant capacity of sour cherries. *Food and Bioprocess Technology*, 7, 829–841. doi: <https://doi.org/10.1007/s11947-013-1130-8>

Xie, L., Mujumdar, A. S., Fang, X. M., Wang, J., Dai, J. W., Du, Z. L., Xiao, H. W., Liu, Y., & Gao, Z. J. (2017). Far-infrared radiation heating assisted pulsed vacuum drying (FIR-PVD) of Wolfberry (*Lycium barbarum* L.): effects on drying kinetics and quality attributes. *Food and Bioprocess Technology*, 102, 320–331. doi: <https://doi.org/10.1016/j.fbp.2017.01.012>

Zaghdoudi, K., Pontvianne, S., Framboisier, X., Achard, M., Kudaibergenova, R., Ayadi-Trabelsi, M., Kalthoum-cherif, J., Vanderesse, R., Frochot, C., & Guiavarc'h, Y. (2015). Accelerated solvent extraction of carotenoids from: Tunisian Kaki (*Diospyros kaki* L.), peach (*Prunus persica* L.) and apricot (*Prunus armeniaca* L.). *Food chemistry*, 184, 131–139. doi: <https://doi.org/10.1016/j.foodchem.2015.03.072>

Figure 1. Experimental design-flow chart.

Figure 2. Total phenolic (a), flavonoid (b) and monomeric anthocyanin (c) content observed in raw, convective dried, vacuum dried and lyophilised samples.

Figure 3. Antioxidant capacity of raw, convective dried, vacuum dried and lyophilised samples obtained by FRAP (a), DPPH (b) and ABTS (c) test.

Figure 4. Bi-plot distribution of PC1 and PC2 (a) and PC1 and PC3 (b) for grouping investigated physical, chemical and biological parameters.

Figure 5. Score plot of PC1 and PC2 for grouping raw, convective dried, vacuum dried and lyophilised samples with prominent sour cherry (a), sweet cherry (b), apricot (c), plum (d) and peach (e) samples.

Table 1. Fresh and dried stone fruit-abbreviations

| Number | Fruit type | Fruit variety | State | Abbreviation |
|--------|--------------|---------------|------------------|--------------|
| 1 | Sour cherry | Feketicka | Fresh | SCF-F |
| 2 | | Erdi Botermo | | SCEB-F |
| 3 | Sweet cherry | Lapins | | SCL-F |
| 4 | | Sweet Heart | | SCSH-F |
| 5 | Apricot | Buda | | AB-F |
| 6 | | NS4 | | ANS4-F |
| 7 | Plum | Anna Spath | | PAS-F |
| 8 | | Toptaste | | PTT-F |
| 9 | Peach | Lela | | PL-F |
| 10 | | Fairtime | | PFT-F |
| 11 | Sour cherry | Feketicka | Convective dried | SCF-C |
| 12 | | Erdi Botermo | | SCEB-C |
| 13 | Sweet cherry | Lapins | | SCL-C |
| 14 | | Sweet Heart | | SCSH-C |
| 15 | Apricot | Buda | | AB-C |
| 16 | | NS4 | | ANS4-C |
| 17 | Plum | Anna Spath | | PAS-C |
| 18 | | Toptaste | | PTT-C |
| 19 | Peach | Lela | | PL-C |
| 20 | | Fairtime | | PFT-C |
| 21 | Sour cherry | Feketicka | Vacuum dried | SCF-V |
| 22 | | Erdi Botermo | | SCEB-V |
| 23 | Sweet cherry | Lapins | | SCL-V |
| 24 | | Sweet Heart | | SCSH-V |
| 25 | Apricot | Buda | | AB-V |
| 26 | | NS4 | | ANS4-V |
| 27 | Plum | Anna Spath | | PAS-V |
| 28 | | Toptaste | | PTT-V |
| 29 | Peach | Lela | | PL-V |
| 30 | | Fairtime | | PFT-V |

| | | | | |
|----|--------------|--------------|-------------|--------|
| 31 | | Feketicka | | SCF-L |
| 32 | Sour cherry | Erdi Botermo | | SCEB-L |
| 33 | | Lapins | | SCL-L |
| 34 | Sweet cherry | Sweet Heart | | SCSH-L |
| 35 | | Buda | | AB-L |
| 36 | Apricot | NS4 | Lyophilised | ANS4-L |
| 37 | | Anna Spath | | PAS-L |
| 38 | Plum | Toptaste | | PTT-L |
| 39 | | Lela | | PL-L |
| 40 | Peach | Fairtime | | PFT-L |

Table 2. Conditions of convective drying, vacuum drying and lyophilisation.

| Number | Fruit | Sample | Pressure [mbar] | Temperature [°C] | Drying time [h] |
|--------|----------------------------|--------|--------------------|---------------------|--------------------|
| 11 | sour cherry Feketicka | SCF-C | 1000 | 78 | 24 |
| 12 | sour cherry Erdi Botermo | SCEB-C | 1000 | 78 | 24 |
| 13 | sweet cherry Lapins | SCL-C | 1000 | 78 | 25 |
| 14 | sweet cherry Sweet Heart | SCSH-C | 1000 | 78 | 23 |
| 15 | apricot Buda | AB-C | 1000 | 78 | 17 |
| 16 | apricot NS4 | ANS4-C | 1000 | 78 | 18 |
| 17 | plum Anna Spath | PAS-C | 1000 | 78 | 24 |
| 18 | plum Toptaste | PTT-C | 1000 | 78 | 24 |
| 19 | peach Lela | PL-C | 1000 | 78 | 22 |
| 20 | peach Fairtime | PFT-C | 1000 | 78 | 21 |
| 21 | sour cherry Feketicka | SCF-V | 20 | 60 | 11 |
| 22 | sour cherry Erdi Botermo | SCEB-V | 20 | 60 | 11 |
| 23 | sweet cherry Lapins | SCL-V | 20 | 60 | 13 |
| 24 | sweet cherries Sweet Heart | SCSH-V | 20 | 60 | 16 |
| 25 | apricot Buda | AB-V | 20 | 60 | 19 |
| 26 | apricot NS4 | ANS4-V | 20 | 60 | 19 |
| 27 | plum Anna Spath | PAS-V | 20 | 60 | 29 |
| 28 | plum Toptaste | PTT-V | 20 | 60 | 28 |
| 29 | peach Lela | PL-V | 20 | 60 | 27 |
| 30 | peach Fairtime | PFT-V | 20 | 60 | 27 |
| 31 | sour cherry Feketicka | SCF-L | 0.01 | -30 | 72 |
| 32 | sour cherry Erdi Botermo | SCEB-L | 0.01 | -30 | 72 |
| 33 | sweet cherry Lapins | SCL-L | 0.01 | -30 | 72 |
| 34 | sweet cherries Sweet Heart | SCSH-L | 0.01 | -30 | 72 |
| 35 | apricot Buda | AB-L | 0.01 | -30 | 72 |
| 36 | apricot NS4 | ANS4-L | 0.01 | -30 | 72 |
| 37 | plum Anna Spath | PAS-L | 0.01 | -30 | 72 |
| 38 | plum Toptaste | PTT-L | 0.01 | -30 | 72 |
| 39 | peach Lela | PL-L | 0.01 | -30 | 72 |
| 40 | peach Fairtime | PFT-L | 0.01 | -30 | 72 |

Table 3. Experimentally obtained values of MC, a_w and ΔE of dried samples.

| Analyses | MC (%) | | a_w | | ΔE | |
|--------------|---------------------------|---------------------------|----------------------------|----------------------------|---------------------------|----------------------------|
| Sample | SCF | SCEB | SCF | SCEB | SCF | SCEB |
| SCF-F/SCEB-F | 83.37±0.06 ^{A,b} | 86.64±0.36 ^{A,a} | 0.929±0.007 ^{A,a} | 0.936±0.001 ^{A,a} | 0.00* | 0.00* |
| SCF-C/SCEB-C | 13.41±0.17 ^{C,a} | 11.20±0.51 ^{D,b} | 0.278±0.004 ^{D,b} | 0.350±0.000 ^{C,a} | 3.15±0.58 ^{B,a} | 4.73±1.70 ^{B,a} |
| SCF-V/SCEB-V | 15.60±0.88 ^{B,b} | 21.29±0.76 ^{B,a} | 0.376±0.003 ^{B,b} | 0.540±0.003 ^{B,a} | 23.44±5.28 ^{A,a} | 7.83±2.26 ^{B,b} |
| SCF-L/SCEB-L | 14.71±0.15 ^{B,b} | 15.11±0.10 ^{C,a} | 0.309±0.006 ^{C,a} | 0.255±0.000 ^{D,b} | 27.52±4.99 ^{A,a} | 26.80±4.93 ^{A,a} |
| Sample | SCL | SCSH | SCL | SCSH | SCL | SCSH |
| SCL-F/SCSH-F | 83.17±0.17 ^{A,a} | 82.00±0.03 ^{A,b} | 0.949±0.003 ^{A,a} | 0.939±0.003 ^{A,a} | 0.00* | 0.00* |
| SCL-C/SCSH-C | 9.05±0.36 ^{D,a} | 9.12±0.004 ^{C,a} | 0.406±0.001 ^{C,a} | 0.349±0.003 ^{C,b} | 23.17±0.14 ^{B,a} | 21.06±1.74 ^{A,b} |
| SCL-V/SCSH-V | 15.36±0.30 ^{B,a} | 14.72±0.24 ^{B,b} | 0.465±0.001 ^{B,a} | 0.419±0.003 ^{B,b} | 17.93±1.44 ^{B,a} | 15.90±5.75 ^{AB,a} |
| SCL-L/SCSH-L | 12.05±0.05 ^{C,b} | 14.48±0.02 ^{B,a} | 0.292±0.000 ^{D,a} | 0.268±0.004 ^{D,b} | 29.07±5.46 ^{A,a} | 11.07±6.32 ^{B,b} |
| Sample | AB | ANS4 | AB | ANS4 | AB | ANS4 |
| AB-F/ANS4-F | 87.77±0.16 ^{A,b} | 88.77±0.08 ^{A,a} | 0.903±0.003 ^{A,b} | 0.948±0.006 ^{A,a} | 0.00* | 0.00* |
| AB-C/ANS4-C | 17.11±0.67 ^{B,a} | 14.15±0.98 ^{B,b} | 0.368±0.001 ^{D,b} | 0.411±0.001 ^{C,a} | 36.21±3.26 ^{B,b} | 47.76±1.97 ^{A,a} |
| AB-V/ANS4-V | 16.23±0.39 ^{B,a} | 14.36±0.24 ^{B,b} | 0.670±0.001 ^{B,a} | 0.544±0.000 ^{B,b} | 13.67±1.84 ^{C,b} | 18.28±1.50 ^{B,a} |
| AB-L/ANS4-L | 16.55±0.22 ^{B,a} | 7.66±0.44 ^{C,b} | 0.425±0.000 ^{C,a} | 0.255±0.001 ^{D,b} | 43.50±1.47 ^{A,b} | 49.43±1.85 ^{A,a} |
| Sample | PTT | PAS | PTT | PAS | PTT | PAS |
| PTT-F/PAS-F | 77.98±0.30 ^{A,a} | 77.81±0.12 ^{A,a} | 0.975±0.007 ^{A,a} | 0.973±0.001 ^{A,a} | 0.00* | 0.00* |
| PTT-C/PAS-C | 7.99±0.05 ^{C,b} | 13.11±0.05 ^{C,a} | 0.404±0.001 ^{C,a} | 0.395±0.003 ^{C,a} | 4.76±0.77 ^{B,b} | 11.72±0.52 ^{A,a} |
| PTT-V/PAS-V | 19.03±0.40 ^{B,b} | 48.71±0.19 ^{B,a} | 0.564±0.000 ^{B,b} | 0.791±0.001 ^{B,a} | 5.25±2.06 ^{B,a} | 7.22±1.72 ^{B,a} |

| PTT-L/PAS-L | 6.54±0.04 ^{D,b} | 11.65±0.13 ^{D,a} | 0.256±0.006 ^{D,a} | 0.258±0.003 ^{D,a} | 8.36±1.57 ^{A,a} | 5.76±3.07 ^{B,a} |
|-------------|---------------------------|---------------------------|----------------------------|----------------------------|---------------------------|---------------------------|
| Sample | PL | PFT | PL | PFT | PL | PFT |
| PL-F/PFT-F | 88.23±0.21 ^{A,a} | 87.95±0.08 ^{A,a} | 0.973±0.001 ^{A,b} | 0.988±0.001 ^{A,a} | 0.00* | 0.00* |
| PL-C/PFT-C | 10.48±0.02 ^{C,b} | 14.96±0.04 ^{B,a} | 0.471±0.001 ^{C,a} | 0.451±0.003 ^{B,b} | 42.91±3.46 ^{A,a} | 34.23±2.65 ^{A,b} |
| PL-V/PFT-V | 12.45±0.06 ^{B,b} | 14.99±0.24 ^{B,a} | 0.522±0.003 ^{B,a} | 0.339±0.000 ^{C,b} | 7.39±2.65 ^{C,b} | 36.92±3.22 ^{A,a} |
| PL-L/PFT-L | 9.14±0.06 ^{D,b} | 11.65±0.29 ^{C,a} | 0.266±0.000 ^{D,b} | 0.287±0.004 ^{D,a} | 22.62±0.28 ^{B,a} | 24.82±3.13 ^{B,a} |

Means that do not share a letter are significantly different ($p < 0.05$)

Uppercase letters A, B, C and D - differences between raw, convective dried, vacuum dried and lyophilised samples within one variety of fruit

Lowercase letters a and b - differences between two varieties of the same type of fruit within one drying technique

Table 4. Experimentally obtained values of dried samples texture.

| Analyses | Shear force (g) | | Penetration force (g) | | Hardness (g) | |
|--------------|-------------------------------|------------------------------|------------------------------|-----------------------------|------------------------------|-------------------------------|
| Sample | SCF | SCEB | SCF | SCEB | SCF | SCEB |
| SCF-F/SCEB-F | 168.6±55.3 ^{C,b} | 310.5±151.0 ^{B,a} | 92.4±11.2 ^{C,a} | 94.6±20.1 ^{D,a} | 210.2±65.9 ^{C,b} | 401.5±67.7 ^{B,a} |
| SCF-C/SCEB-C | 1602.7±1085.0 ^{B,b} | 2723.5±906.7 ^{A,a} | >5000.0 ^A | >5000.0 ^A | 2263.2±810.2 ^{A,b} | 3102.3±894.5 ^{A,a} |
| SCF-V/SCEB-V | 2680.3±1032.5 ^{AB,a} | 988.0±319.9 ^{B,b} | 1730.7±1030.9 ^{B,a} | 664.7±247.9 ^{C,b} | 1129.5±311.7 ^{B,a} | 341.5±136.5 ^{B,b} |
| SCF-L/SCEB-L | 3569.4±1430.6 ^{A,a} | 2867.8±1242.7 ^{A,a} | 1171.6±398.1 ^{B,b} | 1747.8±549.1 ^{B,a} | 654.5±151.3 ^{BC,a} | 673.6±450.5 ^{B,a} |
| Sample | SCL | SCSH | SCL | SCSH | SCL | SCSH |
| SCL-F/SCSH-F | 228.7±124.4 ^{B,a} | 216.3±72.21 ^{B,a} | 135.3±45.0 ^{D,b} | 191.1±38.2 ^{D,a} | 728.6±124.9 ^{B,a} | 810.0±118.8 ^{C,a} |
| SCL-C/SCSH-C | 3194.6±998.1 ^{A,a} | 3372.0±1853.2 ^{A,a} | >5000.0 ^A | >5000.0 ^A | 1800.5±1121.1 ^{A,a} | 2237.9±814.7 ^{A,a} |
| SCL-V/SCSH-V | 777.4±443.1 ^{B,a} | 911.0±159.6 ^{B,a} | 675.6±224.6 ^{C,a} | 696.7±202.0 ^{C,a} | 405.9±144.5 ^{B,a} | 396.1±126.3 ^{C,a} |
| SCL-L/SCSH-L | 3718±1337.7 ^{A,a} | 2705.9±692.6 ^{A,b} | 1096.5±233.1 ^{B,a} | 1368.6±576.2 ^{B,a} | 786.8±235.0 ^{B,b} | 1454.8±340.8 ^{B,a} |
| Sample | AB | ANS4 | AB | ANS4 | AB | ANS4 |
| AB-F/ANS4-F | 10.9±3.10 ^{C,a} | 11.2±4.59 ^{C,a} | 351.2±75.0 ^{C,a} | 148.4±41.2 ^{C,b} | 1107.1±749.7 ^{AB,a} | 549.9±181.5 ^{C,b} |
| AB-C/ANS4-C | 1986.2±1198.6 ^{A,b} | 3535.1±1059.1 ^{A,a} | >5000.0 ^A | >5000.0 ^A | 1851.3±1562.7 ^{A,a} | 2109.1±1742.2 ^{B,a} |
| AB-V/ANS4-V | 896.1±595.2 ^{B,a} | 1205.5±291.7 ^{B,a} | 884.1±383.4 ^{B,a} | 921.5±293.8 ^{B,a} | 478.9±19.6 ^{B,a} | 422.6±174.9 ^{C,a} |
| AB-L/ANS4-L | 1764.1±523.9 ^{A,a} | 147.4±76.5 ^{C,b} | 1102.7±533.9 ^{B,a} | 790.4±274.9 ^{B,a} | 213.9±151.7 ^{B,b} | 3466.0±1337.5 ^{A,a} |
| Sample | PTT | PAS | PTT | PAS | PTT | PAS |
| PTT-F/PAS-F | 442.4±141.7 ^{B,a} | 506.6±220.4 ^{B,a} | 610.1±130.1 ^{C,a} | 420.5±86.4 ^{C,b} | 1914.2±1194.0 ^{B,a} | 1731.2±1215.2 ^{AB,a} |
| PTT-C/PAS-C | 3800.7±938.9 ^{A,a} | 2787.3±1214.9 ^{A,b} | >5000.0 ^A | >5000.0 ^A | 6377.8±3549.0 ^{A,a} | 3427.3±2874.7 ^{A,b} |
| PTT-V/PAS-V | 332.7±137.9 ^{B,b} | 528.2±186.1 ^{B,a} | 509.3±163.2 ^{C,a} | 611.4±358.5 ^{C,a} | 277.4±215.3 ^{B,a} | 252.4±128.0 ^{B,a} |


| | | | | | | |
|--------------|------------------------------|------------------------------|------------------------------|-----------------------------|------------------------------|------------------------------|
| PTT-L/PAS-L | 3856.7±1979.3 ^{A,a} | 2827.6±1425.8 ^{A,a} | 2556.7±844.9 ^{B,a} | 2852.9±713.9 ^{B,a} | 960.5±419.1 ^{B,a} | 805.6±716.9 ^{B,a} |
| Sample | PL | PFT | PL | PFT | PL | PFT |
| PL-F/PFT-F | 127.8±44.3 ^{B,a} | 117.6±36.1 ^{C,a} | 480.4±67.6 ^{C,b} | 749.1±160.2 ^{C,a} | 1691.1±488.8 ^{A,a} | 2041.7±717.1 ^{B,a} |
| PL-C/PFT-C | 2734.1±1226.9 ^{A,a} | 2726.6±1529.9 ^{A,a} | >5000.0 ^A | >5000.0 ^A | 2604.9±1711.3 ^{A,a} | 719.3±538.0 ^{B,b} |
| PL-V/PFT-V | 2800.1±1324.5 ^{A,a} | 1165.1±1081.5 ^{B,b} | 2710.4±1603.0 ^{B,a} | 1384.0±757.9 ^{B,b} | 2425.0±1809.8 ^{A,a} | 787.6±660.6 ^{B,b} |
| PL-L/PFT-L | 369.2±189.6 ^{B,a} | 424.5±266.1 ^{BC,a} | 1181.3±370.7 ^{C,a} | 825.5±237.2 ^{C,b} | 1062.1±548.8 ^{A,b} | 6153.2±3015.0 ^{A,a} |
| Analyses | Springiness | | Cohesiveness | | Chewiness (g) | |
| Sample | SCF | SCEB | SCF | SCEB | SCF | SCEB |
| SCF-F/SCEB-F | 0.94±0.24 ^{A,a} | 0.79±0.03 ^{A,b} | 0.63±0.05 ^{B,a} | 0.62±0.04 ^{A,a} | 124.2±45.6 ^{B,b} | 196.79±34.1 ^{B,a} |
| SCF-C/SCEB-C | 0.80±0.06 ^{A,a} | 0.75±0.04 ^{A,b} | 0.76±0.06 ^{A,a} | 0.69±0.04 ^{AB,b} | 1390.1±548.9 ^{A,a} | 1637.5±546.4 ^{A,a} |
| SCF-V/SCEB-V | 0.61±0.05 ^{B,b} | 0.70±0.10 ^{A,a} | 0.54±0.04 ^{B,b} | 0.61±0.04 ^{A,a} | 352.7±67.6 ^{B,a} | 143.4±50.9 ^{B,b} |
| SCF-L/SCEB-L | 0.42±0.13 ^{C,a} | 0.48±0.13 ^{B,a} | 0.73±0.16 ^{A,a} | 0.61±0.15 ^{A,a} | 189.5±50.2 ^{B,a} | 196.1±127.2 ^{B,a} |
| Sample | SCL | SCSH | SCL | SCSH | SCL | SCSH |
| SCL-F/SCSH-F | 0.76±0.03 ^{A,a} | 0.75±0.07 ^{A,a} | 0.45±0.05 ^{B,a} | 0.45±0.08 ^{C,a} | 252.0±52.3 ^{B,a} | 271.16±141.1 ^{B,a} |
| SCL-C/SCSH-C | 0.34±0.13 ^{C,b} | 0.72±0.13 ^{A,a} | 0.72±0.26 ^{A,a} | 0.73±0.05 ^{A,a} | 475.2±197.6 ^{A,b} | 1199.6±588.3 ^{A,a} |
| SCL-V/SCSH-V | 0.78±0.10 ^{A,a} | 0.74±0.07 ^{A,a} | 0.60±0.02 ^{AB,a} | 0.56±0.03 ^{B,b} | 193.6±82.4 ^{B,a} | 162.4±53.9 ^{B,a} |
| SCL-L/SCSH-L | 0.47±0.09 ^{B,b} | 0.57±0.03 ^{B,a} | 0.63±0.10 ^{A,a} | 0.50±0.02 ^{BC,b} | 224.6±73.5 ^{B,b} | 412.45±85.1 ^{B,a} |
| Sample | AB | ANS4 | AB | ANS4 | AB | ANS4 |
| AB-F/ANS4-F | 0.35±0.10 ^{A,a} | 0.29±0.03 ^{C,b} | 0.27±0.07 ^{C,a} | 0.18±0.05 ^{C,b} | 197.2±252.0 ^{B,a} | 26.5±15.5 ^{B,b} |
| AB-C/ANS4-C | 0.46±0.17 ^{A,a} | 0.49±0.16 ^{B,a} | 0.74±0.10 ^{B,a} | 0.66±0.11 ^{A,a} | 690.5±601.2 ^{A,a} | 685.6±570.0 ^{A,a} |
| AB-V/ANS4-V | 0.37±0.05 ^{A,b} | 0.68±0.12 ^{A,a} | 0.92±0.06 ^{A,a} | 0.53±0.04 ^{B,b} | 162.2±23.2 ^{B,a} | 154.8±76.8 ^{B,a} |

| | | | | | | |
|-------------|---------------------------|---------------------------|---------------------------|--------------------------|------------------------------|-----------------------------|
| AB-L/ANS4-L | 0.40±0.13 ^{A,a} | 0.31±0.06 ^{C,b} | 0.86±0.09 ^{A,a} | 0.24±0.06 ^{C,b} | 79.4±76.1 ^{B,b} | 239.3±82.8 ^{B,a} |
| Sample | PTT | PAS | PTT | PAS | PTT | PAS |
| PTT-F/PAS-F | 0.66±0.08 ^{A,b} | 0.79±0.04 ^{A,a} | 0.59±0.09 ^{B,b} | 0.71±0.04 ^{A,a} | 806.8±655.2 ^{B,a} | 623.1±216.5 ^{B,a} |
| PTT-C/PAS-C | 0.53±0.10 ^{AB,a} | 0.59±0.19 ^{C,a} | 0.50±0.18 ^{B,b} | 0.64±0.12 ^{A,a} | 1588.7±1088.0 ^{A,a} | 1219.1±953.0 ^{A,a} |
| PTT-V/PAS-V | 0.49±0.07 ^{B,b} | 0.64±0.15 ^{BC,a} | 0.76±0.36 ^{AB,a} | 0.63±0.06 ^{A,a} | 89.0±63.3 ^{C,a} | 107.0±60.3 ^{B,a} |
| PTT-L/PAS-L | 0.34±0.20 ^{C,b} | 0.77±0.10 ^{AB,a} | 0.95±0.31 ^{A,a} | 0.66±0.06 ^{A,b} | 321.1±240.5 ^{BC,a} | 436.7±466.2 ^{B,a} |
| Sample | PL | PFT | PL | PFT | PL | PFT |
| PL-F/PFT-F | 0.58±0.38 ^{A,a} | 0.47±0.14 ^{B,a} | 0.13±0.03 ^{C,b} | 0.17±0.03 ^{D,a} | 141.5±109.6 ^{C,a} | 157.9±66.1 ^{B,a} |
| PL-C/PFT-C | 0.67±0.18 ^{A,a} | 0.74±0.26 ^{A,a} | 0.62±0.09 ^{A,b} | 0.91±0.27 ^{A,a} | 1054.5±692.6 ^{A,a} | 513.5±368.7 ^{A,b} |
| PL-V/PFT-V | 0.67±0.18 ^{A,a} | 0.70±0.15 ^{A,a} | 0.53±0.18 ^{AB,a} | 0.60±0.15 ^{B,a} | 672.4±370.3 ^{AB,a} | 322.4±284.6 ^{AB,b} |
| PL-L/PFT-L | 0.71±0.62 ^{A,a} | 0.28±0.05 ^{C,b} | 0.45±0.16 ^{B,a} | 0.39±0.13 ^{C,a} | 400.4±400.4 ^{BC,a} | 616.2±342.3 ^{A,a} |

Means that do not share a letter are significantly different ($p < 0.05$)


Uppercase letters A, B, C and D - differences between raw, convective dried, vacuum dried and lyophilised samples within one variety of fruit


Lowercase letters a and b - differences between two varieties of the same type of fruit within one drying technique



jfpp_15158_f1.tif

Figure 2.







Means that do not share a letter are significantly different ($p < 0.05$)

Uppercase letters A, B, C and D - differences between raw, convective dried, vacuum dried and lyophilised samples within one variety of fruit

Lowercase letters a and b - differences between two varieties of the same type of fruit within one drying technique

Figure 3.







Means that do not share a letter are significantly different ($p < 0.05$)

Uppercase letters A, B, C and D - differences between raw, convective dried, vacuum dried and lyophilised samples within one variety of fruit

Lowercase letters a and b - differences between two varieties of the same type of fruit within one drying technique



jfpp_15158_f4.tif

