This is the final peer-reviewed accepted manuscript of:

Mescolini G, Lupini C, Franzo G, Quaglia G, Legnardi M, Cecchinato M, Tucciarone CM, Blanco A, Turblin V, Biarnés M, Tatone F, Falchieri M, Catelli E. What is new on molecular characteristics of Avian metapneumovirus strains circulating in Europe? Transbound Emerg Dis. 2020 Aug 13. doi: 10.1111/tbed.13788. Epub ahead of print. PMID: 32794302.

The final published version is available online at:
<https://onlinelibrary.wiley.com/doi/abs/10.1111/tbed.13788>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

1 **What is new on molecular characteristics of Avian metapneumovirus strains circulating in**
2 **Europe?**

3

4 Molecular epidemiology of aMPV-B in Europe

5

6 Giulia Mescolini¹, Caterina Lupini^{1,*}, Giovanni Franzo², Giulia Quaglia¹, Matteo Legnardi², Mattia
7 Cecchinato², Claudia Maria Tucciarone², Angela Blanco³, Vincent Turblin⁴, Mar Biarnés³, Fabrizio
8 Tatone⁵, Marco Falchieri⁶, Elena Catelli¹

9

10 ¹ Department of Veterinary Medical Sciences, University of Bologna, Ozzano dell'Emilia (BO),
11 Italy

12 ² Department of Animal Medicine, Production and Health, University of Padua, Legnaro (PD), Italy

13 ³ CESAC - Centre de Sanitat Avícola de Catalunya i Aragó, Reus, Spain.

14 ⁴ MC VET Conseil – RESEAU CRISTAL, Sablé sur Sarthe, France

15 ⁵ RESALAB – RESEAU CRISTAL, Les Herbiers, France

16 ⁶ Field Veterinarian, United Kingdom

17

18 *Corresponding author: Caterina Lupini, Department of Veterinary Medical Sciences -University of
19 Bologna, Via Tolara di Sopra, 50; 40064 – Ozzano dell'Emilia (BO) Italy; Phone +39 051 2097559;
20 e-mail: caterina.lupini@unibo.it

21

22 **Summary**

23 In the present study one hundred and sixteen partial G gene sequences of Avian metapneumovirus
24 (aMPV) subtype B, obtained during routine diagnostics in different European Countries in the last
25 few years (2014-2019), were analysed by sequence and phylogenetic analyses in order to draw an
26 updated picture of the molecular characteristics of circulating strains. Nucleotide sequences were
27 compared with other sequences of European and non-European aMPV-Bs collected prior to that
28 period or retrieved from GenBank. Phylogenetic relationships among the aMPV-B strains,
29 reconstructed using the Maximum Likelihood method implemented in MEGA X, demonstrated that
30 aMPV-B has evolved in Europe from its first appearance, frequently displaying a clear relation with
31 the geographic area of detection. The 40% of aMPV-B viruses analysed were classified as vaccine-
32 derived strains, being phylogenetically related, and showing high nucleotide identity with live
33 commercial vaccine strains licensed in Europe. The remaining 60% were classified as field strains
34 since they clustered separately and showed a low nucleotide identity with vaccines and vaccine-
35 derived strains. The phylogenetic tree showed that the virus has continued to evolve from its first
36 appearance in the '80s since more recently detected strains belonged to clades phylogenetically
37 distant from the older strains. Unlike vaccine-derived strains, field strains tended to cluster
38 according to their geographic origin and irrespective of the host species where the viruses had been
39 detected. In conclusion, the molecular characterization of aMPV-B and the differentiation between
40 vaccines and field strains through G gene sequence analysis can be a useful tool towards correct
41 diagnosis and should be routinely applied in order to better address the control strategies.

42

43 ***Key words:* Avian metapneumovirus, subtype B, Europe, molecular characterization**

44

45

46 **Introduction**

47 *Avian metapneumovirus* (aMPV) infections in Europe have been an issue since its first appearance
48 both in turkeys, chickens and in other minor species (Toquin, Bayon-Auboyer, Eterradossi, & Jestin,
49 1999; Catelli et al., 2001; Cecchinato, Ferreira, Munir, & Catelli, 2017; Cecchinato et al., 2018),
50 causing economic losses mainly due to respiratory or reproductive problems (Cecchinato et al., 2012),
51 often exacerbated by secondary bacterial infections (Giovanardi et al., 2014).

52 First aMPV isolations in Europe date back to the second half of the 1980s, when Turkey
53 Rhinotracheitis (TRT) outbreaks appeared in the United Kingdom (McDougall and Cook, 1986) and
54 France (Giraud, Bennejean, Guittet, & Toquin, 1986). As the virus was spreading all over Europe and
55 the clinical problems in poultry farms became increasingly serious, during the early 1990s (Cook et
56 al. 1989a e 1989b) live attenuated vaccines were developed and became commercially available. By
57 analysing strains circulating in Europe, Juhasz and Easton (1994) reported differences in the G gene
58 between aMPV isolates and proposed the classification into A and B subtypes and confirmed the co-
59 circulation of both subtypes. After this initial study, and due to the increasing use of sequence
60 analysis, two further subtypes, named C and D, were identified in France (Bayon-Auboyer, Arnauld,
61 Toquin, & Eterradossi, 2000; Toquin et al., 2006).

62 Despite these reports, molecular data on aMPV strains circulating in Europe are still poor and
63 scattered (Catelli et al., 2004; Cecchinato et al., 2013a; Listorti et al., 2014; Franzo et al., 2017;
64 Tucciarone et al., 2017 a 2018; Ball, Forrester, & Ganapathy, 2018; Andreopoulou et al., 2019), as
65 most of them originate from few countries. In the last decades, subtype B has been generally the most
66 frequently encountered subtype in Europe, although subtype A has been sporadically reported (Lupini
67 et al., 2011).

68 In order to update the epidemiological picture of circulating strains, the present study was designed
69 to molecularly characterize, by partial G gene sequencing, aMPV subtype B strains detected in
70 Europe in the last years.

71 **Material and Methods**

72 **Sample collection**

73 The survey covered the time period from 2014 to 2019 and targeted aMPV subtype B detected
74 during routine diagnostic activity performed in Italy by the Universities of Bologna and Padua or in
75 Spain by the Centre de Sanitat Avícola de Catalunya i Aragó (CESAC).

76 Samples originated from different European countries and were usually collected during outbreaks
77 of respiratory disorders referable to aMPV infections in turkey, chicken or guinea fowl flocks.
78 Sampling was also performed for epidemiological purposes in absence of clear clinical signs.

79 As a rule, samples consisted of pools of 10 rhino-pharyngeal swabs per flock as previously
80 suggested (Catelli et al., 2004).

81

82 **RNA extraction and PCR analysis**

83 RNA was extracted from each pool of swabs and subjected to a nested RT-PCR targeting the G
84 gene (Cavanagh, Mawditt, Britton, & Naylor, 1999, slightly modified) which allows the
85 simultaneous detection and differentiation of subtypes A and B. The reverse transcription (RT)
86 reaction was primed using oligonucleotide G6- (5'- CTGACAAATTGGTCCTGATT- 3').
87 Subsequently the same oligonucleotide was used together with oligonucleotide Gstart+ (5'-
88 CAAGTATCCAGATGGGGTC- 3') for the first PCR round. The obtained PCR product was then
89 subjected to a second PCR round with oligonucleotide G5- (5'- CAAAGAGCCAATAAGCCCA-
90 3') in conjunction with oligonucleotides G8+A (5'-CACTCACTGTTAGCGTCATA-3') and G9+B
91 (5'-TAGTCCTCAAGCAAGTCCTC-3'). The predicted amplicon length is of 268 bp for subtype A
92 and 361 bp for subtype B. Depending on the laboratory, this step was preceded by a multiplex real-
93 time RT-PCR (qRT-PCR) screening test targeting the SH gene (Cecchinato et al., 2013b) and only
94 positive samples were subsequently amplified by RT-nested PCR.

95

96 **Partial G gene sequencing**

97 The partial G gene amplicons (361 bp-long) obtained were purified using ExoSAP-IT™ *Express*
98 PCR Product Cleanup (Thermo Fisher Scientific, Massachusetts, USA) according to the
99 manufacturer's instructions and sequenced in both directions using RT-nested PCR primers G5- (5'-
100 CAAAGAGCCAATAAGCCCA-3') and G9+B (5'-TAGTCCTCAAGCAAGTCCTC-3'), by a
101 commercial sequencing service (Macrogen Europe, Amsterdam, The Netherlands). The obtained
102 sequences were named using the following nomenclature: aMPV/B/Country of origin/Host species
103 (Turkey: Ty, Chicken: Ck or Guinea fowl)/sample ID number/year of detection.

104

105 **Sequence and phylogenetic analysis**

106 Nucleotide sequences were edited and assembled using BioEdit software, then, using Clustal W,
107 aligned against and compared with G gene sequences of:

108 - The most commonly used subtype B vaccines (i.e. aMPV/B vaccine - strain VCO3; aMPV/B
109 vaccine - Strain 11/94; aMPV/B vaccine - strain 1062 and aMPV/B vaccine - strain PL21)

110 - 82 already available European aMPV B strains (Table S1)

111 - 59 non-European aMPV-B strains, retrieved from *GenBank* (Table S2).

112 Phylogenetic relationships among the aMPV-B strains were reconstructed using the Neighbor-
113 Joining method and Kimura 2-parameter model implemented in MEGA X (Kumar et al., 2018). The
114 substitution model was selected based on Bayesian information criterion (BIC), calculated using the
115 same software. The branch support was calculated by performing 1000 bootstrap replicates; only
116 branches supported by bootstrap values $\geq 70\%$ were considered reliable. Complete deletion option
117 was selected before the analysis begins, to remove sites containing missing data or alignment gaps.

118 Within-group mean pairwise genetic p-distance was estimated using MEGA X in order to evaluate
119 the genetic heterogeneity of aMPV population. Overall nucleotide similarity was further estimated
120 by calculating the arithmetic mean of the sequence identity values, obtained using the Sequence
121 Identity Matrix tool on BioEdit software, between field strains or vaccine strains belonging to the
122 same phylogenetic cluster.

123

124 **Results**

125 116 samples tested aMPV positive, all belonging to subtype B (n. 27 from France, n. 18 from Italy,
126 n. 27 from Romania, n. 36 from Spain and n. 8 from the United Kingdom) and 116 partial G gene
127 sequences were obtained. Sequence data have been submitted to the GenBank database under
128 accession numbers MT432826-MT432923 and MT436220-MT436237 (Tables S3 to S7).

129 Phylogenetic analysis was carried out by reconstructing both a comprehensive tree (Figure 1) and
130 five Country- specific trees (Figures S1 to S5). Furthermore, a phylogenetic tree including selected
131 European and non-European aMPV-B sequences was generated (Figure 2). The number of nucleotide
132 positions included in the final datasets subjected to phylogenetic analysis is specified in the legends
133 of the figures.

134 Regardless of the Country of origin, aMPV strains detected from 2001 to 2019, herein referred to as
135 “recent strains”, were distinguished from those clustering with vaccines and field strains detected
136 prior to the 1990s, referred to as “older strains” (from which the vaccines were established by
137 attenuation) and those forming independent clades.

138 As a rule, aMPV strains were referred to as “vaccine-derived strains” if the maximum nucleotide
139 sequence identity with a reference vaccine strain was greater than or equal to 99 % (2 or less
140 nucleotide differences) and if they fell into the same phylogenetic cluster. The remaining strains were
141 classified as “field strains”.

142 On the above basis, 70 out of 116 aMPV-B strains were labelled as field strains (60%); the remaining
143 46 strains (40%) were identified as vaccine-derived strains.

144 A certain degree of geographic clustering was observed among Romanian, Spanish, French and
145 Italian field strains (Figure 1). Some Italian strains also clustered together with some recent Greek
146 strains.

147 French aMPV field strains (Figure S1) were divided in two clusters. The largest cluster displayed a
148 within-group mean p-distance of 0.003 (overall nucleotide similarity of 99.7%) and included chicken,
149 turkey and guinea fowl strains detected from 2017 to 2019, while the smallest had a within-group
150 mean p-distance of 0.000 (overall nucleotide similarity of 100%) and contains turkey strains detected
151 in 2018. Vaccine-derived strains clustered respectively with vaccine strain PL21, 11/94 or VCO3 and
152 were mostly detected in homologous-vaccinated turkeys or chickens, but also in unvaccinated guinea
153 fowls. The cluster containing vaccines, vaccine-derived strains and older French strains displayed a
154 within-group mean p-distance of 0.006 (overall nucleotide similarity of 99.4%)

155 Italian field strains were grouped in two main clusters (Figure S2): the larger one was composed of
156 several sub-clusters including field strains detected from 2010 to 2019 in turkey, chicken, and guinea
157 fowl flocks and showed a within-group mean p-distance of 0.010 (overall nucleotide similarity of
158 99.0%). The smaller one included field strains detected from 2001 to 2009 and in 2016 in turkeys and
159 chickens, with a within-group mean p-distance of 0.004 (overall nucleotide similarity of 99.6%).
160 Vaccine-derived strains clustering with aMPV-B vaccine strains VCO3 or 1062 were detected from
161 vaccinated turkeys. The cluster containing vaccines, vaccine-derived strains and older Italian strains
162 displayed a within-group mean p-distance of 0.007 (overall nucleotide similarity of 99.3%).

163 Romanian aMPV-B field strains (Figure S3), detected from turkeys or chickens, fell within a single
164 phylogenetic group showing a within-group mean p-distance of 0.001 (overall nucleotide similarity
165 of 99.9%). The vaccine-derived strains, all detected in vaccinated turkey flocks, clustered with

166 vaccine strains PL21, 11/94 or VCO3. The cluster containing vaccines and vaccine-derived strains
167 displayed a within-group mean p-distance of 0.007 (overall nucleotide similarity of 99.3%).

168 In the Spanish tree (Figure S4) field strains fell within two main clusters: a larger one (within-group
169 mean p-distance: 0.006 and overall nucleotide similarity of 99.4%) containing strains detected from
170 2014 to 2017; a smaller one (within-group mean p-distance: 0.003 and overall nucleotide similarity
171 of 99.7%) containing viruses detected from 2014 to 2015. Spanish vaccine-derived strains clustered
172 with vaccine strain 11/94 or PL21. The cluster containing vaccines, vaccine-derived strains and older
173 strains displayed a within-group mean p-distance of 0.008 (overall nucleotide similarity of 99.2%).

174 The British phylogenetic tree (Figure S5) exclusively displayed vaccine-derived strains, clustering
175 with vaccine strain PL21, 11/94 or VCO3 (within-group mean p-distance: 0.007 and overall
176 nucleotide similarity of 99.3%).

177 The tree reconstructed with representative aMPV-B sequences from all over the world (Figure 2)
178 confirmed the Country-specific clustering trend seen for the European strains. Well-defined clusters
179 were identified for field strains circulating in Brazil, Iran, Israel and Turkey. Vaccines, vaccine-
180 derived and older strains clustered together and were showed in the figure compressed in two
181 subtrees. Within-group mean p-distance and overall nucleotide similarity of the cluster containing
182 vaccine PL21 were respectively 0.007 of 99.3%,

183

184 **Discussion**

185 In the present study, one hundred and sixteen aMPV-B partial G gene sequences, obtained during
186 routine diagnostics in different European Countries in the last few years, were analysed by sequence
187 and phylogenetic analyses in order to molecularly characterize them.

188 The G gene, which harbours mutations at variable positions between aMPV-B strains, was
189 conveniently amplified by the routine diagnostic RT-PCR protocol employed in the study, and its

sequencing proved useful for epidemiological purposes. Furthermore, its variability can give an indication of whether vaccine or field strains are present, making the differentiation relatively easy and cheap to perform.

Live attenuated aMPV vaccines are widely administered to prevent disease in turkeys and chickens, and the recovery of vaccine-derived strains is not unusual both in vaccinated or in unvaccinated flocks (Banet-Noach et al., 2009; Lupini et al., 2011; Chacon et al., 2011; Cecchinato et al., 2013a; Listorti et al., 2014; Arafa et al., 2015; Bayraktar et al., 2019; Andreopoulou et al., 2019).

The 40% of the aMPV-B strains detected in the present study were classified as vaccine-derived strains, being phylogenetically related and showing high nucleotide identity with live commercial vaccine strains licensed in Europe. As expected, vaccine-derived strains formed separate clusters depending on the vaccine strain of origin and were detected in all tested European countries.

A large part of the vaccine-derived strains analysed in the present study was detected in homologous-vaccinated birds, from approximately two to four weeks after vaccination, and only occasionally in unvaccinated birds. Reversion to virulence of aMPV subtype A or B has been previously demonstrated (Catelli et al., 2006; Brown et al., 2011, Cecchinato et al., 2014).

Therefore, the detection of vaccine-derived strains closely related to the applied vaccine, concurrently with respiratory signs, could be reliably linked to the vaccine reversion to virulence.

The detection of the strain aMPV/B/France/GuineaFowl/1060/18 in unvaccinated guinea fowls is noteworthy, since its partial G gene sequence shared 100% nucleotide identity with the vaccine strain 11/94. It could be speculated that the vaccine virus could have been introduced as a contaminant by personnel, fomites, vehicle movement or airborne from neighbouring premises.

The field veterinarian reported the presence of a turkey farm at approximately 500 meters from the guinea fowl one. Vaccinal strain 11/94 was applied in the turkey flock and the use of the litter from the turkey farm for fertilization of the surrounding crops was reported. The ability of vaccine-derived aMPVs to spread beyond the administration site is well known and it has been proven by

215 Lupini et al. (2011) following a turkey rhinotracheitis outbreak caused by aMPV subtype A in
216 unvaccinated turkeys.

217 The remaining 60% aMPV-B viruses analysed in the present study were classified as field strains
218 since they clustered separately and showed a low nucleotide identity with vaccines and vaccine-
219 derived strains.

220 The phylogenetic tree reconstructed with European sequences showed that the virus has continued
221 to evolve from its first appearance in the '80s. In fact, more recently detected field strains belonged
222 to clades phylogenetically distant from the older field strains, confirming the previously-reported
223 aMPV tendency to evolve over time (Cecchinato et al., 2010).

224 Unlike vaccine-derived strains, field strains tended to cluster according to their geographic origin,
225 with few exceptions. Distinct clusters were observed for French, Italian, Romanian and Spanish
226 strains, yet some Italian and Greek field isolates clustered closely together, indicating a potential
227 transmission route between these two countries (Tucciarone et al., 2017; Andreopoulou et al.,
228 2019).

229 The molecular epidemiology of aMPV within each country was analysed in detail reconstructing
230 country-related phylogenetic trees. Heterogeneous field strain populations seemed to co-exist within
231 single European countries, with the only exception of Romania, where all identified strains were
232 part of just one clade. This last finding could be explained by keeping into account that all the
233 processed samples came from different sites within the same company in which viral circulation
234 could be compartmentalized (Franzo et al., 2020).

235 French, Italian, and Spanish trees showed a rather heterogeneous field strain population, as the
236 strains fell into more than one cluster, divided in several sub-clusters. The heterogeneity was further
237 deduced from the within-group mean p-distance values. The highest value, indicating the highest
238 heterogeneity of nucleotide sequence, was observed for Italian field strains, presumably because of

239 the larger number of sequences available over a broader time period and coming from different
240 commercial poultry companies.

241 Furthermore, time-related clustering was visible for Italian and Spanish field strains, as strains
242 detected in the last few years tend to form separate clusters or sub-clusters.

243 As a common finding, recent field aMPV-B strains, belonging to different and distant phylogenetic
244 clades, were detected in the presence of respiratory signs in vaccinated flocks. Due to the G gene
245 variability observed in recent aMPV-B strains, and knowing that the encoded surface G
246 glycoprotein is a key antigen for vaccine-induced immune protection (Naylor et al., 2007) and can
247 evolve in order to avoid vaccine-induced immunity (Catelli et al., 2010; Cecchinato et al., 2010),
248 re-assessment of protection conferred by commercially available vaccines against currently
249 circulating aMPV field strains might be necessary to improve disease control strategies.

250 The obtained phylogenetic data revealed that aMPV-B strains clustered together irrespective of the
251 host species where the viruses had been detected. A recent experimental challenge study showed
252 that both chickens and turkeys are susceptible to aMPV-B infection with the same virus isolate
253 (Brown et al., 2019). Moreover Cecchinato et al. (2018) reported an outbreak of respiratory disease
254 in guinea fowls caused by aMPV-B strain identical to the ones circulating in the surrounding turkey
255 flocks. Therefore, no evidences are currently available to support a host-specific adaptation of
256 aMPV variants.

257 As a final remark, the tree reconstructed on European and non-European strains further confirmed
258 the geographic clustering of field strains. As a rule, field strains grouped together for country
259 location and time period, suggesting a local evolution tendency of the virus that might have taken
260 place after a single introduction event. The majority of recently detected European field strains were
261 located in a big subtree indicating a certain genetic similarity and supporting the wide circulation of
262 a quite homogeneous aMPV subtype B clade in European countries, with the only exception of two
263 clusters of phylogenetically-distant French and Spanish strains.

264 In conclusion, the molecular characterization of aMPV subtype B and the differentiation between
265 vaccines and field strains through G gene sequence analysis can be a useful tool towards a correct
266 diagnosis and should be routinely applied in order to better address the control strategies. In this
267 respect, current vaccine research is focused on reducing the issues connected to live attenuated
268 vaccine reversion to virulence or the selection of potentially virulent subpopulations (Franzo et al.,
269 2015). Therefore, considering the associated risks, further efforts should be directed at improving
270 administration and biosecurity measures, in order to reduce their prolonged circulation and
271 spreading.

272 Several research groups have attempted to develop efficacious, more stable and safer next-
273 generation vaccines but, despite this effort, conventional live attenuated vaccines still provide the
274 greatest protection after homologous challenge (Qingzhong et al., 1994; Tarpey et al., 2001;
275 Kapczynski and Sellers, 2003; Kapczynski, 2004; Chary, Njenga, & Sharma, 2005; Yu et al., 2013;
276 Hu, Roth, Zsak, & Yu, 2017).

277 Promising reverse genetics systems for aMPV have been developed in recent years for A, B and C
278 subtypes (Naylor et al., 2004; Yu et al., 2010; Laconi et al., 2016) other than being an exceptional
279 tool for the study of the virus properties (Brown et al., 2011), can be used for the development of
280 rationally modified aMPV vaccines (Naylor, Lupini, & Brown, 2010) or recombinant vaccines
281 (Falchieri et al., 2013) expressing foreign genes.

282

283 **Conflict of Interest Statement**

284 The authors declare no conflict of interest.

285

286 **Ethical approval**

287 The authors confirm that the ethical policies of the journal, as noted on the journal's author guidelines
288 page, have been adhered to. No ethical approval was required as the original research data of this

289 article was obtained from oro-pharyngeal swabs collected for routine diagnostics purpose by field
290 veterinarians.

291

292 **References**

- 293 Andreopoulou, M., Franzo, G., Tucciarone, C. M., Prentza, Z., Koutoulis, K. C., Cecchinato, M.,
294 & Chaligianni, I. (2019). Molecular epidemiology of infectious bronchitis virus and avian
295 metapneumovirus in Greece. *Poultry Science*, 98(11), 5374–5384.
296 <https://doi.org/10.3382/ps/pez360>
- 297 Arafa, A.-S., Mady, W., Hussein, A., Tamam, S., & Madbouly, H. (2015). Molecular
298 Characterization of Vaccine-Derived Mutants of Avian Meta-Pneumoviruses Isolated from
299 Turkeys in Egypt. *American Journal of Virology*, 4(1), 1–11.
300 <https://doi.org/10.3844/ajvsp.2015.1.11>
- 301 Ball, C., Forrester, A., & Ganapathy, K. (2018). Co-circulation of genetically diverse population
302 of vaccine related and unrelated respiratory mycoplasmas and viruses in UK poultry flocks
303 with health or production problems. *Veterinary Microbiology*, 225, 132–138.
304 <https://doi.org/10.1016/j.vetmic.2018.09.009>
- 305 Banet-Noach, C., Simanov, L., Laham-Karam, N., Perk, S., & Bacharach, E. (2009). Longitudinal
306 survey of avian metapneumoviruses in poultry in Israel: infiltration of field strains into
307 vaccinated flocks. *Avian Diseases*, 53(2), 184–189. [https://doi.org/10.1637/8466-090408-](https://doi.org/10.1637/8466-090408-Reg.1)
308 [Reg.1](https://doi.org/10.1637/8466-090408-Reg.1)
- 309 Bayon-Auboyer, M. H., Arnauld, C., Toquin, D., & Eterradossi, N. (2000). Nucleotide sequences
310 of the F, L and G protein genes of two non-A/non-B avian pneumoviruses (APV) reveal a
311 novel APV subgroup. *Journal of General Virology*, 81(11), 2723–2733.
312 <https://doi.org/10.1099/0022-1317-81-11-2723>

313 Bayraktar, E., Umar, S., Yilmaz, A., Turan, N., Franzo, G., Tucciarone, C. M., ... Yilmaz, H.
314 (2019). First Molecular Characterization of Avian Metapneumovirus (aMPV) in Turkish
315 Broiler Flocks. *Avian Diseases*, 62(4), 425–430. [https://doi.org/10.1637/11915-061818-](https://doi.org/10.1637/11915-061818-resnote.1)
316 [resnote.1](https://doi.org/10.1637/11915-061818-resnote.1)

317 Brown, P.A., Allée, C., Courtillon, C., Szerman, N., Lemaitre, E., Toquin, D., Mangart, J.M., ...
318 Eterradossi, N. (2019). Host specificity of avian metapneumoviruses. *Avian Pathology*,
319 48(4), 311–318. <https://doi.org/10.1080/03079457.2019.1584390>

320 Brown, P. A., Lupini, C., Catelli, E., Clubbe, J., Ricchizzi, E., & Naylor, C. J. (2011). A single
321 polymerase (L) mutation in avian metapneumovirus increased virulence and partially
322 maintained virus viability at an elevated temperature. *Journal of General Virology*, 92(Pt
323 2), 346–354. <https://doi.org/10.1099/vir.0.026740-0>

324 Catelli, E., De Marco, M. A., Delogu, M., Terregino, C., & Guberti, V. (2001). Serological
325 evidence of avian pneumovirus infection in reared and free-living pheasants. *Veterinary*
326 *Record*, 149(2), 56–58. <https://doi.org/10.1136/vr.149.2.56>

327 Catelli, E., Cecchinato, M., Cassandro, M., Delogu, M., De Matteo, P., Franciosi, C., ... John
328 Naylor, C. (2004). Avian Pneumovirus infection in turkey and broiler farms in Italy: a
329 virological, molecular and serological field survey. *Italian Journal of Animal Science*,
330 3(3), 287–292. <https://doi.org/10.4081/ijas.2004.287>

331 Catelli, E., Cecchinato, M., Savage, C. E., Jones, R. C., & Naylor, C. J. (2006). Demonstration of
332 loss of attenuation and extended field persistence of a live avian metapneumovirus
333 vaccine. *Vaccine*, 24(42-43), 6476–6482. <https://doi.org/10.1016/j.vaccine.2006.06.076>

334 Cavanagh, D., Mawditt, K., Britton, P., & Naylor, C. J. (1999). Longitudinal field studies of
335 infectious bronchitis virus and avian pneumovirus in broilers using type-specific

336 polymerase chain reactions. *Avian Pathology*, 28(6), 593–605.

337 <https://doi.org/10.1080/03079459994399>

338 Cecchinato, M., Catelli, E., Lupini, C., Ricchizzi, E., Prosperi, S., & Naylor, C. J. (2014).

339 Reversion to virulence of a subtype B avian metapneumovirus vaccine: Is it time for

340 regulators to require availability of vaccine progenitors? *Vaccine*, 32(36), 4660–4664.

341 <https://doi.org/10.1016/j.vaccine.2014.06.030>

342 Cecchinato, M., Ferreira, H. L., Munir, M., & Catelli, E. (2017). Avian metapneumovirus. In

343 Mononegaviruses of veterinary importance. Volume 2: molecular epidemiology and

344 control. <https://doi.org/10.1079/9781780644172.0127>

345 Cecchinato, M., Lupini, C., Silveira, F., Listorti, V., Mescolini, G., Morandini, E., ... Catelli, E.

346 (2018). Molecular characterization of avian metapneumovirus from Guinea fowls (*numida*

347 *meleagris*). *Pakistan Veterinary Journal*, 38(4), 419–423.

348 <https://doi.org/10.29261/pakvetj/2018.088>

349 Cecchinato, M., Drigo, M., Lupini, C., Martini, M., Listorti, V., Franzo, G., ... Catelli, E. (2013a).

350 Field survey of avian metapneumovirus in Northern Italy. *Large Animal Review*, 19(6),

351 267–270.

352 Cecchinato, M., Catelli, E., Lupini, C., Ricchizzi, E., Clubbe, J., Battilani, M., & Naylor, C. J.

353 (2010). Avian metapneumovirus (AMPV) attachment protein involvement in probable

354 virus evolution concurrent with mass live vaccine introduction. *Veterinary Microbiology*,

355 146(1–2), 24–34. <https://doi.org/10.1016/j.vetmic.2010.04.014>

356 Cecchinato, M., Lupini, C., Munoz Pogoreltseva, O. S., Listorti, V., Mondin, A., Drigo, M., &

357 Catelli, E. (2013b). Development of a real-time RT-PCR assay for the simultaneous

358 identification, quantitation and differentiation of avian metapneumovirus subtypes A and

359 B. *Avian Pathology*, 42(3), 283–289. <https://doi.org/10.1080/03079457.2013.788130>

- Cecchinato, M., Lupini, C., Ricchizzi, E., Falchieri, M., Meini, A., Jones, R. C., & Catelli, E. (2012). Italian Field Survey Reveals a High Diffusion of Avian Metapneumovirus Subtype B in Layers and Weaknesses in the Vaccination Strategy Applied. *Avian Diseases*, 56(4), 720–724. <https://doi.org/10.1637/10202-041312-reg.1>
- Chacón, J. L., Mizuma, M., Vejarano, M. P., Toquín, D., Eterradosi, N., Patnayak, D. P., ... Piantino Ferreira, A. J. (2011). Avian Metapneumovirus Subtypes Circulating in Brazilian Vaccinated and Nonvaccinated Chicken and Turkey Farms. *Avian Diseases*, 55(1), 82–89. <https://doi.org/10.1637/9501-081310-reg.1>
- Chary, P., Njenga, M. K., & Sharma, J. M. (2005). Protection by recombinant viral proteins against a respiratory challenge with virulent avian metapneumovirus. *Veterinary Immunology and Immunopathology*, 108(3-4), 427–432. <https://doi.org/10.1016/j.vetimm.2005.06.012>
- Cook, J.K.A., Ellis, M.M., Dolby, C.A., Huggins, M.B., Holmes, H.C., Finney, P.M. (1989a). A Live Attenuated Turkey Rhinotracheitis Virus Vaccine. 1. Stability of the Attenuated Strain. *Avian Pathology*, 18(3), 511–522. <https://doi.org/10.1080/03079458908418623>
- Cook, J. K. A., Dolby, C. A., Ellis, M. M., Huggins, M. B., Holmes, H. C., & Finney, P. M. (1989b). A Live Attenuated Turkey Rhinotracheitis Virus Vaccine. 2. the Use of the Attenuated Strain as an Experimental Vaccine. *Avian Pathology*, 18(3), 523–534. <https://doi.org/10.1080/03079458908418624>
- Falchieri, M., Lupini, C., Cecchinato, M., Catelli, E., Kontolaimou, M., & Naylor, C. J. (2013). Avian metapneumoviruses expressing Infectious Bronchitis virus genes are stable and induce protection. *Vaccine*, 31(22), 2565–2571. <https://doi.org/10.1016/j.vaccine.2013.03.055>

- Franzo, G., Naylor, C. J., Drigo, M., Croville, G., Ducatez, M. F., Catelli, E., ... Cecchinato, M. (2015). Subpopulations in aMPV vaccines are unlikely to be the only cause of reversion to virulence. *Vaccine*, 33(21), 2438–2441. <https://doi.org/10.1016/j.vaccine.2015.03.092>
- Franzo, G., Tucciarone, C. M., Enache, M., Bejan, V., Ramon, G., Koutoulis, K. C., & Cecchinato, M. (2017). First Report of Avian Metapneumovirus Subtype B Field Strain in a Romanian Broiler Flock During an Outbreak of Respiratory Disease. *Avian Diseases*, 61(2), 250–254. <https://doi.org/10.1637/11557-121216-ResNote.1>
- Franzo, G., Tucciarone, C.M., Moreno, A., Legnardi, M., Massi, P., Tosi, G., ..., Cecchinato, M. (2020). Phylodynamic analysis and evaluation of the balance between anthropic and environmental factors affecting IBV spreading among Italian poultry farms. *Scientific Reports*, 10, 7289. <https://doi.org/10.1038/s41598-020-64477-4>
- Giovanardi, D., Lupini, C., Pesente, P., Rossi, G., Ortali, G., & Catelli, E. (2014). Longitudinal field studies of Avian metapneumovirus and Turkey hemorrhagic enteritis virus in turkeys suffering from colibacillosis associated mortality. *Veterinary Research Communications*, 38(2), 129–137. <https://doi.org/10.1007/s11259-014-9596-z>
- Giraud, P., Bennejean, G., Guittet, M., & Toquin, D. (1986). Turkey rhinotracheitis in France: preliminary investigations on a ciliostatic virus. *Veterinary Record*, 119(24), 606–607. <https://doi.org/10.1136/vr.119.24.606-a>
- Hu, H., Roth, J. P., Zsak, L., & Yu, Q. (2017). Engineered Newcastle disease virus expressing the F and G proteins of AMPV-C confers protection against challenges in turkeys. *Scientific Reports*, 7(1), 4025. <https://doi.org/10.1038/s41598-017-04267-7>
- Juhasz, K., & Easton, A. J. (1994). Extensive sequence variation in the attachment (G) protein gene of avian pneumovirus: Evidence for two distinct subgroups. *Journal of General Virology*, 75(11), 2873–2880. <https://doi.org/10.1099/0022-1317-75-11-2873>

407 Kapczynski, D. R. (2004). Development of a Virosome Vaccine Against Avian Metapneumovirus
 408 Subtype C for Protection in Turkeys. *Avian Diseases*, 48(2), 332–343.
 409 <https://doi.org/10.1637/7115>

410 Kapczynski, D. R., & Sellers, H. S. (2003). Immunization of Turkeys with a DNA Vaccine
 411 Expressing Either the F or N Gene of Avian Metapneumovirus. *Avian Diseases*, 47(4),
 412 1376–1383. <https://doi.org/10.1637/7033>

413 Kumar, S., Stecher, G., Li, M., Knyaz, C., & Tamura, K. (2018). MEGA X: Molecular
 414 evolutionary genetics analysis across computing platforms. *Molecular Biology and*
 415 *Evolution*, 35(6), 1547–1549. <https://doi.org/10.1093/molbev/msy096>

416 Laconi, A., Clubbe, J., Falchieri, M., Lupini, C., Cecchinato, M., Catelli, E., ... Naylor, C. J.
 417 (2016). A comparison of AMPV subtypes A and B full genomes, gene transcripts and
 418 proteins led to reverse-genetics systems rescuing both subtypes. *Journal of General*
 419 *Virology*, 97(6), 1324–1332. <https://doi.org/10.1099/jgv.0.000450>

420 Listorti, V., Lupini, C., Cecchinato, M., Pesente, P., Rossi, G., Giovanardi, D., ... Catelli, E.
 421 (2014). Rapid detection of subtype B avian metapneumoviruses using RT-PCR restriction
 422 endonuclease digestion indicates field circulation of vaccine-derived viruses in older
 423 turkeys. *Avian Pathology*, 43(1), 51–56. <https://doi.org/10.1080/03079457.2013.866212>

424 Lupini, C., Cecchinato, M., Ricchizzi, E., Naylor, C. J., & Catelli, E. (2011). A turkey
 425 rhinotracheitis outbreak caused by the environmental spread of a vaccine-derived avian
 426 metapneumovirus. *Avian Pathology*, 40(5), 525–530.
 427 <https://doi.org/10.1080/03079457.2011.607428>

428 McDougall, J. S., & Cook, J. K. (1986). Turkey rhinotracheitis: preliminary investigations. *The*
 429 *Veterinary Record*, 118(8), 206–207. <https://doi.org/10.1136/vr.118.8.206>

Naylor, C. J., Brown, P. A., Edworthy, N., Ling, R., Jones, R. C., Savage, C. E., & Easton, A. J. (2004). Development of a reverse-genetics system for Avian pneumovirus demonstrates that the small hydrophobic (SH) and attachment (G) genes are not essential for virus viability. *Journal of General Virology*, 85(Pt 11), 3219–3227. <https://doi.org/10.1099/vir.0.80229-0>

Naylor, C. J., Ling, R., Edworthy, N., Savage, C. E., & Easton, A. J. (2007). Avian metapneumovirus SH gene end and G protein mutations influence the level of protection of live-vaccine candidates. *Journal of General Virology*, 88(Pt 6), 1767–1775. <https://doi.org/10.1099/vir.0.82755-0>

Naylor, C. J., Lupini, C., & Brown, P. A. (2010). Charged amino acids in the AMPV fusion protein have more influence on induced protection than deletion of the SH or G genes. *Vaccine*, 28(41), 6800–6807. <https://doi.org/10.1016/j.vaccine.2010.07.015>

Qingzhong, Y., Barrett, T., Brown, T. D. K., Cook, J. K. A., Green, P., Skinner, M. A., & Cavanagh, D. (1994). Protection against turkey rhinotracheitis pneumovirus (TRTV) induced by a fowlpox virus recombinant expressing the TRTV fusion glycoprotein (F). *Vaccine*, 12(6), 569–573. [https://doi.org/10.1016/0264-410X\(94\)90319-0](https://doi.org/10.1016/0264-410X(94)90319-0)

Tarpey, I., Huggins, M. B., Davis, P. J., Shilleto, R., Orbell, S. J., & Cook, J. K. A. (2001). Cloning, expression and immunogenicity of the avian pneumovirus (Colorado isolate) F protein. *Avian Pathology*, 30(5), 471–474. <https://doi.org/10.1080/03079450120078653>

Toquin, D., Bayon-Auboyer, M.H., Eterradosi, N. & Jestin, V. (1999). Isolation of a pneumovirus from a Muscovy duck. *The Veterinary Record*, 23, 680.

Toquin, D., Guionie, O., Jestin, V., Zwingelstein, F., Allee, C., & Eterradosi, N. (2006). European and American subgroup C isolates of avian metapneumovirus belong to different genetic lineages. *Virus Genes*, 32(1), 97–103. <https://doi.org/10.1007/s11262-005-5850-3>

- Tucciarone, C.M., Andreopoulou, M., Franzo, G., Prentza, Z., Chaligiannis, I., & Cecchinato, M. (2017). First Identification and Molecular Characterization of Avian metapneumovirus Subtype B from Chickens in Greece. *Avian Diseases*, 61(3), 409–413. <https://doi.org/10.1637/11631-032017-CaseR>
- Tucciarone, C. M., Franzo, G., Lupini, C., Alejo, C. T., Listorti, V., Mescolini, G., ... Cecchinato, M. (2018). Avian Metapneumovirus circulation in Italian broiler farms. *Poultry Science*, 97(2), 503–509. <https://doi.org/10.3382/ps/pex350>
- Yu, Q., Estevez, C., Song, M., Kapczynski, D., & Zsak, L. (2010). Generation and biological assessment of recombinant avian metapneumovirus subgroup C (aMPV-C) viruses containing different length of the G gene. *Virus Research*, 147(2), 182–188. <https://doi.org/10.1016/j.virusres.2009.10.021>
- Yu, Q., Roth, J. P., Hu, H., Estevez, C. N., Zhao, W., & Zsak, L. (2013). Protection by Recombinant Newcastle Disease Viruses (NDV) Expressing the Glycoprotein (G) of Avian Metapneumovirus (aMPV) Subtype A or B against Challenge with Virulent NDV and aMPV. *World Journal of Vaccines*, 03(04), 130–139. <https://doi.org/10.4236/wjv.2013.34018>

470 **Figure legends**

471 **Figure 1.** Phylogenetic tree based on G gene nucleotide sequences of 60 selected European aMPV-
472 B strains detected in the present study (identical sequences within a country were removed),
473 including the most commonly used subtype B vaccines and European aMPV-B strains previously
474 obtained or retrieved from GenBank. The evolutionary history was inferred using the Neighbor-
475 Joining method in MEGA X. All nucleotide positions containing gaps and missing data were
476 eliminated, there were a total of 246 positions in the final dataset. The two subtrees containing
477 vaccines and vaccine-derived strains have been compressed. Field aMPV-B strains clusters are
478 included in square brackets and coloured by country of origin. Only bootstrap values ≥ 70 are
479 reported.

480 **Figure 2.** Phylogenetic tree based on G gene nucleotide sequences of selected European and non-
481 European aMPV-B strains, previously obtained or retrieved from GenBank. The evolutionary
482 history was inferred using the Neighbor-Joining method in MEGA X. All nucleotide positions
483 containing gaps and missing data were eliminated, there were a total of 171 positions in the final
484 dataset. The two subtrees containing vaccines and vaccine-derived strains have been compressed.
485 Field aMPV-B strains clusters are coloured by country of origin. The country of origin was added in
486 brackets next to the strain name when it was not clearly specified in the very name. Only bootstrap
487 values ≥ 70 are reported.

488

489 **Supplementary figure legends**

490 **Figure S1.** Phylogenetic tree based on G gene nucleotide sequences of 27 French aMPV-B strains
491 detected in the present study, including the most commonly used subtype B vaccines (marked with
492 a green triangle) and 2 French aMPV-B strains previously obtained. The evolutionary history was
493 inferred using the Neighbor-Joining method in MEGA X. All nucleotide positions containing gaps

494 and missing data were eliminated, there were a total of 253 positions in the final dataset. Field
495 aMPV-B strains clusters are included in square brackets. Only bootstrap values ≥ 70 are reported.

496

497 **Figure S2.** Phylogenetic tree based on G gene nucleotide sequences of 18 Italian aMPV-B strains
498 detected in the present study, including the most commonly used subtype B vaccines (marked with
499 a green triangle) and 42 Italian aMPV -B strains, previously obtained or retrieved from GenBank.
500 The evolutionary history was inferred using the Neighbor-Joining method in MEGA X. All
501 nucleotide positions containing gaps and missing data were eliminated, there were a total of 252
502 positions in the final dataset. Field aMPV-B strains clusters are included in square brackets. Only
503 bootstrap values ≥ 70 are reported.

504

505 **Figure S3.** Phylogenetic tree based on G gene nucleotide sequences of 27 Romanian aMPV-B
506 strains detected in the present study and the most commonly used subtype B vaccines (marked with
507 a green triangle). The evolutionary history was inferred using the Neighbor-Joining method in
508 MEGA X. All nucleotide positions containing gaps and missing data were eliminated, there were a
509 total of 282 positions in the final dataset. Field aMPV-B strains clusters are included in square
510 brackets. Only bootstrap values ≥ 70 are reported.

511

512 **Figure S4.** Phylogenetic tree based on G gene nucleotide sequences of 36 Spanish aMPV-B strains
513 detected in the present study and the most commonly used subtype B vaccines (marked with a green
514 triangle). The evolutionary history was inferred using the Neighbor-Joining method in MEGA X.
515 All nucleotide positions containing gaps and missing data were eliminated, there were a total of 303
516 positions in the final dataset. Field aMPV-B strains clusters are included in square brackets. Only
517 bootstrap values ≥ 70 are reported.

518

519 **Figure S5.** Phylogenetic tree based on G gene nucleotide sequences of 8 British aMPV-B strains
520 detected in the present study, including the most commonly used subtype B vaccines (marked with
521 a green triangle) and 10 British aMPV -B strains, previously obtained or retrieved from GenBank.
522 The evolutionary history was inferred using the Neighbor-Joining method in MEGA X. All
523 nucleotide positions containing gaps and missing data were eliminated, there were a total of 302
524 positions in the final dataset. Only bootstrap values ≥ 70 are reported.