

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Causes of stiffness after total knee arthroplasty: a systematic review

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Zaffagnini, S., Di Paolo, S., Meena, A., Alesi, D., Zinno, R., Barone, G., et al. (2021). Causes of stiffness after total knee arthroplasty: a systematic review. *INTERNATIONAL ORTHOPAEDICS*, 45(8), 1983-1999 [10.1007/s00264-021-05023-3].

Availability:

This version is available at: <https://hdl.handle.net/11585/818999> since: 2026-01-23

Published:

DOI: <http://doi.org/10.1007/s00264-021-05023-3>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

Causes Of Stiffness After Total Knee Arthroplasty: A Systematic Review

Stefano Zaffagnini, M.D., Full Professor ^{1, 2}	stefano.zaffagnini@unibo.it
Stefano Di Paolo, Eng ²	stefano.dipaolo@ior.it
Amit Meena, MBBS, MS, DNB ³	dr.meenaamit1624@gmail.com
Domenico Alesi, M.D. ¹	domenicoalesi@ymail.com
Raffaele Zinno, MSc ²	raffaele.zinno2@unibo.com
Giuseppe Barone, MSc ²	giuseppe.barone8@unibo.it
Nicola Pizza, M.D. ¹	nicola.pizza@ior.it
Laura Bragonzoni, Ph.D., Senior Assistant Professor ²	laura.bragonzoni4@unibo.it

¹ 2nd Orthopaedic and Traumatologic Clinic, IRCCS Istituto Ortopedico Rizzoli, Via G.B. Pupilli 1, 40136, Bologna, Italy

² University of Bologna, Via Zamboni, 33, 40126 Bologna, Italy

³ Central Institute of Orthopaedics, Vardhman Mahavir Medical College & Safdarjung Hospital, 110029, New Delhi, India

Corresponding Author: Domenico Alesi, M.D.

IRCCS Istituto Ortopedico Rizzoli, via G.B Pupilli 1, 40136, Bologna, Italy

phone: +39 3358071822 - mail: domenicoalesi@ymail.com

Declaration

Ethics Approval: Not applicable

Consent to Participate: Not applicable

Consent to Publish: Not applicable

Authors Contribution: S.Z and L.B conceived of the presented idea and were in charge of overall direction and planning, S.DP, A.M, D.A and N.P wrote the manuscript with input from all authors, R.Z and G.B did the bibliographical research and drew up tables and figure.

Funding: This research was supported by Invibio Knees Limited: Resolution n.194 of June 12th 2020.

Conflicts of interest/Competing interests: Authors declare that they have no conflict of interest.

Availability of data and material: Not applicable

Causes Of Stiffness After Total Knee Arthroplasty: A Systematic Review

Abstract

Background: Knee stiffness after Total Knee Arthroplasty (TKA) often leads to pain and discomfort, failing to meet patients' expectations on the surgical procedure. Despite the growing debate on the topic, a comprehensive literature analysis of stiffness causes has never been conducted. Thus, the purpose of the present study was to systematically review the literature regarding the main causes of stiffness after TKA.

Methods: Pubmed Central, Scopus, and EMBASE databases were systematically reviewed according to the Preferred Reporting Items for Systematic Reviews and Meta-analysis (PRISMA) guidelines for studies on stiffness and pain or discomfort after TKA through November 2020. Overall, 25 articles matched the selection criteria and were included in the study. Clinical relevance and strength of evidence of the included studies were graded using the risk of bias and the methodological index for non-randomized studies quality assessment tools.

Results: The main causes of pain and discomfort due to stiffness were surgery-related issues, i.e., component malpositioning and over-voluming, implant loosening, psychological distress, obesity, which could be considered "modifiable" factors, and expression of profibrotic markers, high material hypersensitivity-related cytokines level, male gender, previous contralateral TKA and high preoperative pain, which could be considered "non-modifiable" factors.

Conclusion: The use of alternative technologies such as surgical robots, anatomy-based devices, and more inert and less stiff component materials could help reducing stiffness caused by both modifiable and even some non-modifiable factors. Furthermore, early diagnostic detection of stiffness onset could consistently support surgeons in patient-specific decision making.

25

26 **Keywords:** Pain; Stiffness; Total Knee Arthroplasty

27

28 **List of Abbreviations**

29 Magnetic resonance images (MRI)

30 Knee Society Score (KSS)

31 Pain Catastrophizing Scale Score (PCS)

32 Oxford Knee Score (OKS)

33 Infrapatellar fat pad (FP)

34 Synovial membrane (SM)

35 Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC)

36 Quality of Well-Being scale (QWB-7)

37 Short Form (SF) 12 scores

38 The Knee injury and Osteoarthritis Outcome Score (KOOS)

39 External Rotation (ER)

40 Internal Rotation (IR)

41 Normal Rotation (NR)

42 **Introduction**

43 Total Knee Arthroplasty (TKA) is a well-established treatment for end-stage osteoarthritis (OA) that
44 effectively reduces pain and improves knee functionality [10]. The primary patient's expectations are
45 usually related to pain and joint mobility improvement [36].

46 However, about 20% of patients remain "unsatisfied" by the procedure [6]. The main complaints are
47 discomfort or pain referred to the operated joint. A heterogeneous list of possible causes has been
48 proposed. Excluding the most frequent and evident causes, i.e., aseptic or septic loosening and
49 implant failure [16, 17], there are several other causes with less clear categorization and treatment
50 path.

51 Knee stiffness consists of a decreased range of motion after TKA and has been investigated both as
52 a cause of and as a complication due to pain and discomfort [4, 9, 33].

53 Although there is a growing interest in this topic, the literature is fragmented: it is not clear whether
54 the reduced knee range of motion could be defined as a flexion contracture, a reduced flexion, or
55 both, and no consensus about pathogenesis has been reached [31]. Moreover, no comprehensive
56 analysis of stiffness causes after TKA has been conducted: previous literature reviews were mainly
57 focused on the treatment of stiffness to increase the knee range of motion.

58 Hence, the purpose of this study was to systematically review the main causes of stiffness after TKA.

59

60 **Material and method**

61 A literature review was performed using a strategy search design to collect articles regarding stiffness
62 and pain or discomfort after TKA. Inclusion criteria adopted were: (1) studies of patients who
63 underwent primary TKA; (2) articles published in the last 10 years; (3) articles published in English;
64 (4) involving human species. Review articles were excluded from the search.

65 *Article Selection*

66 The search was conducted according to the Preferred Reporting Items for Systematic Reviews and
67 Meta-analysis (PRISMA) Guidelines [24] by two independent reviewers on PubMed Central, Scopus
68 and EMBASE, for studies available until November 2020. The keywords used for initial screening
69 were ((TKA[Title/Abstract]) OR (TKR[Title/Abstract]) OR (Total Knee
70 Replacement[Title/Abstract]) OR (Total Knee Arthroplasty[Title/Abstract])) AND
71 (stiff*[Title/Abstract]) AND ((Pain[Title/Abstract]) OR (discomfort*[Title/Abstract])).

72 Two authors (A.M., N.P.) independently reviewed each article's title and abstract from the literature
73 search. The assessors were not blinded to the authors of the publications. The full text was obtained
74 and evaluated when eligibility could not be assessed from the first screening. Any disagreements
75 were resolved via a consensus discussion between the reviewers, and a third reviewer (S.Z.) was
76 consulted if the disagreement could not be resolved.

77 The Cochrane risk of bias (ROB) tool was used for the quality assessment of randomized controlled
78 trials, while the methodological index for non-randomized studies (MINORS) was used for quality
79 assessment of observational studies. Two authors (G.B., R.Z.) independently performed the quality
80 assessment for each article. A third author (L.B.) was consulted in case of disagreement.

81 **Results**

82 After the exclusion process detailed above (Figure 1), the strategy search generated a systematic
83 review of 25 articles. Among these articles, 7 were related to surgical issues. Six of them identified
84 component malpositioning as a cause for stiffness [1, 2, 16, 28, 34, 37], reporting extension deficit,
85 and poor subjective and objective clinical outcomes, such as the Knee Society Score (KSS).
86 Component malpositioning was mainly identified as internal rotation of both the femoral and tibial
87 components and sagittal plane malalignment. One article [23] detected component over-voluming in
88 the 24% of retrospectively evaluated knees and identified it as a main cause of stiffness.

89 Three articles were related to stiffness due to TKA components loosening [13, 15, 32], identified as
90 a major cause of pain and discomfort, and related to higher odds of revision rate.

91 One article [22] found that the genetic expression of profibrotic markers was a factor influencing the
92 limited range of motion and subsequent patients' pain and discomfort, evaluated through the Oxford
93 Knee Scores (OKS) and Pain Catastrophizing Scores (PCS).

94 Three articles were related to immunological and allergy [12, 26, 38] issues. The authors detected
95 high levels of inflammatory cytokines in painful knees presenting stiffness [26, 38], as well as lower
96 postoperative outcomes in poly-allergic patients [12].

97 Three articles were related to psychological issues [3, 19, 30]. Patients with anxiety, depressive
98 symptoms, functional limitation, and severe preoperative pain were more likely to show stiffness and
99 become more dissatisfied after TKA than other patients.

100 Eight studies focused on demographic and other factors [5, 7, 14, 18, 20, 21, 27, 29]. Young patients
101 reported residual symptoms and limitations after modern TKAs, as well as male patients and patients
102 with lung disease, diabetes, and preoperative stiffness. TKA performed after a knee injury (fracture
103 or ligament injury) or a previous osteotomy and preoperative radiologic minimal joint space width
104 were related to poorer outcomes after TKA.

105 The characteristics and the results of the included articles are reported in Table 1.

106

107 *Methodological quality assessment*

108 No randomized controlled trial study was retrieved. Therefore, the MINORS tool was used to
109 perform the quality assessment of all the included articles. Of the 25 articles included in this analysis,
110 8 were comparative studies and 17 were non-comparative studies. The average score for the
111 comparative studies was 18.4 (range 14 – 22), while the average score for the non-comparative
112 studies was 11.0 (range 5 – 16) (Appendix A) [35].

113

114 **Discussion**

115 The purpose of the present study was to review the main causes of stiffness after TKA. Multiple

causes with different incidence emerged from the literature review (Table 2). A distinction could be made between “modifiable” factors, i.e., surgical issues, loosening, psychological distress, obesity, and “non-modifiable” factors, i.e., male gender, young age, material hypersensitivity, high preoperative pain, and gene expression of profibrotic markers .

Among the modifiable factors, malposition and over-voluming were revealed to be critically leading to stiffness. Indeed, the prosthetic components' positioning is fundamental to allow correct kinematics and guarantee the long-term survival of modern TKAs. A mistake in component positioning can trigger chronic inflammation, leading to pain and functional limitation until implant failure [1, 2]. Thus, the prosthetic components' appropriate positioning is mandatory to avoid pain, discomfort, and, eventually, reduce the revision rate after TKA. To face this problem, in the last years, the use of computer-assisted surgery (CAS) and patient-specific instrumentation (PSI) has been developed to improving implant positioning. Both these technologies showed promising results concerning correct femoral positioning, perioperative blood loss, and operative time. However, the clinical relevance of such improvement remains uncertain [11].

The aseptic loosening was detected as a major cause for patients' dissatisfaction leading to implant revision. Several aspects can contribute to this fatal condition, e.g., cementation technique, bone quality, component design. In the last years, a significant effort has been put into improving bone-implant fixation, e.g., the tibial baseplate design and the cementation process. Moreover, the tribology of the material used in modern TKA has been questioned: pre-clinical studies have been conducted on alternative materials (ceramic and polymers) to demonstrate the influence of material stiffness on implant failure. In particular, a lower stress-shielding phenomenon was hypothesized using a less stiff material [25]. If confirmed in clinical studies, the use of less stiff materials could represent a promising alternative to standard TKA implant components.

The patient's psychological conditions have been detected as decisive factors influencing pain and discomfort after TKA. Since the rehabilitation phase needs active patient engagement to reach satisfying results, a positive attitude is highly required. During the patient's selection for this

142 procedure, the surgeon should consider psychological aspects, and if any abnormalities are detected,
143 the patient should be aware of the higher risk of poorer postoperative outcomes. Education programs
144 have been introduced to prepare TKA patients from both physical and psychological sides. Recent
145 studies [8] demonstrated the positive effects of these programs on postoperative patients' conditions
146 during the rehabilitation phase.

147 Among the non-modifiable factors, the expression of profibrotic markers correlated with the onset of
148 postsurgical stiffness, and high preoperative inflammatory cytokines level correlated with poorer
149 postoperative outcomes. Even though these factors are a-specific, a substantial contribution to
150 reducing the risk of inflammatory marker expression and the onset of stiffness in predisposed subjects
151 could come from the biocompatibility of TKA components material. The use of more inert materials
152 than classical cobalt-chrome alloy could indeed limit the alteration of intraarticular homeostasis and
153 the inflammatory response.

154 Two main clinical implications could be deduced from the present review. First, the use of alternative
155 technological solutions, e.g., surgical robots, PSI, and less stiff and more inert materials, could be
156 crucial to act both on some modifiable (surgical issues and loosening) and non-modifiable factors
157 (expression of pro-fibrotic markers or allergy). Such tools might induce a further reduction in the
158 percentage of patients experiencing knee stiffness and subsequent pain and discomfort. However,
159 clinical evidence is still missing and needs to be addressed.

160 Second, some of the factors leading to stiffness are difficult to be detected in routine clinical practice.
161 Indeed, even this condition can be quite easily detected in the postoperative outpatient setting, a
162 comprehensive and well-structured methodology to preventively identify the cohort of patients at risk
163 to develop post-TKA stiffness is still lacking. Such clinical diagnosis could consistently support the
164 surgeons in decision making on the type of implant and the surgical procedures to adopt per each
165 patient.

166 The present review has some limitations. First, the heterogeneous approaches adopted in evaluating
167 stiffness for the different factors did not allow a meta-analysis of the retrieved data. Second, the

168 possibility of falling into article selection bias could have affected the validity and generalizability of
169 this study. However, the authors rigorously followed the PRISMA guidelines [24] and decided to
170 include both prospective and retrospective studies to provide the most comprehensive view on such
171 a specific topic. Furthermore, the MINORS tool was used to assess the quality of the included articles.
172 Although the literature is well-populated with studies regarding the postoperative treatment of
173 stiffness after TKA, the present review confirmed a lack of consistent investigations on the causes of
174 this clinical expression.

175

176 **Conclusion**

177 Stiffness after TKA might occur for different and, sometimes, multifactorial causes. Possible
178 solutions should be looked for into alternative aspects, e.g., different component material properties
179 and anatomy-based TKA designs. Moreover, further studies are needed to investigate the
180 development of diagnostic strategies to detect stiffness onset as early as possible.

181

182 **References**

- 183 1. Abdelnasser MK, Adi MM, Elnaggar AA, Tarabichi S (2020) Internal rotation of the tibial
184 component in total knee arthroplasty can lead to extension deficit. *Knee Surg Sports Traumatol*
185 *Arthrosc* 28:2948–2952
- 186 2. Bédard M, Vince KG, Redfern J, Collen SR (2011) Internal rotation of the tibial component
187 is frequent in stiff total knee arthroplasty. *Clin Orthop Relat Res* 469:2346–2355
- 188 3. Belford K, Gallagher N, Dempster M, Wolfenden M, Hill J, Blaney J, O’Brien S, Smit A-M,
189 Botha P, Molloy D, Beverland D (2020) Psychosocial predictors of outcomes up to one year
190 following total knee arthroplasty. *Knee* 27:1028–1034
- 191 4. Cheuy VA, Foran JRH, Paxton RJ, Bade MJ, Zeni JA, Stevens-Lapsley JE (2017)
192 Arthrofibrosis Associated With Total Knee Arthroplasty. *J Arthroplasty* 32:2604–2611
- 193 5. Clement ND, Bardgett M, Weir D, Holland J, Deehan DJ (2019) Increased symptoms of
194 stiffness 1 year after total knee arthroplasty are associated with a worse functional outcome and
195 lower rate of patient satisfaction. *Knee Surg Sports Traumatol Arthrosc* 27:1196–1203
- 196 6. Clement ND, Bardgett M, Weir D, Holland J, Gerrand C, Deehan DJ (2018) Three groups of
197 dissatisfied patients exist after total knee arthroplasty: early, persistent, and late. *Bone Joint J* 100-
198 B:161–169
- 199 7. Clement ND, Merrie KL, Weir DJ, Holland JP, Deehan DJ (2019) Asynchronous Bilateral
200 Total Knee Arthroplasty: Predictors of the Functional Outcome and Patient Satisfaction for the
201 Second Knee Replacement. *J Arthroplasty* 34:2950–2956
- 202 8. Ditton E, Johnson S, Hodyl N, Flynn T, Pollack M, Ribbons K, Walker FR, Nilsson M
203 (2020) Improving Patient Outcomes Following Total Knee Arthroplasty: Identifying Rehabilitation
204 Pathways Based on Modifiable Psychological Risk and Resilience Factors. *Front Psychol* 11:1061
- 205 9. Donaldson JR, Tudor F, Gollish J (2016) Revision surgery for the stiff total knee
206 arthroplasty. *Bone Joint J* 98-B:622–627
- 207 10. Evans JT, Walker RW, Evans JP, Blom AW, Sayers A, Whitehouse MR (2019) How long
208 does a knee replacement last? A systematic review and meta-analysis of case series and national
209 registry reports with more than 15 years of follow-up. *Lancet* 393:655–663
- 210 11. Gong S, Xu W, Wang R, Wang Z, Wang B, Han L, Chen G (2019) Patient-specific
211 instrumentation improved axial alignment of the femoral component, operative time and
212 perioperative blood loss after total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc*
213 27:1083–1095
- 214 12. Graves CM, Otero JE, Gao Y, Goetz DD, Willenborg MD, Callaghan JJ (2014) Patient
215 reported allergies are a risk factor for poor outcomes in total hip and knee arthroplasty. *J*
216 *Arthroplasty* 29:147–149
- 217 13. Gu A, Malahias M-A, Cohen JS, Richardson SS, Stake S, Blevins JL, Sculco PK (2020)
218 Prior Knee Arthroscopy Is Associated With Increased Risk of Revision After Total Knee
219 Arthroplasty. *J Arthroplasty* 35:100–104
- 220 14. Gungor S, Fields K, Aiyer R, Valle AGD, Su EP (2019) Incidence and risk factors for
221 development of persistent postsurgical pain following total knee arthroplasty: A retrospective
222 cohort study. *Medicine (Baltimore)* 98:e16450
- 223 15. Hagman DS, Granade CM, Smith LS, Yakkanti MR, Malkani AL (2020) Results of
224 Cemented Posterior-Stabilized Total Knee Arthroplasty in Obese Patients With an Average 10-Year
225 Follow-Up. *J Arthroplasty* 35:2097–2100
- 226 16. Hirschmann MT, Amsler F, Rasch H (2015) Clinical value of SPECT/CT in the painful total
227 knee arthroplasty (TKA): a prospective study in a consecutive series of 100 TKA. *Eur J Nucl Med*
228 *Mol Imaging* 42:1869–1882
- 229 17. Hirschmann MT, Konala P, Iranpour F, Kerner A, Rasch H, Friederich NF (2011) Clinical
230 value of SPECT/CT for evaluation of patients with painful knees after total knee arthroplasty- a
231 new dimension of diagnostics? *BMC Musculoskelet Disord* 12:36

18. Järvenpää J, Kettunen J, Soininvaara T, Miettinen H, Kröger H (2012) Obesity has a negative impact on clinical outcome after total knee arthroplasty. *Scand J Surg* 101:198–203
19. Lavernia CJ, Villa JM, Iacobelli DA (2015) What is the role of mental health in primary total knee arthroplasty? *Clin Orthop Relat Res* 473:159–163
20. Liebensteiner M, Wurm A, Gamper D, Oberaigner W, Dammerer D, Krismer M (2019) Patient satisfaction after total knee arthroplasty is better in patients with pre-operative complete joint space collapse. *Int Orthop* 43:1841–1847
21. Liebs TR, Herzberg W, Roth-Kroeger AM, Rütther W, Hassenpflug J (2011) Women recover faster than men after standard knee arthroplasty. *Clin Orthop Relat Res* 469:2855–2865
22. Mann CJ, Bradley AM, Jones ER, McNamara IR, Smith TO, Riley GP (2019) Stiffness post-total knee replacement: A proof of principle study investigating the effect of gene expression analysis of markers of fibrosis. *Knee* 26:914–922
23. Marmor S, Renault E, Valluy J, Saffarini M (2019) Over-voluming predicted by pre-operative planning in 24% of total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc* 27:1544–1551
24. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group (2009) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol* 62:1006–1012
25. Nakahara I, Takao M, Bandoh S, Bertollo N, Walsh WR, Sugano N (2012) Novel surface modifications of carbon fiber-reinforced polyetheretherketone hip stem in an ovine model. *Artif Organs* 36:62–70
26. Paish HL, Baldock TE, Gillespie CS, Del Carpio Pons A, Mann DA, Deehan DJ, Borthwick LA, Kalson NS (2019) Chronic, Active Inflammation in Patients With Failed Total Knee Replacements Undergoing Revision Surgery. *J Orthop Res* 37:2316–2324
27. Parvizi J, Nunley RM, Berend KR, Lombardi AV, Ruh EL, Clohisy JC, Hamilton WG, Della Valle CJ, Barrack RL (2014) High level of residual symptoms in young patients after total knee arthroplasty. *Clin Orthop Relat Res* 472:133–137
28. Planckaert C, Larose G, Ranger P, Lacelle M, Fuentes A, Hagemeister N (2018) Total knee arthroplasty with unexplained pain: new insights from kinematics. *Arch Orthop Trauma Surg* 138:553–561
29. Putman S, Argenson J-N, Bonnevalle P, Ehlinger M, Vie P, Leclercq S, Bizot P, Lustig S, Parratte S, Ramdane N, Colmar M, Société française de chirurgie orthopédique et traumatologie (SOFOT) (2018) Ten-year survival and complications of total knee arthroplasty for osteoarthritis secondary to trauma or surgery: A French multicentre study of 263 patients. *Orthop Traumatol Surg Res* 104:161–164
30. Razmjou H, Boljanovic D, Wright S, Murnaghan J, Holtby R (2015) Association between Neuropathic Pain and Reported Disability after Total Knee Arthroplasty. *Physiother Can* 67:311–318
31. Rodríguez-Merchán EC (2019) The stiff total knee arthroplasty: causes, treatment modalities and results. *EFORT Open Rev* 4:602–610
32. Sadoghi P, Leithner A, Weber P, Friesenbichler J, Gruber G, Kastner N, Pohlmann K, Jansson V, Wegener B (2011) Radiolucent lines in low-contact-stress mobile-bearing total knee arthroplasty: a blinded and matched case control study. *BMC Musculoskelet Disord* 12:142
33. Sanders TL, Kremers HM, Bryan AJ, Kremers WK, Stuart MJ, Krych AJ (2017) Procedural intervention for arthrofibrosis after ACL reconstruction: trends over two decades. *Knee Surg Sports Traumatol Arthrosc* 25:532–537
34. Scott CEH, Clement ND, Yapp LZ, MacDonald DJ, Patton JT, Burnett R (2019) Association Between Femoral Component Sagittal Positioning and Anterior Knee Pain in Total Knee Arthroplasty: A 10-Year Case-Control Follow-up Study of a Cruciate-Retaining Single-Radius Design. *J Bone Joint Surg Am* 101:1575–1585
35. Slim K, Nini E, Forestier D, Kwiatkowski F, Panis Y, Chipponi J (2003) Methodological

283 index for non-randomized studies (minors): development and validation of a new instrument. ANZ
 284 J Surg 73:712–716
 285 36. Tekin B, Unver B, Karatosun V (2012) Expectations in patients with total knee arthroplasty.
 286 Acta Orthop Traumatol Turc 46:174–180
 287 37. White PB, Ranawat AS (2016) Patient-Specific Total Knees Demonstrate a Higher
 288 Manipulation Rate Compared to “Off-the-Shelf Implants.” J Arthroplasty 31:107–111
 289 38. Yang S, Dipane M, Lu CH, Schmalzried TP, McPherson EJ (2019) Lymphocyte
 290 Transformation Testing (LTT) in Cases of Pain Following Total Knee Arthroplasty: Little
 291 Relationship to Histopathologic Findings and Revision Outcomes. J Bone Joint Surg Am 101:257–
 292 264
 293
 294

295 **Figure Legend**

296

297 Fig.1: Flow diagram, describing the number of studies identified, included and excluded with
298 relative reasons.

299

300

301 **Appendix A:** Methodological quality assessment. Individual MINORS criteria [35].

302

Study	A clear stated aim	Inclusion of consecutive patients	Prospective collection of data	Endpoints appropriate to the aim of the study	Unbiased assessment of the study endpoint	Follow-up period appropriate to the aim of the study	Loss to follow up less than 5%	Prospective calculation of the study size	An adequate control group	Contemporary groups	Baseline equivalence of groups	Adequate statistical analyses	Total
Abdelnasser et al. [1]	2	1	2	1	1	0	0	2	N/A	N/A	N/A	N/A	9
Bedard et al. [2]	2	2	2	1	1	1	0	2	N/A	N/A	N/A	N/A	11
Belford et al. [3]	2	2	2	1	1	2	1	2	N/A	N/A	N/A	N/A	13
Clement, Bardgett et al. [5]	2	2	2	1	1	2	1	2	2	1	2	2	20
Clement, Merrie et al. [7]	2	2	2	1	1	2	1	2	N/A	N/A	N/A	N/A	13
Graves et al. [12]	2	2	2	1	1	2	1	0	N/A	N/A	N/A	N/A	11
Gu et al. [13]	2	2	2	2	1	1	2	2	2	2	2	2	22
Gungor et al. [14]	2	2	2	2	2	2	2	2	N/A	N/A	N/A	N/A	16
Hagman et al. [15]	2	2	2	2	0	2	0	0	N/A	N/A	N/A	N/A	10
Hirschmann et al. [16]	1	1	2	1	0	0	0	0	N/A	N/A	N/A	N/A	5
Järvenpää et al. [18]	1	2	2	2	0	1	0	0	2	2	0	2	14
Lavernia et al. [19]	2	1	2	1	0	2	0	2	2	2	2	2	18
Liebensteiner et al. [20]	2	2	2	1	0	1	2	2	N/A	N/A	N/A	N/A	12
Liebs et al. [21]	2	2	2	2	0	2	1	2	N/A	N/A	N/A	N/A	13
Mann et al. [22]	1	2	2	1	2	1	0	2	N/A	N/A	N/A	N/A	11
Marmor et al. [23]	2	1	2	1	0	0	1	2	N/A	N/A	N/A	N/A	9
Paish et al. [26]	1	1	2	2	0	0	0	2	2	2	2	2	16
Parvazi et al. [27]	1	1	2	1	0	1	1	0	N/A	N/A	N/A	N/A	7
Planckaert et al. [28]	2	2	2	2	1	2	1	0	2	1	1	2	18
Putman et al. [29]	2	2	2	2	0	2	1	2	N/A	N/A	N/A	N/A	13
Razmjou et al. [30]	2	0	2	2	0	2	1	2	N/A	N/A	N/A	N/A	11
Sadoghi et al. [32]	2	2	2	2	2	2	2	0	2	2	2	2	22
Scott et al. [34]	2	1	2	2	0	2	1	0	2	2	1	2	17
White et al. [37]	2	2	2	2	0	2	2	0	N/A	N/A	N/A	N/A	12
Yang et al. [38]	2	1	2	2	0	2	2	0	N/A	N/A	N/A	N/A	11

303

Table 1: Characteristics and results of the included studies

Category	Authors	Year	Type of study	Level of evidence	Aim	Study subjects	Instruments	Results	Conclusions
Surgery-related issues	Abdelnasser et al.[10]	2020	Observational study	III	To investigate whether induced IR of the tibial component intraoperatively would change in vivo knee kinematics	31 TKA patients (14 M, 17 F) CR: 22; PS: 9	Radiographs	- Tibial NR: CR -1.6 ± 3.4; PS 1.6 ± 3.9 - Tibial IR: CR 3.7 ± 2.6; PS 8.8 ± 4.4	IR of the tibial component during TKA can lead to postoperative knee extension deficit, pain and limited motion.
	Bedard et al.[11]	2011	Observational study	IV	1. Incidence of IR of the femoral and tibial components in stiff TKAs; 2. If revision surgery that included correction of rotational positioning improved pain, ROM, and patellar tracking; 3. If revision altered nonrotational radiographic parameters	52 TKA patients 34 scanned Pre Revision; 18 scanned After Revision	- CT scans - KSS	- Femoral IR: Pre 3.1 N24; Post 2.1 (n=15) - Femoral ER: Pre 1.3 N10 ; Post 1.1 (n=3) - Tibial IR: Pre 13.7 N33 ; Post 3.3 (n=3) - Tibial ER: Pre 1.0 N1; Post 8.5 15 - KSS - Pain: Pre 10,3 ; Post 33,7 - Knee: Pre 41,6 ; Post 77,3 - Function: Pre 48 ; Post 65,7	The incidence of IR was 24 of 34 femoral and 33 of 34 tibial components; Revision arthroplasty improved Knee Society function, knee, and pain scores; Post-revision nonrotational parameters were unchanged.
	Hirschmann et al.[4]	2015	Observational study	IV	To evaluate the clinical value of SPECT/CT in patients with knee pain after primary TKA	23 TKA patients	Hybrid SPECT/CT	- SPECT/CT imaging changed the suspected diagnosis and the proposed treatment in 19/23 (83%) knees - Progression of patellofemoral OA (n = 11), loosening of the tibial (n = 3) and loosening of the femoral component (n = 2) were identified as the leading causes of pain after TKA	SPECT/CT was very helpful in establishing the diagnosis particularly in patients with patellofemoral problems and malpositioned or loose TKA.

	Planck et al.[12]	2018	Observational study	IV	To analyze knee 3D kinematics of TKA patients with unexplained anterior knee pain and compare them to those of an asymptomatic TKA group, and a healthy control group	62 knees: 24 asymptomatic, 21 painful, and 17 control	▪ CT scans ▪ Radiographs	▪ LEFS: 66.6 ± 8.2 Asympt; 43.6 ± 13.7 Pain; N/A Control ▪ KOOS - Pain 90.0 ± 11.2 Asympt; 59.5 ± 17.7 Pain; 93.1 ± 15.1 Control - Symptom 84.4 ± 16.0 Asympt; 62.1 ± 21.1 Pain; 91.7 ± 11.6 Control - AVQ 90.0 ± 10.3 Asympt; 60.9 ± 16.3 Pain; 94.0 ± 14.7 Control - Sport 55.0 ± 23.2 Asympt; 23.3 ± 19.1 Pain; 87.2 ± 22.3 Control - QoL 82.3 ± 21.1 Asympt; 43.5 ± 29.3 Pain; 86.7 ± 24.4 Control ▪ Component rotation - Tibial rotation: $7.8^\circ \pm 5.4^\circ$; $0.8^\circ \pm 6.8^\circ$ Pain - Femoral rotation: $-0.5^\circ \pm 2.3^\circ - 2.1^\circ \pm 2.6^\circ$ Pain - Combined rotation: $7.3^\circ \pm 6.1^\circ - 1.4^\circ \pm 7.0^\circ$ Pain	Painful TKA patients presented three well-known characteristics: a stiff knee gait, a valgus alignment when walking, and combined TKA components slightly internally rotated.
--	-------------------	------	---------------------	----	---	--	---	---	---

	Scott et al.[13]	2019	Case-Control Follow-up Study	IV	To investigate sagittal femoral component position as a predictor of anterior knee pain at long-term follow-up after cruciate-retaining single-radius TKA without routine patellofemoral resurfacing	162 TKA patients	▪ Radiographs ▪ SF-12 ▪ OKS ▪ PROMs ▪ HSS	▪ Femorotibial angle: 175.1° Pain; 177.9° No Pain ▪ Coronal plane - Medial proximal tibial angle: 89.7° Pain; 88.9° No Pain - Lateral distal femoral angle: 85.7° Pain; 85.7° No Pain ▪ Sagittal plane - Posterior tibial slope: 4.5° Pain; 5.3° No Pain - Femoral component flexion: 20.6 Pain; 1.4° No Pain - Posterior condylar offset ratio: 94.0% Pain; 97.3% No Pain - Anterior femoral offset ratio: 17.2% Pain; 13.3% No Pain ▪ PCS 35.5 (11.6) Pain; 43.4 (10.6) No Pain ▪ MCS 48.5 (9.4) Pain; 51.5 (9.7) No Pain ▪ OKS 29.6 (10.9) Pain; 40.1 (7.1) No Pain ▪ Dissatisfied 14 (19) Pain; 4 (4) No Pain	Sagittal plane positioning and alignment of the femoral component were associated with long-term anterior knee pain, with femoral component extension being a major risk factor
	White et al.[14]	2016	Prospective study	IV	To restore a patient's anatomical geometry of the knee	74 TKA patients Patient-Specific TKA: 21 (7 M, 14 F) PFC PS: 42 (14 M, 28 F) PFC NC-	▪ WOMAC	Pre-op ▪ Patient-Specific TKA: 11.5 Pain; 4.6 Stiffness; 35.3 Function; 3.5 Satisfaction ▪ PFC PS: 11.1 Pain; 5.1 Stiffness; 36.2 Function; 2.1 Satisfaction ▪ PFC NC-CR: 7.8 Pain; 3.4 Stiffness; 30.1 Function; 0.8 Satisfaction 2 Years post-op ▪ Patient-Specific TKA: 4.8 Pain; 3.0 Stiffness; 15.2 Function; 7.0 Satisfaction	Patient-specific TKAs were associated with higher manipulation rate and lower satisfaction scores

						CR: 11 (10 M. 1 F)		▪ PFC PS: 2.8 Pain; 2.2 Stiffness; 10.4 Function; 8.3 Satisfaction ▪ PFC NC-CR: 1.2 Pain; 1.3 Stiffness 5.2 Function; 8.9 Satisfaction	
	Marmor et al.[15]	2019	Observational study	IV	1) To investigate differences in volumes of resected bone compared to implanted components in TKA 2) To determine the incidence and factors related to ‘over-voluming’	100 TKA patients (45 M, 55 F)	▪ MRI	▪ HKA: $178.3^{\circ} \pm 4.1^{\circ}$ ▪ Mechanical lateral femoral angle: $88.2^{\circ} \pm 2.9^{\circ}$ ▪ Mechanical medial proximal tibial angle: $83.3^{\circ} \pm 3.5^{\circ}$ ▪ Over-voluming: 24% of the TKA patients	Over-voluming could be associated with prosthetic overhang or excessive tensions within the joint capsule, and contribute to unexplained pain and stiffness following TKA
Loosening	Gu et al.[16]	2020	Observational study	III	To determine the relationship between prior Knee arthroscopy (KA) within 2 years of TKA on revision rates after TKA	138,019 TKA patients 3357 previous KA vs 134,662 No KA	▪ Current Procedural Terminology ▪ International Classification of Diseases	PREVIOUS KA ▪ Frequency of revision TKA: 2.77% ▪ Frequencies of aseptic loosening: 1.25% ▪ MUA: 4.28% ▪ PJI: 2.65% ▪ Periprosthetic fractures 0.15% ▪ Postoperative stiffness: 17.78% NO KA ▪ Frequency of revision TKA: 1.79% ▪ Frequencies of aseptic loosening: 0.75% ▪ MUA: 3.27% ▪ PJI: 1.83% ▪ Periprosthetic fractures: 0.34% ▪ Postoperative stiffness: 11.53%	Knee arthroscopy before TKA increases the rates of revision, PJI, aseptic loosening, and stiffness

	Hagman et al.[17]	2020	Retrospective study	IV	To establish long-term survival and complication rates in obese patients undergoing cemented, posterior-stabilized TKA with an average 10-year follow-up.	181 TKA patients (body mass index >35)	▪ Radiographs ▪ Blood analysis	▪ Aseptic loosening - Total: 25 (14%) - Tibial component only: 11 (6%) - Tibial and femoral components: 13 (7%) ▪ Aseptic patellar component loosening 1 (1%) ▪ Polyethylene wear 9 (5%) ▪ Prosthetic joint infection 2 (1%) ▪ Stiffness/immobilization syndrome 1 (1%) ▪ Instability 1 (1%) ▪ Patellofemoral pain 1 (1%) ▪ Total failures leading to revision total knee arthroplasty 39 (22%)	Aseptic loosening is the leading cause of failure following TKA in obese; Survivorship with aseptic loosening as the endpoint was 91.2% at 10 years and 86.7% at 15 years.
	Sadoghi et al.[18]	2011	Retrospective study	IV	To test for a correlation between knee pain and tibial radiolucent lines	56 TKA patients	▪ WOMAC ▪ KSS ▪ Radiographs	▪ Pain: 28 patients; 28 No pain: ▪ Radiolucent lines: 27 Pain patients; 6 No pain patients	Radiolucency should still be suspected as the possible cause of pain
Gene expression	Mann et al.[19]	2019	Observational study	III	To establish proof of principle of a link between phenotypic expression and stiffness after TKR	100 TKA patients	▪ Radiographs ▪ PCS ▪ OKS	Preoperative ▪ Extension: $-6.47^{\circ} \pm 6.16^{\circ}$ Stiff; $-3.87^{\circ} \pm 4.98^{\circ}$ No Stiff ▪ Flexion: $95.00^{\circ} \pm 9.78^{\circ}$ Stiff; $95.53^{\circ} \pm 15.97^{\circ}$ No Stiff ▪ OKS: 17.73 ± 5.48 Stiff; 15.80 ± 7.88 No Stiff Postoperative ▪ Extension: $-10.07^{\circ} \pm 7.33^{\circ}$ Stiff; $-2.07^{\circ} \pm 2.37^{\circ}$ No Stiff ▪ Flexion: $78.00^{\circ} \pm 13.34^{\circ}$ Stiff; $105.67^{\circ} \pm 6.78^{\circ}$ No Stiff ▪ OKS: 23.46 ± 8.62 Stiff;	Joint stiffness is a multifactorial problem after TKR but biological factors may play a significant role in stiffness after TKR

								21.47 ± 7.18 No Stiff ▪ PCS - Total Score: 18.60 ± 10.52 Stiff; 19.00 ± 16.60 No Stiff - Rumination): 6.73 ± 5.27 Stiff; 6.33 ± 6.00 No Stiff - Magnification: 2.87 ± 1.41 Stiff; 4.00 ± 3.68 No Stiff - Helplessness: 9.00 ± 5.32 Stiff; 8.73 ± 7.57 No Stiff	
Immunology & Allergy	Graves et al.[20]	2014	Observational study	III	To compare patient reported outcomes following total joint arthroplasty in patients with and without multiple reported allergies	459 THA or TKA patients	▪ WOMAC ▪ SF-36	Preoperative ▪ PCS: 30.7 ± 0.4 (0-3 allergies); 29.3 ± 1.1 (4+ allergies) ▪ MCS: 49.4 ± 0.6 (0-3 allergies); 48.3 ± 1.5 (4+ allergies) ▪ WOMAC - Pain: 46.9 ± 1.1 (0-3 allergies); 43.4 ± 3.2 (4+ allergies) - Stiffness: 41.4 ± 1.1 (0-3 allergies); 35.8 ± 2.7 (4+ allergies) - Function: 45.5 ± 1.0 (0-3 allergies); 39.8 ± 2.4 (4+ allergies)	Patients with multiple reported allergies belong to a group with a statistically significant lower functional status after hip and knee arthroplasty per WOMAC and SF36
	Paish et al.[21]	2019	Retrospective cohort study	III	To investigate whether pro-inflammatory mediators are elevated in revision knee patients, indicating an active, ongoing inflammatory process that may contribute to pain	62 patients (29 Primary; 33 Revision) with TKA	▪ WOMAC ▪ SF ▪ FP ▪ SM	▪ WOMAC pain: 30 Primary TKA; 52 Revision TKA ▪ GM-CSF: 0.5 ± 0.2 Primary; 1.3 ± 0.3 Revision ▪ IL-5: 0.4 ± 0.1 Primary; 2.2 ± 0.8 Revision ▪ IL-8: 80 ± 31 Primary; 3,147 ± 1,552 Revision	A number of biologically active markers involved in inflammation and matrix deposition were significantly elevated many years

								▪ IL-10: 0.8 ± 0.5 Primary; 1.6 ± 0.4 Revision ▪ CCL2: 417 ± 38 Primary; 5,285 ± 1,004 Revision ▪ CCL3: 15.6 ± 3.3 Primary; 177 ± 51 Revision ▪ CCL4: 69.6 ± 8 Primary; 258 ± 54 Revision ▪ CCL13: 42.5 ± 20 Primary; 220 ± 76 Revision ▪ Flt--1: 97.6 ± 12 Primary; 1575 ± 612 Revision ▪ VEGF: 396.1 ± 46 Primary; 1724 ± 268 Revision ▪ VCAM--1: 105,613 ± 7,290 Primary; 134,770 ± 7,434 Revision ▪ ICAM--1: 862,890 ± 12,439 Primary; 150,915 ± 24118 Revision	following primary TKA
	Yang et al.[22]	2019	Retrospective cohort study	IV	To analyze the relationships between a positive LTT for metal, the histopathology from the revision surgery, and pre-revision and post-revision clinical and functional outcomes	27 TKA patients (6 M, 21 F)	▪ Radiograph ▪ KSS ▪ [3H]-thymidine	▪ LTT stimulation index: 9.4 ± 8.2 Pre-revision KSS ▪ Clinical: 49.8 ± 16.7 ▪ Functional: 44.8 ± 16.7 ▪ ROM: 93.2° ± 25.2° Post-revision KSS ▪ Clinical 75.6 ± 14.6 ▪ Functional 57.4 ± 15.5 ▪ ROM: 116.6° ± 19.9°	Positive LTT result may not indicate that an immune reaction is the cause of pain and stiffness post-TKA
Psychology	Belford et al.[23]	2020	Prospective study	IV	To examine the psychosocial factors that predicted pain, stiffness and physical	102 TKA patients	▪ OKS ▪ WOMAC	▪ WOMAC - Pain: Pre 12.04 (3.68); 1y after 5.68 (5.10)	The findings indicate that preoperative psychosocial factors

					functioning up to one year following TKA			- Function: Pre 41.62 (12.80); 1y after 21.83 (17.01) - Stiffness: Pre 5.19 (1.76); 1y after 3.15 (2.01) ▪ OKS: Pre 15.99 (7.95); 1y after 33.62 (10.40)	are important in understanding outcomes of TKA
	Laverni a et al.[24]	2015	Observational study	III	1. To assess patients outcomes, including Knee Society (KS) scores, of those patients who developed arthrofibrosis and underwent manipulation and compare them with a control group; 2. In patients with arthrofibrosis, to determine outcomes and KS scores of those with psychological distress and those without it	53 TKA patients (57 knees) with arthrofibrosis and underwent manipulation under anesthesia; 58 TKA patients (63 knees) without arthrofibrosis;	▪ QWB-7 ▪ SF-36 ▪ WOMAC ▪ KSS	Arthrofibrosis (group1) vs no Arthrofibrosis (group2): ▪ QWB-7 total Pre: Group1 0.522 (0.05); Group2 0.530 (0.06); Post: Group1 0.587 (0.10); Group2 0.609 (0.13) ▪ KSS knee score Pre: Group1 42.59 (22.9); Group2 43.27 (23.7) Post: Group1 81.18 (22.9); Group2 86.58 (22.7) ▪ KSS function score Pre: Group1 26.84 (20.5); Group2 37.06 (19.3) Post: Group1 51.30 (22.1); Group2 59.76 (21.4) ▪ KSS ROM: Pre: Group1 103.07 (18.7); Group2 104.92 (22.0) Post: Group1 101.63 (21.5); Group2 109.76 (17.0) ▪ WOMAC stiffness Pre: Group1 3.96 (2.3); Groups2 3.61 (2.2)	Patients who developed arthrofibrosis had worse preoperative KS function than patients without postoperative arthrofibrosis. Patients with arthrofibrosis and psychological distress perceived themselves as having worse quality of wellbeing and more stiffness before the surgical intervention and their worse quality of well-being perception was sustained after surgery

								Post: Groups1 1.23 (2.2); Group2 0.56 (1.2) - WOMAC total Pre: Groups1 61.16 (18.2); Group2 57.06 (15.8) Post: Group1 17.00 (26.2); Group2 7.22 (13.6) - Knee active flexion Pre: Group1 102.39 (19.1); Group2 102.44 (21.2) Post: Group1 96.71 (17.3); Group2 104.19 (16.1) Distressed (group1) vs nonDistressed (group2): - QWB-7 total Pre: Group1 0.490 (0.04);Group20.547 (0.05) Post: Group1 0.537 (0.09); Group2 0.627 (0.10) - KSS knee score: Pre: Group1 41.00 (22.7); Group2 43.78 (23.4) Post: Group1 80.32 (22.3); Group2 81.86 (23.8) - KSS function score Pre: Group1 19.40 (18.5); Group2 32.66 (20.3) Post: Group1 50.00 (24.4); Group2 52.32 (20.6) - KSS ROM:	
--	--	--	--	--	--	--	--	---	--

								Pre: Group1 105.40 (15.6); Group2 101.25 (21.0) Post: Group1 102.73 (21.0); Group2 100.83 (22.2) - WOMAC stiffness Pre: Group1 4.92 (2.1); Groups2 3.22 (2.1) Post: Groups1 1.52 (2.2); Group2 1.0 (2.1) - WOMAC total: Pre: Groups1 66.96 (17.0); Group2 56.81 (18.0) Post: Group1 20.64 (27.5); Group2 14.16 (25.2) - Knee active flexion: Pre Group1 102.68 (16.4); Group2 102.16 (21.3) Post: Group1 96.23 (18.4); Group2 97.07 (16.8)	
	Razmjou et al.[25]	2015	Observational study	III	To determine whether reporting neuropathic pain (NP) at an average of 5 years after total knee arthroplasty (TKA) was related to patient age, sex, preoperative comorbidity, arthritis self-efficacy, or disability before surgery and at 1 year after surgery	63 TKA patients (16 M, 47 F)	- S-LANSS scores - WOMAC - PHQ-9	- NP: 9 patients - No NP: 54 patients	Participants with NP reported higher levels of pain, stiffness, and physical dysfunction as early as 1 year after surgery that continued up to an average of 5 years after surgery
Demographics and others	Clement , Bardgett et al.[26]	2019	Retrospective study	III	1. To compare the WOMAC and SF12 outcome of patients with increased symptoms of stiffness 1 year after TKA with those who had no change or improvement in symptoms;	2589 TKA patients Increased Stiffness group (IS)	- WOMAC - SF-12	Pre-Surgery WOMAC - Pain: IS 42.4 (15.9) ; NIS 35.1 (17.7) - Physical function: IS 45.3 (16.9); NIS 36.3 (16.9)	Patients with increased symptoms of stiffness after TKA have a worse functional outcome

					2. To identify independent predictors of increased symptoms of stiffness 1 year following TKA	vs Non Increased Stiffness group (NIS)		- Stiffness: IS 61.0 (22.4) ; NIS 36.0 (19.6) ▪ SF12 PCS: IS 28.4 (7.6); NIS 27.5 (7.4); ▪ SF12 MCS:IS 46.3 (13.3) ; NIS 47.1 (13.6) 1 year after ▪ WOMAC - Pain: IS 63.1 (24.0) ; NIS 81.0 (19.8) - Physical function: IS 55.4 (23.6); NIS 74.5 (20.7) - Stiffness: IS 41.0 (22.4) ; NIS 73.2 (21.3) ▪ SF12 PCS: IS 31.8 (8.8) ; NIS 37.9 (11.1) ▪ SF12 MCS:IS 44.0 (14.7) ; NIS 49.5 (12.7);	and a lower rate of patient satisfaction. Pain, function, and the total WOMAC scores were poor predictors of increasing stiffness after TKA
	Clemen, Merrie et al.[27]	2019	Retrospective study	II	1. To identify independent preoperative predictors of outcome and patient satisfaction for the second total knee arthroplasty 2. To compare the predictors of outcome and patient satisfaction of the first to the second TKA and whether dissatisfaction with the first TKA predicts satisfaction with the second TKA	454 asynchrono us bilateral primary TKA patients First TKA vs Second TKA	▪ WOMAC	Pre Surgery ▪ WOMAC - Pain: FirstTKA 35.3 (17.4); SecondTKA 38.8 (16.8) - Physical function: First 36.0 (16.9); Second 40.4 (16.7) - Stiffness: First 36.7 (20.0); Second 41.2 (20.1) 1 year after ▪ WOMAC - Pain: First 83.7 (19.7) ; Second 80.2 (21.9) - Physical function: First 76.8 (19.9); Second 76.6 (19.5); - Stiffness: First 73.7 (22.1); Second 75.5 (20.8);	Improvement in pain and function is less with the second TKA, but the satisfaction rate remains similar. There are common independent predictors for change in the WOMAC score for the first and second TKA; however, the predictors of satisfaction were different with no common factors

	Gungor et al.[28]	2019	Retrospective study	IV	1.To estimate the risk of developing moderate-to-severe Persistent Postsurgical Pain (PPP) after primary TKA; 2. To explore potential predictors of the development of PPP in this clinical setting	578 TKA patients	Collaborative Orthopedic Replacement Registry	The risk of PPP after TKA was 31.3%	This study estimated a high risk of moderate-to-severe PPP after primary TKA. Racial differences and types of peripheral nerve blocks might be associated with greater odds of developing moderate-to-severe PPP after TKA surgery
	Järvenpää et al.[29]	2012	Observational study	III	To analyze clinically and radiologically the results of total knee arthroplasty in obese patients	48 TKA patients (52 knees); 29 non-obese vs 23 obese	- KSS - WOMAC - TUG - Radiographs	- KSS Knee score: 88.9 ± 16.3 Non-obese; 83.6 ± 13.0 Obese - KSS Function score: 76.3 ± 31.5 Non-obese; 63.6 ± 30.1 Obese - ROM: 109.6 ± 8.1 Non-obese; 104.6 ± 9.2 Obese - WOMAC - Pain: 11.6 ± 17.9 Non-obese; 20.7 ± 24.7 Obese - Stiffness: 13.4 ± 21.2 Non-obese; 26.9 ± 24.3 Obese - Physical function: 14.4 ± 22.4 Non-obese; 26.5 ± 10.7 Obese - Walking distance: 2935 ± 2303 Non-obese; 1894 ± 1852 Obese - TUG: 12.1 ± 6.3 Non-obese; 12.7 ± 4.9 Obese	Obese patients had poorer clinical success at final follow-up. WOMAC scores were significantly higher in obese patients.

								▪ Deviation from straight mech.axis: 3.3 ± 3.3 Non-obese;3.3 ± 2.4 Obese	
	Liebenst einer et al.[30]	2019	Retrospective comparative study	II	To determine if preoperative radiologic minimal joint space width (mJSW) is related to the outcome ofTKA	377 TKA patients (141 M, 236 F)	▪ Minimal Joint Space Width (mJSW) ▪ Radiograph ▪ WOMAC	Preoperative ▪ WOMAC - Pain: 50% (mJSW: 0–1 mm); 50% (mJSW: ≥2 mm) - Function: 49.5% (mJSW: 0–1 mm); 52% (mJSW: ≥2 mm) - Stiffness: 55% (mJSW: 0–1 mm); 55% (mJSW: ≥2 mm) - Total: 52% (mJSW: 0–1 mm); 53% (mJSW: ≥2 mm) Postoperative (1 year) ▪ WOMAC - Pain: 6% (mJSW: 0–1 mm); 14% (mJSW: ≥2 mm) - Stiffness: 15% (mJSW: 0–1 mm); 25% (mJSW: ≥2 mm) - Function: 9.5% (mJSW: 0–1 mm); 20% (mJSW: ≥2 mm) - Total: 10% (mJSW: 0–1 mm); 19% (mJSW: ≥2 mm)	Patients with preoperative complete joint space collapse (0 to 1 mm mJSW) achieve a significantly better WOMAC result from TKA than do those with a mJSW equal to or greater than 2 mm

	Liebs et al.[31]	2011	Retrospective cohort study	II	To establish whether (1) women had worse knee function and health-related quality of life after TKA compared with men, (2) lower improvements in scores, and (3) slower recovery after surgery	494 TKA patients (141 M, 353 F)	▪ WOMAC ▪ SF-36	After 2 years ▪ WOMAC - Pain: 11.9 ± 16.2 Male ; 13.6 ± 17.7 Female - Physical function: 15.9 ± 16.5 Male; 19.8 ± 20.3 Female - Stiffness: 21.0 ± 23.3 Male; 22.1 ± 23.1 Female ▪ SF12 PCS: 43.0 ± 9.5 Male; 40.5 ± 10.0 Female ▪ SF12 MCS: 52.8 ± 9.6 Male; 50.2 ± 10.9 Female	Women had faster recovery at 3- and 6-month follow-ups than men. It could be attributable to the lower preoperative health-related quality of life in women
	Parvizi et al.[32]	2014	Multicenter study	III	To quantify the degree of residual symptoms and specific functional deficits in young patients who had undergone TKA	661 TKA patients	▪ Interview	▪ Patients satisfied of function: 90% ▪ Patients satisfied performing normal daily life activity: 89% ▪ Patients satisfied of pain relief: 91%	Many patients continue to experience knee symptoms after full recovery from TKA
	Putman et al.[33]	2018	Retrospective study	IV	To assess survival, functional outcomes, and the nature and frequency of complications	263 TKA patients (141 M, 122 F)	▪ KOOS ▪ Radiographs	▪ Survival rate: 89% ▪ KOOS: 159 little or no discomfort, 157 little or no pain, 41 extreme or severe distress ▪ HKA: $179^\circ \pm 3.2^\circ$ (range 171°-188°)	Although TKA improved knee range of motion, only 60% of patients were satisfied with the procedure

Table 2: Summary of incidence and causes of stiffness in TKA patients. The percentage of patients reporting stiffness was pooled from the different scores of each study.

Category	N° of studies	N° of TKA patients	% of patients with stiffness	Main Causes
Surgery-related issues	7	487	47%	• IR of tibial or femoral component (n=3) • Malalignment (n=3) • Over-voluming (n=1)
Loosening	3	138256	12%	• Cementation technique (n=1) • Bone quality (n=1) • Component design (n=1)
Gene expression	1	100	50%	• Expression of profibrotic markers (n=1)
Immunology & Allergy	3	332	25%	• High material hypersensitivity-related cytokines level (n=3)
Psychology	3	343	36%	• Anxiety (n=2) • Depressive symptoms (n=2) • Functional limitation (n=2) • Severe preoperative pain (n=1)
Demographics and others	8	5473	17%	• Young age (n=1) • Male gender (n=1) • Obesity (n=1) • Previous surgery (n=1) • Diabetes (n=1) • Preoperative stiffness (n=2)

