



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Benzodiazepines' role in managing gastrointestinal disorders

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Balon, R., Sonino, N., Rafanelli, C. (2021). Benzodiazepines' role in managing gastrointestinal disorders. *PSYCHOTHERAPY AND PSYCHOSOMATICS*, 90(2), 81-84 [10.1159/000510686].

Availability:

This version is available at: <https://hdl.handle.net/11585/837227> since: 2022-11-24

Published:

DOI: <http://doi.org/10.1159/000510686>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

Benzodiazepines' role in managing gastrointestinal disorders

Richard Balon, ¹

Nicoletta Sonino, ^{2,3}

Chiara Rafanelli, ⁴

¹ Departments of Psychiatry and Behavioral Neurosciences and Anesthesiology, Wayne State University School of Medicine, Detroit, Michigan, USA

² Department of Statistical Sciences, University of Padova, Padova, Italy

³ Department of Psychiatry, University at Buffalo, State University of New York, Buffalo, NY, USA

⁴ Department of Psychology, University of Bologna, Bologna, Italy

Address all correspondence to: Richard Balon, M.D., Department of Psychiatry, Tolan Park Bldg. #A, 3901 Chrysler Service Dr., Detroit, MI 48201, USA; phone 1-313-9933416; fax: 1-313-9933422; e-mail: rbalon@wayne.edu

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29

Benzodiazepines have been used broadly in gastroenterology, for instance in treating anxiety symptomatology associated with gastrointestinal (GI) illness, or in treating some underlying conditions (e.g., addressing alcohol dependence in liver disease), or in various diagnostic endoscopy procedures, and in preparation for surgery. It is known that anxiety and GI symptomatology/disorders frequently co-exist, with anxiety exacerbating GI disorders, and some GI symptoms provoking anxiety.

Association of anxiety and GI symptomatology

The complex relationship between anxiety and GI symptomatology has attracted limited attention over the years. In a study by Esler and Goulston (1) patients with the irritable-colon syndrome who had predominantly diarrhea were significantly more anxious and more neurotic than the control population of general medical patients. When analyzing results of the Epidemiological Catchment Area study, Walker and colleagues (2) compared subjects reporting unexplained GI symptoms to those not reporting any GI symptoms. Subjects reporting at least one unexplained GI symptoms were significantly more likely to have experienced lifetime episodes of panic disorder (2.5% vs. 0.7%), agoraphobia (10.0% vs. 3.6%) and major depression (7.5% vs. 2.9%) when compared to subjects with no unexplained GI symptoms. The lifetime prevalence rates differences were even higher for subjects with two unexplained GI symptoms: 5.2% for panic disorder, 17.8% for agoraphobia and 13.4% for major depression. In her review (3), Spinelli concluded that approximately 40-60% of patients with irritable bowel syndrome (IBS) have one or more concomitant psychiatric disorder, with anxiety, depression and somatoform disorders being the most common, and that psychosocial stress exacerbates GI symptoms, including symptoms of IBS. In a study on anxiety and depressive symptoms and medical illness among adults with anxiety disorders, Niles and colleagues (4) found that anxiety was independently associated with having ulcer and that severity of anxiety and depressive symptoms was strongly associated with having more medical conditions. They felt that bidirectional relation between anxiety and various diseases is likely. Goodwin and Stein (5) found a strong relationship between generalized anxiety disorder (GAD) and peptic ulcer disease (PUD) (five-fold increase over people without GAD). In his review on anxiety and medical

1 illness, Schuckit (6) suggested that three GI problems (colitis, ulcer, IBS) may be exacerbated
2 by anxiety.

3 **Potential mechanisms**

4 The mechanism of the bidirectional relationship between anxiety and GI
5 illness/symptomatology is not understood and has not been **sufficiently investigated**. In his
6 review of this relationship, Clouse (7) postulated interaction of emotional or other stressful, non-
7 intestinal stimuli on gastrointestinal tract function and possible direct relation of these stimuli to
8 some gastrointestinal illnesses, with mechanisms being both neural and humoral. He wrote that
9 “The substrate for modulation of the gastrointestinal tract by anxiety and other psychiatric
10 disorders is provided by our knowledge of central nervous system control mechanisms of
11 gastrointestinal functions:” (7, p.414). Other investigators (8) emphasized the role of stress in
12 peptic ulcer disease. However, since the discovery of the role of *Helicobacter pylori* in the
13 etiology of peptic ulcer disease, the interest in studying the role of stress in this entity decreased
14 though many recognize its multicausal etiology, including stress and unhealthy lifestyle.
15 Interestingly, some indirect evidence of the impact of anxiety on GI symptomatology comes
16 from studies examining the impact of benzodiazepines (BZs) on various aspects of GI
17 functioning.

18 Birnbaum and colleagues (9) found that oral administration of 10 mg of diazepam
19 markedly reduced nocturnal human basal gastric secretion in duodenal ulcer patients when
20 compared to placebo. **They postulated that, since the decrease in secretion was achieved by a**
21 **drug that has no anticholinergic effect in vitro, the mechanism should refer to regulatory**
22 **autonomic functions in the central nervous system (9).** One may wonder whether the decrease of
23 anxiety and/or the improved sleep were the causative factors.

24 It is well known that anxiety, especially chronic one, increases smooth muscles tension
25 throughout the human body, including the GI system. It is commonly believed that this is
26 centrally mediated though we do not have full understanding of this process. Benzodiazepine
27 tetrazepam (used as muscle relaxant) relieved spasm of isolated rat duodenum and guinea pig
28 ileum (10), which suggests peripheral involvement. On the other hand, intravenous diazepam
29 decreased the pressure of lower esophageal sphincter in humans (11) that may suggest both
30 peripheral (myogenic influence) and central (anxiety) mechanism. These findings suggest the

1 possibility of using BZs for relieving spasm of musculature within the GI system. **It would**
2 **probably make sense to use longer acting BZs such as clonazepam rather than short-acting ones**
3 **(e.g., alprazolam), considering constant and continuous changes in GI motility and tone.**

4 Increasingly important areas to consider in the involvement of anxiety in GI
5 symptomatology are the regulatory function of the brain-gut axis (12) and the impact of the
6 microbiota (13), especially in view of the mentioned (4) bi-directional relationship between
7 anxiety and GI symptomatology/illness. The brain receives constant stream of interoceptive input
8 from the GI tract and after integrating them sends response back to different GI tract cells,
9 including the mentioned GI smooth muscle cells (12). The brain can modulate the afferent
10 signals from the GI system by endogenous pain facilitation or reduced endogenous pain
11 inhibition (12), and may play a role in painful GI symptomatology, e.g. in the irritable bowel
12 syndrome. Clearly, alteration of homeostatic reflexes could be associated with alteration in
13 various GI functions, such as secretion and motility, thus leading to abdominal pain, discomfort,
14 and alteration of bowel habits (12). The regulation of the gut includes sympathetic and
15 parasympathetic nervous system, the HPA axis, and involves various neurotransmitters,
16 including the gamma-aminobutyric acid (GABA) (13). Various gut signaling (in both directions)
17 could be influenced by numerous neurotransmitters, and the release of some of them, such as
18 GABA, could be influenced by BZs.

19 The mechanism of anxiety involvement in GI symptoms/disorders though widely known
20 is not well understood. It seems, however, that medications such as benzodiazepines, may
21 alleviate GI symptoms, possibly both centrally and peripherally.

22 **BZ use and efficacy**

23 **Most of the studies examining the use of BZs in the treatment of GI symptomatology**
24 **were done during the 1970s and 1980s.**

25 In 1973, Baume and Cuthbert (14) noted that medazepam was significantly better than
26 placebo in relieving major symptom complexes of functional bowel disease, e.g. aerophagy,
27 nervous dyspepsia and gastrointestinal pain of 30 patients in a double-blind cross-over clinical
28 trial. Expanding their research on BZs in GI symptoms, Baume and colleagues (15) compared
29 diazepam (5 mg BID) and lorazepam (1 mg BID) to placebo in a double-blind, cross-over study

1 of 28 patients with functional GI symptoms. Both BZs were significantly better than placebo.
2 The authors noted that lorazepam (at that time it was fairly new medication) was well tolerated.

3 Lorazepam was found to be useful in the treatment of GI illness with anxiety component
4 in several other studies. Kasich and colleagues (16) reported that lorazepam (0.5 mg to 4.0
5 mg/day) was better than placebo and comparable to diazepam (2.5 to 30 mg/day) in a double-
6 blind parallel treatment study of 120 patients with chronic gastrointestinal disease associated
7 with anxiety. They concluded that lorazepam is an effective, safe and useful adjunct in the
8 management of functional and organic GI disorders where anxiety has negative influence.
9 Caplan and Vanov (17) found lorazepam significantly more helpful than placebo in 61 anxious
10 patients with demonstrable “organic” or “functional” GI diseases in a double-blind placebo-
11 controlled study of lorazepam (mean daily dose 3 mg). Lorazepam was helpful not just in
12 relieving anxiety, but also in some GI symptoms, such as heartburn, constipation, aerophagia,
13 bad taste and others. Finally, Stokes (18) reported that lorazepam was effective in a double-blind
14 placebo comparison of lorazepam. The initial dose of lorazepam was 3 mg/day; 21% of patients
15 were on a higher dose at the end of the study. The improvement was significantly greater with
16 lorazepam than with placebo on all measures (anxiety, global, GI symptomatology).

17 The place of BZs in the management of GI symptoms/illness was also examined in
18 studies combining BZs with other medications used in treatment of GI problems. Huscher and
19 colleagues (19) studied ranitidine and prazepam or ranitidine and placebo in a double-blind study
20 of 50 patients with an active, endoscopically proven duodenal ulcer. At the end of their 4 -week
21 study, duodenal ulcer healed in 95.6% of patients on ranitidine + prazepam and 75% of patients
22 on ranitidine + placebo ($p = 0.03$) (there were 2 dropouts in prazepam group and one in placebo
23 group). Combination of octatropine methyl-bromide (anticholinergic agent) and diazepam (40
24 mg/2.5 mg twice daily) in patients with IBS was not significantly different from placebo in a
25 study by Pace and colleagues (20).

26 It is important to note that the side effects in all mentioned studies were usually mild,
27 mostly sedation and sleepiness.

28 The evidence from these studies suggests an important role of BZs in managing various
29 GI symptoms and disorders.

1 **BZs versus antidepressants**

2 Considering their favorable side effects profile, BZs should be preferred to
3 antidepressants in the management of GI symptoms with anxiety. BZs are usually not associated
4 with GI side effects, while antidepressants are frequently associated with a number of adverse
5 events. Tricyclic antidepressants, effective in the treatment of functional GI disorders (21, 22),
6 also in conjunction with cognitive behavioral therapy (23) may cause constipation. Second
7 generation antidepressants, such as SSRI and SNRI, may have a more profound unfavorable
8 impact (24, 25), causing nausea, vomiting, dyspepsia, diarrhea and abdominal pain (24). Even
9 short-term use of SSRI may be significantly associated with GI bleeding, suggesting that the
10 same precautions that are used with non-steroidal anti-inflammatory drugs and aspirin are
11 appropriate (25). As a result, newer generation antidepressants do not seem to be suitable for the
12 treatment of anxiety associated with GI symptomatology, and BZs should be preferred.

13

14 **Conclusion**

15 The role of BZs in the management of various troublesome gastrointestinal symptoms,
16 (e.g., aerophagia and gastrointestinal pain), or some acute and chronic GI disorder (e.g., peptic
17 ulcer disease, colitis, and IBS) has not been fully appreciated. Their use should be expanded. In
18 addition to anxiety frequently accompanying these conditions, BZs may, through either
19 decreased gastric secretion and/or relaxation of smooth muscles tone, alleviate GI
20 symptomatology, and contribute to patient well-being. The use of BZs in GI disorders may
21 facilitate a transition of patients from pharmacotherapy to psychotherapy, which is an effective
22 treatment particularly for functional symptoms (26, 27), and may provide more enduring results.

23

24

25

26

27

1 **References**

- 2 1. Esler MD, Goulston KJ: Levels of anxiety in colonic disorders. *N Engl J Med* 1973; 288: 16-
3 20.
- 4 2. Walker EA, Katon WJ, Jemelka RP, Roy-Byrne PP: Comorbidity of gastrointestinal
5 complaints, depression, and anxiety in the Epidemiological Catchment Area (ECA) study. *Am J*
6 *Med* 1992; 92 (suppl 1A): 26S-30S.
- 7 3. Spinelli A: Irritable bowel syndrome. *Clin Drug Invest* 2007; 27: 15-33.
- 8 4. Niles AN, Dour HJ, Stanton AL, Roy-byrne PP, Stein MB, Sherbourne C, Rosw RD, Craske
9 MG: Anxiety and depressive symptoms and medical illness among adults with anxiety disorders.
10 *J Psychosom Res* 2005; 78: 109-115.
- 11 5. Goodwin RD, Stein MB: Generalized anxiety disorder and peptic ulcer disease among adults
12 in the United States. *Psychosom Med* 2002; 64: 862-866.
- 13 6. Schuckit MA: Anxiety related to medical disease. *J Clin Psychiatry* 1983; 44 [11, Sec 2]: 31-
14 36,
- 15 7. Clouse RE: Anxiety and gastrointestinal illness. *Psychiatr Clin North Am* 1988; 11(2) 399-417
- 16 8. Wolf S, Almy TP, Bachrach WH, Spiro HM, Sturdevant RAL, Weiner H: The role of stress in
17 peptic ulcer disease. *J Human Stress* 1979; 5(2): 27-37.
- 18 9. Birnbaum D, Karmeli F, Tefera M: The effect of diazepam on human gastric secretion. *Gut*
19 1971; 12: 616-618.
- 20 10. Perez-Guerrero C, Suarez J, Herrera MD, Marhuenda E: Spasmolytic effects of tetrazepam
21 on rat duodenum and guinea-pi ileum. *Pharmacol Res* 1997; 35: 493-497.
- 22 11. Rushnak MJ, Leevy CM: Effect of diazepam on the lower esophageal sphincter. *Am J*
23 *Gastroenterol* 1980; 73: 127-130.
- 24 12. Mayer EA, Tillisch K: The brain-gut axis in abdominal pain syndromes. *Ann Rev Med* 2011;
25 62: 381-396.

- 1 13. Mayer EA, Tillisch K, Gupta A: Gut/brain axis and the microbiota. *J Clin Invest* 2015; 125:
2 926-938.
- 3 14. Baume P, Cuthbert J: The effect of medazepam in relieving symptoms of functional
4 gastrointestinal distress. *Aust N Z J Med* 1973; 3 (5): 457-460.
- 5 15. Baume P, Tracey M, Dawson L: Efficacy of two minor tranquilizers in relieving symptoms
6 of functional gastrointestinal distress. *Aust N Z J Med* 1975; 5: 503-506.
- 7 16. Kasich AM, Richard DJ, Vanov SK: Lorazepam in the management of anxiety associated
8 with chronic gastrointestinal disease: a double-blind study. *Curr Ther Res Clin Exp* 1976; 19:
9 292-306.
- 10 17. Chaplan A, Vanov SK: GI illness: treatment of the anxiety component. *Psychosomatics*
11 1977; 18: 49-54.
- 12 18. Stokes DK: Lorazepam in anxiety associated with chronic enteritis and ulcerative colitis. *J*
13 *Clin Psychiatry* 1978; 39 (Suppl.10, Pt 2): 53-57.
- 14 19, Huscher C, Magni G, Salmi A, Bossini S, Felini C, Besozzi F, de Leo D: Ranitidine versus
15 ranitidine and prazepam in the short-term treatment of duodenal ulcer – a double-blind controlled
16 trial. *Eur J Clin Pharmacol* 1985; 28: 177-180.
- 17 20. Pace F, Maurano A, Ciacci C, Savarino V, Attili A, Iaquinto G, Magni E, Bianchi Porro G:
18 Octatropine methyl bromide and diazepam combination (Valpinax) in patients with irritable
19 bowel syndrome: a multicenter, randomized, placebo-controlled trial. *Eur Rev Med Pharmacol*
20 *Sci* 2010; 14: 155-162.
- 21 21. Jackson JL, O'Malley PG, Tomkins G, Balden E, Santoro L, Kroenke K: Treatmentn of
22 functional gastrointestinal disorders with antidepressant medications: a meta-analysis. *Am J med*
23 2000; 108: 65-72.
- 24 22. Clouse RE: Antidepressants for functional gastrointestinal syndromes. *Dig Dis Sci* 1994; 39:
25 2352-2363.
- 26 23. Drossman DA, Toner BB, Whitehead WE, Diamant NE, Dalton CB, Duncan S, Emmott S,
27 Proffitt V, Akman D, Fruecianta K, Le T, Meyer K, Brandshaw B, Mikula K, Morris CB,

- 1 Blackman CJ, Hu Y, Jia Hm Li JZ, Koch GG, Bangdiwala SI: Cognitive -behavioral therapy
2 versus education and desipramine versus placebo for moderate to severe functional bowel
3 disorders. *Gastroenterology* 2003; 125: 19-31.
- 4 24. Carvalho AF, Sharma MS, Brunoni AR, Vieta E, Fava GA: The safety, tolerability and risks
5 associated with the use of newer generation antidepressant drugs: a critical review of the
6 literature. *Psychother Psychosom* 2016; 85: 270-288.
- 7 25. Wang YP, Chen YT, Tsai CF, Li Sy, Suo JC, Wang CJ, et al. Short-term use of serotonin
8 reuptake inhibitors and risk of upper gastrointestinal bleeding. *Am J Psychiatry* 2014; 171:54-61
- 9 26. Laird KT, Tanner-Smith EE, Russel AC, Hollon SD, Walker LS: Short-term and long-term
10 efficacy of psychological therapies for irritable bowel syndrome: a systematic review and meta-
11 analysis. *Clin Gastroenterol Hepatol* 2016; 14: 937-947.
- 12 27. Laird KT, Tanner-Smith EE, Russel AC, Hollon SD, Walker LS: Comparative efficacy of
13 psychological therapies for improving mental health and daily functioning in irritable bowel
14 syndrome: a systematic review and meta-analysis. *Clin Psychol Rev* 2017;51: 142-152.
- 15
- 16