

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Allelopathic interactions between phytobenthos and meiofaunal community in an Adriatic benthic ecosystem:
Understanding the role of aldehydes and macroalgal structural complexity

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Lenzo, D., Pezzolesi, L., Samori', C., Rindi, F., Pasteris, A., Pistocchi, R., et al. (2022). Allelopathic interactions between phytobenthos and meiofaunal community in an Adriatic benthic ecosystem: Understanding the role of aldehydes and macroalgal structural complexity. SCIENCE OF THE TOTAL ENVIRONMENT, 807(Pt 2), 1-14 [10.1016/j.scitotenv.2021.150827].

Availability:

This version is available at: <https://hdl.handle.net/11585/839536> since: 2024-05-17

Published:

DOI: <http://doi.org/10.1016/j.scitotenv.2021.150827>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Lenzo D.; Pezzolesi L.; Samori' C.; Rindi F.; Pasteris A.; Pistocchi R.; Colangelo M.A.: *Allelopathic interactions between phytobenthos and meiofaunal community in an Adriatic benthic ecosystem: Understanding the role of aldehydes and macroalgal structural complexity*

SCIENCE OF THE TOTAL ENVIRONMENT, vol.807, ISSN 0048-9697

DOI: 10.1016/j.scitotenv.2021.150827

The final published version is available online at:

<https://dx.doi.org/10.1016/j.scitotenv.2021.150827>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Allelopathic interactions between phytobenthos and meiofaunal community in an Adriatic benthic ecosystem: understanding the role of aldehydes and macroalgal structural complexity

Denise Lenzo¹, Laura Pezzolesi^{1*}, Chiara Samori², Fabio Rindi³, Andrea Pasteris¹, Rossella Pistocchi¹, Marina Antonia Colangelo¹

¹ Dipartimento di Scienze Biologiche, Geologiche e Ambientali - Università di Bologna, via Sant'Alberto 163, 48123 Ravenna (Italy)

² Dipartimento di Chimica "Giacomo Ciamician" - Università di Bologna, via Selmi 2, 40126 Bologna (Italy)

³ Dipartimento di Scienze della Vita e dell'Ambiente - Università Politecnica delle Marche, via Brecce Bianche, 60131 Ancona (Italy)

*Corresponding Author

Dr. Laura Pezzolesi

Tel. +39 (0)544 937373; fax: +39 (0)544 937411

laura.pezzolesi@unibo.it

25 **Abstract**

26 Macroalgae produce several allelopathic substances, including polyunsaturated aldehydes (PUAs),
27 which may inhibit photosynthesis and growth rates of other algal species, and grazing. Additionally,
28 macroalgal structural complexity is an important factor in determining abundance patterns and size
29 structure of epiphytic organisms.

30 In this study the PUAs production of two Mediterranean macroalgae, *Dictyopteris polypodioides*,
31 (DP, Phaeophyceae, Dictyotales) and *Cystoseira compressa* (CC, Phaeophyceae, Fucales), was
32 characterized to clarify the relationships between the meiobenthic and microphytobenthic
33 communities. Results showed a higher PUAs production and a diverse qualitative profile for DP,
34 which reported long-chain compounds (i.e. C14-C16) as main aldehydes, than CC, with the short-
35 chain C6:2 as main compound, as well as variability among sampling times. A clear separation of the
36 meiofauna and microphytobenthos assemblages was found for the macroalgae, but with different
37 temporal trends. Dissimilarities were due to five microalgal orders, namely Naviculales, Lyrellales,
38 Gonyaulacales (i.e. *Ostreopsis*), Bacillariales, and Lichnophorales, and to the meiofaunal groups
39 nematodes, copepods, and copepod nauplii, which were more abundant on DP than on CC. Results
40 indicate that macroalgal complexity is a major determinant of the meiofaunal community structure
41 (accounting to 26% of the variation), rather than PUAs production itself (17%). PUAs effects seem
42 species-specific, thus affecting some grazers instead of the entire community. Conversely,
43 microphytobenthos affected the meiofauna assemblages, particularly harpacticoids, confirming the
44 role of these organisms as primary food source of all marine food chain producers. Since PUAs are
45 produced also by several epiphytic diatoms, the understanding of their effects on the community
46 structure and on the relationships among taxa in the field is complicated and requires further in-depth
47 investigations in simplified systems (i.e. microcosms).

48

49 **Keywords**

50 Brown macroalgae; PUAs; microalgae; meiofauna; harpacticoid copepods; chemical ecology.

51

52 **1. Introduction**

53 Chemical compounds produced by aquatic organisms and microorganisms have received increasing
54 attention for the important role that may have in modulating the interactions in marine ecosystems
55 (Paffenhöfer et al., 2005; Ianora et al., 2012). Many organisms produce allelochemicals, secondary
56 metabolites that affect survival and growth of other species and may give a competitive advantage to
57 their producers (Legrand et al., 2003; Fistarol et al., 2004; Tillmann, 2004; Allen et al., 2016).

58 Several studies regarding chemical interactions between various marine organisms, including
59 copepods, urchins, sea stars, algae, mollusks, and polychaetes, were focused on direct responses of
60 target species to allelochemicals and have been performed mainly under laboratory conditions (e.g.
61 Ianora et al., 2004; Adolph et al., 2004; Caldwell et al., 2004; Vardi et al., 2006 Ribalet et al., 2007;
62 Taylor et al., 2007).

63 Algae produce a variety of these allelopathic substances, such as phenolic compounds, alkaloids,
64 peptides, oxoacids, and polyketides, among which there are toxins synthesized by several species
65 (Snyder et al., 2003; Agostini-Costa et al., 2012; Pistocchi et al., 2012), polyunsaturated fatty acid
66 (PUFAs) (Grima et al., 1995; Tonon et al., 2002; Patil et al., 2007) and their derivatives, such as
67 polyunsaturated aldehydes (PUAs) (Wichard et al., 2005; Ianora et al., 2012; Pezsolesi et al., 2017).

68 Consequences of allelopathic interactions between different algal species can be of various nature
69 and magnitude, including loss of motility, cell deformation (Tang and Gobler, 2010; Pichierri et al.,
70 2017), pigmentation loss, cytoplasm aggregation, formation of vesicles and cellular lysis (Fistarol et
71 al., 2004). However, the majority of allelochemicals have milder effects, for example inhibition of
72 photosynthesis, reduction of growth rate, and inhibition of grazing (Legrand et al., 2003).

73 The effect of PUAs have been mostly investigated for planktonic microalgae and only a few studies
74 have evidenced PUAs production by benthic species, such as diatoms (Jüttner et al., 2010; Scholz
75 and Liebezeit, 2012; Pezsolesi et al., 2017), and macroalgae (Kajiwara et al., 1996; Akakabe et al.,
76 2003; Pezsolesi et al., 2021). Macroalgae are among the most important components of marine coastal

77 ecosystems because they are highly productive (Pinckney and Zingmark, 1993), have a high
78 taxonomic diversity, and may act as foundation species providing habitat for different organisms
79 (Cacabelos et al., 2010). The shape and structural complexity of macroalgae are important factors in
80 determining the abundance patterns and size structure of the epiphytic organisms (McAbendroth et
81 al., 2005). The more structurally complex macroalgal species show abundant and various populations
82 of invertebrates because they provide greater surface area for the colonization of epifaunal
83 assemblages and epiphytic microalgae (Chemello and Milazzo, 2002). Among epifaunal
84 assemblages, macrofauna is the most investigated, while meiofaunal communities are overlooked.
85 Meiofauna represent the most abundant and taxonomically diversified metazoans on Earth (Giere,
86 2009) and on hard substrates can overcome macrofauna in terms of abundance (Gibbons and Griffiths,
87 1986). Meiofauna plays a dominant role in the exchange of organic matter (Sandulli et al., 2014;
88 Semprucci et al., 2016) as part of the "small food web" (size class 45-1000 μm). Moreover, it supports
89 most of the higher trophic levels (Giere, 2009), being an important food resource for macrofauna,
90 small fish, juveniles and other epibenthic predators (Chardy and Dauvin, 1992). In phytal
91 environments, harpacticoid copepods are the dominant meiofauna group (Hicks, 1977; Coull and
92 Wells, 1983; Hall and Bell, 1993) and show high diversity (Sarmiento and Santos, 2012). They feed
93 mainly on diatoms and, for this reason, they have a high content of fatty acids and play a key
94 nutritional role for fish, carnivorous crustaceans (prawns and their larvae), and polychaetes (Coull,
95 1999; Giere, 2009).

96 Indeed, few studies have focused on the benthic environment and included the potential role of
97 macroalgae both in terms of production of allelochemical compounds and regulation of interactions
98 between various organisms (Kajiwara et al., 1996; Akakabe et al., 2003), while only one laboratory
99 study analyzed the effects of PUAs on a species of harpacticoid (i.e. *Tisbe holothuriae*) (Taylor et al.,
100 2007). PUAs strongly impair the reproduction of various potential grazers in *in vitro* studies (Poulet
101 et al., 1994; Ianora et al., 2004b), while in the field the relationship between aldehyde production and
102 reproductive failure of other higher-level organisms (i.e. copepods) remains unclear (Leflaive and

103 Ten-Hage, 2009). Some studies performed to investigate the allelopathic activity of fresh thalli or
104 extracts of *Ulva* spp. reported inhibitory effects on the growth of several microalgae, including
105 harmful species (Tang and Gobler, 2011). The production of aldehydes and their potential
106 consequences on biotic interactions may thus be explored, especially in macroalgae. This would help
107 to determine whether these compounds, which are present across phylogenetic and environmental
108 barriers, may play a role in the ecology of their producers (Leflaive and Ten-Hage, 2009), perhaps
109 affecting the development of epiphytic organisms, such as meiofauna or microalgae.

110 The main aims of this study were to i) analyze the qualitative and quantitative production of aldehydes
111 by two different macroalgae over a period of some months (Mediterranean spring and summer); ii)
112 understand the relationship between meiobenthic communities, with particular interest to
113 harpacticoid copepods, and the microphytobenthos present on the two macroalgal species,
114 considering the structural complexity of macroalgae and the potential role of aldehydes in regulating
115 their interaction.

116 Specifically, this study was performed in a benthic environment of the North-western Adriatic Sea
117 (Piscinetta del Passetto, Ancona, Italy), characterized by the presence of a rich phytobenthic
118 assemblage and by annual blooms of a toxic dinoflagellate, *Ostreopsis* cf. *ovata* Fukuyo, 1981. Two
119 macroalgal species, *Cystoseira compressa* (Esper) Gerloff & Nizamuddin and *Dictyopteris*
120 *polypodioides* A.P. De Candolle J.V. Lamouroux, commonly present in this site (Rindi et al., 2020)
121 and representative of the Mediterranean phytobenthic community were selected, based on their
122 different complexity and PUAs composition (Pezzolesi et al., 2021). To our knowledge, this is the
123 first study where macroalgal allelochemistry has been investigated to understand both the
124 microphytobenthos and meiofauna community structure and their interactions, in relation to the algal
125 complexity.

126

127 **2 Material and methods**

128 2.1 Sampling area and sampling procedure

129 The study was performed in a semi-enclosed and shallow (mean depth 1.5 m) inlet called Piscinetta
130 del Passetto (Conero Riviera, Italy, northern Adriatic Sea: 43°37'09" N, 13°31'54" E), described and
131 showed in Pezzolesi et al. (2021). Sampling was carried out at six different times from May to
132 September 2018 (see table S2), with monthly frequency except for September (when sampling was
133 performed twice). In the Mediterranean area, this period corresponds to late spring and summer.
134 Surface temperature and salinity were measured with a multiparameter water probe HQ30d (Hach-
135 Lange GmbH) and a refractometer Atago S-10, respectively.

136 Apical parts (i.e. tips) of the thalli (first 5-8 cm) of two macroalgae, i.e., *Cystoseira compressa*
137 (Phaeophyceae, Fucales) and *Dictyopteris polypodioides* (Phaeophyceae, Dictyotales), were sampled
138 at a depth of approximately 0.5 m. At the study site, during the time of the year in which sampling
139 was carried out, these species are present and mostly occur in a well-developed habit. For each
140 macroalgal species, six replicates were collected by snorkeling at each sampling time using 50 mL
141 polypropylene tubes (VWR International), avoiding the dispersion of the associated epiphytic
142 organisms.

143 Water samples for assessment of dissolved aldehydes and nutrient analysis were collected in 2 L
144 polyethylene bottles (VWR International) in the proximity of the sampled macroalgae. Water was
145 subsequently filtered using GF/F Whatman filters (0.7 µm porosity, 47 mm) and stored at -22 °C until
146 analysis. Macroalgae were treated to remove all associated benthic organisms as described in
147 Pezzolesi et al. (2021) using filtered seawater. Briefly, each tube containing the thallus and their
148 storage water was vigorously shaken to separate the macroalga from the epiphytic microalgae and the
149 meiofauna. Then the tube was rinsed with filtered seawater and vigorously washed several times until
150 epiphytic organisms were completely removed. The seaweed thalli were dried with absorbent paper,
151 then weighed to determine fresh weight (g fw) and photographed; finally, they were stored at -80 °C
152 in new tubes. The total volume of washing seawater of each sample (approximately 150 mL) was
153 measured and then divided into two aliquots, one for the microphytobenthos and the other for the

meiofauna analyses. Aliquots for microphytobenthos (about 75 mL) were fixed with Lugol and stored in 250 ml dark glass bottles. Aliquots for meiofauna analysis were sieved through a 1000- μ m mesh and a 45- μ m mesh; the meiobenthic organisms retained on the finer mesh sieve were fixed with 70% alcohol and stored in a 50 ml falcon until subsequent analyses. Each apical part of macroalga was placed on a white surface with a reference scale and then photographed using a digital camera Canon EOS 750D. To take into account the different morphology and complexity of the two algal species, pictures were processed using the program ImageJ v1.53u. For each image the scale was fixed and transformed to binary format with a pixel width of 1 cm. Then the area, the perimeter, and the fractal complexity (D) were measured based on the image and using a method analogous to the grid method (boundary dimension) proposed by Sugihara and May (1990). In this method, the fractal dimension is the slope of the linear fit of $\log N(s)$ versus $\log (1/s)$; where s represents the scale of analysis and $N(s)$ is the number of objects observed at that scale.

2.2 Aldehydes

2.2.1 Aldehydes (PUAs) produced by macroalgae

The extraction and quantification of PUAs produced by the different macroalgae was carried out as described in Pezzolesi et al. (2021) by gas chromatography-mass spectrometry (GC-MS). Specifically, a portion of the apical part of the thallus (about 0.2-0.8 g f. wt.) was shredded with mortar and pestle, in liquid nitrogen. The powder thus obtained was transferred into 10 mL tubes. Derivatization of the polyunsaturated aldehydes was performed with O-(2,3,4,5,6-pentafluorobenzyl)hydroxylamine hydrochloride solution (PFBHA HCl) and quantification was based on the internal standard (i.e. benzaldehyde). All reagents were purchased from Sigma-Aldrich (Milan, Italy) and used without any further purification.

2.2.2 Dissolved aldehydes (dPUAs) in seawater

179 Dissolved aldehyde concentration was determined following the protocol described by Vidoudez and
180 Pohnert (2008) with slight modifications, using PFBHA HCl in Tris–HCl 100 mM pH 7.2 as
181 derivatizing reagent and benzaldehyde as internal standard, as described in Pezzolesi et al. (2021).

182

183 *2.3 Nutrients*

184 Nitrate and phosphate analyses were performed on filtered seawater aliquots (GF/F Whatman
185 filters) and analyzed spectrophotometrically according to Strickland and Parsons (1972).

186

187 *2.4 Microphytobenthic community*

188 The quali-quantitative analysis of the microphytobenthos associated with the two macroalgae was
189 performed using an inverted optical microscope (Zeiss Axiovert 100) at 320x and 200x magnification.
190 Sub-samples (3-5 ml) of epiphytic communities fixed with Lugol were settled in counting chambers
191 after homogenization, according to the Utermöhl's sedimentation method (Utermöhl, 1958; Edler and
192 Elbrächter, 2010). Counting was performed in different ways. The microphytobenthos community
193 was examined at 320x magnification on 30 random fields or 4–5 transects; then a counting at 200x
194 of the organisms present on the whole sedimentation chamber was performed to obtain a correct
195 evaluation of uncommon taxa. The microphytobenthos was identified and recognized following
196 various manuals and identification keys (e.g. Tomas, 1997; Kraberg et al., 2010). The identification
197 of individuals was based exclusively on observable morphological characters (such as shape, size,
198 number of chloroplasts); the current taxonomic status for the microalgae was confirmed following
199 AlgaeBase (Guiry and Guiry, 2021).

200

201 *2.5 Meiofauna and harpacticoid copepods*

202 Meiofaunal organisms of each sample were counted and identified at higher taxonomic levels. All
203 harpacticoids (Order Harpacticoida Sars G.O., 1903) were collected under a stereomicroscope (Nikon
204 SMZ 1500) and stored in 70% alcohol inside 1.5 mL-Eppendorf labeled for subsequent identification.

205 Harpacticoids were identified to species level using Lang (1948; 1965), and Boxshall and Halsey
206 (2004).

207

208 2.6 Data analyses

209 PUAs produced by the macroalgae and dissolved in the seawater were analyzed and expressed as μg
210 g^{-1} fw and $\mu\text{g L}^{-1}$, respectively. Microphytobenthos was analyzed in terms of total epiphytic cells and
211 main orders or genera; data were expressed as the number of cells per gram of fresh macroalgae
212 weight (cells g^{-1} fw). Meiofauna and harpacticoid communities were analysed according to total
213 density (N), standardized towards the fresh weight of macroalgae (ind. g^{-1} fw), the total number of
214 taxa or species (S), and Shannon diversity (H'). For each variable, the average value of the six
215 replicates $\pm$ standard error is reported. All univariate variables were analyzed by a 2-way crossed
216 ANOVA; the factors considered were the macroalgal species (fixed, two levels: *Cystoseira*
217 *compressa* (CC) and *Dictyopteris polypodioides* (DP)) and time (fixed, 6 levels). Cochran's test was
218 used to check for the homogeneity of variances and data were transformed, if necessary (Underwood,
219 1996). If even after transformation it was not possible to obtain homogeneity of the variances,
220 untransformed data were analysed and results were considered robust if significant at $p < 0.01$ to
221 compensate for the increased probability of type I error (Underwood, 1996). When significant main
222 effects or interactions were detected, the Student–Newman–Keuls (SNK) test was used for pairwise
223 a posteriori comparisons.

224 The community structure of each assemblage (microphytobenthos, meiofauna and harpacticoids) was
225 analyzed by non-metric multidimensional scaling (nMDS) based on Bray-Curtis similarity of square
226 root-transformed data. Differences in community structures were assessed by permutational non-
227 parametric multivariate analysis of variance (PERMANOVA) following the same experimental
228 design adopted for ANOVA (Anderson, 2001; 2005). When significant main effects or interactions
229 were detected, the specific procedure provided within PERMANOVA was used for pairwise a

230 posteriori comparisons. The analyses were performed using unrestricted permutation of the raw data
 231 and 9999 permutations.

232 Taxa that mostly contributed to the dissimilarity/similarity among/within macroalgal species and
 233 times were identified using the SIMPER analysis (Clarke, 1993).

234 Relationships between macroalgae complexity, PUAs, dominant microphytobenthic taxa, and both
 235 meiofauna and harpacticoid copepod assemblages were analyzed by the distance-based linear model
 236 (DistLM) procedure in PERMANOVA+ (Anderson et al., 2008).

237 A total of 14 explanatory variables, grouped in three sets, were considered: macroalgae complexity
 238 descriptors (perimeter/area, D, Area), PUA concentrations (C14:5, C16:4, C16:3, C6:2), and
 239 microphytobenthic taxa abundances (*Navicula* spp. J.B.M. Bory de Saint-Vincent, 1822, *Lyrella* spp.
 240 N.I. Karayeva, 1978, *Cocconeis* spp. C.G. Ehrenberg, 1837, *Cylindrotheca* spp. L. Rabenhorst,
 241 1859/*Nitzschia* spp. A.H. Hassall, 1845, other diatoms, *Ostreopsis* cf. *ovata*, other dinoflagellates).

242 The relationships between meiobenthic taxa data and harpacticoid abundance (square root
 243 transformed data) and the three groups of variables were analyzed by explicitly examining the
 244 proportion of variation in the taxa data (or harpacticoid species abundance) that was explained by
 245 PUAs concentrations and microphytobenthic taxa abundances over and above the amount explained
 246 by macroalgae complexity descriptors.

247 The Akaike Information Criterion with correction (AICc) was used to select the model. Prior to the
 248 DistLM, a draftsman plot and correlation matrices were performed to detect possible skewness of the
 249 variables and/or strong correlation among pairs of variables (Anderson and Robinson, 2001).

250 Variables were not strongly correlated, so all variables were entered in the analysis. Concentrations
 251 of *Lyrella* spp., and *Cocconeis* spp were square root transformed. Finally, Pearson correlation analysis
 252 was used to test which macroalgal attribute (i.e. complexity descriptors such as D and macroalgal
 253 surface (S), and PUAs composition) explained the variation of some representative genera of
 254 microphytobenthos, of the main meiofauna taxa and of the most representative harpacticoid species.

255 Densities of microphytobenthic genera, main taxa and harpacticoid species were square root
256 transformed.

257 Significance level was set at 0.05 (5%) for all tests. ANOVA, Cochran test, SNK test, and correlations
258 were performed by R (version 3.5.3) using packages Lme4; all multivariate analyses were conducted
259 with PRIMER v7 (Clarke and Gorley, 2015) with the PERMANOVA + add-on (Anderson et al.,
260 2008).

261

262 3. Results

263 3.1 PUAs production by macroalgae

264 The interpretation of the chromatograms and relative mass spectra obtained by GC-MS gave the
265 quali-quantitative profile of the main aldehydes produced by *Dictyopteris polypodioides* (DP) and
266 *Cystoseira compressa* (CC).

267 The major aldehydes production was found in T1 (May) for both macroalgae, with values of $225.5 \pm$
268 35.6 and $17.1 \pm 4.0 \mu\text{g g}^{-1}$ fw for DP and CC, respectively. Conversely, the lowest amounts were
269 measured at times T3 (July) for DP and T2 (June) for CC, with a concentration of 44.3 ± 5.3 and 2.0
270 $\pm 0.3 \mu\text{g g}^{-1}$ fw, respectively (Table 1). From a quantitative point of view, ANOVA results (Table S1)
271 showed a higher PUAs production in DP than in CC and variability among different sampling times
272 ($F=17.73$, $P<0.001$).

273 From a qualitative point of view, the main compounds detected were the short-chain PUA hexadienal
274 (C6:2), which was present in both algae and resulted as the main compound in CC, and some long-
275 chain PUAs, namely hexadecatrienal (C16:3), hexadecatetraenal (C16:4), and tetradecapentaenal
276 (C14:5), which were detected exclusively in DP (Fig. S1). Specifically, C14:5 was the most abundant
277 compound in DP, with average concentrations ranging from a maximum of $115.3 \pm 20.8 \mu\text{g g}^{-1}$ fw at
278 time T1 to a minimum of $13.2 \pm 2.4 \mu\text{g g}^{-1}$ fw at time T3, corresponding to the 50% and 32% of the
279 total aldehydes (Table 1; Fig. S1). C16:4 had the maximum average concentration at time T1 with
280 $36.4 \pm 8.8 \mu\text{g g}^{-1}$ fw, decreasing gradually to a minimum of $5.9 \pm 1.5 \mu\text{g g}^{-1}$ (relative abundance of

8-29%) at time T6. C16:3 showed a similar pattern and an average maximum concentration of $30.9 \pm 9.3 \mu\text{g g}^{-1}$ fw at T1 (relative abundance of 12%); then, its amount decreased to a minimum of $8.7 \pm 3.1 \mu\text{g g}^{-1}$ fw at time T4, and it was not detected at all at T6. In DP the most variable compound was C6:2, with a maximum average concentration of $16.1 \pm 2.2 \mu\text{g g}^{-1}$ fw at time T1; this aldehyde was not detected at times T2 and T5. C6:2 was the main aldehyde in CC, accounting for 64-91% of the total amount, and its average concentration had the maximum and minimum values of 15.4 ± 3.5 and $1.0 \pm 0.1 \mu\text{g g}^{-1}$ fw at time T1 and T2, respectively. This short-chain aldehyde resulted not significantly different between the two algae at the different sampling times.

289

290 3.2 PUEs (dPUEs) and nutrients in seawater

291 The total average concentrations of dPUEs detected in the seawater in the proximity of the two
292 macroalgae (Fig. S2) highlighted high values (ranging from 99.0 to $287.8 \mu\text{g L}^{-1}$, corresponding to
293 0.9 and $2.7 \mu\text{M}$) and temporal changes, with concentrations significantly lower at times T2 and T3,
294 and higher (about $200 \mu\text{g L}^{-1}$) in the subsequent times (T4-T6) (Table S1). A maximum value was
295 recorded at time T4 for DP and at time T6 for CC (287.8 ± 34.2 and $261.7 \pm 33.6 \mu\text{g L}^{-1}$, respectively).
296 ANOVA carried out on dPUEs concentrations in seawater showed highly significant differences for
297 the factor time ($F=11.8$, $P<0.001$), while no significant differences were found between the two algal
298 species (Table S1). Seawater temperatures during the sampling times ranged between 23 and 26°C ,
299 with salinity values of about 34-39 which were higher at T4-T5 (August-September). Concerning
300 nutrients, the concentration of nitrates was generally low, with a peak of $3.26 \mu\text{M}$ in T2 (June), while
301 phosphates were about 0.1-0.2 μM , with a peak at T5 (0.24 μM) (Table S2).

302

303 3.3 Macroalgal complexity

304 Surface area (S) and fractal dimension (D) were calculated to evaluate the different morphological
305 complexity of the apical parts of the two macroalgae that were analysed in the present work (Fig. S3).
306 Overall S and D were higher in DP than in CC. In CC the surface remained relatively constant along

the sampling period ($20\text{--}25\text{ cm}^2\text{ g}^{-1}$), while in DP S was higher at T5 and T6 than in the previous times (max value $127\text{ cm}^2\text{ g}^{-1}$). As for D, more marked differences between the two algae were found at times T5 and T6, with higher values in DP than in CC (max values 1.84 and 1.73, respectively).

3.4 Microphytobenthic community

In total 25 genera were identified, belonging to 18 orders, and mostly to diatoms and dinoflagellates. Sporadically, cyanobacteria (Cyanophyceae Schaffner, 1909) and juvenile stages of green algae (Chlorophyta Pascher, 1914) were also observed.

Total density was significantly higher in DP ($2608441 \pm 425527\text{ cells g}^{-1}\text{ fw}$) than in CC ($486309 \pm 77877\text{ cells g}^{-1}\text{ fw}$) (Fig. 1; Table S3). Results of ANOVA showed a significant interaction between algae and times ($F=5.75$, $P<0.001$) (Table S3) due to a different temporal trend of density between the two macroalgae. Generally, diatoms represented the most abundant component of the microphytobenthic community, accounting for average values of 96% and 87% of the organisms in DP and CC, respectively.

Microalgal species belonging to the diatom group were classified into 22 genera and 15 orders. Some species remained unidentified, namely undetermined pennate and centric diatoms (Fig. 2). On both macroalgae the most represented order was Naviculales Bessey, 1907, that in DP showed an increasing trend from T1, with an average density of $198287 \pm 81945\text{ cells g}^{-1}\text{ fw}$, to T6, with an average density of $3489032 \pm 1159137\text{ cells g}^{-1}\text{ fw}$ (Table S4), which represented 58% of the epiphytic diatoms (Fig. 2A). For Bacillariales Hendey, 1937, a constant amount of about 60,000 cells $\text{g}^{-1}\text{ fw}$ was observed in DP from T1 to T4, and a maximum at time T5 (average density of $621096 \pm 213575\text{ cells g}^{-1}\text{ fw}$) was recorded, which corresponded to 15% of the total diatoms (Fig. 2A). In terms of average abundance, Lyrellales D.G. Mann and Cocconeidales E.J. Cox, 2015 represented less than 10% of the diatoms (ranging from 25,8831 to 10,865 cells $\text{g}^{-1}\text{ fw}$), while species belonging to other orders (i.e. Surirellales D.G. Mann, 1990, Thalassiophysales D.G. Mann, 1990, Mastogloiales

332 D.G. Mann, Licmophorales Round, 1990, Striatellales Round, 1990, Tabellariales Round, 1990)
 333 occurred with average densities below 100,000 cells g⁻¹ fw.
 334 On CC almost all genera identified had a peak at T5, while a constant average density was observed
 335 from T1 to T4, except for Licmophorales that reached a maximum at T2 (229178 ± 91470 cells g⁻¹
 336 fw), accounting for 33% of the diatoms (Fig. 2B). Specifically, Naviculales, whose concentration
 337 ranged from 25 to 56% of the entire diatom community, occurred with the lowest density of 70,457
 338 ± 31,328 cells g⁻¹ fw at T1 and a maximum average density at T5 (27,5598 ± 89,219 cells g⁻¹ fw).
 339 Bacillariales were about 10% of the total diatoms and increased consistently from T1 (8,869 ± 2,935
 340 cells g⁻¹ fw) to T5 (79,071 ± 18,396 cells g⁻¹ fw), then they decreased at T6 (13,850 ± 4,154 cells g⁻¹
 341 fw). Cocconeidales had a peak at T1 (33,615 ± 16,335 cells g⁻¹ fw), accounting for the 16% of the
 342 total diatoms. All other orders (i.e. Surirellales, Thalassiophysales, Mastogloiales, Licmophorales,
 343 Striatellales, Tabellariales) represented less than 5% of the entire diatom assemblage.
 344 The average density of dinoflagellates was significantly higher on DP than on CC. This group was
 345 recorded especially from T4 to T6 (Table S4) due to the presence of a bloom of the toxic dinoflagellate
 346 *Ostreopsis* cf. *ovata*, whose concentration reached a maximum average density at T6 (364,167 ±
 347 77,126 cells g⁻¹ fw).
 348 The nMDS plot carried out on the microphytobenthic communities showed a clear separation of the
 349 assemblages associated with the two macroalgae (Fig. 3A), but with a different temporal trend.
 350 PERMANOVA supported this pattern resulting in a significant interaction between algae and times
 351 (pseudo-F= 2,2779, P≤0.001; Table S5), and the post-hoc analysis carried out between the two algae
 352 at the various sampling times confirmed the different structure of the two communities at each time.
 353 SIMPER analysis revealed that average similarities for DP was 36% and for CC was 40%, whereas
 354 dissimilarities between the two macroalgae at each time ranged from 66% and 82% (Table S6).
 355 Results showed that the average dissimilarities between the two algae at each time were largely due
 356 to the higher abundances of five orders, namely Naviculales, Lyrellales, Gonyaulacales Taylor, 1980

(i.e. *Ostreopsis cf. ovata*), Bacillariales, and Licmophorales, that were always more abundant on DP than on CC.

3.5 Meiobenthic community

A total of 12 taxa belonging to the meiobenthos were identified, one represented by larval stage Copepoda nauplii (referred from now on as copepod nauplii) defined as the larvae of the meiobenthic copepod species (mainly harpacticoids). Copepods and their nauplii were counted separately in view of their different ecology (Hicks and Coull 1983). Harpacticoida (33.1%) and Nematoda (31.2%) were the dominant taxa, followed by copepod nauplii (25.2%), Gastropoda Cuvier, 1795 (5.4%), Isopoda Latreille, 1817 (1.6%), Polychaeta Grube, 1850 (1.6%), Amphipoda Latreille, 1816 (0.7%) Halacaroidea Cunliffe, 1954 (0.5%), Ostracoda Latreille, 1802 (0.3%), Kinorhyncha Reinhard, 1885 (0.3%), Chironomidae Newman, 1834 (0.2%) and Bivalvia Linnaeus, 1758 (0.1%). Overall, total density (N) and number of taxa (S) at high taxonomic resolution resulted higher on DP than on CC (Figs 4 and S4). Total density was in mean 371 ± 46 and 90 ± 15 N g⁻¹ fw in DP and CC, respectively (Fig. 4), while the number of taxa was on average 5.6 ± 0.2 in DP and 4.4 ± 0.2 in CC (Fig. S4). ANOVA results showed a significant interaction between the factors alga and time ($F=7.06$, $P<0.001$; $F=5.9$, $P<0.001$) for both total density and number of taxa (Table S7), suggesting that the differences between macroalgae were not consistent over time. For total density (Fig. 5), significantly higher values occurred on DP than on CC at corresponding times, except for T1 (Table S7). Moreover, for DP densities increased with a peak at T6 (828 ± 126 N g⁻¹ fw), while for CC the highest density occurred at T5 (217 ± 63 N g⁻¹ fw) and then decreased at T6.

The number of taxa showed the highest value on DP at T2 (7 ± 1) and the lowest on CC at T6 (3 ± 0). It resulted significantly higher on DP than on CC at times T2, T4, and T6, while was lower at T5 (5 ± 1) (Fig. S4; Table S7).

The nMDS analysis of meiofauna communities showed a clear separation between samples belonging to the two macroalgae but, for each alga, the composition of the meiofaunal assemblage changed

383 following a different temporal pattern (Fig. 3B). PERMANOVA supported this pattern with a
384 significant interaction between algae and times (pseudo-F= 2,606, $P \leq 0.001$; Table S8). The
385 assemblages on the two algae resulted significantly different at corresponding sampling times, with
386 the exception of T5 and, within each alga community, the structure changed among times as shown
387 by post hoc comparisons (Table S8).

388 SIMPER analysis revealed that average similarities for each macroalga were 42.8% for DP and 43.7%
389 for CC. Moreover, the dissimilarities between macroalgae at each time ranged from 52% to 82%.
390 Both similarities and dissimilarities were largely due to the variations in abundance of the three
391 dominant taxa: nematodes, copepods, and copepod nauplii (Table S9).

392 In DP average densities of nematodes (Fig. 5A; Table S10) increased from T1 ($57 \pm 22 \text{ N g}^{-1} \text{ fw}$) to
393 a maximum in T6 ($287 \pm 46 \text{ N g}^{-1} \text{ fw}$), as well as the average values of the density of the copepods
394 (Fig. 5B) that showed a similar trend to that of nematodes, with a pick at T6 ($285 \pm 69 \text{ N g}^{-1} \text{ fw}$). The
395 densities of copepod nauplii (Fig. 5C), defined as the larvae of the meiobenthic Copepod species
396 (harpacticoids), showed two peaks at times T2 ($215 \pm 52 \text{ N g}^{-1} \text{ fw}$) and T6 ($217 \pm 34 \text{ N g}^{-1} \text{ fw}$).

397 Conversely, in CC nematodes, copepods and copepod nauplii had lower mean densities at each time
398 when compared with those found in DP (Fig. 5). Both nematodes and copepods densities were very
399 low from time T1 (14 ± 6 and $8 \pm 2 \text{ N g}^{-1} \text{ fw}$) to T4 (10 ± 2 and $14 \pm 3 \text{ N g}^{-1} \text{ fw}$); only at T5 an
400 increase occurred (105 ± 32 and $77 \pm 27 \text{ N g}^{-1} \text{ fw}$) and then a decrease took place at T6 (19 ± 4 and
401 $15 \pm 7 \text{ N g}^{-1} \text{ fw}$). Instead, the mean abundance of copepod nauplii slowly increased until T6 (43 ± 15
402 $\text{N g}^{-1} \text{ fw}$).

403

404 3.6 Harpacticoid community

405 Twelve harpacticoid species were identified from the two macroalgae; they belonged to 11 genera
406 and 11 families. The dominant species was *Heterolaophonte minuta* Boeck, 1873 (32.5%), followed
407 by *Amphiascus parvulus* Claus, 1866 (11.9%), *Harpacticus gracilis* Claus, 1863 (11.6%),
408 *Paradactylopodia brevicornis* Claus, 1866 (10.8%), *Parastenhelia spinosa* Fischer, 1860 (10.2%),

409 and *Ectinosoma melaniceps* Boeck, 1865 (9.6%). The other species ranged from 6.8% (*Porcellidium*
410 *viride* Philippi, 1840) to 0.2% (*Scutelledium longicaudum longicaudum* Philippi, 1840). Results of
411 ANOVA for total density (N), and number of species (S) are shown in table S11. Generally, total
412 densities (N) were significantly higher in DP than in CC at T2, T3, T4, and T6 (Fig. 5B), while the
413 number of harpacticoid species (S) was higher on DP compared to CC at T2 (7 vs 3) and T6 (7 vs 3)
414 (Table S12).

415 The nMDS plot of harpacticoid species showed a clear separation between algae and a different
416 temporal pattern of assemblages on each macroalga (Fig. 3C). These results were supported by the
417 significant interaction between the factors alga and time (PERMANOVA pseudo-F=19.084; P<0.01).
418 In particular, harpacticoid assemblages resulted significantly different between the two algae at each
419 time, with the exception of T1 and T5 (Table S13). A significant gradual change in the community
420 structure took place in DP, whereas on CC significant community structure changes were evident at
421 T5. The details of pairwise comparisons among times for each alga are shown in Table S13.

422 SIMPER results revealed that average dissimilarity between the two algae at each time was largely
423 due to the high abundances of five species, namely *Heterolaophonte minuta*, *Parasthenelia spinosa*,
424 *Paradactylopodia brevicornis*, *Harpacticus gracilis*, and *Porcellidium viride*, that resulted more
425 abundant in DP than in CC (Table S14).

426 The mean abundance over time of these six species is shown in figure 6. In DP, the densities of
427 *Heterolaophonte minuta* showed two peaks at times T4 ($68 \pm 17 \text{ N g}^{-1} \text{ fw}$) and T6 ($82 \pm 20 \text{ N g}^{-1} \text{ fw}$),
428 whereas on CC the densities were always low, with a peak at T3 ($17 \pm 7 \text{ N g}^{-1} \text{ fw}$). In DP the density
429 of *Parasthenelia spinosa* increased at T5 ($17 \pm 4 \text{ N g}^{-1} \text{ fw}$) and T6 ($27 \pm 9 \text{ N g}^{-1} \text{ fw}$), while in CC
430 there was an increase at T5 ($12 \pm 5 \text{ N g}^{-1} \text{ fw}$), then a decrease at T6 ($4 \pm 1 \text{ N g}^{-1} \text{ fw}$). *Paradactylopodia*
431 *brevicornis* showed higher densities at times T3 ($11 \pm 6 \text{ N g}^{-1} \text{ fw}$), T5 ($16 \pm 6 \text{ N g}^{-1} \text{ fw}$), and T6 (31
432 $\pm 11 \text{ N g}^{-1} \text{ fw}$) in DP, while on CC this species was almost always absent, except at T5 ($16 \pm 6 \text{ N g}^{-1}$
433 fw). The average densities of *Harpacticus gracilis* resulted always higher on DP. The densities of
434 *Porcellidium viride* showed the highest values on DP at T3 ($31 \pm 11 \text{ N g}^{-1} \text{ fw}$), then decreased and

435 finally disappeared at T6; conversely, the same pattern occurred for CC, but with lower densities. On
436 DP the density of *Ectinosoma melaniceps* increased at T6 ($19 \pm 6 \text{ N g}^{-1} \text{ fw}$), while on CC this species
437 was almost always absent, except at T5 ($7 \pm 3 \text{ N g}^{-1} \text{ fw}$).

438

439 *3.7 Relationship between the meiofauna and harpacticoid assemblages with macroalgae complexity,* 440 *PUAs, and microphytobenthos*

441 The results of DistLM (Table S15) carried out to analyze the relationships between all the three sets
442 of variables and the meiofauna assemblage showed that for the marginal test the three macroalgae
443 complexity descriptors alone accounted for 26% of the variation in the meiofauna abundance, PUAs
444 for 17% and microphytobenthic taxa for 38%.

445 After fitting complexity descriptors, PUAs and microphytobenthos taxa explained an additional 7%
446 and 13% respectively, resulting in a total variation of 47%. These additional amounts were significant
447 according to the sequential test.

448 The same analysis carried out on the harpacticoid assemblages showed that the three complexity
449 descriptors alone accounted for 19% of the variation in the harpacticoid abundance, PUAs for 13%,
450 and microphytobenthic taxa for 23%.

451 After fitting complexity, PUAs and microphytobenthos taxa explained an additional 9% and 10%
452 respectively, resulting in a total variation of 38%. Only the additional amount of PUAs resulted
453 significant according to the sequential test.

454

455 **4. Discussion**

456 Qualitative and quantitative differences were observed in the PUAs produced by the two macroalgae
457 considered (*Dictyopteris polypodioides*, DP, and *Cystoseira compressa*, CC), in agreement with
458 previous results obtained for the same species (Pezzolesi et al., 2021). DP produced, in fact, higher
459 average concentrations than CC, but also a variety of long-chain compounds, such as hexadecatrienal
460 (C16:3), hexadecatetraenal (C16:4) and tetradecapentaenal (C14:5), while *C. compressa* produced

the short-chain hexadienal (C6:2) as main compound, demonstrating the ability of macroalgae to produce short, middle and/or long-chain aldehydes (Kajiwara et al., 1996; Akakabe et al., 2003; Pezzolesi et al., 2021). These results highlighted that PUAs profile could be a fingerprint for each algal species, since the same compounds were consistently detected regardless of the sampling period, while their relative and total amount may vary depending on environmental conditions or morphotypes (Alsufyani et al., 2014; Pezzolesi et al., 2021). In addition, since the apical parts of both algae are those in which growth actively takes place, they are the most metabolically active parts, and therefore presumably also those in which the greatest production of PUAs takes place. Dissolved PUAs (dPUAs) concentrations in proximity of the macroalgae was high (in the order of μM), especially when compared with results of previous studies carried out in an Adriatic planktonic community (Ribalet et al., 2014), but are in accordance with values previously recorded in the same site in other studies (Bartual et al., 2020; Pezzolesi et al., 2021) or hypothesized to occur in proximity of the PUA producers (Ribalet et al., 2008; Bartual et al., 2018). These high concentrations are reasonable considering that the sampling area is colonized by a well structured phytobenthic community (macro- and microalgal), thus dPUAs derive from the contribution of the different algal species. Additionally, the reduced hydrodynamism of this benthic site causes a low dispersion of any secondary metabolites produced by the various organisms.

Seawater parameters (temperature, salinity and nutrients) were within the range previously reported for the Piscinetta site (e.g. Accoroni et al., 2012), and confirmed the seasonal variability associated to this shallow inlet, that is subjected to a moderate anthropic impact (mainly in the form of summer tourism, as it is a popular site for swimming in the summer months). Results showed a predominance of diatoms in the microphytobenthic community, as usually reported in the Adriatic Sea, even in planktonic communities (e.g. Accoroni et al., 2016). Dinoflagellates were present at low densities, except during the bloom of *Ostreopsis* cf. *ovata*, which was the main dinoflagellate and showed its typical blooming trend, with maximum abundances recorded in late summer (September) and within ranges previously observed (e.g. Gémin et al., 2020).

487 The main diatom genera found in the present study (e.g. *Navicula*, *Cylindrotheca*, *Lyrella*, *Cocconeis*,
 488 *Gyrosigma* A.H. Hassall, 1845, *Licmophora* C. Agardh, 1827, *Nitzschia*, *Mastogloia* Thwaites ex
 489 W.Smith, 1856, *Striatella* C. Agardh, 1832, *Coscinodiscus* Ehrenberg, 1839) are among the most
 490 common on Mediterranean macroalgae and in the microphytobenthos, particularly in the Adriatic Sea
 491 (Carnicer et al., 2015; Accoroni et al., 2016; Rogelja et al., 2016; Pennesi and Danovaro, 2017;
 492 Ternon et al., 2020). Species belonging to the orders Naviculales and Lyrellales on DP, to the
 493 Licmophorales on CC and, generally, to centric diatoms on both algae, showed an inverse trend (low
 494 density at T1 and high at subsequent times) in relation to the production of PUAs by macroalgae.
 495 These results could be ascribed to a negative effect of these compounds, as also demonstrated by
 496 Ribalet et al. (2007) for some planktonic species. Long-chain aldehydes, such as C14-C16, can induce
 497 a stronger growth inhibition than short-chain PUAs, probably due to longer alkyl chains that increase
 498 the reactivity of the molecules (Adolph et al., 2003).
 499 Since many diatom species recorded in the present study, as those belonging to the genus
 500 *Cylindrotheca* spp. and *Nitzschia* spp., are themselves among the main PUAs producers (Wichard et
 501 al., 2005a; Lavrentyev et al., 2015; Pezzolesi et al., 2017; C3zar et al., 2018), they may have
 502 developed different sensitivities and/or tolerances to these compounds in a species-specific way.
 503 Studies have also shown a different susceptibility based on the life cycle, with more resistant juvenile
 504 cells and more sensitive stationary phase cells (Ribalet et al., 2007; Leflaive and Ten-Hage, 2009)
 505 and based also on other factors, such as cell size, wall properties and lipid content. In particular,
 506 species with a well-structured and mineralized cell wall, a low surface-volume ratio and a certain
 507 lipid content can limit the ability of these compounds to penetrate the cell. Centric diatoms could
 508 potentially be more sensitive to PUAs, as those found in the present study (*Chaetoceros* spp.
 509 Ehrenberg, 1844, *Coscinodiscus* spp., *Guinardia* spp. H. Peragallo, 1892 and *Rhizosolenia* spp.
 510 Brightwell, 1858) are not listed among the species able to produce these compounds (Wichard et al.,
 511 2005).

512 *Ostreopsis* cf. *ovata* does not seem to be negatively influenced, in terms of abundance, by the
513 production of PUAs by macroalgae. Conversely, macroalgal complexity seems to explain the different
514 abundances found on CC and DP. The presence of a rigid cell wall and the high biovolume could
515 partially explain the apparent lower sensitivity of *Ostreopsis* cells to PUAs compared to other species,
516 in addition to the protective role that can be offered by the mucilaginous layer produced by *O.* cf.
517 *ovata*, which provides an additional barrier against the substances dissolved in the water column
518 (Allen et al., 2016). Similarly, benthic dinoflagellates have shown a higher resistance than planktonic
519 ones to potential allelochemicals (Ben Gharbia et al., 2017).

520 Regarding the meiofauna, 12 major taxa associated with the two macroalgae were found, with a total
521 density recorded in some samples of over 1300 individuals per gram of alga and within the range
522 found in previous studies (Jarvis and Seed, 1996). Harpacticoid copepods, together with their nauplii,
523 were the most abundant taxon, comprising 58% of the total meiofauna, followed by Nematodes that
524 were 33%, as reported in previous studies (Carlo Heip, Magda Vincx, 1985; De Troch et al., 2005).

525 By comparing the results obtained for the two macroalgae, the average abundances of total meiofauna
526 and the number of taxa resulted higher on DP than CC, as well as the abundances of copepods and
527 nematodes. As a result, the community structure resulted different between the two macroalgae and
528 showed a different temporal pattern, although they were subjected to the same environmental
529 characteristics in terms, for instance, of hydrodynamism and tides. DP and CC are different for life
530 cycle, thallus structure and production of compounds. In particular, DP is formed by ribbon-like
531 fronds, with very irregular and proliferating edges, on which numerous meiobenthic organisms can
532 settle. In the area where sampling was conducted, the species persists in its fully developed habit for
533 most of spring and summer. CC is a highly branched, leathery macrophyte; when fully grown, in the
534 study area it may reach 1-1.5 m in height. For a large-sized, habitat-forming species, its growth is
535 relatively fast, especially if compared with other Mediterranean fucalean brown algae. However, in
536 the area of the Passetto its full development is limited to a quite restricted period of the year (from
537 May to mid July); by mid summer this species loses most of its branches and persists in a more

538 reduced habit, with a few short branches. This means that most of its fronds occur in the field for a
539 shorter time and are not available long enough to allow the settlement of a very diverse epiphytic
540 community. The shorter temporal availability of this substrate also means that there will be a lower
541 accumulation of sediment and detritus, a potential source of food for meiofaunal organisms (Hicks,
542 1980; Gibbons, 1988; Frame et al., 2007). Both species have apical growth, determined by divisions
543 of a group of meristematic cells in DP and a single apical cell in CC. Therefore, the apical parts are
544 the youngest parts of the thallus, on which sediment and epiphytes had less time to settle.

545 Taking into consideration all these aspects, our results suggest that the morphological complexity of
546 DP may affect total number of individuals, but also the associated species that have evolved
547 morphological adaptations necessary for the adhesion to a thallus of this type (Taylor and Cole, 1994;
548 Chemello and Milazzo, 2002). Although macrophytes are not true fractal objects, estimates of
549 complexity using tools of fractal geometry have proved to be a useful approach for quantifying and
550 separating effects of habitat architecture from those of habitat quantity (Gee and Warwick, 1994;
551 McAbendroth et al., 2005; Hooper and Davenport, 2006). However, fractal measures performed for
552 not truly fractal objects are just an estimate of complexity for a given scale.

553 The putative role of PUAs on structuring meiofauna may be reflected by the low density of nauplii,
554 copepods, and nematodes at time T1 recorded on both macroalgae, which could be related to the high
555 concentration of PUAs in this period, as observed also for microphytobenthos. Moreover, host
556 specificity may be supposed, according to what determined by Bates and DeWreede (2007), that
557 found specific chemical, structural and morphological characteristics of the algal species.

558 Harpacticoids have a number of features that make them an attractive group of benthic organisms in
559 which PUA toxicity responses could be investigated, such as abundance, ecological importance, and
560 short generation cycles (Raisuddin et al., 2007). In this study, harpacticoids associated with the two
561 macroalgae consisted of 12 species. Even if the number of species was relatively low compared to
562 results of previous studies (Hicks, 1977b; Arroyo et al., 2006), it is interesting to note that the 12
563 species belonged to 12 genera and 11 families, so showing a high taxonomic distinctness. Moreover,

no species was present exclusively on a single macroalga; only the relative abundance of individual species was different between the two macroalgae, with *Heterolaophonte minuta* as the dominant species on both. Almost all species showed low abundance at T1, potentially suggesting a role of PUAs, as previously postulated for meiofauna and microphytobenthos. To our knowledge, only one study was carried out in laboratory to analyze the effects of these compounds on the harpacticoid *Tisbe holothuriae* (Taylor et al., 2007); while, other studies carried out on planktonic ecosystem have shown deleterious effects of PUAs produced by diatoms on the reproduction of calanoid copepods (e.g. *Temora stylifera*, *Calanus helgolandicus*), which feed on them (Ianora et al., 2003, 2012; Miralto et al., 1999), as well as apoptosis in maturing oocytes (Poulet et al., 2007a) during embryo development (Romano et al., 2003) and in newly hatched nauplii (Ianora et al., 2004b).

It has to be considered that the present study was carried out in the field, so the link between PUAs effects on the community structure is more difficult to evaluate, due to the variability of PUAs production both by diatoms (Wichard et al., 2005b) and macroalgae (Pezzolesi et al., 2021), of the copepod sensitivity (Ianora et al., 2003; Sommer, 2009) and to the detoxification ability developed by certain species of copepods (e.g. Taylor et al., 2007; Wichard et al., 2008). As attested by Taylor et al. (2007), the benthos tends to be a more stressful environment compared to planktonic, where rapid fluctuations in physical conditions occur and both natural and anthropogenic toxins can accumulate at high levels within the sediments between the algal fronds. Since benthic organisms must be highly adapted to survive in such a harsh environment, it is not unreasonable to speculate that harpacticoids may have a more developed detoxification system than planktonic calanoid copepods, thus being better equipped to resist to the toxic effect of oxylipins. Indeed, a number of candidate detox genes were found in an analysis of 686 sequence tags expressed by *Tigriopus japonicus* (Lee et al., 2005, 2008). Therefore, it is possible that the species found during the present study have developed effective and efficient detoxification strategies to survive in an environment such as the Passetto area. The different community structure between the two macroalgae and the temporal changes could be also explained by the ecological and trophic role of the various species,

590 and by the different morphological evolution of the two macroalgae during the sampling time. This
 591 is an important aspect to consider in the case of this study, as we sampled apical parts of thalli of CC
 592 and DP, that may be not representative of the entire thalli, as are the youngest parts on which epiphytes
 593 had less time to settle and also those most exposed to light. In particular, four species (*Porcellidium*
 594 *viride*, *Parastenhelia spinosa*, *Heterolaophonte minuta*, and *Harpacticus gracilis*) belonged to the
 595 phytal group sensu strictu, and two (*Paradactylopodia brevicornis*, and *Ectinosoma melaniceps*)
 596 belonged to migrator and cosmopolitan group (Mascart et al., 2015). *E. melaniceps* is a tolerant
 597 eurytopic species, which presumably is not affected by the biochemical compounds produced by
 598 macroalgae, as for example reported for the green alga *Ulva lactuca* (Hicks, 1980). *P. brevicornis* is
 599 cosmopolitan species that have a wide distribution range and was found in different habitats, therefore
 600 able to adapt to a large number of different environmental conditions (Hicks, 1980). Conversely, *P.*
 601 *viride*, *H. minuta* and *H. gracilis* are endemic species of the Mediterranean Sea with certain
 602 morphological characteristics that allow them to live adhering to macroalgal surfaces; therefore, they
 603 could either be affected by the effect of the various compounds produced by both macroalgae and
 604 microphytobenthos, or adapt to the various environmental conditions.

605 The DistLm procedure highlighted that the observed differences in the meiofauna and harpacticoid
 606 community structure between the two macroalgal species could be mainly explained by
 607 microphytobenthic main taxa, after fitting macroalgal complexity.

608 Microalgae, in fact, are at the base of the food web and provide energy for all the trophic levels above
 609 them; thus, these results confirm their important role as primary food source of essentially all marine
 610 food chains producers. In fact, most of the epifaunal taxa directly graze upon the macroalgae or the
 611 epiphytes for their food source, thus variations in the epiphytic algal density and composition between
 612 two macroalgal species may influence the abundance of associated epifauna (Gestoso et al., 2010).

613 Additionally, the macroalgal complexity, rather than PUAs production alone, resulted an important
 614 component influencing the community structure. The role of the shape and structural complexity of
 615 macroalgae in determining the abundance patterns and size structure of the epiphytic organisms is

616 supported in literature, either for epifaunal and epiphytic assemblages (e.g. Chemello and Milazzo,
617 2002; McAbendroth et al., 2005; Cacabelos et al., 2010). To better investigate the effect of macroalgal
618 complexity and PUAs composition on variations of the microphytobenthos at some representative
619 genera levels, on main meiofauna taxa, and on harpacticoid species Pearson correlation analysis was
620 used (Fig. 7). Among PUAs, long-chain compounds (i.e. C14-C16), when compared with the short
621 one (i.e. C6:2), showed higher effects on the abundances of some representative microalgae (i.e.
622 *Navicula* spp., *Lyrella* spp., *Cocconeis* spp., *Cylindrotheca* spp./*Nitzschia* spp., and *Ostreopsis* cf.
623 *ovata*), on the main meiobenthos taxa (i.e. Nematoda, Copepoda, copepod nauplii) and on
624 harpacticoid species (i.e. *H. minuta*, *H. gracilis*, *P. viride*). Results are in agreement with laboratory
625 studies performed to investigate the responses of allelochemicals, such as PUAs, on target organisms,
626 including copepods and algae (Ianora et al., 2004a; Taylor et al., 2007; Adolph et al., 2004; Caldwell
627 et al., 2004; Ribalet et al., 2007; Pichierri et al., 2016). Macroalgal complexity, either in terms of
628 fractal dimension and area, is highly correlated and showed significant relationships with almost all
629 the taxa considered, except with *P. viride*, potentially due to its ecological role, being a phytal species
630 able to colonize a variety of macroalgae thanks to morphological adaptations that has evolved to
631 attach to morphologically diverse thalli (Hicks, 1980).

632 It is important to point out that several metabolites, as well as PUAs, are known to be produced by
633 macroalgae, including some (e.g. diterpenoids, polyphenols) that inhibit the growth of microalgae,
634 bryozoan or other benthic species (Ternon et al., 2020), thus their contribution to the observed
635 dynamics could not be excluded. Additionally, the epiphytic community that colonize the macroalgal
636 surface may contribute either to the surface metabolome, adding inhibitory effects on the co-occurring
637 species (Monti and Cecchin, 2012) or to the higher trophic levels, and directly interacting with the
638 epifaunal, thus adding complexity to the understanding of the relationships among these organisms
639 and to the role of PUAs. Finally, toxic microalgae such as *O. cf. ovata*, that was found to bloom
640 during the sampling period, have been reported to affect copepods, and particularly nauplii (Guidi-
641 Guilvard et al., 2012), with different sensitivities among species (Pavaux et al., 2019).

642 Taking into consideration all these aspects, PUAs production by macroalgae, together with their
643 complexity, resulted one of the main factors involved in the benthic community structure dynamics,
644 but it is not enough to explain the differences in the microphytobenthos and meiofana assemblages.

645

646 5. Conclusion

647 Epiphytic communities on the two macroalgae highlighted a clear separation of the meiofauna and
648 microphytobenthos assemblages with different temporal trends. The average dissimilarities were due
649 to several microalgal orders, namely Naviculales, Lyrellales, Gonyaulacales (i.e. *Ostreopsis* cf.
650 *ovata*), Bacillariales, and Licmophorales, and to the three meiofauna dominant taxa (nematodes,
651 copepods, and copepod nauplii) that were always more abundant on DP than on CC. Particularly,
652 average dissimilarities of harpacticoid copepods were largely due to the abundances of five species,
653 namely *Heterolaophonte minuta*, *Parastenhelia spinosa*, *Paradactylopodia brevicornis*, *Harpacticus*
654 *gracilis*, *Porcellidium viride*, and *Ectinosoma melaniceps*. Generally, variations in the meiofauna and
655 harpacticoid abundances were mainly due to macroalgal complexity variables and
656 microphytobenthos, while a minor contribution was due to PUAs. Results documented that i)
657 microphytobenthos resulted to affect the meiofauna population dynamics, in particular the
658 harpacticoid assemblages, attesting the role of these organisms as primary food source of essentially
659 all marine food chains producers, being at the base of the food web and providing energy for all the
660 trophic levels above them, ii) the macroalgal complexity rather than PUAs production alone could be
661 a major trigger of the community structure. PUAs effects, in fact, resulted species-specific, thus
662 affecting some grazers instead of the entire community structure, as demonstrated also by Pearson's
663 ρ correlations between taxa abundances and several macroalgal parameters. Among PUAs, long-
664 chain compounds (i.e. C14-C16), with respect to the short one (i.e. C6:2), showed higher effects on
665 the abundances of some representative microalgal genera, harpacticoid species and on the main
666 meiobenthos taxa.

667 Since several of the epiphytic diatom species found, in addition to macroalgae, can produce PUAs,
668 the understanding of the effects of these compounds on the community structure and on the
669 relationships among taxa in field studies are complicated, thus opening to further in-depth
670 investigations in simplified systems (i.e. microcosms).

671

672 **Acknowledgements**

673 This research did not receive any specific grant from funding agencies in the public, commercial, or
674 not-for-profit sectors. The authors wish to thank Elisa Bombardiere for help with microphytobenthos
675 analysis as part of her master's thesis; Prof. Victor Ugo Ceccherelli for help with harpacticoid species
676 identification; Mara Simonazzi and Dr. Franca Guerrini for help during some samplings, and Prof.
677 Cecilia Totti for advices about taxonomy of diatoms. The authors would like to thank the anonymous
678 reviewers for their careful reading of the manuscript and the insightful comments.

679

680 **Note**

681 The authors declare no competing financial interest.

682

683 **CRedit author statement**

684 Denise Lenzo: Data curation, Investigation, Formal analysis, Writing- Original draft preparation.
685 Laura Pezzolesi: Conceptualization, Methodology, Data curation, Resources, Writing- Original draft
686 preparation, Funding acquisition. Chiara Samorì: Data curation, Writing - Review & Editing. Fabio
687 Rindi: Resources, Reviewing and Editing. Andrea Pasteris: Supervision, Reviewing and Editing.
688 Rossella Pistocchi: Funding acquisition, Reviewing and Editing. Marina Antonia Colangelo:
689 Conceptualization, Methodology, Formal analysis, Software, Writing- Original draft preparation,
690 Supervision.

691

692 **References**

693 Accoroni, S., Colombo, F., Pichierri, S., Romagnoli, T., Marini, M., Battocchi, C., Penna, A., Totti,
694 C., 2012. Ecology of *Ostreopsis cf. ovata* Blooms in the Northwestern Adriatic Sea .
695 Cryptogam. Algal. 33, 191–198. <https://doi.org/10.7872/crya.v33.iss2.2011.191>

696 Accoroni, S., Romagnoli, T., Pichierri, S., Totti, C., 2016. Effects of the bloom of harmful benthic
697 dinoflagellate *Ostreopsis cf. ovata* on the microphytobenthos community in the northern
698 Adriatic Sea. Harmful Algae 55, 179–190. <https://doi.org/10.1016/j.hal.2016.03.003>

699 Adolph, S., Bach, S., Blondel, M., Cueff, A., Moreau, M., Pohnert, G., Poulet, S.A., Wichard, T.,
700 Zuccaro, A., 2004. Cytotoxicity of diatom-derived oxylipins in organisms belonging to
701 different phyla. J. Exp. Biol. 207, 2935–2946. <https://doi.org/10.1242/jeb.01105>

702 Adolph, S., Poulet, S.A., Pohnert, G., 2003. Synthesis and biological activity of $\alpha,\beta,\gamma,\delta$ -unsaturated
703 aldehydes from diatoms. Tetrahedron 59, 3003–3008. [https://doi.org/10.1016/S0040-](https://doi.org/10.1016/S0040-4020(03)00382-X)
704 [4020\(03\)00382-X](https://doi.org/10.1016/S0040-4020(03)00382-X)

705 Agostini-Costa, T.S., Vieira, R.F., Bizzo, H.R., Silveira, D., Gimenes, M.A., 2012. Secondary
706 metabolites, in: Dhanarasu, D.S. (Ed.), Chromatography and Its Applications. pp. 131–164.
707 <https://doi.org/10.1017/s026972700000614x>

708 Akakabe, Y., Matsui, K., Kajiwara, T., 2003. 2,4-Decadienals are produced via (R)-11-HPITE from
709 arachidonic acid in marine green alga *Ulva conglobata*. Bioorganic Med. Chem. 11, 3607–
710 3609. [https://doi.org/10.1016/S0968-0896\(03\)00364-X](https://doi.org/10.1016/S0968-0896(03)00364-X)

711 Allen, J.L., Ten-Hage, L., Leflaive, J., 2016. Allelopathic interactions involving benthic
712 phototrophic microorganisms. Environ. Microbiol. Rep. 8, 752–762.
713 <https://doi.org/10.1111/1758-2229.12436>

714 Alsufyani, T., Engelen, A.H., Diekmann, O.E., Kuegler, S., Wichard, T., 2014. Prevalence and
715 mechanism of polyunsaturated aldehydes production in the green tide forming macroalgal
716 genus *Ulva* (Ulvales, Chlorophyta). Chem. Phys. Lipids 183, 100–109.
717 <https://doi.org/10.1016/j.chemphyslip.2014.05.008>

718 Anderson, M., Gorley, R., Clarke, K., 2008. PERMANOVA for PRIMER: guide to software and

719 statistical methods. PRIMER–E Ltd., Plymouth.

720 Anderson, M.J., 2001. A new method for non-parametric multivariate analysis of variance. *Austral*

721 *Ecol.* 26, 32–46. <https://doi.org/10.1046/j.1442-9993.2001.01070.x>

722 Anderson, M.J., Connell, S.D., Gillanders, B.M., Diebel, C.E., Blom, W.M., Saunders, J.E.,

723 Landers, T.J., 2005. Relationships between taxonomic resolution and spatial scales of

724 multivariate variation. *J. Anim. Ecol.* 74, 636–646. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2656.2005.00959.x)

725 [2656.2005.00959.x](https://doi.org/10.1111/j.1365-2656.2005.00959.x)

726 Anderson, M.J., Robinson, J., 2001. Permutation tests for linear models. *Aust. New Zeal. J. Stat.* 43,

727 75–88. <https://doi.org/10.1111/1467-842X.00156>

728 Arroyo, N.L., Maldonado, M., Walters, K., 2006. Within- and between-plant distribution of

729 harpacticoid copepods in a North Atlantic bed of *Laminaria ochroleuca*. *J. Mar. Biol. Assoc.*

730 *United Kingdom* 86, 309–316. [https://doi.org/DOI: 10.1017/S0025315406013166](https://doi.org/DOI:10.1017/S0025315406013166)

731 Bartual, A., Hernanz-Torrijos, M., Sala, I., Ortega, M.J., González-García, C., Bolado-Penagos, M.,

732 López-Urrutia, A., Romero-Martínez, L., Lubián, L.M., Bruno, M., Echevarría, F., García,

733 C.M., 2020. Types and distribution of bioactive polyunsaturated aldehydes in a gradient from

734 mesotrophic to oligotrophic waters in the Alborán Sea (Western Mediterranean). *Mar. Drugs*

735 18, 159. <https://doi.org/10.3390/md18030159>

736 Bartual, A., Morillo-García, S., Ortega, M.J., Cózar, A., 2018. First report on vertical distribution of

737 dissolved polyunsaturated aldehydes in marine coastal waters. *Mar. Chem.* 204, 1–10.

738 <https://doi.org/10.1016/j.marchem.2018.05.004>

739 Bates, C.R., DeWreede, R.E., 2007. Do changes in seaweed biodiversity influence associated

740 invertebrate epifauna? *J. Exp. Mar. Bio. Ecol.* 344, 206–214.

741 <https://doi.org/10.1016/j.jembe.2007.01.002>

742 Ben Gharbia, H., Yahia, O.K.D., Cecchi, P., Masseret, E., Amzil, Z., Herve, F., Rovillon, G., Nouri,

743 H., M’Rabet, C., Couet, D., Triki, H.Z., Laabir, M., 2017. New insights on the species-specific

744 allelopathic interactions between macrophytes and marine HAB dinoflagellates. *PLoS One* 12.

745 <https://doi.org/10.1371/journal.pone.0187963>

746 Boxshall, G.A., Halsey, S.H., 2004. An introduction to copepod diversity. Ray Society, Andover.

747 Cacabelos, E., Olabarria, C., Incera, M., Troncoso, J.S., 2010. Effects of habitat structure and tidal
748 height on epifaunal assemblages associated with macroalgae. *Estuar. Coast. Shelf Sci.* 89, 43–
749 52. <https://doi.org/10.1016/j.ecss.2010.05.012>

750 Caldwell, G.S., Bentley, M.G., Olive, P.J.W., 2004. First evidence of sperm motility inhibition by
751 the diatom aldehyde 2E, 4E-decadienal. *Mar. Ecol. Prog. Ser.* 273, 97–108.
752 <https://doi.org/10.3354/meps273097>

753 Carlo Heip, Magda Vincx, G.V., 1985. The ecology of marine nematodes. *Oceanogr,Mar.Biol.*
754 <https://doi.org/10.1111/j.1432-1033.1975.tb20978.x>

755 Carnicer, O., Guallar, C., Andree, K.B., Diogène, J., Fernández-Tejedor, M., 2015. *Ostreopsis* cf.
756 *ovata* dynamics in the NW Mediterranean Sea in relation to biotic and abiotic factors. *Environ.*
757 *Res.* 143, 89–99. <https://doi.org/10.1016/j.envres.2015.08.023>

758 Chardy, P., Dauvin, J.C., 1992. Carbon flows in a subtidal fine sand community from the western
759 English Channel: a simulation analysis. *Mar. Ecol. Prog. Ser.* 81, 147–161.
760 <https://doi.org/10.3354/meps081147>

761 Chemello, R., Milazzo, M., 2002. Effect of algal architecture on associated fauna: Some evidence
762 from phytal molluscs. *Mar. Biol.* 140, 981–990. <https://doi.org/10.1007/s00227-002-0777-x>

763 Clarke, K.R., 1993. Non-parametric multivariate analyses of changes in community structure. *Aust.*
764 *J. Ecol.* 18, 117–143. <https://doi.org/10.1111/j.1442-9993.1993.tb00438.x>

765 Clarke, K.R., Gorley, R.N., 2015. Getting started with PRIMER v7. PRIMER-E. Prim. Plymouth
766 2–20.

767 Coull, B.C., 1999. Role of meiofauna in estuarine soft-bottom habitats. *Austral Ecol.* 24, 327–343.
768 <https://doi.org/10.1046/j.1442-9993.1999.00979.x>

769 Coull, B.C., Wells, J.B.J., 1983. Refuges from Fish Predation : Experiments with Phytal Meiofauna
770 from the New Zealand Rocky Intertidal Published by : Wiley on behalf of the Ecological

771 Society of America Stable URL : <https://www.jstor.org/stable/1937513> REFERENCES Linked
 772 references are av. Ecology 64, 1599–1609.

773 Cózar, A., Morillo-García, S., Ortega, M.J., Li, Q.P., Bartual, A., 2018. Macroecological patterns of
 774 the phytoplankton production of polyunsaturated aldehydes. Sci. Rep. 8, 1–10.
 775 <https://doi.org/10.1038/s41598-018-29787-8>

776 De Troch, M., Vandepitte, L., Raes, M., Suárez-Morales, E., Vincx, M., 2005. A field colonization
 777 experiment with meiofauna and seagrass mimics: Effect of time, distance and leaf surface area.
 778 Mar. Biol. 148, 73–86. <https://doi.org/10.1007/s00227-005-0062-x>

779 Edler, L., Elbrächter, M., 2010. No Title, in: B. Karlson, C. Cusack, E.B. (Ed.), Microscopic and
 780 Molecular Methods for Quantitative Phytoplankton Analysis. UNESCO, Paris, pp. 13–20.

781 Fistarol, G.O., Legrand, C., Selander, E., Hummert, C., Stolte, W., Granéli, E., 2004. Allelopathy in
 782 *Alexandrium* spp.: Effect on a natural plankton community and on algal monocultures. Aquat.
 783 Microb. Ecol. 35, 45–56. <https://doi.org/10.3354/ame035045>

784 Frame, K., Hunt, G., Roy, K., 2007. Intertidal meiofaunal biodiversity with respect to different algal
 785 habitats: A test using phytal ostracodes from Southern California. Hydrobiologia 586, 331–
 786 342. <https://doi.org/10.1007/s10750-007-0707-5>

787 Gee, J.M., Warwick, R.M., 1994. Metazoan community structure in relation to the fractal
 788 dimensions of marine macroalgae. Mar. Ecol. Prog. Ser. 103, 141–150.
 789 <https://doi.org/10.3354/meps103141>

790 Gémin, M.P., Réveillon, D., Hervé, F., Pavaux, A.S., Tharaud, M., Séchet, V., Bertrand, S., Lemée,
 791 R., Amzil, Z., 2020. Toxin content of *Ostreopsis* cf. *ovata* depends on bloom phases, depth and
 792 macroalgal substrate in the NW Mediterranean Sea. Harmful Algae 92, 101727.
 793 <https://doi.org/10.1016/j.hal.2019.101727>

794 Gestoso, I., Olabarria, C., Troncoso, J.S., 2010. Variability of epifaunal assemblages associated
 795 with native and invasive macroalgae. Mar. Freshw. Res. 61, 724–731.
 796 <https://doi.org/10.1071/MF09251>

797 Gibbons, M.J., 1988. The impact of sediment accumulations, relative habitat complexity and
 798 elevation on rocky shore meiofauna. *J. Exp. Mar. Bio. Ecol.* 122, 225–241.
 799 [https://doi.org/10.1016/0022-0981\(88\)90125-6](https://doi.org/10.1016/0022-0981(88)90125-6)

800 Gibbons, M.J., Griffiths, C.L., 1986. A comparison of macrofaunal and meiofaunal distribution and
 801 standing stock across a rocky shore, with an estimate of their productivities. *Mar. Biol.* 93,
 802 181–188. <https://doi.org/10.1007/BF00508255>

803 Giere, O., 2009. Meiobenthology. *The Microscopic Motile Fauna of Aquatic Sediments*.
 804 Meiobenthology Microsc. Motile Fauna Aquat. Sediments 1–527. [https://doi.org/10.1007/978-](https://doi.org/10.1007/978-3-540-68661-3)
 805 [3-540-68661-3](https://doi.org/10.1007/978-3-540-68661-3)

806 Grima, E.M., Pérez, J.A.S., Camacho, F.G., Medina, A.R., Giménez, A.G., López Alonso, D., 1995.
 807 The production of polyunsaturated fatty acids by microalgae: from strain selection to product
 808 purification. *Process Biochem.* 30, 711–719. [https://doi.org/10.1016/0032-9592\(94\)00047-6](https://doi.org/10.1016/0032-9592(94)00047-6)

809 Guidi-Guilvard, L.D., Gasparinl, S., Lemée, R., 2012. The negative impact of *Ostreopsis cf. ovata*
 810 on phytal meiofauna from the coastal NW mediterranean. *Cryptogam. Algal.* 33, 121–128.
 811 <https://doi.org/10.7872/crya.v33.iss2.2011.121>

812 Guiry, M.D., Guiry, G.M., 2021. *AlgaeBase*. – World-wide electronic publication, National
 813 University of Ireland, Galway [WWW Document].
 814 <<https://www.algaebase.org/search/species/>.

815 Hall, M.O., Bell, S.S., 1993. Meiofauna on the seagrass *Thalassia testudinum*: population
 816 characteristics of harpacticoid copepods and associations with algal epiphytes. *Mar. Biol. Int.*
 817 *J. Life Ocean. Coast. Waters* 116, 137–146. <https://doi.org/10.1007/BF00350741>

818 Hicks, G.R.F., 1980. Structure of phytal harpacticoid copepod assemblages and the influence of
 819 habitat complexity and turbidity. *J. Exp. Mar. Bio. Ecol.* 44, 157–192.
 820 [https://doi.org/10.1016/0022-0981\(80\)90151-3](https://doi.org/10.1016/0022-0981(80)90151-3)

821 Hicks, G.R.F., 1977a. Observations on substrate preference of marine phytal harpacticoids
 822 (Copepoda). *Hydrobiologia* 56, 7–9.

823 Hicks, G.R.F., 1977b. Species associations and seasonal population densities of marine phytal
824 harpacticoid copepods from Cook Strait. *New Zeal. J. Mar. Freshw. Res.* 11, 621–643.
825 <https://doi.org/10.1080/00288330.1977.9515702>

826 Hooper, G.J., Davenport, J., 2006. Epifaunal composition and fractal dimensions of intertidal
827 marine macroalgae in relation to emersion. *J. Mar. Biol. Assoc. United Kingdom* 86, 1297–
828 1304. <https://doi.org/10.1017/S0025315406014329>

829 Ianora, A., Miralto, A., Poulet, S.A., Carotenuto, Y., Buttino, I., Romano, G., Casotti, R., Pohnert,
830 G., Wichard, T., Colucci-D'Amato, L., Terrazzano, G., Smetacek, V., 2004a. Aldehyde
831 suppression of copepod recruitment in blooms of a ubiquitous planktonic diatom. *Nature* 429,
832 403–407. <https://doi.org/10.1038/nature02526>

833 Ianora, A., Miralto, A., Romano, G., 2012. Antipredatory Defensive Role of Planktonic Marine
834 Natural Products, in: Fattorusso, E., Gerwick, W.H., Tagliatela-Scafati, O. (Eds.), *Handbook*
835 *of Marine Natural Products*. Springer Netherlands, Dordrecht, pp. 711–748.
836 https://doi.org/10.1007/978-90-481-3834-0_13

837 Ianora, A., Poulet, S.A., Miralto, A., 2003. The effects of diatoms on copepod reproduction: A
838 review. *Phycologia* 42, 351–363. <https://doi.org/10.2216/i0031-8884-42-4-351.1>

839 Ianora, A., Turner, J.T., Esposito, F., Carotenuto, Y., D'Ippolito, G., Romano, G., Fontana, A.,
840 Guisande, C., Miralto, A., 2004b. Copepod egg production and hatching success is reduced by
841 maternal diets of a non-neurotoxic strain of the dinoflagellate *Alexandrium tamarense*. *Mar.*
842 *Ecol. Prog. Ser.* 280, 199–210. <https://doi.org/10.3354/meps280199>

843 Jarvis, S.C., Seed, R., 1996. The meiofauna of *Ascophyllum nodosum* (L.) Le Jolis:
844 Characterization of the assemblages associated with two common epiphytes. *J. Exp. Mar. Bio.*
845 *Ecol.* 199, 249–267. [https://doi.org/10.1016/0022-0981\(95\)00184-0](https://doi.org/10.1016/0022-0981(95)00184-0)

846 Jüttner, F., Messina, P., Patalano, C., Zupo, V., 2010. Odour compounds of the diatom *Cocconeis*
847 *scutellum*: Effects on benthic herbivores living on *Posidonia oceanica*. *Mar. Ecol. Prog. Ser.*
848 400, 63–73. <https://doi.org/10.3354/meps08381>

- 849 Kajiwara, T., Matsui, K., Akakabe, Y., 1996. Biogenesis of volatile compounds via oxylipins in
850 edible seaweeds, in: Takeoka, G.R., Teranishi, R., Williams, P.J., Kobayashi, A. (Ed.), ACS
851 Symposium Series. Washington D. C., pp. 146–166. [https://doi.org/10.1021/bk-1996-](https://doi.org/10.1021/bk-1996-0637.ch014)
852 0637.ch014
- 853 Kraberg, A., Baumann, M., Dürselen, C.-D., 2010. Coastal Phytoplankton: Photo Guide for
854 Northern European Seas, Verlag Dr. Friedrich Pfeil.
- 855 Lang, K., 1965. Copepoda Harpacticoida from the Californian Pacific coast. K. Sven.
856 Vetenskapsakademiens Handl. 1–560. 10, 1–560.
- 857 Lang, K., 1948. Monographie der Harpacticiden. Håkan Ohlsson, Lund.
- 858 Lavrentyev, P.J., Franzè, G., Pierson, J.J., Stoecker, D.K., 2015. The effect of dissolved
859 polyunsaturated aldehydes on microzooplankton growth rates in the Chesapeake Bay and
860 Atlantic coastal waters. *Mar. Drugs* 13, 2834–2856. <https://doi.org/10.3390/md13052834>
- 861 Lee, K.W., Raisuddin, S., Rhee, J.S., Hwang, D.S., Yu, I.T., Lee, Y.M., Park, H.G., Lee, J.S., 2008.
862 Expression of glutathione S-transferase (GST) genes in the marine copepod *Tigriopus*
863 *japonicus* exposed to trace metals. *Aquat. Toxicol.* 89, 158–166.
864 <https://doi.org/10.1016/j.aquatox.2008.06.011>
- 865 Lee, Y.M., Kim, I.C., Jung, S.O., Lee, J.S., 2005. Analysis of 686 expressed sequence tags (ESTs)
866 from the intertidal harpacticoid copepod *Tigriopus japonicus* (Crustacea, Copepoda). *Mar.*
867 *Pollut. Bull.* 51, 757–768. <https://doi.org/10.1016/j.marpolbul.2005.02.014>
- 868 Leflaive, J., Ten-Hage, L., 2009. Chemical interactions in diatoms: Role of polyunsaturated
869 aldehydes and precursors. *New Phytol.* 184, 794–805. [https://doi.org/10.1111/j.1469-](https://doi.org/10.1111/j.1469-8137.2009.03033.x)
870 8137.2009.03033.x
- 871 Legrand, C., Rengefors, K., Fistarol, G.O., Granéli, E., 2003. Allelopathy in phytoplankton -
872 Biochemical, ecological and evolutionary aspects. *Phycologia* 42, 406–419.
873 <https://doi.org/10.2216/i0031-8884-42-4-406.1>
- 874 Mascart, T., Agosto, L., Lepoint, G., Remy, F., De Troch, M., 2015. How do harpacticoid copepods

colonize detrital seagrass leaves? *Mar. Biol.* 162, 929–943. <https://doi.org/10.1007/s00227-015-2632-x>

McAbendroth, L., Ramsay, P.M., Foggo, A., Rundle, S.D., Bilton, D.T., 2005. Does macrophyte fractal complexity drive invertebrate diversity, biomass and body size distributions? *Oikos* 111, 279–290. <https://doi.org/10.1111/j.0030-1299.2005.13804.x>

Miralto, A., Barone, G., Romano, G., Poulet, S.A., Ianora, A., Russo, G.L., Buttino, I., Mazzarella, G., Laablr, M., Cabrini, M., Glacobbé, M.G., 1999. The insidious effect of diatoms on copepod reproduction. *Nature* 402, 173–176. <https://doi.org/10.1038/46023>

Monti, M., Cecchin, E., 2012. Comparative growth of three strains of *ostreopsis ovata* at different light intensities with focus on inter-specific allelopathic interactions. *Cryptogam. Algal.* 33, 113–119. <https://doi.org/10.7872/crya.v33.iss2.2011.113>

Paffenhöfer, G.A., Ianora, A., Miralto, A., Turner, J.T., Kleppel, G.S., Ribera D’Alcalà, M., Casotti, R., Caldwell, G.S., Pohnert, G., Fontana, A., Müller-Navarra, D., Jónasdóttir, S., Armbrust, V., Båmstedt, U., Ban, S., Bentley, M.G., Boersma, M., Bundy, M., Buttino, I., Calbet, A., Carlotti, F., Carotenuto, Y., D’Ippolito, G., Frost, B., Guisande, C., Lampert, W., Lee, R.F., Mazza, S., Mazzocchi, M.G., Nejstgaard, J.C., Poulet, S.A., Romano, G., Smetacek, V., Uye, S., Wakeham, S., Watson, S., Wichard, T., 2005. Colloquium on diatom-copepod interactions. *Mar. Ecol. Prog. Ser.* 286, 293–305. <https://doi.org/10.3354/meps286293>

Patil, V., Källqvist, T., Olsen, E., Vogt, G., Gislerød, H.R., 2007. Fatty acid composition of 12 microalgae for possible use in aquaculture feed. *Aquac. Int.* 15, 1–9. <https://doi.org/10.1007/s10499-006-9060-3>

Pavaux, A.-S., Rostan, J., Guidi Guilvard, L., Marro, S., Ternon, E., Olivier P., T., Lemée, R., Gasparini, S., 2019. Effects of the toxic dinoflagellate *Ostreopsis cf. ovata* on survival, feeding and reproduction of a phytal harpacticoid copepod. *J. Exp. Mar. Bio. Ecol.* 516, 103–113. <https://doi.org/10.1016/j.jembe.2019.05.004>

Pennesi, C., Danovaro, R., 2017. Assessing marine environmental status through

901 microphytobenthos assemblages colonizing the Autonomous Reef Monitoring Structures
 902 (ARMS) and their potential in coastal marine restoration. *Mar. Pollut. Bull.* 125, 56–65.
 903 <https://doi.org/10.1016/j.marpolbul.2017.08.001>

904 Pezzolesi, L., Accoroni, S., Rindi, F., Samorì, C., Totti, C., Pistocchi, R., 2021. Survey of the
 905 allelopathic potential of Mediterranean macroalgae: production of long-chain polyunsaturated
 906 aldehydes (PUAs). *Phytochemistry* 189. <https://doi.org/10.1016/j.phytochem.2021.112826>

907 Pezzolesi, L., Pichierri, S., Samorì, C., Totti, C., Pistocchi, R., 2017. PUFAs and PUAs production
 908 in three benthic diatoms from the northern Adriatic Sea. *Phytochemistry* 142, 85–91.
 909 <https://doi.org/10.1016/j.phytochem.2017.06.018>

910 Pichierri, S., Accoroni, S., Pezzolesi, L., Guerrini, F., Romagnoli, T., Pistocchi, R., Totti, C., 2017.
 911 Allelopathic effects of diatom filtrates on the toxic benthic dinoflagellate *Ostreopsis cf. ovata*.
 912 *Mar. Environ. Res.* 131, 116–122. <https://doi.org/10.1016/j.marenvres.2017.09.016>

913 Pichierri, S., Pezzolesi, L., Vanucci, S., Totti, C., Pistocchi, R., 2016. Inhibitory effect of
 914 polyunsaturated aldehydes (PUAs) on the growth of the toxic benthic dinoflagellate *Ostreopsis*
 915 *cf. ovata*. *Aquat. Toxicol.* 179, 125–133. <https://doi.org/10.1016/j.aquatox.2016.08.018>

916 Pinckney, J., Zingmark, R.G., 1993. Biomass and production of benthic microalgal communities in
 917 estuarine habitats. *Estuaries* 16, 887–897. <https://doi.org/10.2307/1352447>

918 Pistocchi, R., Guerrini, F., Pezzolesi, L., Riccardi, M., Vanucci, S., Ciminiello, P., Dell’Aversano,
 919 C., Forino, M., Fattorusso, E., Tartaglione, L., Milandri, A., Pompei, M., Cangini, M., Pigozzi,
 920 S., Riccardi, E., 2012. Toxin Levels and Profiles in Microalgae from the North-Western
 921 Adriatic Sea—15 Years of Studies on Cultured Species. *Mar. Drugs*.
 922 <https://doi.org/10.3390/md10010140>

923 Poulet, S.A., Ianora, A., Miralto, A., Meijer, L., 1994. Do diatoms arrest embryonic development in
 924 copepods? *Mar. Ecol. Prog. Ser.* 111, 79–86. <https://doi.org/10.3354/meps111079>

925 Raisuddin, S., Kwok, K.W.H., Leung, K.M.Y., Schlenk, D., Lee, J.S., 2007. The copepod
 926 *Tigriopus*: A promising marine model organism for ecotoxicology and environmental

927 genomics. *Aquat. Toxicol.* 83, 161–173. <https://doi.org/10.1016/j.aquatox.2007.04.005>

928 Ribalet, F., Bastianini, M., Vidoudez, C., Acri, F., Berges, J., Ianora, A., Miralto, A., Pohnert, G.,
929 Romano, G., Wichard, T., Casotti, R., 2014. Phytoplankton cell lysis associated with
930 polyunsaturated aldehyde release in the northern Adriatic Sea. *PLoS One* 9.
931 <https://doi.org/10.1371/journal.pone.0085947>

932 Ribalet, F., Berges, J.A., Ianora, A., Casotti, R., 2007. Growth inhibition of cultured marine
933 phytoplankton by toxic algal-derived polyunsaturated aldehydes. *Aquat. Toxicol.* 85, 219–227.
934 <https://doi.org/10.1016/j.aquatox.2007.09.006>

935 Ribalet, F., Intertaglia, L., Lebaron, P., Casotti, R., 2008. Differential effect of three
936 polyunsaturated aldehydes on marine bacterial isolates. *Aquat. Toxicol.* 86, 249–255.
937 <https://doi.org/10.1016/j.aquatox.2007.11.005>

938 Rindi, F., Gavio, B., Díaz-Tapia, P., Di Camillo, C.G., Romagnoli, T., 2020. Long-term changes in
939 the benthic macroalgal flora of a coastal area affected by urban impacts (Conero Riviera,
940 Mediterranean Sea). *Biodivers. Conserv.* 29, 2275–2295. [https://doi.org/10.1007/s10531-020-](https://doi.org/10.1007/s10531-020-01973-z)
941 01973-z

942 Rogelja, M., Cibic, T., Pennesi, C., De Vittor, C., 2016. Microphytobenthic community
943 composition and primary production at gas and thermal vents in the Aeolian Islands
944 (Tyrrhenian Sea, Italy). *Mar. Environ. Res.* 118, 31–44.
945 <https://doi.org/10.1016/j.marenvres.2016.04.009>

946 Romano, G., Russo, G.L., Buttino, I., Ianora, A., Miralto, A., 2003. A marine diatom-derived
947 aldehyde induces apoptosis in copepod and sea urchin embryos. *J. Exp. Biol.* 206, 3487–3494.
948 <https://doi.org/10.1242/jeb.00580>

949 Sandulli, R., Semprucci, F., Balsamo, M., 2014. Taxonomic and functional biodiversity variations
950 of meiobenthic and nematode assemblages across an extreme environment: a study case in a
951 Blue Hole cave. *Ital. J. Zool.* 81, 508–516. <https://doi.org/10.1080/11250003.2014.952356>

952 Sarmiento, V.C., Santos, P.J.P., 2012. Species of Harpacticoida (Crustacea, Copepoda) from the

953 phytal of Porto de Galinhas coral reefs, northeastern Brazil. Check List 8, 936–939.

954 <https://doi.org/10.15560/8.5.936>

955 Scholz, B., Liebezeit, G., 2012. Screening for competition effects and allelochemicals in benthic

956 marine diatoms and cyanobacteria isolated from an intertidal flat (southern North Sea).

957 Phycologia 51, 432–450. <https://doi.org/10.2216/11-80.1>

958 Semprucci, F., Balsamo, M., Sandulli, R., 2016. Assessment of the ecological quality (EcoQ) of the

959 Venice lagoon using the structure and biodiversity of the meiofaunal assemblages. Ecol. Indic.

960 67, 451–457. <https://doi.org/10.1016/j.ecolind.2016.03.014>

961 Snyder, R. V., Gibbs, P.D.L., Palacios, A., Abiy, L., Dickey, R., Lopez, J. V., Rein, K.S., 2003.

962 Polyketide synthase genes from marine dinoflagellates. Mar. Biotechnol. 5, 1–12.

963 <https://doi.org/10.1007/s10126-002-0077-y>

964 Sommer, U., 2009. Copepod growth and diatoms: Insensitivity of *Acartia tonsa* to the composition

965 of semi-natural plankton mixtures manipulated by silicon:nitrogen ratios in mesocosms.

966 Oecologia 159, 207–215. <https://doi.org/10.1007/s00442-008-1193-9>

967 Strickland, J.D., Parsons, T., 1972. A Practical Handbook of Seawater Analysis, 2nd ed. Bull.

968 Fish. Res. Board Can.

969 Sugihara, G., May, M.R., 1990. Applications of fractals in ecology. Trends Ecol. Evol. 5, 79–86.

970 [https://doi.org/10.1016/0169-5347\(90\)90235-6](https://doi.org/10.1016/0169-5347(90)90235-6)

971 Tang, Y.Z., Gobler, C.J., 2011. The green macroalga, *Ulva lactuca*, inhibits the growth of seven

972 common harmful algal bloom species via allelopathy. Harmful Algae 10, 480–488.

973 <https://doi.org/10.1016/j.hal.2011.03.003>

974 Tang, Y.Z., Gobler, C.J., 2010. Allelopathic effects of *Cochlodinium polykrikoides* isolates and

975 blooms from the estuaries of Long Island, New York, on co-occurring phytoplankton. Mar.

976 Ecol. Prog. Ser. 406, 19–31. <https://doi.org/10.3354/meps08537>

977 Taylor, R.B., Cole, R.G., 1994. Mobile epifauna on subtidal brown seaweeds in northeastern New

978 Zealand. Mar. Ecol. Prog. Ser. 115, 271–282. <https://doi.org/10.3354/meps115271>

979 Taylor, R.L., Caldwell, G.S., Dunstan, H.J., Bentley, M.G., 2007. Short-term impacts of
 980 polyunsaturated aldehyde-producing diatoms on the harpacticoid copepod, *Tisbe holothuriae*.
 981 J. Exp. Mar. Bio. Ecol. 341, 60–69. <https://doi.org/10.1016/j.jembe.2006.10.028>
 982 Ternon, E., Paix, B., Thomas, O.P., Briand, J.-F., Culioli, G., 2020. Exploring the Role of
 983 Macroalgal Surface Metabolites on the Settlement of the Benthic Dinoflagellate *Ostreopsis* cf.
 984 *ovata*. Front. Mar. Sci. 7. <https://doi.org/10.3389/fmars.2020.00683>
 985 Tillmann, U., 2004. Interactions between planktonic microalgae and protozoan grazers. J. Eukaryot.
 986 Microbiol. 51, 156–168. <https://doi.org/10.1111/j.1550-7408.2004.tb00540.x>
 987 Tomas, C.R., 1997. Identifying Marine Phytoplankton. Academic Press.
 988 Tonon, T., Harvey, D., Larson, T.R., Graham, I.A., 2002. Long chain polyunsaturated fatty acid
 989 production and partitioning to triacylglycerols in four microalgae. Phytochemistry 61, 15–24.
 990 [https://doi.org/10.1016/S0031-9422\(02\)00201-7](https://doi.org/10.1016/S0031-9422(02)00201-7)
 991 Underwood, A.J., 1996. Experiments in Ecology: Their Logical Design and Interpretation Using
 992 Analysis of Variance. Cambridge University Press, Cambridge. [https://doi.org/DOI:](https://doi.org/DOI:10.1017/CBO9780511806407)
 993 [10.1017/CBO9780511806407](https://doi.org/DOI:10.1017/CBO9780511806407)
 994 Utermöhl, H., 1958. Methods of collecting plankton for various purposes are discussed. SIL
 995 Commun. 1953-1996 9, 1–38. <https://doi.org/10.1080/05384680.1958.11904091>
 996 Vardi, A., Formiggini, F., Casotti, R., De Martino, A., Ribalet, F., Miralto, A., Bowler, C., 2006. A
 997 stress surveillance system based on calcium and nitric oxide in marine diatoms. PLoS Biol. 4,
 998 0411–0419. <https://doi.org/10.1371/journal.pbio.0040060>
 999 Wichard, T., Poulet, S.A., Boulesteix, A.L., Ledoux, J.B., Lebreton, B., Marchetti, J., Pohnert, G.,
 1000 2008. Influence of diatoms on copepod reproduction. II. Uncorrelated effects of diatom-
 1001 derived $\alpha,\beta,\gamma,\delta$ -unsaturated aldehydes and polyunsaturated fatty acids on *Calanus*
 1002 *helgolandicus* in the field. Prog. Oceanogr. 77, 30–44.
 1003 <https://doi.org/10.1016/j.pocean.2008.03.002>
 1004 Wichard, T., Poulet, S.A., Halsband-Lenk, C., Albaina, A., Harris, R., Liu, D., Pohnert, G., 2005a.

1005 Survey of the chemical defence potential of diatoms: Screening of fifty species for $\alpha,\beta,\gamma,\delta$ -
1006 unsaturated aldehydes. J. Chem. Ecol. 31, 949–958. <https://doi.org/10.1007/s10886-005-3615->
1007 z

1008 Wichard, T., Poulet, S.A., Pohnert, G., 2005b. Determination and quantification of $\alpha,\beta,\gamma,\delta$ -
1009 unsaturated aldehydes as pentafluorobenzyl-oxime derivates in diatom cultures and natural
1010 phytoplankton populations: Application in marine field studies. J. Chromatogr. B Anal.
1011 Technol. Biomed. Life Sci. 814, 155–161. <https://doi.org/10.1016/j.jchromb.2004.10.021>

1012

1013

1014

1015

1016

1017

1018 **Figure captions**

1019 **Figure 1** – Total density of microphytobenthos community (cells g⁻¹ fw) in *Dictyopteris*
1020 *polypodioides* (DP) and *Cystoseira compressa* (CC) at the different sampling times (T1-T6).

1021 **Figure 2** – Relative abundance (%) of species belonging to the Bacillariophyceae orders in A)
1022 *Dictyopteris polypodioides* (DP) and B) *Cystoseira compressa* (CC) at the different sampling times
1023 (T1-T6).

1024 **Figure 3** - Two-dimensional nMDS of centroids for A) microphytobenthos orders, B) meiofauna,
1025 and C) harpacticoid community on the two algae *Dictyopteris polypodioides* (DP) and *Cystoseira*
1026 *compressa* (CC) at the different sampling times (T1-T6).

1027 **Figure 4** – Total density (N g⁻¹ fw) of meiobenthos in *Dictyopteris polypodioides* (DP) and
1028 *Cystoseira compressa* (CC) at the different sampling times (T1-T6).

1029 **Figure 5** – Total density (n° organisms g⁻¹ fw) of A) nematodes, B) copepods and C) copepod
 1030 nauplii in *Dictyopteris polypodioides* (DP) and *Cystoseira compressa* (CC) at the different sampling
 1031 times (T1-T6).

1032 **Figure 6** – Total density (N g⁻¹ fw) of harpacticoids in A) *Dictyopteris polypodioides* (DP) and B)
 1033 *Cystoseira compressa* (CC) at the different sampling times (T1-T6).

1034 **Figure 7** – Results of Pearson correlation analysis between substrate attribution (D and S) and PUA
 1035 composition and densities of more representative genera of microphytobenthos, main meiofauna
 1036 taxa and harpacticoid species, after square root transformation. Non-significant relationships are
 1037 indicated by white squares. Signif. codes: *** p < 0.001, ** p < 0.01, * p < 0.05. D, fractal
 1038 dimension; S, macroalgal surface; tot, total PUA; C14:5, tetradecapentaenal; C16:4,
 1039 hexadecatetraenal; C16:3, hexadecatrienal; C6:2, hexadienal.

1040

1041

1042

1043

1044 Table 1: Relative abundance (%) of identified and unknown polyunsaturated aldehydes (PUAs) and
 1045 total concentration of PUAs (µg g⁻¹ fw) in *Dictyopteris polypodioides* (DP) and *Cystoseira*
 1046 *compressa* (CC) at the different sampling times (T1-T6).

1047

Time	Alga	C6:2	C16:3	C16:4	C14:5	Unknown	Tot (µg g ⁻¹ fw)
T1	DP	10%	12%	15%	50%	13%	225.5
T2	DP	0%	39%	20%	29%	14%	73.9
T3	DP	7%	21%	29%	32%	11%	44.3
T4	DP	7%	8%	14%	60%	11%	100.3
T5	DP	0%	12%	19%	61%	7%	82.1
T6	DP	6%	0%	8%	79%	7%	87.4
T1	CC	91%	0%	0%	0%	9%	17.1
T2	CC	64%	0%	0%	0%	36%	2.0
T3	CC	81%	0%	0%	0%	19%	2.3
T4	CC	88%	0%	0%	0%	12%	3.4
T5	CC	83%	0%	0%	0%	17%	5.3
T6	CC	68%	0%	0%	0%	32%	4.8

1048





Figure 1

1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091


Figure 2
A)



1092

1093

1094 B)




1095

1096 Figure 3


1097

A)


1098



B)



C)



1100

1101

1102 Figure 4

1103






Figure 5

1126


1127



1128



1129



1130

1131


1132

1133

1134 Figure 6

1135


1136 A)



1137

1138

1139 B)



1140

1141

1142

1143 Figure 7

1144

