



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE
DELLA RICERCA

Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Primitive permutation IBIS groups

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Lucchini A., Morigi M., Moscatiello M. (2021). Primitive permutation IBIS groups. JOURNAL OF COMBINATORIAL THEORY. SERIES A, 184, 1-17 [10.1016/j.jcta.2021.105516].

Availability:

This version is available at: <https://hdl.handle.net/11585/846160> since: 2022-01-18

Published:

DOI: <http://doi.org/10.1016/j.jcta.2021.105516>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Lucchini, A., Morigi, M., & Moscatiello, M. (2021). Primitive permutation IBIS groups. Journal of Combinatorial Theory. Series A, 184

The final published version is available online at
<https://dx.doi.org/10.1016/j.jcta.2021.105516>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

PRIMITIVE PERMUTATION IBIS GROUPS

ANDREA LUCCHINI, MARTA MORIGI, AND MARIAPIA MOSCATIELLO

ABSTRACT. Let G be a finite permutation group on Ω . An ordered sequence of elements of Ω , $(\omega_1, \dots, \omega_t)$, is an irredundant base for G if the pointwise stabilizer $G_{(\omega_1, \dots, \omega_t)}$ is trivial and no point is fixed by the stabilizer of its predecessors. If all irredundant bases of G have the same size we say that G is an IBIS group. In this paper we show that if a primitive permutation group is IBIS, then it must be almost simple, of affine-type, or of diagonal type. Moreover we prove that a diagonal-type primitive permutation groups is IBIS if and only if it is isomorphic to $\text{PSL}(2, 2^f) \times \text{PSL}(2, 2^f)$ for some $f \geq 2$, in its diagonal action of degree $2^f(2^{2^f} - 1)$.

1. INTRODUCTION

Let $G \leq \text{Sym}(\Omega)$ be a finite permutation group. A subset \mathcal{B} of Ω is a *base* for G if the pointwise stabilizer $G_{(\mathcal{B})}$ is trivial and we denote by $b(G)$ the minimal size of a base for G . An ordered sequence of elements of Ω , $\Sigma := (\omega_1, \dots, \omega_t)$, is *irredundant* for G if no point in Σ is fixed by the stabiliser of its predecessors. Moreover Σ is an *irredundant base* of G if it is a base and it is irredundant. In particular, such an irredundant base provides the the following decreasing chain reaching the identity

$$G > G_{\omega_1} > G_{(\omega_1, \omega_2)} > \dots > G_{(\omega_1, \dots, \omega_{t-1})} > G_{(\omega_1, \dots, \omega_t)} = 1,$$

where the inclusion of subgroups are strict.

Determining base sizes is a fundamental problem in permutation group theory, with a long history stretching back to the nineteenth century. Since the knowledge of how an element $g \in G$ acts on a base \mathcal{B} completely determines the action of g on Ω , bases plays an important role in computational group theory. The smaller the base, the less memory is needed to store group elements, so there are practical reasons for trying to find small bases.

Irredundant bases for a permutation group possess some of the features of bases in a vector space. Indeed, \mathcal{B} is a basis for a vector space V if and only if it is an irredundant base for the general linear group $\text{GL}(V)$ in its natural action on V . However, some familiar properties of bases in vector spaces do not extend to bases for permutation groups: irredundant bases for groups in general are not preserved by re-ordering, and they can have different sizes.

In [2], Cameron and Fon-Der-Flaass showed that all irredundant bases for a permutation group G have the same size if and only if all the irredundant bases for G are preserved by re-ordering. Groups satisfying one of the previous equivalent properties are called *Irredundant Bases of Invariant Size groups*, *IBIS groups* for short. Moreover, Cameron and Fon-Der-Flaass [2] also proved that for a permutation group G to be IBIS is a necessary and sufficient for the irredundant bases of G to be the base of a combinatorial structure known as *matroid*. If this condition hold, then G acts geometrically on the matroid and when G acts primitively, and is not cyclic of prime order, then the matroid is geometric (see [2] for more details). This brought Cameron to ask for a possible classification of the IBIS groups. As explained by Cameron himself in [1, Section 4-14], there is no hope for a complete classification of IBIS groups when the cardinalities of the bases are large. However it might be reasonable to pose this question for primitive groups.

2010 *Mathematics Subject Classification*. 20B15.

Key words and phrases. primitive group; base size; IBIS group .

It is easy to observe that the symmetric group $\text{Sym}(k)$ in its natural action on $[k] := \{1, \dots, k\}$ has base size $k - 1$, and that all subsets of $[k]$ with size $k - 1$ are irredundant bases for $\text{Sym}(k)$. Similarly the base size for the action of the alternating group $\text{Alt}(k)$ acting naturally on $[k]$ has base size $k - 2$ and all subsets of $[k]$ with size $k - 2$ are irredundant bases for $\text{Alt}(k)$. There are only finitely many values of k for which $\text{Sym}(k)$ or $\text{Alt}(k)$ acts as a primitive IBIS group not in its natural representation [2, Theorem 3.5]. In particular if the action of $\text{Sym}(k)$ and $\text{Alt}(k)$ on m -sets of $[k]$ is IBIS, then $k \leq 12$ and $m \leq 5$. In this context, it is worth remembering that $\text{Sym}(6)$ and $\text{Alt}(6)$ act as primitive IBIS groups on 2-sets (with degree 15) and on partitions into 2-sets of size 3 of $[6]$ (with degree 10). There exist other almost simple primitive groups that are IBIS, *e.g.* $\text{PSL}(2, q)$ and $\text{PGL}(2, q)$ on their action of degree $q + 1$, $\text{PSL}(2, 2^f)$ acting on the cosets of a dihedral subgroup of order $2(2^f + 1)$, $\text{PSL}(3, 2)$ in degree 7, and $\text{Alt}(7)$ in degree 15. In another direction, it is not difficult to observe that Frobenius groups are IBIS groups with base size 2. Hence, the affine group $\text{AGL}(1, p)$ is a primitive IBIS group with base size 2 for every prime p . More in general, $\text{AGL}(d, p)$ is a primitive IBIS group with base size $d + 1$ for every prime p , and for every natural number d .

Therefore we have quite a few examples of almost simple and affine-type primitive groups that are IBIS. So before launching in a complete classification of primitive IBIS groups, we decided to start the investigation of primitive permutation groups that can be IBIS in the classes of diagonal-type, product-type and twisted wreath product-type. The main result of this paper is the following theorem, that we prove in Section 6.

Theorem 1.1. *Let G be a primitive permutation IBIS group. Then one of the following holds:*

1. *The group G is of affine type.*
2. *The group G is almost simple.*
3. *The group G is of diagonal type.*

Moreover, G is an IBIS primitive group of diagonal type if and only if it belongs to the infinite family of non-monolithic diagonal type groups $\{\text{PSL}(2, 2^f) \times \text{PSL}(2, 2^f) \mid f \geq 2\}$ having degree $|\text{PSL}(2, 2^f)| = 2^f(2^{2f} - 1)$.

It is not difficult to see that there are no primitive permutation IBIS groups having non-abelian socle and base size 2 (see Lemma 2.3). This could lead to expect that there are not examples of IBIS groups with non-abelian socle and small base size. However, in Proposition 3.4, we prove that $\text{PSL}(2, 2^f) \times \text{PSL}(2, 2^f)$ in its diagonal action of degree $|\text{PSL}(2, 2^f)|$ is an IBIS group with base size 3. In Propositions 3.2, 3.1 and Theorem 3.19, as already mentioned in Theorem 1.1, we show that there are no other diagonal-type primitive permutation groups that are IBIS. During our investigation, we prove some results regarding non-abelian simple groups (see Subsection 2.1) and about the base size of primitive diagonal-type permutation groups (Lemma 3.7 and Lemma 3.14). Finally, in Sections 4, 5, we prove that there are no IBIS primitive groups that are of product-type nor twisted wreath product groups.

2. PRELIMINARIES

The socle of a finite primitive group is a non-trivial direct product of simple groups that are pairwise isomorphic. The essential tool for study primitive permutation groups depending on the group and the action of the socle is the O’Nan-Scott Theorem. In [12], Liebeck, Praeger and Saxl gave a self-contained proof of such theorem. There, precisely five types of primitive groups are defined, namely the *affine-type*, the *almost simple*, the *diagonal-type*, the *product-type*, and the *twisted wreath product-type* and it is shown that, every finite primitive group belongs to exactly one of these types. We refer the reader to [12] for the precise definition of the classes.

Now, we state and prove same preliminary results regarding bases.

Remark 2.1. Since a base of G with minimal size $b(G)$ is clearly irredundant, then a permutation group G is IBIS if and only if all its irredundant bases have size $b(G)$.

The proof of the following lemma is straightforward.

Lemma 2.2. *Let G be a transitive permutation group on a set Ω . Then $\mathcal{B} \subset \Omega$ is an irredundant base if and only if $\mathcal{B}^g \subset \Omega$ is an irredundant base for every $g \in G$.*

Lemma 2.3. *Let $G \leq \text{Sym}(\Omega)$ be a primitive group with non-abelian socle and let M be a normal subgroup of G containing the socle of G . If M is not regular, then M has an irredundant base of size at least 3. In particular there are no IBIS primitive permutation groups G with non-abelian socle and $b(G) = 2$.*

Proof. Let $\omega \in \Omega$. The point stabilizer G_ω normalizes $M_\omega = M \cap G_\omega$, hence $G_\omega \leq N_G(M_\omega)$. Since M is not regular, M_ω is not normal in G , so $G_\omega \leq N_G(M_\omega) < G$. On the other hand G_ω is a maximal subgroup of G , so $G_\omega = N_G(M_\omega)$ and consequently $N_M(M_\omega) = M_\omega$. Now assume by contradiction that $M_\omega \cap M_\omega^x = 1$, for every $x \in M \setminus M_\omega$. Then M would be a Frobenius group with kernel K and complement M_ω (see [13, 8.5.5]). In particular $K = \text{soc}(M)$ would be a nilpotent normal subgroup of G (see [13, 10.5.6]), in contradiction with the fact that $\text{soc}(G)$ is a direct product of isomorphic non-abelian simple groups. \square

2.1. Results on finite non-abelian simple groups. In this subsection, the finite non-abelian simple CT-groups are characterized.

Definition 2.4. A finite group G is said to be a *CT-group* if commutativity is a transitive relation on $G \setminus \{1\}$.

Lemma 2.5. *A group G is a CT-group if and only if for every $x, y \in G \setminus \{1\}$, either $C_G(x) = C_G(y)$ or $C_G(x) \cap C_G(y) = 1$.*

Proof. First, let assume that G is a CT-group. Moreover assume that t is a non-identity element of $C_G(x) \cap C_G(y)$. Since t commutes with both x and y , then every element commuting with x commutes with t and consequently with y . Similarly every element commuting with y commutes with t and consequently with x . That is $C_G(x) = C_G(y)$.

Conversely assume that for every $x, y \in G \setminus \{1\}$, either $C_G(x) = C_G(y)$ or $C_G(x) \cap C_G(y) = 1$. Let $a, b, c \in G \setminus \{1\}$ and assume that a commutes with b and b commutes with c . Then $a \in C_G(a) \cap C_G(b)$ and $b \in C_G(b) \cap C_G(c)$, so $C_G(a) = C_G(b) = C_G(c)$. \square

Lemma 2.6. *The group $T = \text{PSL}_2(2^f)$, with $f \geq 2$ is a CT-group.*

Proof. Let $x \in T$. If x is not the identity, the minimal polynomial of x coincides with its characteristic polynomial. In the latter case, the matrices that commute with x are all polynomials in x , so they all commute with each other. That is $C_T(x)$ is an abelian subgroup of T . Now let a, b , and t be non-trivial elements of T such that a commutes with t and t commutes with b . Since a and b are elements of $C_T(t)$ that is an abelian group, then a and b commute. \square

In 1957, Suzuki [14] showed that if G is a finite non-abelian simple CT-group then G is isomorphic to some $\text{PSL}(2, 2^f)$ with $f \geq 2$. Hence the following result holds true.

Theorem 2.7. *A finite non-abelian simple group is a CT-group if and only if it is isomorphic to $\text{PSL}(2, 2^f)$ for some $f \geq 2$.*

3. DIAGONAL TYPE

Let $k \geq 2$ be an integer, and let T be a non-abelian simple group. We set $[k] := \{1, \dots, k\}$ and

$$\begin{aligned} W(k, T) &:= \{(x_1, \dots, x_k)\pi \in \text{Aut}(T) \wr \text{Sym}(k) \mid x_1 \equiv x_i \pmod T \text{ for all } i \in [k]\}, \\ D(k, T) &:= \{(x, \dots, x)\pi \in \text{Aut}(T) \wr \text{Sym}(k)\}, \\ \Omega(k, T) &:= D(k, T)/W(k, T), \text{ the set of the right cosets of } D(k, T) \text{ in } W(k, T) \\ A(k, T) &:= W(k, T) \cap \text{Aut}(T)^k. \end{aligned}$$

Observe that $W(k, T) = A(k, T) \rtimes \text{Sym}(k) = T^k \cdot (\text{Out}(T) \times \text{Sym}(k))$. The group G is of diagonal type if there exists an integer $k \geq 2$ and a non-abelian simple group T such that $T^k \leq G \leq W(k, T)$ and G acts primitively on $\Omega(k, T)$. Throughout this section we will assume that G is a primitive permutation group of diagonal type. Note that G has socle T^k and degree $|T|^{k-1}$. Moreover G is primitive on $\Omega(k, T)$ when

$$P_G := \{\pi \in \text{Sym}(k) \mid (g_1, \dots, g_k)\pi \in G \text{ for some } (g_1, \dots, g_k) \in A(k, T)\} \leq \text{Sym}(k)$$

is primitive on $[k]$ or $k = 2$ and $P_G = 1$. Recall that a group is *monolithic* if it has a unique minimal normal subgroup. If $k = 2$ and $P_G = 1$ then G has two minimal normal subgroups, hence the group is non-monolithic. In all other cases, the primitive diagonal-type groups are monolithic.

Throughout the section, we shall use the following notation without further comment. Given $t, s_2, \dots, s_k \in T$, we set

$$H := \{x \in \text{Aut}(T) \mid (x, \dots, x) \in G\}, \quad C(t) := C_H(t), \quad \text{and} \quad [1, s_2, \dots, s_k] := D(k, T)(1, s_2, \dots, s_k).$$

From Remark 2.1, it is clear that the study of primitive permutation IBIS groups is intimately related to the base sizes. For primitive diagonal groups, the minimal base size has been studied in [4]. In the following proposition, we summarize the results proved in [4] which we shall use in this section.

Proposition 3.1. [4] *The following hold true.*

- (1) *If the top group P_G does not contain $\text{Alt}(k)$, then $b(G) = 2$.*
- (2) *If $k = 2$, then $b(G) = 3$ when $P_G = 1$, and $b(G) \in \{3, 4\}$ otherwise.*
- (3) *If $P_G = \text{Alt}(k)$, then $b(G) = 3$ when $k = 2$, and $b(G) = 2$ when k is 3 or 4.*
- (4) *If $P_G = \text{Sym}(k)$, then $b(G) \in \{3, 4\}$ when $k = 2$, and $b(G) \in \{2, 3\}$ when k is 3 or 4.*

3.1. Diagonal non-monolithic primitive groups. In this subsection, $k = 2$ and $P_G = 1$.

Here we will prove that G is IBIS if and only if $G \cong (\text{PSL}(2, 2^f))^2$, for some $f \geq 2$.

It is not difficult to observe that the sequence $([1, 1], [1, t_1], [1, t_2], \dots, [1, t_r])$ is an irredundant sequence for G if and only if the following hold true:

- t_1, \dots, t_r are non trivial elements of T ;
- for $1 \leq i \leq r - 1$, the subgroup $\bigcap_{1 \leq j \leq i} C(t_j)$ properly contains $\bigcap_{1 \leq j \leq i+1} C(t_j)$
- $\bigcap_{1 \leq j \leq r} C(t_j) = 1$.

Proposition 3.2. *If $k = 2$, $P_G = 1$ and G is an IBIS group, then $T = (\text{PSL}(2, 2^f))^2$, for some $f \geq 2$.*

Proof. If $\text{soc}(G)$ is not isomorphic to $(\text{PSL}(2, 2^f))^2$, then by Theorem 2.7 we can choose two non trivial elements of T , s and t , such that $C(t) \cap C(s) \neq 1$ and $C(t) \neq C(s)$. Then $([1, 1], [1, s], [1, t])$ is an irredundant sequence that is not a base. In particular we can complete $([1, 1], [1, s], [1, t])$ to an irredundant base of G , thus there exists an irredundant base of G with size at least 4. Since $b(G) = 3$ by Proposition 3.1, it follows from Remark 2.1 that G is not IBIS. Therefore we conclude that $\text{soc}(G) = T^2 = (\text{PSL}(2, 2^f))^2$. \square

Proposition 3.3. *If $k = 2$, $P_G = 1$, $T = \text{PSL}(2, 2^f)$ and G is IBIS, then $G = \text{soc}(G) = T^2$.*

Proof. Let $\varphi \in \text{Aut}(T)$ be the automorphism induced by the Frobenius automorphism of the field \mathbb{F}_{2^f} , so that $\text{Aut}(T) = \langle \varphi \rangle T$. If $G \neq T^2$, then there exists $\varphi^i \in \text{Aut}(T)$ such that $(\varphi^i, \varphi^i) \in G$. That is, $\varphi^i \in H$. Consider the following elements of T :

$$w := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad z := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

Note that $w \in C(w) \setminus C(z)$. Moreover, it is clear that $w^{\varphi^i} = w$ and $z^{\varphi^i} = z$, that is $\varphi^i \in C(w) \cap C(z)$. Hence $([1, 1], [1, w], [1, z])$ is an irredundant sequence that is not a base. Hence there exists an irredundant base of G with size at least 4. Since $b(G) = 3$ by Proposition 3.1, it follows from Remark 2.1 that G is not IBIS. Therefore we conclude that $G = T^2 = (\text{PSL}(2, 2^f))^2$. \square

Proposition 3.4. *Let $G = T^2 = (\text{PSL}_2(2^f))^2$, where $f \geq 2$. Then G is an IBIS group.*

Proof. Since $b(G) = 3$ by Proposition 3.1, it suffices to show that each irredundant base has size 3. Denoting by \mathcal{B} an irredundant base, by Lemma 2.2 we can assume that $[1, 1]$ is the first element of \mathcal{B} . Since $|\mathcal{B}| \geq 3$, there exist two different non-trivial elements s and t of T , such that $[1, t]$ and $[1, s]$ are in \mathcal{B} . Moreover, since \mathcal{B} is irredundant, then $C_T(t) \neq C_T(s)$. Hence from Theorem 2.7, we deduce that $C_T(t) \cap C_T(s) = 1$ and so \mathcal{B} has size 3. \square

3.2. Diagonal monolithic primitive groups. In this section we show that the monolithic primitive groups of diagonal type are not IBIS groups. Recall that if a group G of diagonal type with $T^k \leq G \leq W(k, T)$ is monolithic, then P_G is primitive on its action on k points. Throughout the subsection, we shall assume that G is monolithic without further mentioning.

First we assume $k = 2$. In this case $T^2 \leq G \leq W(2, T)$ and $P_G = \langle \sigma \rangle$, where $\sigma := (1, 2)$. Note that there exists $y \in \text{Aut}(T)$ such that $(y, y)\sigma \in G$. Recall that $H = \{z \in \text{Aut}(T) \mid (z, z) \in G\}$ and consider the coset Hy . Since $((y, y)\sigma)^2 = (y^2, y^2)$, we have that $y^2 \in H$. For any $x \in \text{Aut}(T)$, we get that $(x, x)\sigma \in G$ if and only if $Hx = Hy$. Further, notice that $Hy = H$ if and only if $\sigma \in G$.

For any $t \in T$, define

$$I(t) := \{x \in yH \mid x^{-1}tx = t^{-1}\}.$$

Lemma 3.5. *Let $t_1, \dots, t_r \in T$ and $\mathcal{B} = ([1, 1], [1, t_1], \dots, [1, t_r])$. Then \mathcal{B} is a base for G if and only if*

$$\bigcap_{1 \leq i \leq r} C(t_i) = 1 \quad \text{and} \quad \bigcap_{1 \leq i \leq r} I(t_i) = \emptyset.$$

Moreover, \mathcal{B} is an irredundant sequence if and only if t_1, \dots, t_r are non-trivial elements of T and, for $1 \leq j \leq r - 1$, either the set $\bigcap_{1 \leq i \leq j} C(t_i)$ properly contains $\bigcap_{1 \leq i \leq j+1} C(t_i)$ or the set $\bigcap_{1 \leq i \leq j} I(t_i)$ properly contains $\bigcap_{1 \leq i \leq j+1} I(t_i)$.

Proof. We have $G_{[1,1]} = \{(h, h), (x, x)\sigma \mid h \in H, x \in yH\}$. To conclude it suffices to notice that $(h, h) \in G_{[1, t_i]}$ if and only if $h \in C(t_i)$, while $(x, x)\sigma \in G_{[1, t_i]}$ if and only if $x \in I(t_i)$. \square

The following result will play an important role in investigating the case $k = 2$.

Theorem 3.6. [11, Theorem 1.1] *Let T be a non-abelian finite simple group, that is not $\text{Alt}(7)$, $\text{PSL}(2, q)$, $\text{PSL}(3, q)$ nor $\text{PSU}(3, q)$. Then there exist $x, t \in T$ such that the following hold:*

- (i) $T = \langle x, t \rangle$;
- (ii) t is an involution;
- (iii) there is no involution $\alpha \in \text{Aut}(T)$ such that $x^\alpha = x^{-1}$, $t^\alpha = t$.

Lemma 3.7. *Assume that $T^2 \leq G \leq W(2, T)$, with T as in Theorem 3.6. Then $b(G) = 3$.*

Proof. Let $T = \langle x, t \rangle$, with x, t as in Theorem 3.6 and let $\mathcal{B} := ([1, 1], [1, x], [1, t])$. Note that $C_{\text{Aut}(T)}(x) \cap C_{\text{Aut}(T)}(t) = 1$ and $I(x) \cap I(t) = \emptyset$, so by Lemma 3.5 \mathcal{B} is a base for G , that is $b(G) \leq 3$. By Proposition 3.1, $b(G) \geq 3$, so the result follows. \square

Proposition 3.8. *Assume that $T^2 \leq G \leq W(2, T)$, with $T \neq \text{PSL}(2, 2^f)$ for every $f \geq 2$. If $b(G) \leq 3$, then G is not an IBIS group.*

Proof. By Theorem 2.7, there exist two nonidentity elements s and t in T such that $C_T(t) \cap C_T(s) \neq 1$ and $C_T(t) \neq C_T(s)$. Then $([1, 1], [1, s], [1, t])$ is an irredundant sequence that is not a base. In particular we can complete $([1, 1], [1, s], [1, t])$ to an irredundant base of G , thus there exists an irredundant base of G with size at least 4. Since we are assuming $b(G) \leq 3$, the result follows from Remark 2.1. \square

Combining Lemma 3.7 and Proposition 3.8 we get the following:

Corollary 3.9. *Let $T^2 \leq G \leq W(2, T)$, with T as in Theorem 3.6. Then G is not an IBIS group.*

In what follows, we are going to show that also when $T \in \{\text{Alt}(7), \text{PSL}(2, q), \text{PSL}(3, q), \text{PSU}(3, q)\}$, the group $T^2 \leq G \leq W(2, T)$ is not IBIS.

Although the results for $\text{Alt}(7)$ and $\text{Alt}(5)$ follow straightforwardly from a GAP [6] computation, we prefer to keep them so detailed as we believe that such easy constructions help the reader to handle subsequent arguments more easily.

Proposition 3.10. *If $(\text{Alt}(7))^2 \leq G \leq W(2, \text{Alt}(7))$, then G is not an IBIS group.*

Proof. If $(1, 2) \in yH$ then $((1, 2), (1, 2))\sigma \in G$. In this case, consider $s_1 = (1, 2)(3, 4)$, $s_2 = (1, 2)(3, 5)$ and $s_3 = (1, 2)(3, 6)$. Since $(1, 2) \in I(s_1) \cap I(s_2) \cap I(s_3)$ and $C_{\text{Alt}(7)}(\langle s_1, s_2, s_3 \rangle) < C_{\text{Alt}(7)}(\langle s_1, s_2 \rangle) < C_{\text{Alt}(7)}(s_1)$, then $([1, 1], [1, s_1], [1, s_2], [1, s_3])$ is an irredundant sequence of G that is not a base.

Notice that $(1, 2) \notin yH$ if and only if $H = \text{Alt}(7) = Hy$. In this case, consider $t_1 = (1, 2, 3)$, $t_2 = (1, 2, 4)$, and $t_3 = (1, 2, 5)$. Since $(1, 2)(6, 7) \in I(t_1) \cap I(t_2) \cap I(t_3)$ and $C_{\text{Alt}(7)}(\langle t_1, t_2, t_3 \rangle) < C_{\text{Alt}(7)}(\langle t_1, t_2 \rangle) < C_{\text{Alt}(7)}(t_1)$, then $([1, 1], [1, t_1], [1, t_2], [1, t_3])$ is an irredundant sequence of G that is not a base.

In particular, in both cases, completing the irredundant sequences to a base of G , we find an irredundant base of G with size at least 5. By Proposition 3.1 $b(G) \leq 4$, hence the result follows from Remark 2.1. \square

For $T = \text{PSL}(2, q)$ we need to work differently according to the parity of q . Since $\text{PSL}(2, 4) \cong \text{PSL}(2, 5) \cong \text{Alt}(5)$, in order to simplify the investigation, we deal with $T = \text{Alt}(5)$ separately.

Proposition 3.11. *Let $T = \text{Alt}(5)$ and let $T \leq G \leq W(T, 2)$. Then G is not an IBIS group.*

Proof. We prove the result by considering different cases and, in each of them, we set $a := (1, 2, 3)$.

First, let $yH \neq H$. Under this assumption we have that $H = T = \text{Alt}(5)$ and so, without loss of generality, we can assume that $y = (1, 2)$. Let $a_2 := (1, 2, 3, 4, 5)$ and let $\mathcal{B}_1 := ([1, 1], [1, a], [1, a_2])$. Note that $a \in C(a) \setminus C(a_2)$, $C(a) \cap C(a_2) = 1$, and $I(a) \cap I(a_2) = \emptyset$. By Lemma 3.5 then \mathcal{B}_1 is an irredundant base of size 3. Now, let $b_2 := (1, 2, 4)$. Note that $a \in C(a) \setminus C(b_2)$ and $(1, 2) \in I(a) \cap I(b_2)$. Hence $([1, 1], [1, a], [1, b_2])$ is an irredundant sequence that is not a base. Completing this to an irredundant bases of G , we produce an irredundant base of size at least 4. By Remark 2.1, G is not IBIS in this case.

Let $yH = H = T$, let $c_2 := (1, 2)(3, 4)$, and let $\mathcal{S}_1 := ([1, 1], [1, a], [1, c_2])$. Since $C(a) \cap C(c_2) = 1$ and $I(a) \cap I(c_2) = \emptyset$, then \mathcal{S}_1 is an irredundant base of size 3. Let $d_2 := (1, 2)(4, 5)$. Then $a \in C(a) \setminus C(d_2)$ and $d_2 \in I(a) \cap I(d_2)$. Consequently $([1, 1], [1, a], [1, d_2])$ is an irredundant sequence that is not a base and we can conclude as in the previous paragraph.

Finally, we need to consider the case $yH = H = \text{Sym}(5)$. Here, let $t := (3, 4, 5)$, $s_1 := (1, 2)(3, 4)$, and $s_2 := (1, 2)(4, 5)$. Since $t \in C(t) \setminus C(s_1)$, $s_1 \in (I(t) \cap I(s_1)) \setminus I(s_2)$, and $(1, 2) \in C(t) \cap C(s_1) \cap C(s_2)$, then $([1, 1], [1, t], [1, s_1], [1, s_2])$ is an irredundant sequence of G that is not a base. Thus completing the irredundant sequences to a base of G , we find an irredundant base of G with size at least 5. By Proposition 3.1 $b(G) \leq 4$, hence the result follows from Remark 2.1. \square

Proposition 3.12. *Let $T = \text{PSL}(2, q)$. If $q > 5$ is odd and $T^2 \leq G \leq W(2, T)$, then G is not an IBIS group.*

Proof. The conjugacy classes of subgroups of $T = \text{PSL}(2, q)$ are well-known (see for instance [9]). Among them there are a conjugacy class of dihedral groups of order $q-1$ (in Aschbacher class \mathcal{C}_2) and a conjugacy class of dihedral groups of order $q+1$ (in Aschbacher class \mathcal{C}_3). We can choose $\epsilon \in \{\pm 1\}$ such that $a = \frac{q+\epsilon}{2}$ is odd. Let $b = \frac{q-\epsilon}{2}$. In T there is just one conjugacy class of involutions and the centralizer of an involution is a dihedral group D_{2b} of order $2b$. Note that $|T| = \frac{1}{2}q(q^2 - 1) = 2qab$. Thus the total number of involutions is qa . There are qb subgroups isomorphic to a dihedral group D_{2a} of order $2a$ and each of them contains a involutions. Hence every involution is contained in b different subgroups of type D_{2a} . In particular, there exist two different subgroups M_1 and M_2 , both isomorphic to D_{2a} , such that $M_1 \cap M_2$ contains an involution x .

First, assume $q > 11$. In this case the dihedral groups of order $2a$ are maximal in G . Let $y_i \in M_i$ be an element of order a such that $M_i = \langle y_i, x \rangle$ and $y_i^x = y_i^{-1}$, for $i = 1, 2$. Since there is no maximal subgroup of T containing both y_1 and y_2 , we have that $T = \langle y_1, y_2 \rangle$. The previous argument does not work if $q \in \{7, 9, 11\}$ but it can be easily checked that, also in these cases, T contains two elements y_1, y_2 of order a and an involution x such that $y_1^x = y_1^{-1}, y_2^x = y_2^{-1}$ and $\langle y_1, y_2 \rangle = T$.

Note that if $z \in \text{Aut}(T)$ is such that $y_i^z = y_i^{-1}$, for $i = 1, 2$, then $zx^{-1} \in C_{\text{Aut}(T)}(\langle y_1, y_2 \rangle) = 1$, hence $z = x$.

Assume that $x \notin yH$. Then we have that $I(y_1) \cap I(y_2) = \emptyset$ and $C(y_1) \cap C(y_2) = 1$, thus, by Lemma 3.5, $\mathcal{B} = ([1, 1], [1, y_1], [1, y_2])$ is an irredundant base for G of size 3. Hence, by Proposition 3.8, G is not IBIS.

Now, assume that $x \in yH$, so that $y \in Hx = H$. Note that there exists a subgroup of type D_{2b} containing x and r such that r has order b and $r^x = r^{-1}$. From $q \neq 5$, it follows that $b > 2$ and $r \neq r^{-1}$. Moreover we have that $T = \langle x, y_1, r \rangle$, where $x \in C(x) \setminus C(\langle x, r \rangle)$, $r^{\frac{b}{2}} \in C(\langle x, r \rangle) \setminus C(\langle x, r, y_1 \rangle)$, and $x \in I(x) \cap I(r) \cap I(y_1)$. Hence $([1, 1], [1, x], [1, r], [1, y_1])$ is an irredundant sequence that is not a base, so G has an irredundant base of size at least 5. As usual, the result follows from Proposition 3.1 and Remark 2.1. \square

Lemma 3.13. *Let $T = \text{PSL}(2, 2^f)$ with $f \geq 3$. Then there exists two maximal subgroups K_1, K_2 of T such that $K_1 \cong K_2 \cong D_{2(q-1)}$ and $|K_1 \cap K_2| = 1$.*

Proof. Let $q = 2^f$. In T there is just one conjugacy class of involutions and the centralizer of an involution is a Sylow 2-subgroup of order q . Note that $|T| = q(q^2 - 1)$. Thus the total number of involutions is $q^2 - 1$. There are $q(q+1)/2$ maximal subgroups isomorphic to $D_{2(q-1)}$ (all conjugate in T). Denote by Ω the set of these subgroups. Any element of Ω contains $q-1$ involutions. Hence every involution is contained in $q/2$ different subgroups of Ω . Now, fix $D \in \Omega$. If $K \in \Omega$ and $K \neq D$, then $|D \cap K| \leq 2$. Let $\Omega_D = \{K \in \Omega \mid |D \cap K| = 2\}$. For any involution $z \in D$, there are exactly $q/2 - 1$ subgroups $K \in \Omega_D$ with $D \cap K = \langle z \rangle$, hence $|\Omega_D| = (q-1)(q/2 - 1) < |\Omega| - 1 = q(q+1)/2 - 1$. Thus there exists $K \neq D$ with $K \in \Omega \setminus \Omega_D$. \square

Lemma 3.14. *Assume that $T^2 \leq G \leq W(2, T)$, with $T = \text{PSL}(2, 2^f)$ and $f \geq 3$. If either $yH \neq H$ or $yH = H = T$, then $b(G) = 3$.*

Proof. First, we assume that $yH \neq H$. Since $q = 2^f$, we have that $\text{PSL}(2, q)$ is isomorphic to $\text{SL}(2, q)$, so we may deal with $T = \text{SL}(2, q)$. Let a be an element of multiplicative order $q-1$ in the

field \mathbb{F}_q . Consider the following elements of T :

$$x := \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}, \quad w := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \text{and } z := wx = \begin{pmatrix} 0 & a^{-1} \\ a & 0 \end{pmatrix}.$$

Then $N := \langle w, z \rangle$ is a dihedral group of order $2(q-1)$. As $C_T(x) = \langle x \rangle$, it follows that $C_T(N) = 1$.

We are going to show that $C_{\text{Aut}(S)}(N) = 1$. Denote by $\varphi \in \text{Aut}(T)$ the automorphism induced by the Frobenius automorphism of the field \mathbb{F}_q . Let $s \in T$, let $1 \leq i \leq f$, and let assume that $\varphi^i s$ centralizes N . In particular, $\varphi^i s$ centralizes x , and so $x^{\varphi^i} = x^{s^{-1}}$ and x are conjugated. Therefore $\{a, a^{-1}\} = \{a^{2^i}, a^{-2^i}\}$.

If $a = a^{-2^i}$, then $a^{2^i+1} = 1$ and so $q-1 = 2^f - 1$ divides $2^i + 1$, contradicting the assumption $f \geq 3$. Hence $a = a^{2^i}$, that is φ^i is the identity. Consequently $s \in C_T(N) = 1$. This proves that $C_{\text{Aut}(S)}(N) = 1$ and in particular that $C(z) \cap C(w) = 1$.

If $I(w) \cap I(z) \neq \emptyset$, then there exist $i \in \mathbb{N}$ and $s \in T$ such that $\varphi^i s \in I(w) \cap I(z)$. Since w and z are involutions, then $\varphi^i s \in C_{\text{Aut}(S)}(N) = 1$. Consequently $1 \in yH$, that contradicts $yH \neq H$.

Hence $I(w) \cap I(z) = \emptyset$ and, from Lemma 3.5, $([1, 1], [1, z], [1, w])$ is a base for G . Combining this and Proposition 3.1, we deduce that $b(G) = 3$.

Now, assume that $yH = H = T$. By Lemma 3.13, G contains two subgroups H_1 and H_2 isomorphic to a dihedral group of order $2(q-1)$, with $H_1 \cap H_2 = 1$. Take $x_1 \in H_1, x_2 \in H_2$, both of order $q-1$. There is no maximal subgroup containing both x_1 and x_2 , so $T = \langle x_1, x_2 \rangle$. In particular, $C(x_1) \cap C(x_2) = 1$. Moreover, if z is an involution inverting x_1 , then $z \in H_1$. Hence, there are no involutions in T inverting both x_1 and x_2 . Therefore $([1, 1], [1, x_1], [1, x_2])$ is an irredundant base of size 3. Combining this and Proposition 3.1, we deduce that $b(G) = 3$. \square

Proposition 3.15. *Let $T = \text{PSL}(2, 2^f)$, with $f \geq 3$. If $T^2 \leq G \leq W(2, T)$ then G is not an IBIS group.*

Proof. As above, we work in $T = \text{SL}(2, 2^f)$. First, assume that $yH \neq H$. Without loss of generality, denoting again by φ the automorphism induced by the Frobenius automorphism of the field \mathbb{F}_{2^f} , we can say that $y = \varphi^t$, for some $t \in \mathbb{N}$. Since $y^2 \in H$, then $f = 2r$ is even. Moreover, we can assume that t divides r . Since $2^t + 1$ divides $2^f - 1$, there exists a nonzero element $c \in \mathbb{F}_{2^f}$ of multiplicative order $2^t + 1$. Consider the following elements

$$x := \begin{pmatrix} c & 0 \\ 0 & c^{-1} \end{pmatrix} \quad \text{and } w := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Note that $x \in C(x) \setminus C(\langle x, w \rangle)$. Further, $x^y = x^{\varphi^t} = x^{-1}$ and $w^y = w^{\varphi^t} = w = w^{-1}$, so that $y \in I(x) \cap I(w) \neq \emptyset$. Hence $([1, 1], [1, x], [1, w])$ is an irredundant sequence that is not a base. From Lemma 3.14, we have that $b(G) = 3$, hence the result follows from Remark 2.1.

Therefore we can assume that $yH = H$, so that $\sigma \in G$.

Let assume that there exists $1 \neq \varphi^i \in \text{Aut}(T)$ such that $(\varphi^i, \varphi^i) \in G$, that is $H > T$. Consider the following elements of T :

$$x := \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}, \quad w := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{and } z := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

Note that $N := \langle x, w \rangle$ is a dihedral group of order $2|x|$ and w inverts x . In particular $x \in C(x) \setminus C(w)$, $w \in I(x) \cap I(w)$, and $w \notin I(z)$. Moreover, it is clear that $x^{\varphi^i} = x$, $w^{\varphi^i} = w$, and $z^{\varphi^i} = z$, that is $\varphi^i \in C(x) \cap C(w) \cap C(z)$. Hence $([1, 1], [1, x], [1, w], [1, z])$ is an irredundant sequence that is not a base. This produces a base of G with size at least 5. Since $b(G) \leq 4$ by Proposition 3.1, the result follows from Remark 2.1.

Finally, assume $Hy = H = T$. Notice that T contains two involutions z_1, z_2 generating a dihedral group of order $2(2^f + 1)$. Since $C(z_1) \cap C(z_2) = I(z_1) \cap I(z_2) = 1$, for every $x \in T$ with $|x| >$

2, $([1, 1], [1, z_1], [1, z_2], [1, x])$ is an irredundant base of size 4. Now, the result follows combining Lemma 3.14 and Remark 2.1. \square

Proposition 3.16. *If $T = \text{PSL}(3, q)$ and $T^2 \leq G \leq W(2, T)$, then G is not an IBIS group.*

Proof. Let $q = p^f$, let

$$N := \left\{ \begin{pmatrix} 1 & x & y \\ 0 & 1 & z \\ 0 & 0 & 1 \end{pmatrix} \mid x, y, z \in \mathbb{F}_p \right\} \subseteq \text{SL}(3, q),$$

and consider the following elements of N :

$$a := \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, b := \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, c := \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Since N intersects the center of $\text{SL}(3, q)$ trivially, we may identify N with a subgroup of T . Note that N is non-abelian of order p^3 , $Z(N) = \langle c \rangle$, and N is isomorphic to a dihedral group of order 8 when $p = 2$, and it has exponent p when p is odd. In particular $a \in C_T(c) \setminus (C_T(c) \cap C_T(b))$, $b \in (C_T(c) \cap C_T(b)) \setminus (C_T(c) \cap C_T(b) \cap C_T(a))$ and $c \in (C_T(c) \cap C_T(b) \cap C_T(a))$. Thus $([1, 1], [1, c], [1, b], [1, a])$ is an irredundant sequence that is not a base. By Proposition 3.1 we have that $b(G) \leq 4$, hence by Remark 2.1, G is not IBIS. \square

Proposition 3.17. *Assume that $S^2 \leq G \leq W(2, S)$, with $T = \text{PSU}(3, q)$. Then G is not an IBIS group.*

Proof. Let $q = p^f$, with p prime, and let α be the automorphism of order 2 of the field \mathbb{F}_{q^2} with q^2 elements defined by $\alpha(x) = x^q$. All α -Hermitian forms in a vector space of dimension 3 over \mathbb{F}_{q^2} are equivalent, so we may assume that $\text{GU}(3, q)$ is the isometry group of the form B such that $B((x_1, x_2, x_3), (y_1, y_2, y_3)) = x_1y_3^q + x_2y_2^q + x_3y_1^q$ associated to the matrix

$$\begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

Let

$$M := \left\{ \begin{pmatrix} 1 & -\alpha^q & \beta \\ 0 & 1 & \alpha \\ 0 & 0 & 1 \end{pmatrix} \mid \alpha, \beta \in \mathbb{F}_{q^2}, \beta + \beta^q + \alpha\alpha^q = 0 \right\}.$$

It is not difficult to prove that M is a subgroup of the special unitary group $\text{SU}(3, q)$, and since M intersects the center of $\text{SU}(3, q)$ trivially, it can be identified with a subgroup of $T = \text{PSU}(3, q)$. We shall use without further comment the fact that the trace map from \mathbb{F}_{q^2} to \mathbb{F}_q given by $\text{tr}(\beta) = \beta + \beta^q$ is surjective. Taking a non-zero element $\gamma \in \mathbb{F}_{q^2}$ such that $\gamma + \gamma^q = 0$, then

$$c := \begin{pmatrix} 1 & 0 & \gamma \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

is a non-trivial central element of M . Denote by ω a primitive element of \mathbb{F}_{q^2} , then $\omega^{q+1} \in \mathbb{F}_q$ and consequently there exists $\delta \in \mathbb{F}_{q^2}$ such that $\delta + \delta^q = -\omega\omega^q$. Hence we can consider the following elements of M :

$$a := \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} \text{ and } b := \begin{pmatrix} 1 & -\omega^q & \delta \\ 0 & 1 & \omega \\ 0 & 0 & 1 \end{pmatrix}.$$

Since $\omega^q \neq \omega$, then a and b do not commute. Hence, we have that $a \in C(c) \setminus (C(c) \cap C(b))$, $b \in (C(c) \cap C(b)) \setminus (C(c) \cap C(b) \cap C(a))$, and $c \in (C(c) \cap C(b) \cap C(a))$. So $([1, 1], [1, c], [1, b], [1, a])$ is an

irredundant sequence of size 4 that is not a base. Now, the result follows combining Proposition 3.1 and Remark 2.1. \square

So far, we proved that when G is monolithic primitive, then $T^2 \leq G \leq W(2, T)$ is a non IBIS group. Hence we need to consider the case $k \geq 3$.

Proposition 3.18. *Let $k \geq 3$ and let $T^k \leq G \leq W(k, T)$. Then G is not an IBIS group.*

Proof. First, let $k = 3$ and assume by contradiction that G is an IBIS group. By Proposition 3.1 and Lemma 2.3 we can assume $P_G = \text{Sym}(3)$. In particular $\sigma := (1, 2) \in P_G$, so there exist $y \in \text{Aut}(T)$, $s_2, s_3 \in T$ such that $(y, s_2y, s_3y)\sigma \in G$. That is $(y, y, y)\sigma \in G$ for some $y \in \text{Aut}(T)$. It follows from the Classification of Finite Simple Groups that there exists a non-trivial element t in $C_T(y)$ (see for example ([7, 1.48])). So we have that $(y, y, y)\sigma \in G_{([1,1,1],[1,1,t])}$, while $(y, y, y)\sigma \notin G_{[1,t,t]}$. If $s \notin C_T(t)$, then $(s, s, s) \in G_{[1,1,1] \setminus G_{[1,1,t]}}$. Moreover, $(t, t, t) \in G_{([1,1,1],[1,1,t],[1,t,t])}$. Hence $([1, 1, 1], [1, 1, t], [1, t, t])$ is an irredundant sequence that is not a base. In particular G has an irredundant base of size at least 4. Now, the result follows combining 3.1 and Remark 2.1.

Now we can assume that $k \geq 4$. If P_G does not contain $\text{Alt}(k)$, then $b(G) = 2$ by Proposition 3.1, hence the result follows from Lemma 2.3. So we can assume that $\text{Alt}(k) \leq P_G$. Let $(u_1, \dots, u_k) \in T^k \leq G$. Recall that $g = (y, \dots, y)\sigma$ stabilises $[u_1, \dots, u_k]$ if and only if

$$g^{(u_1, u_2, u_3, \dots, u_k)^{-1}} = (u_1y u_{1\sigma}^{-1}, \dots, u_ky u_{k\sigma}^{-1})\sigma \in D(k, T),$$

i.e. if and only if

$$(3.1) \quad u_1y u_{1\sigma}^{-1} = \dots = u_ky u_{k\sigma}^{-1}.$$

Consider in particular $(u_1, \dots, u_k) = (t_1, t_2, 1, \dots, 1)$ where t_1 and t_2 are chosen so that they are not conjugate in $\text{Aut}(T)$ and $T = \langle t_1, t_2 \rangle$ (choose for instance an involution t_1 , then the existence of t_2 follows from the results in [8]). Let $\tau := [u_1, \dots, u_k]$. We are going to deduce from (3.1) that $g = (y, \dots, y)\sigma$ stabilises τ if and only if 1 and 2 are fixed by σ and $y = 1$. Let $w_i := u_iy u_{i\sigma}^{-1}$ with $w_i = t_i$ if $i \leq 2$, $w_i = 1$ otherwise. Assume $1\sigma^{-1} = j > 2$. Then $w_j = yt_1^{-1}$, and $w_i \in \{y, yt_2^{-1}\}$, when $i \geq 3$ and $i \neq j$. This would imply $w_i \neq w_j$ in contradiction with (3.1). So we must have $1\sigma^{-1} \in \{1, 2\}$. A similar argument shows that $2\sigma^{-1} \in \{1, 2\}$. It follows that $w_1 = w_2 = w_3 = \dots = w_k = y$. If $1\sigma = 2$, then $t_1yt_2^{-1} = y$, i.e. $t_2 = t_1^y$ in contradiction with the assumption that t_1 and t_2 are not conjugate in $\text{Aut}(T)$. So we must have $1\sigma = 1$ and $2\sigma = 2$. This implies $t_1yt_1^{-1} = t_2yt_2^{-1} = y$, hence $y \in C_{\text{Aut}(T)}(\langle t_1, t_2 \rangle) = C_{\text{Aut}(T)}(T) = 1$.

Let $\iota := [(1, \dots, 1)]$. We have proved that $G_{(\iota, \tau)} = \{\sigma \in G \mid 1\sigma = 1, 2\sigma = 2\}$. Since $\text{Alt}(k) \leq P_G$, there exists $a \in \text{Aut}(T)$ such that $z = (a, \dots, a)(1, 2)(3, 4) \in G$. Let $1 \neq s \in C_T(a)$ and consider $\eta := [s, s, 1, \dots, 1]$. Clearly $G_{(\iota, \tau)} \leq G_{(\iota, \eta)}$ and $z \in G_{(\iota, \eta)} \setminus G_{(\iota, \tau)}$. Further, there exists $b \in \text{Aut}(T)$ such that $v := (b, \dots, b)(1, 2, 3) \in G$. It follows from (3.1) and the fact that $s \neq 1$, that $v \in G_\iota \setminus G_\eta$. So we have $G_\iota > G_{(\iota, \eta)} > G_{(\iota, \eta, \tau)} = G_{(\iota, \tau)}$ and therefore G is not IBIS. \square

Combining Propositions 3.9, 3.10, 3.11, 3.12, 3.15, 3.16, 3.17, 3.18, we finally deduce the following result.

Theorem 3.19. *Every monolithic primitive group of diagonal type is a non IBIS group.*

4. PRODUCT TYPE

Let $H \leq \text{Sym}(\Gamma)$ be a primitive group of almost simple type or of diagonal type with socle T . For $k \geq 2$, we consider $W = H \wr \text{Sym}(k)$ acting on $\Omega = \Gamma^k$, with its natural product action. Then G is of product type if $T^k \leq G \leq W$ and the group P of the elements $\sigma \in \text{Sym}(k)$ such that $(h_1, \dots, h_k)\sigma \in G$ for some $h_i \in H$ is transitive. Write $\Omega = \Gamma_1 \times \dots \times \Gamma_k$ where $\Gamma_i = \Gamma$ for each i . By a result of Kovács [10] we may assume that G induces H on each of the k factors Γ_i of Ω . In the whole section we will assume that G is a primitive permutation group of product type.

Theorem 4.1. *The group G is not an IBIS group.*

Proof. To show this result, let assume by contradiction that G is an IBIS group. We are going to construct two irredundant bases of G and we will compare their lengths.

Let $\mathcal{S} := (\gamma_0, \gamma_1, \dots, \gamma_r)$ be an irredundant base of $T \leq \text{Sym}(\Omega)$. By Lemma 2.3 we deduce that

$$(4.1) \quad r \geq 2.$$

Let $\alpha_i = (\gamma_i, \dots, \gamma_i)$ be a k -tuple with $0 \leq i \leq r$ and let $\beta_{i,j}$ be the k -tuple having the i -th component equal to γ_j and the others equal to γ_0 , for $1 \leq i \leq k$ and $1 \leq j \leq r$. The following sequences

$$\mathcal{B}_s = (\alpha_0, \alpha_1, \dots, \alpha_r) \quad \text{and} \quad \mathcal{B}_l = (\alpha_0, \beta_{1,1}, \dots, \beta_{1,r}, \beta_{2,1}, \dots, \beta_{2,r}, \dots, \beta_{k,1}, \dots, \beta_{k,r})$$

are irredundant sequences of G , and \mathcal{B}_s has size $r + 1$, while \mathcal{B}_l has size $1 + kr$. Indeed, from the choice of \mathcal{S} , for every $1 \leq a \leq r - 1$, there exists $u_a \in T$ that stabilizes $\gamma_0, \dots, \gamma_a$ but does not stabilize γ_{a+1} , and consequently $(u_a, u_a, \dots, u_a) \in G$ stabilises $\alpha_0, \alpha_1, \dots, \alpha_a$ but not α_{a+1} . Moreover $(u_a, 1, \dots, 1) \in G$ stabilises $\alpha_0, \beta_{1,1}, \dots, \beta_{1,a}$ but not $\beta_{1,a+1}$, $(1, u_a, \dots, 1) \in G$ stabilises $\alpha_0, \beta_{1,1}, \dots, \beta_{1,r}, \beta_{2,1}, \dots, \beta_{2,a}$ but not $\beta_{2,a+1}$ and proceeding with this argument we finally get that $(1, 1, \dots, u_a) \in G$ stabilises $\alpha_0, \beta_{1,1}, \dots, \beta_{1,r}, \beta_{2,1}, \dots, \beta_{2,r}, \dots, \beta_{k,1}, \dots, \beta_{k,a}$, but not $\beta_{k,a+1}$.

Let $X := G \cap H^k$. It is not difficult to observe that

$$X_{(\mathcal{B}_s)} = X_{(\mathcal{B}_l)} = \{(h_1, \dots, h_k) \in X \mid h_1 \in H_{(\mathcal{S})}, \dots, h_k \in H_{(\mathcal{S})}\}.$$

Now, let $\mathcal{A} = (\delta_1, \dots, \delta_b)$ be a base of minimal size of $X_{(\mathcal{B}_s)} = X_{(\mathcal{B}_l)}$. Hence $\mathcal{B}_s \sqcup \mathcal{A}$ and $\mathcal{B}_l \sqcup \mathcal{A}$ are still irredundant sequences for G , where \sqcup denotes the concatenation of the two sequences.

Here, let $\delta_1, \dots, \delta_c$, and $\nu_1, \dots, \nu_{c'}$ be elements of G such that the sequences $\mathcal{B}_1 = \mathcal{B}_s \sqcup \mathcal{A} \sqcup \{\delta_1, \dots, \delta_c\}$ and $\mathcal{B}_2 = \mathcal{B}_l \sqcup \mathcal{A} \sqcup \{\nu_1, \dots, \nu_{c'}\}$ are irredundant bases of G . Since G is IBIS, then $|\mathcal{B}_1| = r + 1 + b + c = 1 + rk + b + c' = |\mathcal{B}_2|$. Hence

$$(4.2) \quad r(k - 1) = c - c' \leq c.$$

Note that $G_{(\mathcal{B}_s \sqcup \mathcal{A})} \cap X = 1$, then $G_{(\mathcal{B}_s \sqcup \mathcal{A})} \cong G_{(\mathcal{B}_s \sqcup \mathcal{A})} / (G_{(\mathcal{B}_s \sqcup \mathcal{A})} \cap X) \cong G_{(\mathcal{B}_s \sqcup \mathcal{A})} X / X \leq \text{Sym}(k)$. So, denoting by $\ell(\text{Sym}(k))$ the maximal length of a chain of subgroups of $\text{Sym}(k)$, we can estimate c with $\ell(\text{Sym}(k))$. From (4.2) and (4.1) we deduce that

$$(4.3) \quad 2(k - 1) \leq \ell(\text{Sym}(k)).$$

By [3] we have that $\ell(\text{Sym}(k)) < 3k/2$, hence (4.3) yields $k < 4$. Since $\ell(\text{Sym}(2)) = 1$ and $\ell(\text{Sym}(3)) = 2$, (4.3) allows us to exclude also $k = 2$ and $k = 3$. This contradicts the assumption $k \geq 2$ and the result follows. \square

5. TWISTED WREATH PRODUCT TYPE

Let T be a non-abelian simple group, and let k be an integer that is at least 2. A group G of twisted wreath product type with socle T^k acts primitively on a set $|\Omega|$ with degree $|T^k|$ and is also a twisted wreath product, which is a split extension of T^k by a transitive subgroup P of $\text{Sym}(k)$. In the whole section we will assume that G is a primitive permutation group of twisted wreath product type.

Theorem 5.1. *The group G is not an IBIS group.*

Proof. We may identify G with a subgroup of the wreath product $W = \text{Aut}(S) \wr P$, with the property that $\text{soc}(G) = T^k$ and T^k has a maximal complement, say H , in G , which is isomorphic to P . In particular for every $\sigma \in P$ there exists a unique element $(a_1, \dots, a_k) \in (\text{Aut}(T))^k$ such that $(a_1, \dots, a_k)\sigma \in H$.

By [5, Theorem 5.0.1], if P is a primitive subgroup of $\text{Sym}(k)$, then $b(G) = 2$. In this case, since $\text{soc}(G)$ is non-abelian, the result follows from Lemma 2.3. So we may assume that k is not a prime.

Since P is transitive, it contains a fixed-point free permutation σ . Either σ or one of its powers (in the case when σ is a k -cycle) admits at least two orbits of size at least 2. So it is not restrictive to assume that P contains an element

$$(5.1) \quad \rho = (1, \dots, r)(r+1, r+2, \dots, r+s)\tau$$

for some $r, s \geq 2$ and $\tau \in \text{Sym}(k)$ which pointwise fixes the set $\{1, \dots, r+s\}$. There exists a unique element $(a_1, \dots, a_k) \in (\text{Aut}(T))^k$ such that $h := (a_1, \dots, a_k)\rho \in H$. It follows from the Classification of Finite Simple Groups that $C_T(a_1 \cdots a_r) \neq 1$ ([7, 1.48]). In particular there exists $t \in C_T(a_1 \cdots a_r)$ of prime order, say p . Choose $u \in T$ of prime order q , with $q \neq p$, and consider the following elements of T^k : $y_1 = (t, t^{a_1}, t^{a_1 a_2}, \dots, t^{a_1 a_2 \cdots a_{r-1}}, 1, \dots, 1)$, $y_2 = (1, \dots, 1, u, 1, \dots, 1)$, where u is the entry in position $r+1$, and $y = y_1 y_2$. We have that $h \in C_H(y_1) = C_H(y^q)$ but $h \notin C_H(y)$. Hence $H \cap H^{y^q} \cap H^y = H \cap H^y = C_H(y) < C_H(y^q) = H \cap H^{y^q} < H$ and therefore G is not IBIS. \square

6. PROOF OF THEOREM 1.1

We work through the non-affine and non-almost simple classes of the O’Nan-Scott Theorem. If G is of diagonal type, as showed in Propositions 3.4, 3.2, 3.1 and Theorem 3.19, there is a unique infinite family of IBIS groups. If G is of product action type or twisted wreath product, then the result follows immediately from Sections 4 and 5 respectively. Hence the proof is complete.

ACKNOWLEDGMENTS. All authors are members of GNSAGA.

REFERENCES

- [1] P. J. Cameron, *Permutation Groups*, London Mathematical Society Student Texts. Cambridge University Press, 1999.
- [2] P. J. Cameron and D. G. Fon-Der-Flaas, Bases for Permutation Groups and Matroids, *Europ. J. Combin.* **16** (1995), 537–544.
- [3] P. J. Cameron, R. Solomon and A. Turull, Chains of Subgroups in Symmetric Groups, *J. Algebra* **127** (1989), 340–352.
- [4] J. B. Fawcett, The base size of a primitive diagonal group, *J. Algebra* **375** (2013), 302–321.
- [5] J. B. Fawcett, Bases of primitive permutation groups, Phd dissertation.
- [6] GAP, The GAP Group, <https://www.gap-system.org>
- [7] D. Gorenstein, *Finite simple groups. An introduction to their classification*. University Series in Mathematics. Plenum Publishing Corp., New York, 1982.
- [8] R. M. Guralnick and W. M. Kantor, Probabilistic generation of finite simple groups. Special issue in honor of Helmut Wielandt. *J. Algebra* 234 (2000), 743–792.
- [9] B. Huppert, *Endliche Gruppen I*, Springer, Berlin, 1967.
- [10] L.G. Kovács, Primitive subgroups of wreath products in product action, *Proc. London Math. Soc.* **58** (1989), 306–322.
- [11] D. Leemans and M. W. Liebeck, Chiral polyhedra and finite simple groups, *Bull. Lond. Math. Soc.* **49** (2017), no. 4, 581–592.
- [12] M. W. Liebeck, C. E. Praeger and J. Saxl, On the O’Nan-Scott theorem for finite primitive permutation groups. *J. Australian Math. Soc. (A)* **44** (1988), 389–396.
- [13] D. J. S. Robinson, *A course in the theory of groups*, Springer Science & Business Media, 1996.

- [14] M. Suzuki, The nonexistence of a certain type of simple groups of odd order, *Proc. Amer. Math. Soc.* **8** (1957), 686–695.

ANDREA LUCCHINI, DIPARTIMENTO DI MATEMATICA “TULLIO LEVI-CIVITA”, UNIVERSITÀ DEGLI STUDI DI PADOVA, VIA TRIESTE 63, 35121, PADOVA, ITALY
Email address: `lucchini@math.unipd.it`

MARTA MORIGI, UNIVERSITÀ DI BOLOGNA, PIAZZA DI PORTA SAN DONATO 5, 40126 BOLOGNA, ITALY
Email address: `marta.morigi@unibo.it`

MARIAPIA MOSCATIELLO, DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI BOLOGNA, PIAZZA DI PORTA SAN DONATO 5, 40126 BOLOGNA, ITALY
Email address: `mariapia.moscatiello@unibo.it`