

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Detection and identification of a 16SrIII-J subgroup phytoplasma associated with faba bean in Peru

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Torres Suarez G., Gamarra Gamarra D., Villar C.M., Llacza Munive S.L., Satta E., Carrasco Lozano E.C., et al. (2021). Detection and identification of a 16SrIII-J subgroup phytoplasma associated with faba bean in Peru. JOURNAL OF PHYTOPATHOLOGY, 169(4), 203-208 [10.1111/jph.12977].

Availability:

This version is available at: <https://hdl.handle.net/11585/846401> since: 2022-01-19

Published:

DOI: <http://doi.org/10.1111/jph.12977>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the peer reviewed version of the following article:

Gilberto Torres Suarez, Delia Gamarra Gamarra, Charo M. Villar, Sally L. Llacza Munive, Eleonora Satta, Emerson C. Carrasco Lozano, Assunta Bertaccini
Detection and identification of a 16SrIII-J subgroup phytoplasma associated with faba bean in Peru

which has been published in final form at

JOURNAL OF PHYTOPATHOLOGY Volume 169, Issue 4 April 2021

Pages 203-208

<https://dx.doi.org/10.1111/jph.12977>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions. This article may not be enhanced, enriched or otherwise transformed into a derivative work, without express permission from Wiley or by statutory rights under applicable legislation. Copyright notices must not be removed, obscured or modified. The article must be linked to Wiley's version of record on Wiley Online Library and any embedding, framing or otherwise making available the article or pages thereof by third parties from platforms, services and websites other than Wiley Online Library must be prohibited.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)
When citing, please refer to the published version.

1 **Detection and identification of a 16SrIII-J subgroup phytoplasma associated**
2 **with faba bean in Peru**

3
4 Gilberto Torres Suarez¹, Delia Gamarra Gamarra¹, Charo Milagros Villar¹, Sally.
5 Lorena Llacza Munive¹, Eleonora Satta², Emerson Clovis Carrasco Lozano¹,
6 Assunta Bertaccini²

7
8 ¹ Center for Plant Molecular Biology Research, Universidad Nacional del Centro del
9 Peru, Av. Mariscal Castilla N° 3909, El Tambo, Huancayo, Peru.

10 ² Department of Agricultural and Food Sciences, *Alma Mater Studiorum* –
11 University of Bologna, viale G. Fanin, 40, 40127, Bologna, Italy.

12
13 Correspondence

14 Emerson Clovis Carrasco Lozano, Center for Plant Molecular Biology Research,
15 Universidad Nacional del Centro del Peru, Av. Mariscal Castilla N° 3909, El
16 Tambo, Huancayo, Peru.

17 E-mail: ecarrasco@uncp.edu.pe; eccarrascol@unal.edu.co

18
19 **KEYWORDS:** *Vicia faba*, 16SrIII phytoplasma, dwarfism, abnormal leaf sprouts,
20 Peru.

25 **Abstract**

26 Faba bean samples with symptoms of yellowing, dwarfism, shoot proliferation,
27 internode shortening, leaf sprouts and lack of pod and seed production were
28 collected from Huancayo and Chupaca provinces, Junin-Peru, and analyzed to
29 verify phytoplasma presence and identity. After total nucleic acid extraction, the
30 amplification with universal phytoplasma primers, using nested polymerase chain
31 reactions, on the 16S ribosomal gene followed by restriction fragment length
32 polymorphism analysis and sequence analysis allowed the classification of the
33 detected phytoplasma in the subgroup 16SrIII-J. The alignment of 30 16S
34 ribosomal gene sequences from 15 faba bean symptomatic samples from single
35 plants allowed verifying the consistent presence of 5 single nucleotide
36 polymorphisms that are however not modifying the phytoplasma classification. The
37 phytoplasma identity was also corroborated by the amplification and the restriction
38 fragment length polymorphism analyses carried out on the ribosomal protein gene
39 amplicons obtained with primers specific for the phytoplasmas enclosed in the
40 16SrIII group. This is the first description in Peru of a disease associated with
41 phytoplasmas in faba beans.

42

43 **2 | INTRODUCTION**

44 *Vicia faba* L., known as faba bean, is a leguminous species originated in the
45 Mediterranean region and grown worldwide for its high content of carbohydrate,
46 protein, vitamins, minerals (Zn, Fe, Mg, Ca) and omega-3 (Etemadi et al. 2019). In
47 2017, Peru had a cropping area of 65,985 ha, with an average yield of 11,478
48 kg/ha and total seed production of 75,736 tons (FAOSTAT 2019). Faba bean is

49 worldwide infected by various fungi, bacteria, chromista, viruses, and nematodes.
50 The main diseases reported are chocolate spot (caused by *Botrytis fabae*), leaf
51 blight (*Xanthomonas campestris*), bacterial spot (*Pseudomonas syringae*), root rot
52 (*Fusarium solani*), powdery mildew (*Erysiphe cichoracearum*), and infections by
53 viral complexes (Etemadi et al. 2019). During the last decades *V. faba* was also
54 reported to be infected by phytoplasmas (Castro and Romero 2004; Al-Saleh and
55 Amer 2014; Salehi et al. 2016; Omar 2017), that are bacteria discovered in 1967
56 (Doi et al. 1967) lacking the cell wall and having a diameter of 200 to 800 nm. Their
57 genomes are of small size and possess variable numbers of extrachromosomal
58 fragments (Nishigawa et al. 2003). Phytoplasmas are hosted in the plant phloem
59 and in the hemolymph of their insect vectors that are mainly those having siphon-
60 like mouth systems, as leafhoppers and psyllids (Weintraub and Beanland 2006).
61 They infect hundreds of plant species in which different symptoms can be
62 observed; moreover, different phytoplasmas could be associated with
63 indistinguishable symptomatology in diverse plant host species (Bertaccini 2007,
64 Namba 2019). These phytoplasma-associated symptoms include leaf yellowing,
65 growth reduction, virescence, and witches' broom, shortening of internodes, small
66 leaves, phyllody, floral malformation, and plant death (Salehi et al. 2016; Kumari et
67 al. 2019).

68 Since their discovery it was difficult to detect and identify the phytoplasmas
69 in symptomatic plants since they are present in a reduced concentration in the
70 infected tissues where are unevenly distributed (Berges et al. 2000). Phytoplasmas
71 belong to the class *Mollicutes* provisionally enclosed in the '*Candidatus*
72 *Phytoplasma*' genus (IRPCM 2004). PCR/RFLP analyses on their 16S ribosomal

73 gene allowed the phytoplasma classification in ribosomal groups and subgroups
74 that is integrating the provisional classification for further differentiation among the
75 detected strains (Bertaccini and Lee 2018).

76 In recent years in the faba bean fields located in the Junin region, symptoms
77 related to phytoplasma presence have been observed with 8% incidence, however,
78 to date the possible presence and identity of phytoplasmas was not studied.
79 Therefore, the objective of this work was to identify the phytoplasma associated
80 with the disease of the faba bean in the provinces of Huancayo and Chupaca in the
81 Junin region of Peru.

82

83 **2 | MATERIALS AND METHODS**

84 **2.1| Sample collection and DNA extraction**

85 Faba bean samples with symptoms of yellowing, dwarfism, shoot proliferation,
86 internode shortening, leaf sprouts and lack of pod and seed production (Figure 1a-
87 b) were collected from Huancayo (15 samples) and Chupaca provinces (16
88 samples), Junin-Peru, and analyzed to verify phytoplasma presence and identity.
89 The 31 samples each collected from each from a separate plant, were processed
90 and dehydrated with silica gel, then leaf tissues were sprayed with liquid nitrogen,
91 and 25 mg were weighed for the extraction of DNA with innuPREP Plant DNA Kit
92 (Analytik Jena, Germany) according to the instructions of the manufacturer. For the
93 negative control, DNAs of four symptomless plants were also extracted following
94 the same procedure. As positive control, the DNA of a strain from *Euscelidius*
95 *variegatus* (16SrIII-B, API) maintained in the micropropagated collection
96 (Bertaccini, 2014) was used. The quality of the obtained DNA was verified by

97 means of a 1% agarose gel and quantified with a spectrophotometer NanoDrop™
98 (ThermoScientific™).

99 **2.2| PCR detection and RFLP analysis**

100 Phytoplasma detection was achieved using PCR assays with P1/P7 (Deng and
101 Hiruki 1991; Schneider et al. 1995), R16mF2/mR1 primers followed in nested PCR
102 reaction by R16F2n/R2 (Gundersen and Lee 1996) and fU5/rU3 (Lorenz et al.
103 1995) primers. The total nucleic acid used was about 20-40 ng for the PCR
104 reactions and amplicons, diluted in sterile distilled water 1:30, were used as
105 template for the nested PCR reactions. Cycling conditions reported (Schaff et al.
106 1992) were employed in all the PCR reactions. The amplicon visualization was
107 obtained by electrophoresis in 1% agarose gel followed by ethidium bromide (10
108 ng/μl) staining. The R16F2n/R2 and fU5/rU3 amplicons were analyzed by RFLP
109 using *Tru1* and *HpaII* (Fermentas, Vilnius, Lithuania) restriction enzymes and
110 compared with the corresponding profile of the phytoplasma strain API in 6.7%
111 polyacrylamide gels stained and visualized as for the agarose gels reported above.
112 Further molecular characterization was achieved in nested PCR on the ribosomal
113 protein (rp) operon with primer pairs rpL2F3/rp(I)R1A and rp(III)F1/rp(III)R1,
114 specifically amplifying about 800 bp in the rp coding gene of the phytoplasmas
115 enclosed in ribosomal group 16SrIII, under the reported conditions (Martini et al.
116 2007). The following phytoplasma strains enclosed in the ribosomal group 16SrIII
117 and maintained in periwinkle (*Catharanthus roseus*) as a collection of
118 micropropagated shoots (Bertaccini 2014) were used as controls: X disease
119 phytoplasma (16SrIII-A, CX), a strain from *Euscelidius variegatus* (16SrIII-B, API),
120 *Vaccinium* witches' broom (16SrIII-F, VAC), spirea stunt (16SrIII-E, SPI), and

121 poinsettia branching (16SrIII-H, JRI). The amplicons were analyzed by RFLP with
122 the restriction enzyme *Tru1I*, observed after electrophoresis and visualized as for
123 the agarose gels reported above.

124 **2.3| Sequencing and phylogenetic analysis of 16S rDNA**

125 The extraction from agar of R16mF2/R16mR1 and fU5/rU3 amplified fragments of
126 approximately 1,400 and 876 bp respectively, was performed with the PCR Clean-
127 Up System Kit (Promega, USA). Amplicons were then sent for both directions
128 sequencing using the primer pairs employed in the amplification at Macrogen
129 (Seoul, Korea). The sequence alignment and manual revision were done with the
130 BLASTn program, followed by the comparison with phytoplasma sequences in the
131 GenBank. The consensus sequence obtained was then subjected to virtual RFLP
132 analysis with the *iPhyClassifier* using all the 17 available restriction enzymes with
133 the 16SrIII subgroup strains (Zhao et al. 2009). Phylogenetic analysis with the
134 Minimum Evolution system (Rzhetsky and Nei 1992) was also carried out using
135 MEGA 6 (Tamura et al. 2013) with the 43 described 'Ca. Phytoplasma' species and
136 with a number of phytoplasma strains enclosing those classified in 16SrIII-J
137 subgroup or reported from Peru and other South American agricultural areas.

138

139

140 **3| RESULTS**

141 From the 31 symptomatic samples collected from individual plants no or weak
142 amplification was obtained with P1/P7 primers while two samples were amplified
143 with primers R16mF2/R16mR1. By using primers amplifying shorter fragments in
144 nested PCR the detection of phytoplasma expected length bands was obtained in

145 all the samples from symptomatic plants; in particular primers fU5/rU3 (amplicons
146 of approximately 850 bp) provided 100% of amplification from all the samples from
147 symptomatic plants and primers R16F2n/R2 (amplicons of approximately 1,200 bp)
148 provided 80% of amplification (data not shown). The samples from asymptomatic
149 plants and the sterile water used as negative controls did not produce bands in any
150 PCR reaction. From the fU5/rU3 amplified fragments 10 samples were selected for
151 direct sequencing, the remaining fragments were subjected to RFLP analysis with
152 *Tru1I* and *HpaII* (Figure 2a). The profiles obtained resulted identical to those of the
153 strain from *Euscelidius variegatus* (16SrIII-B, API) (data not shown). The two 1,400
154 bp amplicons obtained with R16mF2/R16mR1 were sequenced in both directions
155 and the consensus sequence of 1,320 bp (4 times coverage for each nucleotide in
156 each position) was deposited in the GenBank under the accession number
157 MT358307; this strain is referred here as faba bean yellows and witches' broom
158 (FB-YWB) phytoplasma. The R16F2n/R2 sequence (1,240 bp) analyzed *in silico*
159 RFLP by the *iPhyClassifier* (Figures 2b, c and d) was having a 1.00 as similarity
160 coefficient to the phytoplasma strains enclosed in the subgroup 16SrIII-J. The
161 phylogeny allowed clustering the FB-YWB sequence with phytoplasma strains
162 enclosed in this ribosomal subgroup detected in other geographic areas of South
163 America (Figures 3a and b). The alignment of 30 16Sr sequences amplified with
164 fU5/rU3 primers (about 800 bp) from the faba bean phytoplasma strains detected
165 in Peru showed the consistent presence of three SNPs (*i.e.* present in more than
166 three sequences) at positions 384 (A/G), 442 (C/T), 831 (C/A) and of three
167 insertions T at 492, C at 827, G at 1015 counted 3' to 5' in the deposited
168 consensus sequence (data not shown). The specific amplification of the

169 phytoplasmas with rp 16SrIII group-specific primers followed by RFLP analyses
170 with *Tru1I* (Figure 4), corroborated the close genetic relationship of this
171 phytoplasma with those classified in the 16SrIII subgroup.

172

173 **4| DISCUSSION**

174 The detection of 16SrIII-J in symptomatic faba bean samples is the first
175 identification of phytoplasmas enclosed in this ribosomal subgroup in Peru; they
176 were reported first in Brazil (Montano et al. 2000) and later infecting several other
177 crop species in South American states such as Argentina (Galdeano et al. 2013;
178 Fernández et al. 2020), Brazil (Munhoz et al. 2019), and Chile (Quiroga et al.,
179 2020). The identification of variability at the 16Sr level is not unusual in
180 phytoplasmas classified in the 16SrIII group where the frequent presence of
181 interoperon heterogeneity is reported in several strains (Davis et al. 2013).

182 The first studies of phytoplasmas in Peru date back to the late 1970s (Nault
183 et al. 1979). Based on serological and microscopic techniques, the presence of
184 phytoplasmas, spiroplasmas, and viruses in corn crops was verified. After that, a
185 survey of phytoplasmas in different regions and crops allowed identifying
186 phytoplasmas enclosed in ribosomal group 16SrI in diverse plants species, of
187 group 16SrII in potato (*Solanum tuberosum*), while phytoplasmas in the 16SrIII
188 group were detected in tomato (*Solanum lycopersicum*) and dandelion (*Taraxacum*
189 sp.) (Hodgetts et al. 2009). More recently, a phytoplasma classified in the 16SrXV-
190 A was detected in grapevine (Wei et al. 2017).

191 Phytoplasmas have not been identified in faba bean in Peru previously,
192 however, a phytoplasma enclosed in the 16SrIII group was reported in Spain

193 (Castro and Romero 2004), while in Saudi Arabia (Al-Saleh and Amer 2014, Omar,
194 2017) and Iran (Salehi et al. 2016) 16SrII phytoplasma strains were identified in
195 symptomatic plants. The identified phytoplasma is largely reported in South
196 American countries it is therefore necessary to continue the studies to verify the
197 presence of insect vectors following the recent demonstration of *Paratanus*
198 *exitiosus* and *Bergallia valdiviana* in Chile (Quiroga et al. 2019). Furthermore,
199 assays to verify a possible seed transmission, reported for other phytoplasmas in
200 tomato, corn (*Zea mays*) and carrot (*Daucus carota*) (Calari et al. 2011; Satta et al.
201 2020) should be planned.

202

203 **ACKNOWLEDGEMENTS**

204 This research was financed by the Vice-Rector's Office for Research (CANON
205 Projects, resolution No. 1565-R-2017) of the Universidad Nacional del Centro del
206 Peru. We are also thankful to the members of the research team of the Center for
207 Plant Molecular Biology Research. Research data are not shared.

208

209 **ORCID**

210 Gilberto Torres Suarez: <https://orcid.org/0000-0003-2179-4922>

211 Delia Gamarra Gamarra: <https://orcid.org/0000-0002-6262-237X>

212 Charo Milagros Villar: <https://orcid.org/0000-0002-1267-3697>

213 Sally Lorena Llacza Munive: <https://orcid.org/0000-0003-1055-9951>

214 Eleonora Satta: <https://orcid.org/0000-0002-2566-5928>

215 Emerson Clovis Carrasco Lozano: <https://orcid.org/0000-0001-6075-6457>

216 Assunta Bertaccini: <https://orcid.org/0000-0002-5650-1512>

217

218 **REFERENCES**

- 219 Al-Saleh, M.A., Amer M.A. (2014) Molecular characterization of the 16SrII group of
220 phytoplasma associated with faba bean (*Vicia faba* L.) in Saudi Arabia.
221 Journal of Animal and Plant Sciences, 24:221–228
- 222 Berges, R., Rott, M., Seemüller. E. (2000) Range of phytoplasma concentration in
223 various plant hosts as determined by competitive polymerase chain reaction.
224 Phytopathology 90:1145–1152
- 225 Bertaccini, A. (2007) Phytoplasmas: diversity, taxonomy, and epidemiology.
226 Frontiers in Bioscience 12:673–689
- 227 Bertaccini, A. (2014) Internet Resource: <http://www.ipwgnet.org/collection> (verified
228 March 10, 2020)
- 229 Bertaccini, A., Lee, I-M. (2018) Phytoplasmas: an update. In: Rao GP, Bertaccini A,
230 Fiore N, Liefting LW. (eds) Phytoplasmas: Plant Pathogenic Bacteria - I
231 Characterisation and epidemiology of phytoplasma-associated diseases.
232 Singapore, Springer, pp 1–29.
- 233 Calari, A., Paltrinieri, S., Contaldo, N., Sakalieva, D., Mori, N., Duduk, B.,
234 Bertaccini, A. (2011) Molecular evidence of phytoplasmas in winter oilseed
235 rape, tomato and corn seedlings. Bulletin of Insectology, 64(Supplement)
236 S157–S158
- 237 Castro, S., Romero, J. (2004) First detection of a phytoplasma infecting faba bean
238 (*Vicia faba* L.) in Spain. Spanish Journal of Agricultural Research 2:253–256
- 239 Davis, R.E., Zhao, Y., Dally, E.L., Lee, I-M., Jomantiene, R., Douglas, S.M. (2013)
240 ‘*Candidatus* Phytoplasma pruni’, a novel taxon associated with X-disease of

241 stone fruits, *Prunus* spp.: multilocus characterization based on 16S rRNA,
242 secY, and ribosomal protein genes. International Journal of Systematic and
243 Evolutionary Microbiology 63:766–776

244 Deng, S.J., Hiruki, C. (1991) Amplification of 16S ribosomal-RNA genes from
245 culturable and nonculturable Mollicutes. Journal of Microbiological Methods
246 14:53–61

247 Doi, Y., Teranaka, M., Yora, K., Asuyama, H. (1967) Mycoplasma or PLT group
248 like microorganisms found in the phloem elements of plants infected with
249 mulberry dwarf, potato witches' broom, aster yellows or paulownia witches'
250 broom. Japanese Journal of Phytopathology 33:259–266

251 Etemadi, F., Hashemi, M., Barker, A.V., Zandvakili, O.R., Liu, X. (2019) Agronomy,
252 nutritional value, and medicinal application of faba bean (*Vicia faba* L.).
253 Horticultural Plant Journal 5:170–182

254 FAOSTAT (2019) Internet Resource: <http://www.fao.org/faostat/en/#data/QC>
255 (verified Dec 10, 2019)

256 Felsenstein, J. (1985) Confidence limits on phylogenies: an approach using the
257 bootstrap. Evolution 39:783–791

258 Fernández, F.D., Guzmán, F.A., Baffoni, P., Reinoso, L., Kiehr, M., Delhey,
259 Favere, V.R., Galdeano, E., Conci, L.R., (2020) Phytoplasmas of subgroup
260 16SrIII-J associated with *Beta vulgaris* in Argentina. Tropical Plant Pathology
261 45:143–147.

262 Galdeano, E., Guzmán, F.A., Fernández, F.D., Conci, L.R. (2013) Genetic diversity
263 of 16SrIII group phytoplasmas in Argentina. Predominance of subgroups
264 16SrIII-J and -B and two new subgroups 16SrIII-W and X. European Journal

265 of Plant Pathology 137:753–764

266 Gundersen, D.E., Lee, I-M. (1996) Ultrasensitive detection of phytoplasmas by
267 nested-PCR assays using two universal primer pairs. *Phytopathologia*
268 *Mediterranea* 35:114–151

269 Hodgetts, J., Chuquillanqui, C., Gamarra, D., Muller, G., Arocha, Y., Pinillos, O.,
270 Velit, E., Lozada, P., Boa, E., Boonham, N., Mumford, R., Barker, I.,
271 Dickinson, M. (2009) Surveys reveal the occurrence of phytoplasmas in
272 plants at different geographical locations in Peru. *Annals of Applied Biology*
273 155:15–27

274 IRPCM (2004) '*Candidatus* Phytoplasma', a taxon for the wall-less, non-helical
275 prokaryotes that colonize plant phloem and insects. *International Journal of*
276 *Systematic and Evolutionary Microbiology* 54:1243–1255

277 Kumari, S., Nagendran, K., Rai, A.B., Singh, B., Rao, G.P., Bertaccini, A. (2019)
278 Global status of phytoplasma diseases in vegetable crops. *Frontiers in*
279 *Microbiology* 10:1349

280 Lorenz, K-H., Schneider, B., Ahrens, U., Seemiiller, E. (1995) Detection of the
281 apple proliferation and pear decline phytoplasmas by PCR amplification of
282 ribosomal and nonribosomal DNA. *Phytopathology* 85:771–776

283 Martini, M., Lee, I-M., Bottner, K.D., Zhao, Y., Botti, S., Bertaccini, A., Harrison,
284 N.A., Carraro, L., Marcone, C., Khan, A.J., Osler, R. (2007) Ribosomal protein
285 gene-based phylogeny for finer differentiation and classification of
286 phytoplasmas. *International Journal of Systematic Bacteriology* 57:2037–
287 2051

288 Montano, H.G., Davis, R.E., Dally, E.L., Pimentel, J.P., Brioso, P.S.T. (2000)

289 Identification and phylogenetic analysis of a new phytoplasma from diseased
290 chayote in Brazil. *Plant Disease* 84:429–436

291 Munhoz, E., Pereira, T., Bedendo, I. (2019) Molecular characterization of a
292 phytoplasma associated with a commercial variety of *Momordica charantia*.
293 *Scientia Agricola* 76:232-236

294 Nishigawa, H., Oshima, K., Miyata, S., Ugaki, M., Namba, S. (2003) Complete set
295 of extrachromosomal DNAs from three pathogenic lines of onion yellows
296 phytoplasma and use of PCR to differentiate each line. *Journal of General*
297 *Plant Pathology* 69:194-198.

298 Namba, S. (2019) Molecular and biological properties of phytoplasmas.
299 *Proceedings of the Japan Academy Serie B* 95:401–418

300 Nault, L.R., Gordon, D.T., Gingery, R.E., Bradfute, O.E., Castillo-Loayza, J. (1979)
301 Identification of maize viruses and mollicutes and their potential insect vectors
302 in Peru. *Phytopathology* 69:824–828

303 Omar, A.F. (2017) Detection and molecular characterization of phytoplasmas
304 associated with vegetable and alfalfa crops in Qassim region. *Plant-*
305 *Microorganism Interactions* 12:58–66

306 Quiroga, N., Longone, V., González, X., Zamorano, A., Pino, A.M., Picciau, L.,
307 Alma, A., Paltrinieri, S., Contaldo, N., Bertaccini, A., Fiore, N. (2019)
308 Transmission of 16SrIII-J phytoplasmas by *Paratanus exitiosus* (Beamer)
309 and *Bergallia valdiviana* Berg 1881 leafhoppers. *Phytopathologia*
310 *Mediterranea* 58:231–237

311 Quiroga, N., Gamboa, C., Soto, D., Pino, A.M., Zamorano, A., Campodonico, J.,
312 Alma, A., Bertaccini, A., Fiore, N. (2020) Update and new epidemiological

313 aspects about grapevine yellows in Chile. *Pathogens* 9:933.

314 Rzhetsky, A., Nei, M. (1992) A simple method for estimating and testing minimum
315 evolution trees. *Molecular Biology and Evolution* 9:945–967

316 Salehi, M., Rasoulpour, R., Izadpanah, K. (2016) Molecular characterization, vector
317 identification and partial host range determination of phytoplasmas
318 associated with faba bean phyllody in Iran. *Crop Protection* 89:12–20

319 Satta, E., Carminati, G., Bertaccini, A. (2020) Phytoplasma presence in carrot
320 seedlings. *Australasian Plant Disease Notes* 15:11

321 Schaff, D.A., Lee, I-M., Davis, R.E. (1992) Sensitive detection and identification of
322 mycoplasmalike organisms by polymerase chain reactions. *Biochemical and*
323 *Biophysical Research Communications* 186:1503–1509

324 Schneider, B., Seemüller, E., Smart, C.D., Kirkpatrick, B.C. (1995) Phylogenetic
325 classification of plant pathogenic mycoplasmalike organisms or
326 phytoplasmas. In: Razin S, Tully JG. (eds) *Molecular and diagnostic*
327 *procedures in Mycoplasmaology*. San Diego, CA, USA, Academic press, pp
328 369–380.

329 Tamura, K., Nei, M. (1993) Estimation of the number of nucleotide substitutions in
330 the control region of mitochondrial DNA in humans and chimpanzees.
331 *Molecular Biology and Evolution* 10:512–526

332 Tamura, K., Peterson, D., Stecher, G., Filipski, A., Kumar, S. (2013) MEGA6:
333 molecular evolutionary genetics analysis version 6.0. *Molecular Biology and*
334 *Evolution* 30:2725–2729

335 Wei, W., Pérez-López, E., Bermúdez-Díaz, L., Davis, R.E., Granda-Wong, C.,
336 Zhao, Y. (2017) First report of a new grapevine yellows disease in Peru and

337 its association with infection by a '*Candidatus* Phytoplasma brasiliense'
338 related phytoplasma strain. *Plant Disease* 101:502

339 Weintraub, P.G., Beanland, L. (2006) Insect vectors of phytoplasmas. *Annual*
340 *Review of Entomology* 51:91–111

341 Zhao, Y., Wei, W., Lee, I-M., Shao, J., Suo, X., Davis, R.E. 2009. Construction of
342 an interactive online phytoplasma classification tool, *iPhyClassifier*, and its
343 application in analysis of the peach X-disease phytoplasma group (16SrIII).
344 *International Journal of Systematic and Evolutionary Microbiology* 59:2582–
345 2593.

346

347

348

349

350

351


352



353


354 **Figure 1** *Vicia faba* plants with symptoms of yellowing, dwarfism, leaf distortion,
355 and abnormal leaf sprouts near to healthy plants (a) and in a close up (b) in a
356 production field in the province of Chupaca, Peru.

357



358


359 **Figure 2** a) Polyacrylamide 6.7% gels showing the restriction profiles of 16S
 360 ribosomal amplicons of faba bean phytoplasma strains from Peru: from left
 361 digestion with *Tru1I* of R16F2n/R16R2 amplicons followed by digestion of the
 362 same samples with the same enzyme of fU5/rU3 amplicons; after the second
 363 marker (P) the same amplicons digested with *HpaII*. Samples 1, 2, and 4, broad
 364 bean samples from the diverse localities. b), c) and d) *In silico* RFLP patterns of
 365 the R16F2n/R2 sequence of the FB-YWB strain (GenBank accession number
 366 MT358307) in comparison with the 16S rIII available phytoplasma subgroups using
 367 the *iPhyClassifier*. Ribosomal subgroups are listed at the top; lane with no name is
 368 strain FB-YWB. *Tru1I* and *MseI* are isoschizomer enzymes. P and MW, marker
 369 Φ X174 digested with *HaeIII* and having the fragment sizes from top to bottom of
 370 1,353; 1,078; 872; 603; 310; 281; 271; 234; 194; 118 and 72 base pairs,
 371 respectively.



372

373 **Figure 3** a) and b) Phylogenetic trees showing relationships at the 16S ribosomal
 374 RNA gene level of the faba bean yellows and witches' broom phytoplasma strain
 375 from Peru (FB-YWB) (in red) compared with the classified '*Ca. Phytoplasma*'
 376 species and with some phytoplasmas classified in the group 16SrIII (reported
 377 ribosomal subgroups are listed on the right in blue) using the Minimum Evolution
 378 method. The replicate tree percentage indicating the clustering of the associated
 379 taxa are next to the branches (Felsenstein 1985). The Tamura-Nei method (1993)
 380 was used to calculate the evolutionary distances that are expressed in base
 381 substitutions per site, only values above 50 are reported. The out group employed
 382 is the *Acholeplasma laidlawii*. The numbers of the GenBank of the sequences
 383 employed are on the right for the '*Ca. Phytoplasma*' species and on the left for the
 384 strains in the 16SrIII group.

385



386

387 **Figure 4** Restriction fragment length polymorphism patterns of ribosomal protein
388 amplicons from faba bean yellows and witches' broom phytoplasma from Peru
389 (Fb1, Fb2, and Fb4) and other phytoplasma strains classified in the ribosomal
390 group 16SrIII after polyacrylamide 6.7% gel electrophoresis. The amplification was
391 carried out in nested PCR with rplIIF1/rplIIR1 primers; the enzyme used is listed at
392 the bottom. Lanes 1, X disease (16SrIII-A, CX), 2, strain from *Euscelidius*
393 *variegatus* (16SrIII-B, API), 3, *Vaccinium* witches' broom (16SrIII-F, VAC), 4, spirea
394 stunt (16SrIII-E, SPI), 5, poinsettia branching (16SrIII-H, JRI). P, marker Φ X174
395 digested with *Hae*III and having the fragment sizes from top to bottom of 1,353;
396 1,078; 872; 603; 310; 281; 271; 234; 194; 118 and 72 base pairs.