



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE  
DELLA RICERCA

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Evaluation of plant derivatives of Meliaceae family as a source of nitrogen for trees

This is the submitted version (pre peer-review, preprint) of the following publication:

*Published Version:*

Marcolini G., Giocchini P., Quartieri M., Baldi E., Sorrenti G., Toselli M. (2022). Evaluation of plant derivatives of Meliaceae family as a source of nitrogen for trees. JOURNAL OF PLANT NUTRITION, 45(11), 1642-1653 [10.1080/01904167.2021.2014868].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/849772> since: 2022-01-31

*Published:*

DOI: <http://doi.org/10.1080/01904167.2021.2014868>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is an original manuscript of an article published by Taylor & Francis in *Journal of plant nutrition* on 08 December 2021, available at:  
<https://doi.org/10.1080/01904167.2021.2014868>.

**Evaluation of plant derivatives of Meliaceae family as a source of nitrogen for trees**

|                               |  |
|-------------------------------|--|
| Journal:                      | <i>Journal of Plant Nutrition</i>  |
| Manuscript ID                 | LPLA-2019-0331.R1  |
| Manuscript Type:              | Research Article   |
| Date Submitted by the Author: | n/a  |
| Complete List of Authors:     | Marcolini, Graziella; University of Bologna<br>Gioacchini, Paola; University of Bologna<br>quartieri, maurizio; University of Bologna<br>Baldi, Elena; University of Bologna,<br>Sorrenti, Giovambattista; University of Bologna<br>Toselli, Moreno; University of Bologna, Department of Agricultural and Food Sciences |
| Keywords:                     | Azadiracta indica, C:N ratio, microbial N, soil respiration, 15N   |
|                               |  |

1  
2  
3  
4 Evaluation of plant derivatives of Meliaceae family as a source of nitrogen for  
5  
6 trees  
7  
8

9 Graziella Marcolini, [graziella.marcolini2@unibo.it](mailto:graziella.marcolini2@unibo.it)  
10  
11

12  
13 Paola Gioacchini, [paola.gioacchini@unibo.it](mailto:paola.gioacchini@unibo.it)  
14  
15

16 Maurizio Quartieri, [maurizio.quartieri@unibo.it](mailto:maurizio.quartieri@unibo.it)  
17  
18

19  
20 Elena Baldi, [elena.baldi7@unibo.it](mailto:elena.baldi7@unibo.it), <https://www.unibo.it/sitoweb/elena.baldi7>  
21  
22

23 Giovambattista Sorrenti, [g.sorrenti@unibo.it](mailto:g.sorrenti@unibo.it), <https://www.unibo.it/sitoweb/g.sorrenti>  
24  
25

26  
27 Moreno Toselli\*, corresponding author, [moreno.toselli@unibo.it](mailto:moreno.toselli@unibo.it)  
28  
29 <https://www.unibo.it/sitoweb/moreno.toselli>, ORCID: 0000-0001-5632-6331  
30  
31

32 Department of Agricultural and Food Sciences, University of Bologna, viale Fanin, 46,  
33  
34 40127, Bologna, Italy  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Evaluation of plant derivatives of Meliaceae family as a source of nitrogen for trees

Marcolini G., Gioacchini P., Quartieri M., Baldi E., Sorrenti G., Toselli M. \*

Department of Agricultural and Food Sciences, viale Fanin, 46, 40127, Bologna, Italy

### Abstract

Soil application of fresh organic matter is a way to increase soil organic matter and provide nutrients to trees. The effect of application of organic matter depends on the interaction among soil, root and microbial biomass. The aim of this research was to evaluate the potential release of N for hybrid GF677 (*P. persica* x *P. dulcis*) uptake, of 6 neemcakes available on the Italian market compared with fresh leaves of *Melia azedarach*, an ornamental tree that grows in the area of investigation. The release of N, and consequently root uptake was related to C:N ratio, the lower the ratio the higher the N concentration in plant tissues and plant growth. Using the  $^{15}\text{N}$  isotope technique, we found that up to 30% of the N applied with fresh *Melia* leaves, was accumulated in the tree, however the mineral N concentration in soil and plant and plant growth was not affected by the application of plant derivatives.

**Key words:** *Azadiracta indica*, C:N ratio,  $^{15}\text{N}$ , soil respiration, microbial N

### Introduction

Soil application of organic fertilizers is a way to increase soil organic matter (OM) and provide nutrients to trees. Organic matter represents an important source of C and energy for soil microbial biomass, hence sustaining a microbial population (Sparling, 1992) with consequent considerable amounts of N immobilized. For this reason, plant roots compete with microorganisms in soil for the same available N, and may reduce N immobilization, thus limiting the supply of mineral N to the microflora (Wang and Bakken, 1989 and 1997). Among the large availability of plant derivatives, neemcake represents a byproduct of neem oil production. Neemcake maintains a valuable fertilization potential which has been recently investigated (Marcolini et al., 2016), that is the result of its chemical and biochemical composition. In particular, the interaction between neemcake and soil promotes a release of

1  
2  
3  
4 mineral N that may be up taken by roots and used by trees or metabolized by soil microbial  
5 biomass (Marcolini et al., 2016).

6  
7 The potential effect of OM as a source of nutrients for plants depends on soil  
8 characteristics, root uptake efficiency, microbial community, environment, etc. The use of  
9 isotopic methods can provide useful agronomic information on the amount of N mineralized  
10 from the residue, on the root uptake rate and on the fraction used for plant metabolism (Hood  
11 et al., 2000). Organic fertilizer derived from plant residues uniformly labelled with  $^{15}\text{N}$  (Ladd  
12 et al., 1981) can be employed for this purpose. In Northern latitudes, melia (*Melia azedarach*)  
13 is often used as an ornamental tree for its resistance to pest and diseases. The release of  
14 nutrients from residues of melia, showed a behavior similar to neemcake (Marcolini et al.,  
15 2016; Toselli et al., 2010), consequently, considering its adaptability to grow in temperate  
16 climate, we used plants of melia to obtain uniformly labelled fresh materials (named leaves  
17 enriched with  $^{15}\text{N}$  isotope).

18  
19 The aims of the present study were to 1) evaluate the effectiveness of plant derivatives  
20 obtained from neemcake and fresh leaves of melia tree as fertilizer-N for peach and 2)  
21 estimate the fraction of  $^{15}\text{N}$  released by the mineralization of fresh melia leaves and absorbed  
22 by peach tree. The two aims were pursued in two separate experiments, experiment A:  
23 Comparison of derivatives as tree fertilizers and experiment B: Estimation of the uptake rate  
24 of labelled N.

## 25 26 27 28 29 30 31 32 33 34 35 36 37 38 **Material and Methods**

### 39 40 41 42 Experiment A: Comparison of derivatives as tree fertilizers

43 The following plant material, commercialized in Italy and described in Marcolini et al.,  
44 (2016) were used: 1) Green neem 1, from India (green neem 1), 2) Neem pelleted, origin  
45 unknown (pelleted neem), 3) Neem, origin unknown (neem cake 1), 4) Deoiled neemcake,  
46 from India (deoiled neem), 5) Oiled neem cake, from India (oiled neem), 6) Green neem 2,  
47 from India (green neem 2), 7) *Melia azedarach* L. leaves harvested in June 2010 from  
48 seedlings grown on pots filled with sand and frozen at  $-20^{\circ}\text{C}$  prior to use (melia leaves).

49 The chemical composition of plant derivatives is listed in Table 1. The derivatives  
50 were oven-dried at  $65^{\circ}\text{C}$  and ball milled. Total C and N concentration were determined with  
51 a CHN elemental analyzer (Thermo Fisher, mod. EA 1110, Bremen, Germany). Solid samples  
52 were weighed in tin cups and dropped in a tube where in the presence of external oxygen  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 flash combustion occurred at a temperature of 1800 °C. The gaseous combustion  
4 products, N<sub>2</sub>, NO<sub>x</sub>, CO<sub>2</sub>, were carried by the helium through a column filled with copper  
5 oxide and from there to a Cu-column where NO<sub>x</sub> was reduced to elementary N. With a  
6 programmed temperature raise in the column the gases were released separately. They flew  
7 along a thermal conductivity detector which produced an electrical signal proportional to the  
8 concentration of N and C. Calcium (Ca), potassium (K), magnesium (Mg), iron (Fe),  
9 manganese (Mn), zinc (Zn) and copper (Cu) were determined by atomic absorption  
10 spectrometry (SpectrAA-200, Varian, Mulgrave, Australia) after wet mineralization  
11 according to US EPA Methods 3052 (Kingston, 1988) by treating 0.5 g of dry weight  
12 (DW) material with 8 mL of nitric acid (65%) and 2 mL of hydrogen peroxide (30%) at  
13 180 °C in an Ethos TC microwave lab station (Milestone, Bergamo, Italy). Phosphorus  
14 (P) concentration was spectrophotometrically quantified at 700 nm (Saunders and  
15 Williams, 1955) on the same mineralized samples used for metal determinations. Liquid  
16 samples were neutralized with 5 M NaOH and enriched with 30 mL of a mixture of 0.1 M  
17 ascorbic acid, 32 mM ammonium molybdate, 2.5 M sulphuric acid and 3 μM potassium  
18 antimonyl tartrate to develop a phospho-molybdic blue color.

### 31 32 *Experimental design*

33  
34 The experiment was carried out in 2010 at the experimental station of the University of  
35 Bologna, in Cadriano (44°35'N, 11°27'E). Plant derivatives were incorporated into a clay  
36 loam Bathicalci Eutric Cambisols soil (FAO, 1990) (Table 2), at the application rate of 8  
37 g fresh weight (FW) kg<sup>-1</sup> soil for neemcakes and 16 g FW kg<sup>-1</sup> for melia leaves.  
38 Micro-propagated plants of GF677 (*P. persica* x *P. dulcis*) rootstocks were placed into 1.5  
39 L pots filled with 1 kg of soil previously mixed with plant derivatives and  
40 compared to an unamended control. On day 68 and 113 after planting, 5 pots per treatment  
41 were destructively sampled. Plants were harvested and divided into roots, stem, shoots  
42 (leaves and shoot axis). The length of the new shoots was recorded and leaf green color,  
43 as an estimation of leaf chlorophyll, was determined by a portable SPAD 502 (Minolta  
44 Co., Ramsey, New Jersey, USA). Leaves were washed with a solution of Tween 20, then,  
45 as all the other organs, they were rinsed with tap water and distilled water, oven-dried,  
46 weighed and milled. Roots and shoots macro- and micronutrient concentrations were  
47 determined as previously described for the derivative characterization, in particular C and  
48 N by CHN elemental analyzer (Thermo Fisher, mod. EA 1110, Bremen, Germany), metals  
49 by atomic absorption spectrophotometry

(SpectrAA-200, Varian, Mulgrave, Australia) and P by spectrophotometry (Saunders and Williams, 1955).

For each sampling date, data were statistically analyzed as in a complete randomized design with 5 replicates (pot) and, when analysis of variance showed statistical significance at  $P \leq 0.05$ , means were separated by Student Newman Keuls (SNK) test.

#### Experiment B: Estimation of the uptake rate of labelled N

Uniformly  $^{15}\text{N}$  labelled melia leaves were obtained by feeding seedlings of *Melia azedarach* grown in sand with a nutrient solution containing  $^{15}\text{N}$ -labelled  $^{15}\text{NH}_4^{15}\text{NO}_3$  (10 atom%  $^{15}\text{N}$ ). Leaves were harvested in August 2010 and stored at  $-20^\circ\text{C}$  prior to use. A subsample was oven-dried, ball-milled and analyzed for N and  $^{15}\text{N}$  concentration with an elemental analyzer coupled with an isotope-ratio mass spectrometer (CF-IRMS, Delta Plus Thermo Fisher, Bermen, Germany). Isotope-ratio of N ( $^{15}\text{N}/^{14}\text{N}$ ) was measured after conversion of the molecule to a gas-phase ion, the resultant flux of electrically charged ions was converted into a proportional electrical current. This allowed to measure total N concentration that was 1.87% and  $^{15}\text{N}$  enrichment that was 1.876 atom%.

#### *Experimental design*

The experiment started in April 2011 at the experimental station of the University of Bologna, in Cadriano. The same clay loam soil (FAO, 1990) as described in table 2 was collected from the field, mixed with sand at a soil:sand ratio of 3:1 and sieved to 4 mm. Labelled melia leaves were ground and sieved to 2 mm and incorporated into the soil at the same application rate as for experiment A (16 g FW  $\text{kg}^{-1}$  soil). One-year-old plants of GF677 rootstocks were placed into pots filled with 2 kg FW amended soil. Another set of plants was potted with unamended soil (control treatment). Immediately after potting, plants were pruned and trained to one shoot, placed in the greenhouse and watered daily. After one month, plants were transferred outdoor, on a bench, under a plastic shelter to protect them from rain. At day 30, 90, 131 and 173 after planting, 5 control and 4 treated plants were destructively harvested and divided into roots, stem, new growth (leaves and shoot axis). The length of the new shoots was recorded and leaf green color, as an estimation of leaf chlorophyll, was determined by a portable SPAD 502 (Minolta Co., Ramsey, New Jersey, USA). All organs were carefully washed with tap water and distilled water, oven-dried, weighed, milled and analyzed for total N and  $^{15}\text{N}$  content with the elemental analyzer coupled with an isotope ratio mass

spectrometer as described previously. The percentage of N derived from melia leaves (% NDFML) was calculated as (Hauck and Bremner, 1976):

$$\% \text{ NDFML} = (\text{atom } ^{15}\text{N excess of plant fertilized with labelled melia leaves} / \text{atom } ^{15}\text{N excess of labelled melia leaves}) \times 100;$$

where atom %  $^{15}\text{N}$  excess was obtained by subtracting from values measured in the treated pools, the respective natural abundance measured in control pool (0.3663 atm %).

Then, the amount of N derived from labelled melia leaves (NDFML) in plant organ were calculated as:

$$\text{NDFML (mg)} = \text{total N (mg)} \times (\% \text{ NDFML} / 100)$$

Finally, the amount of N recovered from the labelled melia leaves was calculated as (Hood et al., 2000):

$$\% \text{ N recovery from melia leaves} = \text{NDFML (mg)} / (\text{N added as melia leaves (mg)}) * 100$$

The soil of each pot was collected, uniformly mixed and analyzed for total N with a CHN elemental analyzer. Microbial biomass C and N were determined by the fumigation-extraction method (Vance et al, 1987), from 5 g moist soil, extracted with 20 mL of 0.5 M  $\text{K}_2\text{SO}_4$  and filtered through filter paper S&S 595 (Ahlstrom, Helsinki, Finland). Another sample of 5 g of moist soil was firstly fumigated with chloroform for 24 h and then extracted in the same way. The extracts were frozen at  $-20^\circ\text{C}$  until analysis for total organic C concentrations using a total organic C measuring unit (TOC-Vcpn TNM-1, Shimadzu, Kyoto, Japan). The amount of microbial C and N were determined by the difference between  $\text{K}_2\text{SO}_4$  extractable C and N in fumigated and non-fumigated soils. No correction factor, that is usually applied to take into account the incomplete recovery of microbial constituents extracted from soil after fumigation was used, since the values of microbial C and N are used here to reveal differences between derivatives.

For each sampling day, data of plant DW, N concentration and NDFML were statistically analyzed as in a complete randomized design, when analysis of variance showed

1  
2  
3  
4 statistical significance at  $P \leq 0.05$ , means were separated by SNK test. Data of soil mineral N  
5 and microbial biomass C and N were statistically analyzed as in a factorial design with 2  
6 factors: treatment (2 levels: control and  $^{15}\text{N}$  melia) and time (4 levels: day 30, 90, 131, 173).  
7  
8 When analysis of variance showed statistical significance at  $P \leq 0.05$ , means were separated by  
9 Student Newman Keuls test. When interaction between factors was significant, 2 standard  
10 error of means (SEM) was used as minimum difference between statistically different values.  
11  
12  
13  
14

## 15 16 **Results**

### 17 18 19 Experiment A: Comparison of derivatives as tree fertilizers

20  
21 Table 1 shows the mineral compositions of the derivatives used in this experiment.  
22  
23 With the exception of N, neem cake 1 presented the highest concentration of all the nutrients  
24 analyzed. Phosphorus ranged between  $11.3 \text{ g kg}^{-1}$  in neem cake 1 and  $2.0 \text{ g kg}^{-1}$  in melia  
25 leaves. Potassium varied between  $24.0 \text{ g kg}^{-1}$  in neem cake 1 and  $8.1 \text{ g kg}^{-1}$  in melia leaves;  
26 among the other derivatives, **green neem 1**, **deolid neem** and **oiled neem** presented higher K  
27 than **green neem 2** and **pelleted neem**. Calcium and Mg were higher in melia leaves compared  
28 to **green neem 1**, **pelleted neem**, **deoiled neem**, **oiled neem**, and **green neem 2**. Iron  
29 concentration widely varied between  $8,447 \text{ mg kg}^{-1}$  in neem cake 1 and  $160 \text{ mg kg}^{-1}$  in melia  
30 leaves. Among the others, **deoiled neem** presented the highest Fe concentration, followed by  
31 **oiled neem**, **green neem 1**, **pelleted neem**, and **green neem 2**. Manganese concentrations  
32 ranged between  $125 \text{ mg kg}^{-1}$  in neem cake 1 and  $43 \text{ mg kg}^{-1}$  in **green neem 2** and melia leaves.  
33  
34 Cu concentration was similar in all derivatives, except in neem cake 1, which showed a  
35 concentration more than twice that observed in the others derivatives. Zinc concentration  
36 ranged between  $16 \text{ mg kg}^{-1}$  in melia leaves and  $70 \text{ mg kg}^{-1}$  in neem cake 1.  
37  
38  
39  
40  
41  
42  
43  
44

45 At the end of the experiment, with the exception of **green neem 1**, all derivatives,  
46 increased leaf chlorophyll content, particularly **oiled neem**, **green neem 2** and melia leaves  
47 (Table 3). Shoot length was increased by soil-addition of **green neem 1**, **pelleted neem**, **neem**  
48 **cake 1**, **deoiled neem**, **green neem 2**. **Pelleted neem**, **neem cake 1** and **green neem 2** increased  
49 all organs and, consequently, total plant DW compared to control (Table 3).  
50  
51  
52  
53

54 At the end of the experiment (day 113), no effect of treatment on organ N  
55 concentration was detected (Table 4). **Deoiled neem** treated-plants showed the highest leaf P  
56 concentration, followed by **oiled neem** and **green neem 1**, while the lowest P concentration  
57 was recorded in **green neem 2** treatment (Table 4). No differences between treatments were  
58  
59  
60

found for K concentration that was on average 18.6 mg kg<sup>-1</sup>. Moreover, leaves of control plants presented the highest Ca concentration, significantly different from that of neem cake 1- and green neem 2-treated-plants. Deoiled neem promoted a higher leaf Mg concentration compared to neem cake 1 and green neem 2. At the end of the experiment, leaf microelement concentrations were not affected by treatments, with the exception of Mn, that was higher in leaves of control, deoiled neem-, oiled neem-, and green neem 2-treated-plants compared to those fertilized with green neem 1 and neem cake 1 (Table 4).

Considering the amount of N in plants as a mean to measure the N removal, 68 days after treatment application, plant fertilized with neem cake 1 and deoiled neem removed a higher amount of N than control (Table 5). At the end of the experiment, in addition to plant treated with neem cake 1 and deoiled neem, those fertilized with green neem 1 and green neem 2 removed a higher amount of N than control (Table 5).

#### Experiment B: Estimation of the uptake rate of labelled N

Although soil application of <sup>15</sup>N-enriched melia leaves increased soil N concentration (1,555 mg kg<sup>-1</sup>) compared to untreated control (1,266 mg kg<sup>-1</sup>), soil nitrate- and ammonium-N were not affected by the application of melia leaves. In detail NH<sub>4</sub><sup>+</sup>-N ranged between 2.35 and 2.37 mg kg<sup>-1</sup> while NO<sub>3</sub><sup>-</sup>-N ranged between 5.1 and 5.6 mg kg<sup>-1</sup> for treated and untreated soil, respectively. Nitrogen concentration of plant organs was not affected by treatment (data not tabulated) at any of the sampling times.

Plant organ DW resulted similar in control and <sup>15</sup>N-melia treated plants and increased with time during the experiment. Total plant DW in untreated plant ranged between 1.47 g (at day 30) and 6.07 g (at day 173) and between 1.12 g (at day 30) and 4.58 g (at day 173) in <sup>15</sup>N-melia treatment (data not tabulated).

At day 30, only a little of NDFML were detected in GF677 organs (Figure 1), with a higher amount in roots compared to stem and shoots (Figure 1, inset). Nitrogen derived from melia leaves increased with time and on day 90, shoot showed a higher amount of NDFML than root and stem. On day 131, NDFML in shoot and root was higher than in stem; on day 173, no organ effect was found (Figure 1). At day 131, 9.02 mg of NDFML were found in shoot (Figure 1), corresponding to a NDFML recovery of 4.55% of total N applied with plant derivatives (Table 6). At that day, stem and roots presented a NDFML of 24% and 29.6%, respectively (Table 6). At day 173, NDFML decreased in the different organs investigated, in term of amount (Figure 1) and percentage (Table 6).

1  
2  
3  
4 Soil application of labelled melia leaves immediately increased microbial biomass C  
5 (Figure 2), the highest value was found at day 1, and then it decreased with time to reach  
6 values similar to the control soil at the end of the experiment. In addition, microbial N  
7 increased after the amendment application, in this case, the highest value was found at day  
8 131, and then it decreased (Figure 3). The microbial C/N ratio decreased from day 1 to day  
9 173 in melia leaf treated plants, compared with unamended control that presented a steady  
10 C/N ratio ranging from 4 to 6 throughout the investigation period of time.  
11  
12  
13  
14  
15  
16  
17

## 18 Discussion

19 The *Meliaceae* derivatives showed a different mineral composition in response to their  
20 different origin, species, organ, production processes, year of harvest, etc. (Marcolini et al.,  
21 2016). The mineral composition of neem cake 1 deserves some comments. In fact, neem  
22 cake1 presented, for most of the nutrients (i.e. P, K, Ca, Mg, Fe, Mn, Cu and Zn),  
23 concentrations that were at least twice higher than the other neemcakes here used and reported  
24 in earlier study (Toselli et al., 2010). Patra et al. (2011) reported a N, P, and K neemcake  
25 concentration of 49.3 g kg<sup>-1</sup>, 10.3 g kg<sup>-1</sup> and 13.2 g kg<sup>-1</sup>, respectively, that were similar for P,  
26 higher for N and lower for K compared to the neem cake 1 used in this experiment.  
27  
28  
29  
30  
31  
32

33 The mineral composition of melia leaves resulted different from that reported in other  
34 studies. For example, melia leaves tested by Toselli et al. (2010) presented a concentration of  
35 K, Ca, Cu and Zn two times higher compared to that found here and a Mg concentration  
36 almost 20 times lower. In addition, the chemical composition of melia litter reported by Singh  
37 and Sharma (2007) differed from our material, principally for Fe and Zn concentration, which  
38 resulted about 9 and 3 times higher compared to our fresh leaves, respectively.  
39  
40  
41  
42

43 This different chemical composition was probably the main reason of the different  
44 effect of the incorporation of *Meliaceae* derivatives on plant growth that was stimulated by  
45 green neem 2, pelleted neem and neem cake 1. These cakes had the highest N concentration  
46 and the lowest C:N ratio, that made the N added (37, 44, and 31 mg kg<sup>-1</sup> N for neem cake 1,  
47 green neem 2, and pelleted neem, respectively) promptly available for root uptake (Marcolini  
48 et al., 2016). However, the higher availability of N did not promote a significant increase of  
49 leaf N, this because of the high growth rate induced by the neemcakes that diluted the N  
50 absorbed by root. In fact, the amount of N removed by the plants treated with pelleted neem,  
51 neem cake 1 and green neem 2 was double the amount removed by control plants; at the same  
52 time leaf N concentration was not different.  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The effect of the derivatives on leaf P concentration is not easy to explain, since plant treated with deoiled neem, oiled neem and green neem 1 showed the highest leaf P concentration, although the highest P concentration was found in neem cake 1 and green neem 2 compounds. Probably, also in this case, there was a dilution effect induced by the application of pelleted neem, neem cake 1 and green neem 2 that stimulated plant growth and prevented the increase of concentration of P in the tissues.

At the end of the experiment, green neem 2- and neem cake 1-treated-plants presented lower leaf Ca concentration compared to the control ones. This can be only partly explained by the high concentration of K and Mg in neem cake 1 and green neem 2, respectively. In fact, the possible antagonistic effect of K and Mg on Ca uptake (Marschner, 1995) was not observed in deoiled neem- and green neem 1-treated-plants, despite of their high concentration of K. A similar behavior was found on one-year-old peaches fertilized with organic amendments (Baldi et al., 2010), where leaf K was increased, while Ca and Mg concentration were depressed by soil application of compost.

Although the high concentration of Fe in all the neemcakes, leaf Fe concentration was not increased by treatments, this means that Fe was not available for root uptake. However, the root system of GF677 hybrid is considered one of the most efficient in absorbing Fe (Tagliavini and Rombolà, 2001), and this ability may have reduced the effect of Fe soil supply.

Soil-applied melia leaves was less effective than neemcakes in increasing nutrient status in peach tree. In fact, none of the nutrient leaf concentration was increased and plant growth was not stimulated by fresh melia leaves compared to the untreated control. Only leaf chlorophyll obtained benefit from soil incorporation of leaf of melia, this can be the first appearance of a positive effect that may involve also the nutritional status later. This response was probably related to the high C:N ratio of melia leaves (17.5%) that limited the mineralization of N and delayed the fertilization effect (Marcolini et al., 2016). Previous research on wheat reported a grain yield and a nutrient content (N, P, K) increase after the application of leaf litter of poplar (*Populus deltoides*), eucalypt (*Eucalyptus* hybrid) and dek (*Melia azedarach*), with or without the inoculation of cellulolytic fungus culture of *Aspergillus awamori* to accelerate the decomposition rate of litters (Singh and Sharma, 2007). In addition, yield of sorghum increased significantly with increasing levels of leaf litter application. Nevertheless, in their experiment, Singh and Sharma (2007), applied N, P and K

1  
2  
3  
4 at a rate of 50, 11, 10 mg kg<sup>-1</sup> soil, respectively that could have enhanced the positive effect of  
5 plant litter compared to our study.  
6

7 The N mineralization behavior of fresh melia leaves was studied in detail in  
8 experiment B where, 131 day after application, the amount of N from mineralization of fresh  
9 organic material tilled into the soil accounted for up to 39% of total N present in plant. In  
10 particular, a little amount of <sup>15</sup>N released by melia leaves was available one month after the  
11 amendment and was found mainly in shoots. Thereafter, the amounts of NDFML increased  
12 with time in all organs, with shoot and leaves representing the strongest sink organs for melia  
13 leaf-derived-N. Also, at the end of the experiment (September) shoot and leaves presented a  
14 higher amount of melia-derived N, compared to stem, but it was lower than the previous  
15 sampling days, indicating a possible root uptake of soil-native-N that diluted the <sup>15</sup>N fraction  
16 within the plant. The availability and uptake of <sup>15</sup>N did not increase plant N concentration, so  
17 that treated and untreated trees showed the same N concentration throughout the season. In  
18 addition, the mineral N concentration of both control and treated soil was similar during all  
19 the experiment. This means that the sum of fractions of labelled N from melia leaves and  
20 unlabeled N from soil-native-N was not modified, if one increased, the other decreased and  
21 vice versa.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

33 Microbial biomass was significantly increased by application of melia leaves, as  
34 testified by the increase of both microbial C and N. In particular, microbial N ranged between  
35 5 and 10 mg kg<sup>-1</sup> and between 10 and 20 mg kg<sup>-1</sup> in control and amended soil, respectively.  
36 This difference can be explained considering an increase of N mineralization in the amended  
37 compared to the unamended soil, followed by an immediate microbial immobilization, and a  
38 consequent competition for this N between microbial biomass and plants. However, the exact  
39 origin of this extra amount of N mineralized cannot be defined; it can derive from both soil-  
40 native-N or melia leaf.  
41  
42  
43  
44  
45  
46

47 At the same time a fast increase of microbial biomass C:N ratio (12) was observed at  
48 day 1, followed by a decrease to 3 (at day 173), as a consequence of incorporation of melia  
49 leaves into the soil. Considering that in control soil the C:N ratio was steady and ranged from  
50 4 to 6, this response may indicate a variation in C and N availability throughout the  
51 experiment, with C available at the beginning, and N later. The different C and N flux in the  
52 soil was related to melia leaf C:N ratio of 17.5 that pushed microbial biomass to immobilize  
53 N. This explanation is indirectly supported by Li et al. (2016) who found a significant,  
54 negative correlation of annual rice yield with soil microbial biomass C:N. In fact, in our  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 experiment the high microbial biomass C:N induced a N immobilization, which reduced the  
5 flux of mineral N for peach root. In fact, although soil total N increased after melia leaf  
6 application, mineral N remained stable, meaning that the release of N by melia leaf  
7 mineralization, and detected as  $^{15}\text{N}$  was equalized by soil-native-N microbial immobilization.  
8  
9

10 In conclusion the effectiveness of neemcakes as N fertilizer is variable, and in part  
11 predictable from the C:N ratio, in fact the cakes with highest fertilization aptitude showed a  
12 C:N ratio lower than 17. In this short-term (6 months) experiment, fresh melia leaves showed  
13 a low effectiveness in term of N fertilization, because the rate of N released was compensated  
14 by the same rate of N immobilization by microbial biomass. However, plants showed a  
15 beneficial effect in term of increase of leaf chlorophyll not justified by the increase of leaf N.  
16 Other investigations can possibly explain the physiological reason of this positive response.  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References

Anderson J.P.E., Domsch K.H., 1978. A physiological method for the quantitative measurement of microbial biomass in soils. *Soil Biology and Biochemistry*, 10: 215-221.

Baldi E., Toselli M, Marangoni B., 2010. Nutrient partitioning in potted peaches (*Prunus persica* L.) trees supplied with mineral and organic fertilizers. *Journal of Plant Nutrition*, 33 (14): 2050-2061.

Dijkstra F.A., Bader N.E., Johnson D.W., Cheng W., 2009. Does accelerated soil organic matter decomposition in the presence of plants increase plant N availability? *Soil Biology and Biochemistry*, 41: 1080-1087.

Hauck R.D., Bremner J.M., 1976. Use of tracers for soil and fertiliser nitrogen research. *Advances in Agronomy*, 28: 219-266.

Hood R., Merckx R., Jensen E.S., Powlson D., Matijevec M., Hardason G., 2000. Estimating crop N uptake from organic residues using a new approach to the <sup>15</sup>N isotope dilution technique. *Plant and Soil*, 223: 33-44.

Li Y., Wu J., Shen J., Liu S., Wang C., Chen D., Huang T., Zhang J., 2016. Soil microbial C:N ratio is a robust indicator of soil productivity for paddy fields. *Scientific Reports*, 6:35266, DOI: 10.1038/srep35266.

Ladd J.N., Oades J.M., Amato M., 1981. Microbial biomass from <sup>14</sup>C and <sup>15</sup>N-labelled plant material decomposing in soils in the field. *Soil Biology and Biochemistry*, 13: 119-126.

Marcolini G., Toselli M., Quartieri M., Gioacchini P., Baldi E., Sorrenti G., Mariani S., 2016. Nitrogen and carbon mineralisation of different Meliaceae derivatives. *Plant Soil Environ.*, 62(3): 122-127.

Marschner H., 1995. Mineral nutrition of higher plants. Academic press, London.

1  
2  
3  
4  
5 Patra P.S., Sinha A.C., Mahesh S.S., 2011. Yield, nutrient uptake and quality of groundnut  
6 (Arachis hypogaea) kernels as affected by organic sources of nutrient Indian Journal of  
7 Agronomy 56 (3): 237-241  
8  
9

10  
11  
12 Saunders W.M., Williams E.G., 1955. Observations of the determinations of organic  
13 phosphorous in soils. Journal of Soil Science, 6: 254-267.  
14  
15

16  
17 Singh B., Sharma K.N., 2007. Nutrition and growth of wheat-sorghum rotation in soils  
18 amended with leaf litter of trees before planting of wheat. Agroforest Syst, 71: 25-34.  
19  
20

21  
22 Sparling, G.P., 1992. Ratio of microbial biomass carbon to soil organic carbon as a sensitive  
23 indicator of changes in soil organic matter. Soil Res. 30 (2):195–207.  
24  
25

26  
27 Tagliavini, M., Rombolà, A.D., 2001. Iron deficiency and chlorosis in orchard and  
28 vineyard ecosystems: a review. Eur. J. Agron. 15, 71–92.  
29  
30

31  
32 Toselli M., Baldi E., Sorrenti G., Quartieri M., Marangoni B., 2010. Evaluation of the  
33 effectiveness of soil-applied plant derivatives of Meliaceae species on nitrogen availability to  
34 peach trees. Scientia Horticulturae 124: 183-188.  
35  
36

37  
38 Vance E.D., Brookes P.C., Jenkinson D.S., 1987. An extraction method for measuring soil  
39 microbial biomass C. Soil Biology and Biochemistry, 19: 703-707.  
40  
41

42  
43 Wang J.G. and Bakken L.R., 1997. Competition for nitrogen during mineralization of plant  
44 residue in soil: microbial response to C and N availability. Soil Biology and Biochemistry, 29  
45 (2): 163-170.  
46  
47

48  
49 Wang J.G. and Bakken L.R., 1989. Nitrogen mineralization in rhizosphere and non-  
50 rhizosphere soil, effect of the spatial distribution of N-rich and N-poor plant residues. In:  
51 Nitrogen in Organic Wastes Applied to Soils (J. A. Hansen and K. Henriksen, Eds): pp 81-97.  
52 Academic, London.  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Table 1. Dry weight (DW) and mineral composition of plant derivatives used in experiment A. In brackets standard deviation (n=2).

| <b>Derivative</b>    | <b>DW</b><br>(g kg <sup>-1</sup> ) | <b>C</b><br>(g kg <sup>-1</sup> ) | <b>N</b><br>(g kg <sup>-1</sup> ) | <b>P</b><br>(g kg <sup>-1</sup> ) | <b>K</b><br>(g kg <sup>-1</sup> ) | <b>Ca</b><br>(g kg <sup>-1</sup> ) | <b>Mg</b><br>(%) | <b>Fe</b><br>(mg kg <sup>-1</sup> ) | <b>Mn</b><br>(mg kg <sup>-1</sup> ) | <b>Cu</b><br>(mg kg <sup>-1</sup> ) | <b>Zn</b><br>(mg kg <sup>-1</sup> ) |
|----------------------|------------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|------------------------------------|------------------|-------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|
| <b>Melia leaves</b>  | 290                                | 420                               | 24                                | 1.9 (±0.02)                       | 8.1 (±0.05)                       | 13.6 (±0.01)                       | 3.8 (±0.04)      | 160 (±7.92)                         | 43 (±0.56)                          | 8 (±0.71)                           | 16 (±0.35)                          |
| <b>Green neem</b>    | 890                                | 470                               | 22                                | 3.5 (±0.03)                       | 15.3 (±0.02)                      | 6.2 (±0.007)                       | 1.9 (±0.02)      | 1324 (±241)                         | 56 (±0.35)                          | 11 (±2.05)                          | 22 (±1.2)                           |
| <b>Pelleted neem</b> | 940                                | 500                               | 31                                | 3.0 (±0.08)                       | 9.7 (±0.03)                       | 3.0 (±0.007)                       | 1.7 (±0.007)     | 1096 (±0.21)                        | 26 (±1.70)                          | 7 (±0.92)                           | 30 (-)                              |
| <b>Neem cake 1</b>   | 920                                | 350                               | 37                                | 11.3 (±0.20)                      | 24.0 (±0.09)                      | 14.6 (±0.01)                       | 6.7 (±0.007)     | 8447 (±1.06)                        | 125 (±0.35)                         | 50 (±11.88)                         | 70 (±0.84)                          |
| <b>Deoiled neem</b>  | 960                                | 410                               | 17                                | 2.1 (±0.02)                       | 15.2 (±0.05)                      | 4.9 (±0.02)                        | 2.0 (±0.01)      | 3035 (±158)                         | 72 (±2.26)                          | 19 (±3.53)                          | 23 (±3.11)                          |
| <b>Oiled neem</b>    | 910                                | 440                               | 16                                | 2.0 (±0.01)                       | 16.3 (±0.03)                      | 6.1 (±0.06)                        | 2.1 (±0.01)      | 2414 (±54.7)                        | 61 (±0.28)                          | 10 (±1.70)                          | 19 (±0.56)                          |
| <b>Green neem 2</b>  | 940                                | 490                               | 44                                | 6.5 (±0.02)                       | 12.5 (±0.04)                      | 5.0 (±0.007)                       | 3.5 (±0.007)     | 620 (±59.2)                         | 43 (±1.27)                          | 10 (-)                              | 54 (±2.12)                          |

Table 2. Selected characteristics of the soil used in the 2 studies

| Characteristic                                      | Value | Analytical method                                       |
|---|-------|---|
| Sand (g kg <sup>-1</sup> )                          | 250   |   |
| Silt (g kg <sup>-1</sup> )                          | 500   |   |
| Clay (g kg <sup>-1</sup> )                          | 250   |   |
| pH  | 7.5   | Water   |
| Calcium Carbonate (g kg <sup>-1</sup> )             | 30    | HCl/De Astis  |
| Organic Matter (g kg <sup>-1</sup> )                | 14    | K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub> Oxidation |
| Total N (g kg <sup>-1</sup> )                       | 11    | Kjeldahl  |
| Exchangeable K (mg kg <sup>-1</sup> )               | 219   | Barium chloride   |
| Exchangeable P (mg kg <sup>-1</sup> )               | 12    | Olsen   |
| Cation Exchange Capacity (meq 100 g <sup>-1</sup> ) | 16.1  | Barium chloride   |
| Electric Conductivity μScm <sup>-1</sup> )          | 220   |   |

Table 3. Effect of soil-applied *Meliaceae* derivatives on leaf chlorophyll, shoot length and plant organ dry weight (DW) after 113 days from treatment application (experiment A)

| Treatment           | Leaf chlorophyll | Shoot length | Leaves                      | Shoot  | Roots  | Stem    | Total   |
|---------------------|------------------|--------------|-----------------------------|--------|--------|---------|---------|
|                     | (Spad Unit)      | (cm)         | (g DW plant <sup>-1</sup> ) |        |        |         |         |
| Control             | 35c              | 32c          | 1.0d                        | 0.69c  | 1.78b  | 0.76c   | 4.64c   |
| Melia               | 40a              | 46abc        | 1.69bcd                     | 1.20bc | 2.83ab | 1.08abc | 6.80bc  |
| Green neem 1        | 36bc             | 71a          | 1.60bcd                     | 1.70ab | 3.13a  | 1.21ab  | 7.64abc |
| Pelleted neem       | 37b              | 56ab         | 2.22abc                     | 1.70ab | 3.56a  | 1.23ab  | 8.72ab  |
| Neem cake 1         | 38b              | 62ab         | 2.32ab                      | 1.96ab | 3.08a  | 1.40ab  | 8.76ab  |
| Deoiled neem        | 37b              | 58ab         | 1.11cd                      | 1.31bc | 2.96a  | 1.10abc | 6.49bc  |
| Oiled neem          | 40a              | 42bc         | 1.44bcd                     | 1.18bc | 2.49ab | 1.02bc  | 6.14bc  |
| Green neem 2        | 42a              | 65ab         | 2.92a                       | 2.26a  | 3.50a  | 1.51a   | 10.2a   |
| <i>Significance</i> | **               | ***          | ***                         | ***    | **     | **      | **      |

\*\* , \*\*\* = effect of *Meliaceae* derivatives significant at  $P > 0.01$  and  $0.001$ , respectively. In the same column, values followed by the same letter are not statistically different.

Table 4. Effect of soil-applied *Meliaceae* derivatives on N, P, K, Ca, Mg, Mn, Fe, Cu and Zn leaf concentrations at the end of the experiment A.

| Treatment     | N                     | P      | K    | Ca     | Mg    | Mn                     |     | Cu   |    | Zn |
|---------------|-----------------------|--------|------|--------|-------|------------------------|-----|------|----|----|
|               |                       |        |      |        |       | Fe                     |     |      |    |    |
|               | (g kg <sup>-1</sup> ) |        |      |        |       | (mg kg <sup>-1</sup> ) |     |      |    |    |
| Control       | 13.7                  | 4.3bc  | 15.8 | 11.1a  | 3.1ab | 31a                    | 122 | 5.27 | 22 |    |
| Melia         | 14.6                  | 4.1bc  | 18.6 | 10.5ab | 3.1ab | 29ab                   | 97  | 6.50 | 22 |    |
| Green neem 1  | 15.8                  | 5.8ab  | 20.1 | 9.6ab  | 3.3ab | 24b                    | 99  | 6.2  | 27 |    |
| Pelleted neem | 13.4                  | 5.5abc | 17.1 | 9.1abc | 3.0ab | 29ab                   | 100 | 6.02 | 24 |    |
| Neem cake 1   | 13.4                  | 5.0bc  | 19.4 | 8.7bc  | 2.5b  | 24b                    | 92  | 5.53 | 22 |    |
| Deoiled neem  | 16.2                  | 7.1a   | 20.9 | 10.3ab | 3.8a  | 33a                    | 79  | 6.92 | 23 |    |
| Oiled neem    | 15.8                  | 6.0ab  | 18.9 | 9.7ab  | 3.1ab | 31a                    | 91  | 5.84 | 27 |    |
| Green neem 2  | 14.5                  | 3.6c   | 17.7 | 7.7c   | 2.5b  | 31a                    | 90  | 5.50 | 23 |    |
| Significance  | ns                    | ***    | ns   | ***    | *     | *                      | ns  | ns   | ns |    |

ns, \*, \*\*\* = effect of *Meliaceae* derivatives not significant, significant at  $P \leq 0.05$ , and 0.001, respectively. In the same column, values followed by the same letter are not statistically different.

Table 5. Effect of soil-applied *Meliaceae* derivatives on total N removed by plant at day 68 and 113. Experiment A.

| Treatment           | N (mg) |        |
|---------------------|--------|--------|
|                     | Day    |        |
|                     | 68     | 113    |
| Control             | 20.8b  | 29.2c  |
| Melia               | 38.3ab | 49.9bc |
| Green neem 1        | 45.0ab | 55.7b  |
| Pelleted neem       | 42.4ab | 52.7b  |
| Neem cake 1         | 49.8a  | 60.7ab |
| Deoiled neem        | 39.7ab | 45.2bc |
| Oiled neem          | 37.8ab | 45.5bc |
| Green neem 2        | 51.2a  | 75.9a  |
| <i>Significance</i> | *      | ***    |

\*, \*\*\* = effect of *Meliaceae* derivatives significant at  $P \leq 0.05$ , and 0.001, respectively. In the same column, values followed by the same letter are not statistically different.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

Table 6. Mean  $\pm$ standard deviation (n =4) of the percentage of N derived from labelled melia leaves (NDFML) and correspondent recovery (%) of  $^{15}\text{N}$  in plant organs 30, 90, 131 and 173 day after soil application of  $^{15}\text{N}$ -enriched melia leaves (experiment B).

| Treatment        | NDFML (%)         |                  |                  |                  | $^{15}\text{N}$ Recovery (%) |                   |                   |                   |
|------------------|-------------------|------------------|------------------|------------------|------------------------------|-------------------|-------------------|-------------------|
|                  | day               |                  |                  |                  | day                          |                   |                   |                   |
|                  | 30                | 90               | 131              | 173              | 30                           | 90                | 131               | 173               |
| Shoot and leaves | 7.31( $\pm$ 11.2) | 38.6( $\pm$ 9.1) | 39.1( $\pm$ 7.2) | 30.2( $\pm$ 2.7) | 0.008( $\pm$ 0.01)           | 3.57( $\pm$ 1.9)  | 4.55( $\pm$ 1.9)  | 2.40( $\pm$ 2.47) |
| Stem             | 0.14( $\pm$ 0.02) | 15.7( $\pm$ 3.5) | 24.0( $\pm$ 4.3) | 21.1( $\pm$ 2.6) | 0.003( $\pm$ 0.02)           | 0.29( $\pm$ 0.13) | 0.56( $\pm$ 0.19) | 0.69( $\pm$ 0.43) |
| Roots            | 1.09( $\pm$ 0.37) | 26.2( $\pm$ 4.7) | 29.6( $\pm$ 7.8) | 22.4( $\pm$ 2.9) | 0.04( $\pm$ 0.02)            | 1.43( $\pm$ 0.79) | 2.66( $\pm$ 1.15) | 2.47( $\pm$ 2.29) |




Figure 1. Effect of soil-applied  $^{15}\text{N}$ -enriched melia leaves on N derived from labelled leaves (NDFML) in roots, stem, and shoot+leaves. At each sampling date: ns, \*, \*\* = effect of organ not significant, significant at  $P \leq 0.05$ , and  $P \leq 0.01$ , respectively. Within the same date, values followed by the same letter are not significantly different. In box, the detailed trend at the first sampling time (day 30) is reported.




Figure 2 Effect of soil-applied <sup>15</sup>N-enriched melia leaves on soil microbial biomass C. Interaction time\*treatment significant at  $P \leq 0.001$ . Two standard error of means (SEM) = 16 is minimum difference between two statistically different values.




Figure 3. Effect of soil-applied  $^{15}\text{N}$ -enriched melia leaves on soil microbial biomass N. Interaction time\*treatment not significant. Two standard error of means (SEM) = 3.3 is minimum difference between two statistically different values.