

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Quantification of 2- and 3-isopropylmalic acids in forty Italian wines by UHPLC-MS/MS triple quadrupole and evaluation of their antimicrobial, antioxidant activities and biocompatibility

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Ricciutelli M., Bartolucci G., Campana R., Salucci S., Benedetti S., Caprioli G., et al. (2020). Quantification of 2- and 3-isopropylmalic acids in forty Italian wines by UHPLC-MS/MS triple quadrupole and evaluation of their antimicrobial, antioxidant activities and biocompatibility. FOOD CHEMISTRY, 321, 1-8 [10.1016/j.foodchem.2020.126726].

Availability:

This version is available at: <https://hdl.handle.net/11585/849864> since: 2022-01-31

Published:

DOI: <http://doi.org/10.1016/j.foodchem.2020.126726>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

**Quantification of 2- and 3-isopropylmalic acids in forty Italian wines by
UHPLC-MS/MS triple quadrupole and evaluation of their antimicrobial,
antioxidant activities and biocompatibility**

Massimo Ricciutelli¹, Gianluca Bartolucci², Raffaella Campana³, Sara Salucci³, Serena
Benedetti³, Giovanni Caprioli,^{1*} Filippo Maggi¹, Gianni Sagratini¹, Sauro Vittori¹, Simone
Lucarini^{3*}

¹School of Pharmacy, University of Camerino, Via Sant'Agostino 1, 62032 Camerino,
Italy

²Department of Neurosciences Psychology, Drug Research and Child Health
(NeuroFarBa), University of Florence, Florence, 50139, Italy

³Department of Biomolecular Sciences, Piazza del Rinascimento 6, University of Urbino
Carlo Bo, 61029 Urbino, Italy

***Corresponding authors:**

Giovanni Caprioli, School of Pharmacy, University of Camerino, via S.Agostino 1, 62032
Camerino, Italy. Phone: +390737402238. Fax: +390737637345. Email:
giovanni.caprioli@unicam.it

Simone Lucarini, Department of Biomolecular Sciences, Piazza del Rinascimento 6,
University of Urbino Carlo Bo, 61029 Urbino, Italy. Phone: +390722303333. Fax:
+390722303313. Email: simone.lucarini@uniurb.it

Abstract

2-Isopropylmalic acid (2-IPMA) and 3-isopropylmalic acid (3-IPMA), recently discovered in wines, were simultaneously quantified in forty wines by UHPLC-MS/MS triple quadrupole. Principal component analysis displayed that red wines were more correlated with high amounts of 2-IPMA (average content 31.60 mg/L); white wines were mostly characterized by low levels of both organic acids. No correlation of their levels to other wine features (wine ageing or alcoholic content) were found. 2-IPMA and 3-IPMA showed MICs values of 4096 mg/L and MBCs values of 8192 mg/L or higher against several food borne pathogens. In association, an interesting lower MIC and MBC values (2048 mg/L and 4096 mg/L respectively) were observed against *Y. enterocolitica*. Interestingly, 3-IPMA showed a mild antioxidant activity by DPPH assay ($EC_{50} = 3940$ mg/L), higher than that of 2-IPMA ($EC_{50} > 4800$ mg/L). No toxicity of these compounds against human colorectal and liver cells (TB assay) was observed.

Keywords: wine, 2-isopropylmalic acid, 3-isopropylmalic acid, antimicrobial activity, antioxidant

1. Introduction

The category of wine flavour substances is composed of volatile compounds especially responsible for the odour and non-volatile components particularly contributing to the taste ([Whiting, 1976](#); [Lamikanra, Inyang & Leong, 1995](#)). Organic acids comprise one of the main taste groups, namely, that of sourness. The quantitatively dominating acids of grapes are maleic, tartaric, and citric acids, the first two of which account for over 90% of the total acid content of grapes ([Schreier & Jennings, 2013](#)). Wine is also rich in phenolic compounds, that are important components as they affect organoleptic characteristics, such as colour, astringency and aroma ([Kennedy, 2008](#)). Some years ago, Ginjom et al. ([2011](#)) identified and quantified the individual phenolic compounds present in wine at different wine-making stages but they were not able to identify two compounds, the first one (named U1) with a low molecular mass (176 u) and a large peak area eluting at around 8.66 min, with a maximum absorbance at 275 nm; the second (named U2) with a high molecular mass (508 u) and a large peak area eluting at around 34.52 min, with a maximum absorbance at 360 nm. They noticed that the concentration (mg GAE/l) of U1 increased during ageing, suggesting that it was continuously released either from the oak and/or from the degradation of other phenolics in the wine. The other unknown compound (U2) showed maximum concentration during oaking but decreased as the wine was bottle-aged. Both of these unknowns were independently reported by [Monagas et al. \(2005\)](#) and our group ([Salucci et al., 2017](#)).

In a recent paper, our research group has identified in wines not one but two isomeric compounds, corresponding to U1, with molecular weight of 176 u by using two LC-MS systems, i.e. LC-IT (ion trap) and LC-Q-Orbitrap ([Ricciutelli et al., 2019](#)). The two compounds were not chromatographically separated, as the quantification were performed by extracting the specific fragment of the two co-eluting isomers (i.e. m/z 115 or m/z 73)

and they were determined only in ten among red and white Italian wines. Thus, an improving of the chromatography separation would be desirable to better quantify these two molecules and more wine samples need to be analysed to correlate 2-IPMA and 3-IPMA levels with wine features. The molecular structure of the two molecules is reported in [Fig 1S](#). From literature, 2- and 3-IPMA are intermediates for the biosynthesis of leucine in yeast. In *Saccharomyces cerevisiae*, 2-IPMA is produced in mitochondria from isoketovalerate and then exported to the cytosol where it is isomerized to 3-IPMA. The latter is converted to leucine in two steps. Yeast cells naturally secrete these compounds into their surroundings ([Calvo, Kalyanpur, & Stevens 1962](#); [Dumlao, Hertz, & Clarke 2008](#) and [Marobbio, Giannuzzi, Paradies, Pierri, & Palmieri, 2008](#)). In particular, 2-IPMA secretion chelates aluminium ions and prevents them from entering to the cells, resulting in aluminium tolerance and reducing aluminum toxicity to the yeast cell ([Suzuki, Tamura, Nakanishi, Tashiro, Nishizawa, & Yoshimura, 2007](#)). The literature indicates that the concentrations of the active compounds in wines vary widely among the different grape cultivars from which they are derived. Indeed, it is well-recognized that biological activities and health benefits of wines depend on both the total amounts and the specific amounts of highly bioactive compounds ([Friedman, 2014](#)). To this regard, despite the evidence for the antimicrobial effectiveness of wine ([Moretro & Daeschel, 2004](#); [Carneiro, Couto, Mena, Queiroz, & Hogg, 2008](#); [Daglia, Papetti, Grisoli, Aceti, Dacarro, & Gazzani 2007](#)), the contributions of specific wine components to the wine antimicrobial activity have not been well characterized ([Boban, Tonkic, Budimir, Modun, Sutlovic Punda-Polic &, Boban, 2010](#)). Different components of wine have been proposed to contribute to the antimicrobial activity, but the mechanisms responsible for this activity are not fully understood. In that regard, the reported investigations can be generally divided into two main groups, those giving emphasis to the role of wine phenolics and those accentuating

the role of nonphenolic constituents of wine (Friedman, 2014). The antioxidant activity and cytotoxicity of phenolic compounds in wine were exhaustively reported (Waterhouse, 2002; Salucci et al., 2017 and Prasad, Jeyanthimala, & Ramachandran 2009). On the other hand, fewer reports could be found for acid components (Nelson, Rush, & Wilson, 2016; and Robles, Fabjanowicz, Chmiel, & Plotka-Wasyłka, 2019).

In the light of the scant information available in literature, the aims of the work were a) to develop a new and fast analytical method to simultaneously quantify 2- and 3-IPMA in forty wines (white and red ones) by using ultra-high-performance liquid chromatography–tandem mass spectrometry (UHPLC-MS/MS) triple quadrupole, b) to preliminary characterize the biological activity of these malic acid derivatives. In particular, the antibacterial and bactericidal activities of 2- and 3-IPMA, alone and in associations, against several foodborne pathogens were assessed. Moreover, antioxidant and biocompatibility studies were also carried out.

2. Materials and Methods

2.1 Reagents and standards

The analytical standards of 2-IPMA (CAS Number 3237-44-3) and 3-IPMA (CAS Number 16048-89-8) were purchased from Sigma-Aldrich (Milano, Italy). The stock standard solution was prepared by dissolving 10 mg of each analyte in 10 mL of methanol and stored in a glass-stoppered bottle at 4°C in the dark. Standard working solutions, at various concentrations, were prepared daily by appropriate dilution of aliquots of the stock solutions in methanol. HPLC-grade acetonitrile and methanol were purchased from Sigma-Aldrich (Milan, Italy), while HPLC-grade formic acid 99-100% was bought from J.T. Baker B.V. (Deventer, Holland). For sample preparation and chromatographic analysis, deionized water of 18.2 MΩ/cm resistivity purified with a Milli-Q system (Millipore,

Bedford, USA) was used. All solvents were filtered through a 0.2 μm polyamide filter from Sartorius Stedim (Goettingen, Germany) while all wine samples were filtered through a 0.2 μm single use membrane syringe filter from Phenomenex (Bologna, Italy) before HPLC analysis.

2.2 Wine samples

Red and white wine samples were purchased in the supermarkets of the Camerino area (Camerino, Macerata, Italy).

2.3 Sample preparation

Briefly, 5 mL wine samples were extracted three times with ethyl acetate (5 mL) following and downscaling a previous published method (Ricciutelli et al., 2019; Salucci et al., 2017). The ethyl acetate extracts were pooled together prior to rotary evaporation at 30°C. The residue was re-dissolved in 0.25 mL methanol, filtrated through a 0.2 μm membrane filter from Phenomenex (Bologna, Italy) and then directly injected into the HPLC-MS/MS.

2.4 UHPLC-MS/MS analysis

UHPLC–MS/MS studies were performed using an Agilent 1290 Infinity series and a Triple Quadrupole 6420 from Agilent Technology (Santa Clara, CA) equipped with an electrospray (ESI) source operating in negative ionization mode. The separation of polyphenols was achieved using a Zorbax RRHD C18 analytical column (50 \times 2.10 mm i.d., 1.8 μm), also from Agilent (USA). The mobile phase for UHPLC MS/MS analysis was a mixture of water (A, 97%) and acetonitrile (B, 3%), both with formic acid 0.1% at a flow rate of 0.4 mL min⁻¹ with gradient elution. The solvent composition varied as follows: 0 min, 97% A; 5.5 min, 97% A; 5.5-14 min, 30% A; 14–19 min, 30% A; 19-24 min, 97% A; until the end of the run at 30 min. The injection volume was 1 μL . The temperature of the column was 30°C and the temperature of the drying gas in the ionization source was 300°C. The gas flow was 12 l min⁻¹, the nebulizer pressure was 50

psi and the capillary voltage was 4000 V. Detection was performed by electrospray ionization (ESI)-MS in the “multiple reaction monitoring” (Dynamic-MRM) mode. The selected ion transition and the settings of the mass analyzer are reported in [Table 1](#). Before use, all solvents were filtered through a 0.2 µm filter from Sartorius Stedim (Goettingen, Germany), and before UHPLC analysis, all samples were filtered through a 0.2 µm single use syringe filter from Phenomenex (Bologna, Italy).

2.5. Principal Component Analysis (PCA) Forty samples of red and white wines were analyzed by PCA in order to identify possible correlation groups based on the content of the two isomers of isopropyl-malic acid. A covariance matrix based on 40 samples x 2 variables was created and data were analyzed by STATISTICA v.7.1 (Stat Soft Italia S.r.l., Vigonza, Italy). Score and loading plots were generated by calculating eigenvalues.

2.6 Bacterial strains and culture conditions

Five reference human pathogens were used in this study: *Escherichia coli* O157:H7 ATCC 35150, *Listeria monocytogenes* ATCC 7644, *Staphylococcus aureus* ATCC 29213, *Salmonella enterica* ATCC 13314 and *Yersinia enterocolitica* ATCC 14053. The strains were routinely grown on Tryptic Soy Agar (TSA, VWR, Milan, Italy) at 37 °C for 24 h. All the stock cultures were kept at -80 °C in Nutrient broth (Oxoid, Italy) with 15% of glycerol.

2.7. Determination of minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC)

MICs of all the tests compounds were determined by standard micro-dilution method. First, for each compound a stock solution was prepared in one mL of distilled water and subsequently sterilized by 0.22 µm pore size filters (VWR). Several colonies of each bacterial strain were inoculated in 10 mL of sterile Mueller-Hinton broth (MHB) (VWR) and incubated at 37 °C for 18 h. At the end of incubation, each bacterial suspension was

adjusted by spectrophotometer to about 10^6 cfu/mL (OD 610nm 0.13-0.15) and 100 μ L was added in wells of the 96-well plate together with the appropriate volumes of the test solutions (final concentration from 256 to 8192 mg/L). In addition, 2-IPMA and 3-IPMA were also tested in association (ratio 10:1 and 1:10). For the association, the analyzed concentrations are the sum of the two acids contribution in the correct ratio. For example, total concentration = 4096 μ g/mL of 2-IPMA:3-IPMA (10:1 ratio) is the sum of 3724 μ g of 2-IPMA (10 parts) and 372 μ g of 3-IPMA (1 part) in 1 mL of solution. Two rows were used for positive (bacteria alone) and negative controls (MHB alone), respectively. Amoxicillin and gentamicin (1-128 mg/L) (Sigma, Italy), were used and internal control. After 24h of incubation at 37°C, the plates were observed and the MIC was determined as the lowest concentration that inhibits the visible growth in comparison with the control samples. The optical density (600 nm) of each well was also assessed using a Multiscan Ex Microplate Reader (Thermo Scientific, Italy). All data were expressed as the mean of three independent experiments performed in duplicate. For MBC determination, ten microliters from the invisible growth wells were inoculated in triplicate on TSA and incubated at 37 °C for 24 h; MBC was defined as the lowest concentration of each compound that completely inhibited growth on TSA (approximately 99.5% of killed microorganisms). All the experiments were performed in duplicate.

2.8. DPPH radical scavenging activity

The antioxidant activity of 2-IPMA, 3-IPMA and Malic acid (MA) was evaluated using the DPPH assay, which measures the ability of a compound to act as free radical scavenger or hydrogen donor (Sagar B. Kedare and R. P. Singh, 2011). Briefly, 0.300 mL of sample diluted in ethanol (range 0.3-9.6 mg/mL) was added to 0.300 mL of 100 μ M DPPH in ethanol. Absorbance decrease at 517 nm was recorded after 30 min at room temperature. The scavenger effect was calculated as $\% = [(\text{Abs } 517 \text{ nm of blank} - \text{Abs } 517 \text{ nm of$

sample /Abs 517 nm of blank] x 100. EC₅₀ (i.e. the concentration required to obtain a 50% antioxidant effect) was also calculated. Ascorbic acid (initial concentration 0.3-9.6 µg/mL) was used as standard to check the correctness of the procedures. Statistical significance was assessed by one-way ANOVA and Tukey's posthoc test for multiple comparisons (PRISM 6, GraphPad Software, USA). Significance level was set at $p < 0.05$.

2.9. Human cell culture and treatments (Trypan Blue exclusion assay)

The human colorectal Caco-2 and the liver HepG2 cancer cells, were seeded on multiwell plates at a cell density of 1×10^6 cells/mL and were cultured in a DMEM high glucose medium supplemented with 10% heat-inactivated fetal bovine serum (FBS), 2 mM glutamine, 1% penicillin-streptomycin, and MEM nonessential amino acid solution (1X).

All culture models were maintained at 37°C in humidified air with 5% CO₂ and cell behavior were monitored by means of an Inverted Microscopy (Eclipse TE2000-S Nikon; objective 10x). Cells were treated with different doses of 2-IPMA, 3-IPMA and MA to verify their potential toxic effect. 2-IPMA, 3-IPMA and MA were added to the culture medium (and monitored until 48h) at the following final concentrations: 1.76, 8.8, 35.2, 88, 176, 880 and 1176 mg/L. Moreover, 2-IPMA e 3-IPMA have been administrated to cell cultures in association (ratio 10:1 and 1:10) using these dosages: 1.76 mg/L + 15.84 mg/L, 35.2 mg/L + 316.8 mg/L, 176 mg/L + 1584 mg/L. The treatments were compared with 50 µM etoposide, a known chemoterapic drug (Salucci et al., 2014). Trypan Blue (TB) exclusion assay was employed to monitor cell viability after all treatments since it represents a useful method to quantify live and dead cell number (Salucci et al., 2018).

In addition, cells have been exposed to 0.5 mM hydrogen peroxide (H₂O₂) for 24 h, a pro-oxidant agent and cell death trigger (Salucci et al., 2014). MA, 2-IPMA, 3-IPMA alone and the association of 2- and 3-IPMA (ratio 10:1 and 1:10) have been administrated to cells for 24 h before H₂O₂ treatment to verify their potential protection against oxidative damage.

Also in this case, TB exclusion assay has been used to monitor cell viability for both human colorectal Caco-2 and the liver HepG2 cancer cell lines.

Data collected from each experimental condition were presented as mean of living cell percentage \pm standard deviation of the mean (SEM).

3 Results and discussion

3.1 Method validation of the HPLC-MS/MS method

The investigated performance characteristics of the developed method were linearity, limit of detection (LOD), limit of quantification (LOQs) (**Table 2**), and matrix effect (as signal suppression/enhancement %) (**Table 1S**). Linearity was tested by injecting 7 different concentrations of authentic standard mixtures of the analytes in solvent (**Table 2**) from 0.5 to 100 mg/L. Calibration curves (concentrations versus peak areas) were determined by least-squares regression analysis obtaining correlation coefficients (R^2) higher than 0.9942. The reproducibility of the chromatographic retention times was examined five times over a five-day period ($n=25$) obtaining high stability ($RSD < 1.3\%$). The LODs and LOQs were estimated on the basis of 3:1 and 10:1 signal-to-noise ratios obtained with standards containing the compounds at low concentration levels. The estimated LODs and LOQs were 0.01 and 0.02 mg/L and 0.03 and 0.06 mg/L for 2-IPMA and 3-IPMA respectively (**Table 2**).

Accuracy was calculated via spike-recovery and it was determined using percentage relative errors at two different concentrations by comparing the true value of the analyte in the sample (true concentration) with the values obtained by analysis (measured concentration). As it can be seen from **Table 2S** in supplementary materials, the % relative error for the standards are between + 2.9 and +10.4 % for the 2 different analytes. The lowest % relative error is (+2.9%) obtained for 2-IPMA at concentration level of 50 mg/L,

meanwhile the highest % relative error is +10.4% obtained for 3-IPMA at concentration level of 5 mg/L. The measure of precision usually is expressed in terms of imprecision and computed as standard deviation of the test result. The precision of this method was determined by monitoring the RSD % of the three replicates. As it can be seen from **Table 2S** reported in supplementary materials, all the RSD % obtained are between 2.8 – 5.9% . The lowest RSD was 2.8% for 2-IPMA at concentration level of 50 mg/L, while the highest RSD was 4.9% for 3-IPMA at concentration level of 5 mg/L.

Matrix effect was studied comparing the slope of a calibration curve in pure solvent and of a matrix-matched curve prepared by adding the standards to a wine extract (after extraction). The calibration curve was obtained by subtracting in each concentration the amount of polyphenols obtained from the blank extract. These curves were obtained after injecting seven concentration levels, between 5 and 250 mg/L. The slope of the standard addition plot was compared with the slope of standard calibration plot to evaluate the matrix effects. The signal suppression/enhancement (SSE) was calculated according to the following equation (Caprioli, Nzekoue, Giusti, Vittori & Sagratini, 2018):

$$\text{SSE \%} = (\text{slope matrix matched curve} / \text{slope pure solvent curve}) \times 100;$$

If SSE (%) is about 100% there is no matrix effect, values < 100% indicate signal suppression, while values > 100% indicate signal enhancement. In our study the SSE % ranged from 46 (2-IPMA) to 59 (3-IPMA) indicating a negative matrix effect (ion suppression) of 54 and 41%, respectively (**Table 1S**). In the following paragraph, the quantification of the two analytes in wine samples have been reported taken into account ME studies.

3.2 Quantification of 2-IPMA and 3-IPMA in Italian wines

The validated UHPLC-MS/MS triple quadrupole method was used to analyse 40 Italian wine samples, specifically 19 red and 21 white wines and the two analytes were found in all samples (**Table 3**). **Fig. 2S** show an HPLC-MS/MS chromatogram of a red wine samples with the TIC (total ion chromatogram) and the multiple reaction monitoring (MRM) transition of the two organic acids. This method offers significant improvement with respect to our previous work ([Ricciutelli et al., 2019](#)). The Zorbax RRHD C18 analytical column (50×2.10 mm i.d., $1.8 \mu\text{m}$) provided good performance in terms of resolution and selectivity as the two isomers were perfectly separated (please see **Fig. 2S**). Moreover, LODs and LOQs obtained are at least 10 times lower than values reported in our previous article ([Ricciutelli et al., 2019](#)). The concentration of the two analytes has been expressed in mg/L in the original wine samples. Concerning red wines, the contents of 2-IPMA ranged from 18.9 mg/L of Montepulciano (sample n°1) to 41.6 mg/L of Merlot (n°19). Nero Buono (n°18), Montepulciano D'Abruzzo (n°17) and Vernaccia di Serrapetrona (n°16) wines displayed a conspicuous content of 2-IPMA with levels of 41.25, 38.25 and 37.63 mg/L, respectively. On the other hand, 3-IPMA was found in low concentration ranging from 1.43 mg/L in Montepulciano (n°1) to 4.13 mg/L of Merlot (n°19). The average content of 2-IPMA in the 19 wines was 31.60 mg/L, whereas that of 3-IPMA was 1.65 mg/L.

Concerning the white wines, the contents of the two derivatives of malic acid are a bit lower respect to red ones. In fact, 2-IPMA ranged from 6.7 mg/L of Primofiore (n°20) to 27.8 mg/L of Verdicchio di Matelica (sample n°40). Noteworthy levels of 2-IPMA were found also in Bellone Anfora sample n°39 (26.00 mg/L) and Verdicchio di Matelica sample n°38 (25.82 mg/L). As reported for red wines, the amounts of 3-IPMA detected in white wines were lower with respect to 2-IPMA. The content of 3-IPMA

ranged from 0.56 mg/L of Primofiore (n°20) to 2.42 mg/L of Verdicchio di Matelica n°40 (2.42 mg/L). Bellone (n°34), Verdicchio di Matelica (n°38) and Muller Thurgaut (n°37) wines displayed a conspicuous content of 3-IPMA with levels of 1.91, 1.63 and 1.57 mg/L, respectively. The average contents of 2-IPMA and 3-IPMA in white wines were 19.57 mg/L and 0.93 mg/L, respectively.

PCA analysis allowed to visualize two main groups of wine samples as depicted in **Fig. 1**. 2-IPMA was the main variable influencing data variability in the first principal component (99.6%, values of eigenvectors: 8.42; -0.03). They were represented mostly by red wine samples on the right-hand side of the score plot that were more correlated with high amounts of 2-IPMA. On the other hand, white wine samples were mostly in the left-hand side of the score plot and where characterized by low levels of both organic acids.

3.3 Antimicrobial activity of 2-IPMA and 3-IPMA

The assessment of the antibacterial activity of 2-IPMA and 3-IPMA against five human pathogens was performed according to the National Committee for Clinical Laboratory Standards (NCCLS) document M100-S12 method (**Table 4**). As shown, MICs values of 4096 mg/L were determined for both the compounds for both gram-positive (*L. monocytogenes* ATCC 7644 and *S. aureus* ATCC 29213) and gram-negative (*E. coli* O157:H7 ATCC 35150, *S. enterica* ATCC 13314 and *Y. enterocolitica* ATCC 14053) bacteria. As regards the bactericidal activity, most of the bacterial strains were killed by 2-IPMA and 3-IPMA at the concentration of 8192 mg/L, though in some cases higher MBCs values (>8192 mg/L) were observed. The association 2-IPMA:3-IPMA in the ratio 10:1 did not show remarkable differences in MICs values, resulting of 4096 mg/L for all the microorganisms. Similarly, no differences in MICs values were evidenced using 2-

IPMA:3-IPMA in the ratio 1:10 (MIC 4096 mg/L), with the only exception of *Y. enterocolitica* ATCC 14053 for which a lower MIC (2048 mg/L) was noted with an interesting synergistic effect. As regards the bactericidal activity of the mentioned combinations, MBCs >4096 mg/L were observed for *E. coli* O157:H7 ATCC 35150, *L. monocytogenes* ATCC 7644, *S. aureus* ATCC 29213 and *S. enterica* ATCC 13314, while the MBC value for *Y. enterocolitica* ATCC 14053 resulted to be 4096 mg/L. The MICs of the internal controls showed the susceptibility of all the examined bacteria to Gentamicin, with MICs ranging from 4 to 16 mg/L, as previously observed (Campana et al., 2019; Perinelli et al., 2018, Mengist et al., 2018), while a lower susceptibility was detected for Amoxicillin (from 128 to >128 mg/L) (Obaidat and Stringer 2019; Mengist et al., 2018).

3.4. Antioxidant activity of 2-IPMA and 3-IPMA by DPPH radical scavenging assay

Among the test compounds, 3-IPMA showed the highest scavenging activity towards DPPH ($EC_{50} = 3.94 \pm 0.13$ mg/mL), followed by 2-IPMA and MA ($EC_{50} > 4.8$ mg/mL for both compounds) (Figure 2A). Ascorbic acid, used as reference compound, presented an EC_{50} value of 1.64 ± 0.11 μ g/mL (Data not shown). 2-IPMA and MA show almost comparable no anti-oxidant activities, however 3-IPMA shows a very mild anti-oxidant activity. This property could be attributed to the relative stability of the tertiary radical formed with DPPH compared to the secondary radical formed from 2-IPMA (see Figure 1S, supplementary file).

3.5. Toxicity studies of 2-IPMA and 3-IPMA by TB exclusion assay

2-IPMA, 3-IPMA and MA administered to cell models and monitored until 48h, were not cytotoxic for colon and liver human cells and did not affect their viability. In fact, TB assay revealed that the number of living cells after all treatments is comparable to that

quantified in control condition. The same results can be observed by exposing cells to 2-IPMA and 3-IPMA in association. On the contrary, after etoposide exposure an evident decrease of living cells were observed. These findings clearly appear in **Figure 2B** and **Figure 3S** (see supplementary file) that show no toxicity of the tested acids in all the concentration for HepG2 and Caco-2 cells viability after 24 h of treatments, respectively. The absence of cytotoxicity is confirmed when the 2- and 3-IPMA are used in association (10:1 and 1:10 ratio, **Figure 4S** in supplementary file). Furthermore, 2- and 3-IPMA used singly or in combination do not evidence any protection against cell death induced by the oxidative damage of hydrogen peroxyde (data not shown).

4. Conclusions

A new, sensitive and fast analytical method to simultaneously quantify 2-IPMA and 3-IPMA in forty wines was developed and validated by using UHPLC-MS/MS triple quadrupole. The two analytes were found in all wine samples analyzed; the average content of 2-IPMA was 31.60 mg/L in red wines and 19.57 mg/L in white wines; whereas that of 3-IPMA was 1.65 mg/L and 0.93 mg/L, respectively. Statistical analyses displayed that red wine samples seem to be correlated with high amounts of 2-IPMA; on the other hand, white wine samples were mostly characterized by low levels of both organic acids. The two isomeric compounds were tested alone and in association against five representative food borne pathogens for MIC and MBC determination. 2-IPMA and 3-IPMA showed mild antibacterial and bactericidal activity, having MICs values of 4096 mg/L (23.3 mM) and MBCs values of 8192 mg/L (46.5 mM) or higher for both gram-positive (*L. monocytogenes* and *S. aureus*) and gram-negative (*E. coli*, *S. enterica* and *Y. enterocolitica*) bacteria. In association, 2-IPMA and 3-IPMA did not show remarkable

differences in MICs and MBCs values. However, using 2-IPMA together with 3-IPMA in the ratio 1:10, an interesting lower MIC and MBC (2048 mg/L and 4096 mg/L respectively) were observed against *Y. enterocolitica*, highlighting a possible synergistic effect of the two isomeric acids against this gram-negative strain. Interestingly, 3-IPMA showed a mild antioxidant activity by DPPH assay ($EC_{50} = 3940$ mg/L, 22.4 mM) higher than that of its isomer 2-IPMA ($EC_{50} > 4800$ mg/L). No toxicity of these compounds alone and in association was observed at all tested concentrations as evidenced by TB exclusion assay against human colorectal and liver cell lines. The mild antibacterial and mild antioxidant activities together with the good biocompatibility made 2- and 3-IPMA interesting candidates as preservatives for foods.

Conflict of Interests

The authors declare that no competing interests exist.

References

- Boban, N., Tonkic, M., Budimir, D., Modun, D., Sutlovic, D., Punda-Polic, V., & Boban M. (2010). Antimicrobial effects of wine: separating the role of polyphenols, pH, ethanol, and other wine components. *Journal of Food Science*, 75(5), M322-6.
- Calvo, M. J., Kalyampur, M. G., & Stevens C. M. (1962). 2-Isopropylmalate and 3-isopropylmalate as intermediates in leucine biosynthesis. *Biochemistry*, 6, 1157-1161.
- Campana, R., Merli, A., Verboni, M., Biondo, F., Favi, G., Duranti, A., & Lucarini S. (2019). Synthesis and evaluation of saccharide-based aliphatic and aromatic esters as antimicrobial and antibiofilm agents. *Pharmaceuticals*, 12 (4), 186.

383 Caprioli, C., Nzekoue, F., Giusti, F., Vittori, S., & Sagratini G. (2018). Optimization of an
 384 extraction method for the simultaneous quantification of sixteen polyphenols in thirty-one
 385 pulse samples by using HPLC-MS/MS dynamic-MRM triple quadrupole. *Food Chemistry*,
 386 266, 490-497.

387 Carneiro, A, Couto, J.A., Mena, C., Queiroz, J., & Hogg, T. (2008). Activity of wine
 388 against *Campylobacter jejuni*. *Food Control*, 19, 800–805.

389 Daglia, M., Papetti, A., Grisoli, P., Aceti, C., Dacarro, C., & Gazzani G. (2007).
 390 Antibacterial activity of red and white wine against oral streptococci. *Journal of*
 391 *Agricultural and Food Chemistry*, 55, 5038–42.

392 Dumlao, S. D., Papetti, A., Hertz, N., & Clarke G. (2008). Secreted 3-isopropylmalate
 393 methyl ester signals invasive growth during amino acid starvation in *Saccharomyces*
 394 *cerevisiae*. *Biochemistry*, 47, 698–709.

395 Friedman, M. (2014). Antibacterial, antiviral, and antifungal properties of wines and
 396 winery byproducts in relation to their flavonoid content. *Journal of Agricultural and Food*
 397 *Chemistry*, 62, 6025–6042.

398 Ginjom, I., D’Arcy, B., Caffin, N., & Gidley, M. (2011). Phenolic compound profiles in
 399 selected Queensland red wines at all stages of the wine-making process. *Food Chemistry*
 400 125, 823–834.

401 Lamikanra, O., Inyang I. D., & Leong, S. (1995). Distribution and effect of grape maturity
 402 on organic acid content of red muscadine grapes. *Journal of Agricultural and Food*
 403 *Chemistry*, 43, 3026-3028

404 Kennedy, J. A. (2008). Grape and wine phenolics: Observations and recent findings.
 405 *Ciencia E Investigacion Agraria*, 35, 107–120.

406 Marobbio, C. M. T., Giannuzzi, G., Paradies, E., Pierri, C. L., & Palmieri, F. (2008). α -
 407 Isopropylmalate, a leucine biosynthesis intermediate in yeast, is transported by the
 408 mitochondrial oxalacetate carrier. *Journal of Biological Chemistry*, 17, 283, 28445–28453.

409 Mengist, A., Mengistu, G., & Reta, A. (2018). Prevalence and antimicrobial susceptibility
 410 pattern of salmonella and shigella among food handlers in catering establishments at Debre
 411 Markos University, Northwest Ethiopia. *International Journal of Infectious Diseases*, 75,
 412 74-79.

413 Monagas, M., Suarez, R., Gomez-Cordoves, C., & Bartolome, B. (2005). Simultaneous
 414 determination of nonanthocyanin phenolic compounds in red wines by HPLC/DAD/ESI-
 415 MS. *American Journal of Enology and Viticulture*, 56, 139–147.

416 Moretro, T., & Daeschel, MA. (2004). Wine is bactericidal to foodborne pathogens.
 417 *Journal of Food Science*, 69, 251–7.

418 Nelson, H. N., Rush, K. L., & Wilson, T. (2016). Functions of common beverage
 419 ingredients. In Wilson T., Temple N. (Eds.), *Beverage Impacts on Health and Nutrition*.
 420 *Nutrition and Health*. (pp. 317–329). New York: Humana Press, Cham.

421 Obaidat, M.M., & Stringer, A.P. (2019). Prevalence, molecular characterization, and
 422 antimicrobial resistance profiles of *Listeria monocytogenes*, *Salmonella enterica*, and
 423 *Escherichia coli* O157:H7 on dairy cattle farms in Jordan. *Journal of Dairy Science*, 102,
 424 8710-8720.

425 Perinelli, D.R., Lucarini, S., Fagioli, L., Campana, R., Vllasaliu, D., Duranti, A., &
 426 Casettari, L. (2018). Lactose oleate as new biocompatible surfactant for pharmaceutical
 427 applications. *European Journal of Pharmaceutics and Biopharmaceutics*, 124, 55-62.

428 Prasad, N. R, Jeyanthimala, K, & Ramachandran, S. (2009). Caffeic acid modulates
 429 ultraviolet radiation-B induced oxidative damage in human blood lymphocytes. *J.*
 430 *Photochem. Photobiol. B*, 95, 196–203.

431 Ricciutelli, M., Moretti, S., Galarini, R., Sagratini, G., Mari, M., Lucarini, S., Vittori, S.,
 432 Caprioli, G. (2019). Identification and quantification of new isomers of isopropyl-malic
 433 acid in wine by LC-IT and LC-Q-Orbitrap. *Food Chemistry*, 294, 390-396.

434 Robles, A., Fabjanowicz, M., Chmiel, T., & Plotka-Wasyłka, J. (2019). Determination and
 435 identification of organic acids in wine samples. Problems and challenges. *Trends in*
 436 *Analytical Chemistry*, 120, 115630.

437 Sagar B. Kedare and R. P. Singh (2011). Genesis and development of DPPH method of
 438 antioxidant assay. *J Food Sci Technol.*, 48, 412–422.

439 Salucci S., Burattini S, Battistelli M., Baldassarri V., Curzi D., Valmori A., & Falcieri E.
 440 (2014). Melatonin prevents chemical-induced haemopoietic cell death. *International*
 441 *Journal of Molecular Sciences*, 15 (4), 6625-6640.

442 Salucci S., Burattini S., Giordano F. M., Lucarini S., Diamantini G., & Falcieri E. (2017).
 443 Further highlighting on the prevention of oxidative damage by polyphenol-rich wine
 444 extracts. *Journal of Medicinal Food*, 20, 410–419.

445 Salucci S., Burattini S, Buontempo F., Orsini E., Furiassi L., Mari M., Lucarini S., Martelli
 446 A. M., & Falcieri E. (2018). Marine bisindole alkaloid: A potential apoptotic inducer in
 447 human cancer cells. *European Journal of Histochemistry*, 62 (2), 2881.

448 Schreier, P., & Jennings, W. G. (1979). Flavor composition of wines: A review, C R C
 449 *Critical Reviews in Food Science and Nutrition*, 12, 59-111.

450 Suzuki, T., Tamura, S., Nakanishi, H., Tashiro, M., Nishizawa, N. K., & Yoshimura, E.
 451 (2007). Reduction of aluminum toxicity by 2-IPMA acid in the budding yeast
 452 *Saccharomyces cerevisiae*. *Biological Trace Element Research*, 120, 257-263.

453 Waterhouse, A. L. (2002). Wine phenolics. *Ann NY Acad Sci*, 957, 21-36.

454 Whiting, G. C. (1976). Organic acid metabolism of yeasts during fermentation of alcoholic
 455 beverages. A review. *Journal of the Institute of Brewing*, 82, 84-92.