

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Comparing effects and action mechanisms of BPA and BPS on HTR-8/SVneo placental cells

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Profita, M., Fabbri, E., Spisni, E., Valbonesi, P. (2021). Comparing effects and action mechanisms of BPA and BPS on HTR-8/SVneo placental cells. *BIOLOGY OF REPRODUCTION*, 105(5), 1355-1364 [10.1093/biolre/ioab139].

Availability:

This version is available at: <https://hdl.handle.net/11585/854113> since: 2024-05-20

Published:

DOI: <http://doi.org/10.1093/biolre/ioab139>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Profita M.; Fabbri E.; Spisni E.; Valbonesi P.: *Comparing effects and action mechanisms of BPA and BPS on HTR-8/SVneo placental cells*

BIOLOGY OF REPRODUCTION: Vol. 105 ISSN 0006-3363

DOI: 10.1093/biolre/ioab139

The final published version is available online at:

<https://dx.doi.org/10.1093/biolre/ioab139>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

1

2

3

Comparing effects and action mechanisms of BPA and BPS

4

on HTR-8/SVneo placental cells

5

Marilin Profita¹, Elena Fabbri^{1,2}, Enzo Spisni¹, and Paola Valbonesi¹

6

¹University of Bologna, Department of Biological, Geological, and Environmental Sciences,

7

Ravenna, Italy

² Corresponding author: Elena Fabbri, Department of Biological, Geological and Environmental Science, University of Bologna, Via S. Alberto 163, Ravenna, Italy

E-mail: elena.fabbri@unibo.it

Grant support: This work was supported by MIUR Italy (RFO 2018 and 2019 to EF)

8

9 Running title: Comparing BPA and BPS effects on placental cells

10

11 Summary sentence: Bisphenol A and its analog Bisphenol S, emerging contaminants founded in
12 human serum and placental fluids, significantly affected the biological functions of HTR-8/SVneo
13 cells, through estrogen receptors and MAPK mediated processes.

14

15 Keywords: BPA, BPS, HTR-8/SVneo cells, Endocrine disruptors, Human trophoblast, MAPK

Abstract

Bisphenol A (BPA) is one of the most investigated compound as a suspected endocrine disrupting chemical. It has been found at nM concentrations in the maternal serum, cord serum, and amniotic fluid and also permeates placental tissues. Attempts are being made to replace BPA with the analog Bisphenol S (BPS). Also BPS was found in maternal and umbilical cord serum, and urine samples from a large population of pregnant women. A few studies investigated BPA impact on the placentation process, and even less are available for BPS. This work aimed to elucidate and compare the effects of BPA and BPS on physiological functions of HTR-8/SVneo cells, derived from extravillous trophoblast of first-trimester pregnancy. Proliferation and migration ability of trophoblast cells were assessed in vitro after exposure to BPA or BPS (10^{-13} – 10^{-3} M). Further, induction of the inflammatory response by the bisphenols was studied. To provide insight into the molecular pathways implicated in the responses, experiments were carried out in the presence or absence of tamoxifen as estrogen receptors (ERs) blocker, and U0126 as ERK1/2 phosphorylation inhibitor. Data indicate that BPA significantly affects both proliferation and migration of HTR8/SVneo cells, through ER and ERK1/2 mediated processes. Differently, BPS only acts on proliferation, again through ER and ERK1/2 mediated processes. BPS, but not BPA, induces secretion of interleukins 6 and 8. Such effect is inhibited by blocking ERK1/2 phosphorylation. To the best of our knowledge, these are the first data showing that BPS affects trophoblast functions through ER/MAPK modulation.

1. Introduction

The increasing abnormalities in the female reproductive functions over the years highlighted a strong link between the exposure to industrial chemicals and the effects on human health [1–3]. Within this relationship, the concept of endocrine disrupting chemicals (EDCs) has taken a central role in the field of reproduction. EDCs comprise many different substances from industrial, agricultural, and domestic sources. They have been related with adverse effects on female reproductive system and may cause the development of gynaecological pathologies [4]. Further, exposure to EDCs may interfere with normal growth and development of the fetus and render the placenta unable to support the requested physiological functions [5]. Bisphenol A (BPA) is one of the most investigated substances as a suspected EDC. BPA is a monomer that was first developed as a synthetic estrogen in the 1890s and reported to have the efficacy of estrogens in stimulating the female reproductive system in rats in the 1930s [6]. Later, BPA has widely been used for its cross-linking properties in the manufacture of polycarbonate plastics and epoxy resins [7]. As an additive of plastics [8], BPA is released in food and water from plastics devices and from food packaging materials, as well as from personal care products and thermal receipts/papers, which constitute the main source of oral and dermal intake of BPA in humans. Epidemiological studies have reported that up to 95% of adults have detectable levels of BPA in their serum, saliva, and urine [9], within the nM range. Recent in vivo studies showed that BPA exposure is a known risk factor for the development of type 2 diabetes [10], obesity [11], earlier puberty [12], cardiovascular diseases [13], breast and ovarian cancer [14], altered liver function, oxidative stress and inflammation [15]. As to reproductive tissue, BPA has been found in maternal serum (0–154 ng/ml), amniotic fluid (0–8.38 ng/ml), cord serum (0–62.8 ng/ml), and placental tissue (0–104.9 ng/g) [9, 16–18]. Due to its high lipophilic property, BPA may permeate the placenta [19]. Given

the potential risks posed by BPA on human health and reproduction, attempts are being made to replace it with the analog Bisphenol S (BPS). Structurally similar to BPA, BPS was first synthesized in 1869 as a dye, and is currently used in a variety of industrial applications and introduced into many commercial products available on the market as “BPA-free” [20, 21]. Human exposure to BPS occurs through ingestion, inhalation, and dermal contact [22]. BPS was found in human urine, generally at concentrations and frequencies comparable to BPA [23]. However, the information available on adverse effects by BPS, in particular as a potential EDC, is rather poor. As a consequence of its structural similarity to BPA, BPS could potentially lead to similar endocrine disrupting capacity and effects on the reproductive system [24]. BPS was found in maternal (0.03–0.07 ng/ml) and cord serum samples (0.03–0.12 ng/ml) [25, 26], suggesting its possible ability to cross the placental barrier [27]. BPS was also detected in urine samples from the at-risk population, such as pregnant women [28, 29]. Emerging evidence suggest that BPS is capable of imitating properties of hormones [30], interacting with various physiological receptors, including estrogen receptors (ERs), androgenic and aryl hydrocarbon receptors [31]. Some in vitro studies observed that BPS affects estrogenic and antiandrogenic activities on cells derived from human ovarian and breast cancer, in a similar manner and potency of BPA [32, 33]. Considering its complexity, knowledge on potential EDC effects on placental function is still limited, especially regarding BPA and even more BPS. The development of the human placenta depends entirely on normal differentiation, proliferation, and invasion of trophoblast cells [34]. Impaired function of trophoblasts can lead to severe pregnancy complications [35, 36]. The proliferation and invasion of trophoblast cells represent a complex process, strictly regulated by many factors, including hormones, prostaglandins, cytokines and hypoxia, which either promote or inhibit proliferation and/or invasion [37]. Numerous signalling cascades/proteins are involved in the regulation of

trophoblast cell activity. Among them, mitogen-activated protein kinase (MAPK) pathways play a critical role in a wide range of biological processes [38, 39]. Further, cytokines and growth factors are secreted by trophoblast cells, and regulate the functional activity of the trophoblast via paracrine and autocrine mechanisms, control placenta development, and maintain immunological tolerance in the mother-fetus system [40, 41]. The present study aims to elucidate and compare the effects of BPA and BPS on the proliferation and migration ability of extravillous trophoblast HTR-8/SVneo cells, as well as on the secretion of inflammatory mediators. These cells preserve well-known factors controlling proliferation, migration and invasion, thus representing a good model for the in vitro study of molecular mechanisms at the basis of placentation, as well as of early events modulating placental development [42]. To provide insights into the molecular pathways implicated in cellular responses, involvement of the ERs and modulation of the ERK1/2 dependent pathway by BPA and BPS were investigated.

2. Materials and methods

2.1. Chemicals

Anti-phospho-ERK1/2 (Thr202/Tyr204), anti-ERK1/2, anti-rabbit conjugated with horseradish peroxidase immunoglobulin G (IgG), and U0126 (MEK1,2 inhibitor) were purchased from Cell Signaling Technology, Inc. (Beverly, MA). Roswell Park Memorial Institute RPMI 1640 medium, Fetal Bovine Serum (FBS, EU origin), Charcoal Stripped Fetal Bovine Serum (CS-FBS), glutamine, penicillin/streptomycin, anti β -tubulin, anti-mouse IgG conjugated with horseradish peroxidase, Bisphenol A (BPA), Bisphenol S (BPS), Tamoxifen (TAM) and all other reagents were from Merk Life Science (Milan, Italy). Stock solutions were prepared in DMSO as follows: 0.1 M BPA, 0.1 M BPS, 0.01 M U0126 and 0.01 M TAM.

2.2. Cell culture and treatments

The HTR-8/SVneo human trophoblast cell line was kindly provided by Dr. Charles H. Graham (Queen's University, Ontario, Canada [43]). HTR-8/SVneo cells originated from an explant culture of the human first-trimester placenta; they exhibited the following intrinsic mechanisms: adhesion, migration and invasion [44]. Cells were grown in RPMI 1640 medium supplemented with 10% FBS, 2 mM L-glutamine, and 2% penicillin/streptomycin, and were maintained at 37°C in a standard atmosphere containing 5% CO₂. For the experiments, cells were treated with trypsin and removed from culture flasks, then seeded in 6-well, 12-well or 24-well culture plates at a density of 1.4×10^4 cells/ml, and cultures were incubated overnight. After 24 h culture, cells were exposed to bisphenols (BPA or BPS, dissolved in DMSO) at different concentrations for 3 days. When indicated, cells were incubated with 10 μ M U0126 (ERK1/2 inhibitor) or 0.1 μ M TAM (ER antagonist). Because of the hormonal activity of phenol red and FBS [45], experiments were performed in phenol-red-free RPMI, supplemented with 5% CS-FBS. Dextran treated charcoal is used to selectively remove hormones without nonspecific loss of other serum components. Final DMSO concentration in the medium never exceeded 0.1%. In each experiment, control cells were exposed to the vehicle alone to ascertain absence of effects.

2.3. E-SCREEN assay

The E-screen assay was originally developed to evaluate the proliferative effect of estrogen-like compounds on MCF-7 cells, a human breast cancer cell line, that endogenously overexpresses ERs [46]. Since HTR-8/SVneo cells do possess both nuclear- and membranelocalized- ERs [47], we verified the possibility to use the E-screen assay in the present investigations. Preliminary trials demonstrated that HTR-8/SVneo cells incubated with increasing amounts of 17 β -Estradiol (E2)

are induced to proliferate, starting from the concentration 10^{-12} M, and reaching the maximum response at 10^{-8} M (Supplemental Figure S1). The E-screen assay is then used for the first time on HTR-8/SVneo, with a modification with respect to the original method, i.e. cells are exposed to potential estrogenic compounds for three days instead of five days. In fact, previous observations by our laboratory indicated that HTR-8/SVneo cells replicate at higher rate (~ 22 h) than MCF-7 cells (~ 32 h). Cell proliferation was then determined by a colorimetric assay based on the reduction of a tetrazolium salt, MTT (3(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) by metabolically active cells [48]. Optical density was assessed at 570 nm with background correction at 650 nm, and results are expressed as Proliferative Effect (PE), the ratio of the cell number achieved after BPA or BPS exposure, and the cell number in the control cells.

2.4. Western blotting analysis

Western blotting procedures were carried out as previously reported [49]. After the experimental treatments, cells were lysed in ice-cold lysis buffer containing phosphatase inhibitors (1 mM sodium ortovanadate and 50 mM sodium fluoride) and proteins were electrophoresed, transferred to nitrocellulose membrane and probed with specific antibodies. MAPK activity was assessed by using rabbit anti-phospho-ERK1/2 or anti-total-ERK1/2 as primary antibodies (1:1000) and horseradish-peroxidase-conjugated goat anti-rabbit IgG (1:2000). Immunoblots were developed by enhanced chemiluminescence reagent, and a densitometric analysis of the films was performed by Image Master equipped with TotalLab software ver. 1.0 (Amersham-Pharmacia). Given the inherent variations in film development, densitometry data were normalized to an external reference sample (REF) (unexposed cells) loaded in each immunoblot. MAPK activity was evaluated as phospho/total ERK1/2% of phosphorylation with respect to control. β -tubulin immunodetection was used as the control to assess equal protein loading.

2.5. Transwell cell migration assay

Transwell inserts (PET membrane, 8- μ m pore size, Millicell Hanging, Merck Life Science, Milan, Italy) were used for the cell migration assay, according to manufacturer's instructions. Briefly, after three days of treatment with BPA or BPS, HTR-8/SVneo cells were trypsinized, resuspended in RPMI with 1% FBS and seeded in the upper chamber of each insert at a density of 3×10^4 cells/well; the lower chambers contained 750 μ L RPMI supplemented with 10% FBS. After 24 h of incubation at 37°C the non-migrated cells on the upper surface of the membrane were removed by gentle swabbing. Cells that had reached the lower side of the filter were fixed with methanol for 30 min, stained for 10 min with crystal violet 0.3%, and quantified by visual counting of ten randomly selected fields from each filter under a light microscope. Data were normalized against cell count. Migration capacity was calculated as the percentage of cells that passed through the membranes in the treated- with respect to control- cells.

2.6. Luminex assay

Analysis of inflammatory cytokines and cytokine-induced vascular endothelial growth factor (VEGF) was performed in the medium of trophoblast cell culture, where soluble factors can be released after bisphenols exposure. After the experimental treatments, the medium was collected in 2 mL tubes, centrifuged at $1000 \times g$ for 10 min at 4°C, and supernatant was kept at -80°C until analysis. Secretions of IL-6 (interleukin 6), IL-8 (interleukin 8), IL-1 β (interleukin 1 β), TNF- α (tumor necrosis factor- α), chemokine CCL2 (CC Motif Chemokine Ligand 2), and VEGF were analysed using the immunological kit for Human Magnetic Luminex Screening Assay (Bio-technique s.r.l. Milan, Italy). The analysis was performed using the BioPlex200 equipment (BIO-RAD Laboratories srl, Milan, Italy). Sample concentrations of the different compounds were estimated

through a standard curve using a 5-order polynomial curve and expressed pg/mL (Bio-Plex Manager software 5.0). Data were normalized against cell count.

2.7. Data analysis

Data groups were compared using one-way ANOVA with SigmaPlot software (ver 13, Systat Software Inc.), and followed by Dunnet post-hoc test; a statistical difference was accepted when $P < 0.05$. All experiments were independently repeated at least three times.

3. Results

3.1. BPA and BPS exposure increase HTR-8/SVneo cell proliferation

Cells were exposed to increasing concentrations of BPA or BPS (from 10^{-13} M to 10^{-3} M) for three days. As reported in Figure 1, both BPA and BPS affected the proliferation of HTR-8/SVneo cells. At concentrations ranging from 10^{-11} M to 10^{-7} M BPA or BPS, cell proliferation was significantly increased compared to the control cells. The most active concentration was 10^{-7} M for BPA and 10^{-9} M for BPS, reaching the maximum proliferative effect of 1.89 and 1.62, respectively. BPA at 10^{-5} M concentration significantly increased HTR-8/SVneo cell proliferation, whereas no effect was observed at an equimolar concentration of BPS. At the highest concentration tested (10^{-3} M) of BPA or BPS, the proliferative effect was significantly lower than 1. This result can be ascribed to toxic effects of the compounds on HTR-8/SVneo cells.

3.2. MAPK (ERK1/2) pathway is involved in the stimulatory effects induced by BPA and BPS

The ability of BPA and BPS to modulate the MAPK pathway, and its involvement in the proliferative effect, were investigated by assessing ERK1/2 phosphorylation through western blot analysis. As reported on Figure 2, BPA triggered ERK1/2 phosphorylation in the 10^{-9} – 10^{-5} M

concentration range. A significant increase of ERK1/2 phosphorylation was observed at 10^{-9} M and 10^{-7} M BPS. In the presence of U0126, a selective MAPK inhibitor, BPA and BPS effects on ERK1/2 phosphorylation were significantly reversed.

3.3. BPA but not BPS impairs HTR-8/SVneo cell migration

In vitro effects on the migration capacity of HTR-8/SVneo cells were examined after incubation at different concentrations of BPA or BPS (from 10^{-13} M to 10^{-5} M) for three days. A significant reduction of trophoblast cell migration was observed after treatment with BPA at 10^{-7} M and 10^{-5} M concentrations (-22% and -53% versus control) (Figure 3). BPS exposure did not induce any statistically significant change on HTR-8/SVneo cell migration, at any concentration tested.

3.4. The effects of BPA and BPS on cell proliferation and migration are mediated by ER and MAPK signalling pathways

To reveal the involvement of ER and ERK1/2 signalling pathways in mediating the effects induced by BPA and BPS on HTR-8/SVneo cell proliferation and migration, further experiments were performed in the presence of TAM, a specific antagonist of ER, and U0126. The effects of TAM or U0126 were tested using one of the peak active concentrations of BPA or BPS. Blocking ER with TAM, the effect of both BPA and BPS on cell proliferation (Figure 4A), and the effect of BPA on HTR-8/SVneo cell migration (Figure 4B) were abolished. The proliferative effects induced by BPA and BPS were fully reversed by U0126, which decreased to levels lower than control (Figure 4A). Treatment with U0126 also recovered the migration ability of trophoblast cells, which was inhibited by BPA (Figure 4B).

3.5. BPS but not BPA affects inflammatory cytokines secretion in HTR-8/SVneo cells

Cells were incubated with different concentrations of BPA or BPS (from 10^{-11} M to 10^{-5} M) for three days, then secretion of different inflammatory proteins was measured: IL-6, IL-8, TNF α , IL-1 β , CCL2, and VEGF. Interestingly, IL-6 and IL-8 secretion was affected by BPS exposure (Figure 5A and B), with significant increases at 10^{-11} M that were counteracted by treatment with the U0126 inhibitor. No effect of BPA was detected on interleukin secretion, with respect to the basal levels. Stimulation of HTR8/SVneo cells with BPA or BPS treatment did not affect CCL2 secretion (Figure 5C). In our experimental conditions, basal levels and secretion of IL-1 β TNF- α and VEGF from HTR-8/SVneo cells were always below the sensitivity of the assay.

4. Discussion

BPA and its analog BPS are ubiquitous emerging contaminants of increasing concern. The real-world exposure to BPA and BPS occurs daily at low concentrations, but relevant for human health risk [50]. Emerging evidence suggest high potential of adverse biological effects on reproductive functions after exposure to these chemicals [51–53]. The aim of this work was to elucidate and compare the effects of BPA and BPS on biological activities of HTR-8/SVneo cells, derived from extravillous trophoblast of first-trimester pregnancy. A wide range of concentrations of BPA and BPS (from 10^{-13} M to 10^{-3} M) were used to perform in vitro experiments, which comprises the nM range often documented in human fluids. To the best of our knowledge, these are the first data showing that BPS affects trophoblast functions through ER/MAPK mediated pathways.

Due to the observed effects on humans, particularly those related to hormonal regulation of reproductive processes [54], BPA has been qualified as a xenoestrogen [55]. Based on the chemical similarity to the natural hormone E2, BPA can bind with ERs (ER α and ER β), although displaying 1000 to 2000-fold less affinity with respect to E2 [56]. BPS has been reported to have activity and potency similar to those of BPA [32]. Since the ERs signalling pathways are involved in the cell

proliferation equilibrium, we investigated the possible physiological consequence of BPA and BPS exposure in triggering estrogen-like proliferation on extravillous trophoblast cells. The findings demonstrated that both BPA and BPS increase HTR-8/SVneo cells proliferation, starting from the concentration 10^{-11} M (Supplemental Figure S2), one order of magnitude higher than natural estrogen E2, whose effect on HTR-8/SVneo cells is significant at 10^{-12} M (Supplemental Figure S1). Similar to other EDCs, BPA and BPS displayed non monotonic dose–response curves. More specifically, they showed an inverted U-shaped curve, inducing a significant biological effect at low concentrations, reaching the maximum at 10^{-9} M and 10^{-7} M, and causing cytotoxic responses at the highest doses. The proliferative effect of both BPA and BPS was abolished by the treatment with the ER-blocker TAM, which confirms that both bisphenols mimic the effect of E2. Similar findings, obtained after exposure of cells at different BPA concentrations, were observed on the human choriocarcinoma cell line BeWo [57, 58], JEG-3 cells [59], and human ovarian carcinoma cell line OVCAR3[60]. To the extent of our knowledge, no studies are currently available on BPS effects on reproductive tissues. MAPKs are responsible for converting many cellular and extracellular stimuli into specific responses controlling cell proliferation, differentiation, apoptosis, embryogenesis, and regulation of inflammatory and stress responses [61]. Transduction pathways mediated by phosphorylated ERK1/2 play an essential role in the placenta development [62] and is reported to facilitate trophoblast differentiation [63]. Several data support that BPA-dependent estrogenic activity flows through the ER-mediated extranuclear signals activation, that results in the ERK1/2 phosphorylation [64]. Previous studies have revealed that BPA activates ERK1/2 phosphorylation in human adrenal and breast [65], placenta [66], and also HTR-8/SVneo cells [67]. Present data demonstrate that phosphorylation of ERK1/2 in trophoblast cells is activated by treatment with both BPA and BPS. Since TAM prevented their effect, we concluded

that the ERK1/2 stimulation was induced through the interaction of bisphenols with ERs. Further, ERK1/2 activation is involved in the effect of BPA and BPS on trophoblast proliferation, as the effect was abolished in the presence of the U0126. Overall, BPA and BPS, similarly to E2, activate the ER-dependent signals that culminate with the activation of rapid extra-nuclear pathways in HTR-8/SVneo cells. We conclude that bisphenols effects on trophoblast cells proliferation and migration require the activation of two consequential signals, ERs and ERK1/2 pathways; by blocking one of them the downstream effects are abolished. The present study shows for the first time an ER and MAPK mediated effect of both BPA and BPS on trophoblast cell proliferation. Previous studies reported that BPA did not affect cell proliferation [67–69]. In a further study [70] a concentration-dependent decrease in the proliferation of HTR8/SVneo cells exposed to BPA or BPS was observed. Therefore, additional work is advisable for a better understanding of these modulations. The trophoblast is an embryonic tissue that exerts a crucial role during implantation and placentation, and its migration and invasion capacity represents a pre-requisite for normal embryo implantation.

Numerous studies indicated that exposure to BPA might be associated with severe pregnancy-related complications affecting both mother and fetus [71]. BPA can affect placentation, a significant factor determining pregnancy outcome. It has been associated with implantation failure, miscarriage, premature delivery, and it may also contribute to infertility and subfertility [72, 73]. Few studies about the effect of BPA on the trophoblast migration were performed [67–69, 74], while no information is available yet on the effect of BPS on this process. Present results indicate that only BPA affected cell migration ability. The inhibitory effect of BPA was abolished by TAM, suggesting an estrogen-like effect mediated through ERs. Besides, the migratory ability of cells was recovered by treatment with the inhibitor U0126, demonstrating that ERK1/2 pathways are

relevant for BPA reduction of cell migration. In this study, BPS exposure, unlike BPA, never affected migration of HTR-8/SVneo cells. Similar to cell proliferation, also data on BPA effect on HTR8/SVneo cell migration are conflicting. Some studies [67, 68, 74] observed that migration and invasion were reduced following BPA exposure, while others [69] assessed that BPA enhances cell migration through ERK1/2 stimulation. The present work reports a reduction of migration at micromolar BPA concentrations. The discrepancy among the above studies may be related to the complexity of the migration process modulation. In fact, the control of trophoblast functions is strictly dependent on the balance of various factors [75], which can be influenced by BPA or BPS exposure thus interfering with the intricate role of the trophoblast in human placentation. Among other examples, a very recent investigation reports the important modulation by nutrient transfers, which is affected by BPA [76]. Therefore, further work is advisable to clarify BPA and BPS effects on physiological response of trophoblast cells, possibly performed on ex-vivo preparations thus avoiding possible influence by culture conditions. Previous studies indicated that cytokines may have a critical role in the functions of trophoblast cells; this is particularly true for interleukins [77–79] as key molecules involved in the biological processes related to migration and invasion of trophoblast cells [80]. IL-6 is widely expressed in the endometrium at implantation, and increased levels of placental IL-6 have been associated with preeclampsia, recurrent miscarriage, and infertility [81]. Previous studies provided some evidence that IL-6 enhances migration of HTR8/SVneo cells [82]. In addition, the HTR-8/SVneo cell line secretes IL-8, which is reported to increase cell migration and invasion [83]. However, the critical role of the cytokines on the embryo implantation is not yet understood. Herein, we investigated the effect of BPA and BPS on cytokines secretion by HTR-8/SVneo cells. BPS, only at 10^{-11} M, upregulated interleukins IL-6 and IL-8 secretion. We also assessed that IL-1 β , CCL2, TNF- α and VEGF levels were not affected.

The effect of BPS on IL-6 and IL-8 protein levels was reversed by treatment with U0126. Thus, BPS induction of both IL-6 and IL-8 is mediated by the ERK1/2 signalling. BPA did not cause any effect on cytokines, although a slight but not significant stimulation was observed only at 10^{-11} M BPA on IL-8 secretion. These data are not sufficient to correlate cytokine secretion by HTR-8/SVneo with cell proliferation/migration modulated by BPA or BPS.

5. Conclusion

Our results indicate that BPA, already at sub-nanomolar concentrations, affects both proliferation and migration of HTR-8/SVneo cells through ER/MAPK mediated processes. BPS, at the same concentration range, enhances proliferation, also acting through ER/MAPK pathways. BPS has no effect on HTR-8/SVneo cell migration. Thus, BPA and BPS share estrogen-like activity and modulation of ERK1/2 signaling, while targeting different biological processes in trophoblast cells. Furthermore, BPS while not BPA, increases interleukins secretion through ERK1/2 phosphorylation. The physiological significances of this effect, observed only at a subnanomolar concentration, needs to be further elucidated. In the current state of knowledge, the replacement of BPA with BPS does not seem to ensure the safety of human health.

6. Data availability

The data underlying this article are available in the article and in its online supplementary material. Further information underlying this article will be shared on reasonable request to the corresponding author.

7. Author contributions

MP: running of experiments, graphical representation and writing of first draft. ES:

conceptualization and running of cytokines release measurements; data analysis. EF: conceptualization, supervision, funds, and writing of final draft. PV: theoretical organization of experiments, data analysis and writing of final draft.

8. Conflict of interests

The Authors declare that there are no conflict of interests regarding the publication of this work.

9. References

1. Caserta, D., Maranghi, L., Mantovani, A., Marci, R., Maranghi, F., Moscarini, M. Impact of endocrine disruptor chemicals in gynaecology. *Hum. Reprod. Update* 2007; 14:59–72.
2. Giulivo, M., Lopez de Alda, M., Capri, E., Barceló, D. Human exposure to endocrine disrupting compounds: Their role in reproductive systems, metabolic syndrome and breast cancer. A review. *Environ. Res.* 2016; 151:251–264.
3. Piazza, M.J., Urbanetz, A.A. Environmental toxins and the impact of other endocrine disrupting chemicals in women's reproductive health. *JBRA Assist. Reprod.* 2019; 23:154–164.
4. Brehm, E., Flaws, J.A. Transgenerational Effects of Endocrine-Disrupting Chemicals on Male and Female Reproduction. *Endocrinology* 2019; 160:1421–1435.
5. Yang, C., Song, G., Lim, W. A mechanism for the effect of endocrine disrupting chemicals on placentation. *Chemosphere* 2019; 231:326–336.
6. Dodds EC, Lawson W. Synthetic strogenic Agents without the Phenanthrene Nucleus. *Nature* 1936; 137:996.
7. Björnsdotter, M.K., de Boer, J., Ballesteros-Gómez, A. Bisphenol A and replacements in thermal paper: A Review. *Chemosphere* 2017; 182:691-706.
8. Gunaalan, K., Fabbri, E., Capolupo, M. The hidden threat of plastic leachates: A critical review on their impacts on aquatic organisms. *Water. Res.* 2020; 184:116170.

9. Lee, J., Choi, K., Park, J., Moon, H.-B., Choi, G., Lee, J.J., Suh, E., Kim, H.-J., Eun, S.-H., Kim, G.-H., Cho, G.J., Kim, S.K., et al. Bisphenol A distribution in serum, urine, placenta, breast milk, and umbilical cord serum in a birth panel of mother–neonate pairs. *Sci. Total Environ.* 2018; 626:1494–1501.
10. Sowlat, M.H., Lotfi, S., Yunesian, M., Ahmadkhaniha, R., Rastkari, N. The association between bisphenol A exposure and type-2 diabetes: a world systematic review. *Environ. Sci. Pollut. Res.* 2016; 23:21125–21140.
11. Vom Saal, F.S., Nagel, S.C., Coe, B.L., Angle, B.M., Taylor, J.A. The estrogenic endocrine disrupting chemical bisphenol A (BPA) and obesity. *Mol. Cell. Endocrinol.* 2012; 354:74–84.
12. Leonardi, A., Cofini, M., Rigante, D., Lucchetti, L., Cipolla, C., Penta, L., Esposito, S. The Effect of Bisphenol A on Puberty: A Critical Review of the Medical Literature. *Int. J. Environ. Res. Public Health* 2017; 14:9-1044.
13. Olsén, L., Lind, L., Lind, P.M. Associations between circulating levels of bisphenol A and phthalate metabolites and coronary risk in the elderly. *Ecotoxicol. Environ. Saf.* 2012; 80:179–183.
14. Dumitrascu, M.C., Mares, C., Petca, R.-C., Sandru, F., Popescu, R.-I., Mehedintu, C., Petca, A. Carcinogenic effects of bisphenol A in breast and ovarian cancers. *Oncol. Lett.* 2020; 20:6.
15. Zhang, X., Liu, R. Advances in BPA-induced Oxidative Stress and Related Effects and Mechanisms in Liver, 1991-2017. *Mini Rev. Med. Chem.* 2020; 20:432–443.
16. Vandenberg, L.N., Hauser, R., Marcus, M., Olea, N., Welshons, W.V. Human exposure to bisphenol A (BPA). *Reprod. Toxicol.* 2007; 24:139–177.
17. Jalal, N., Surendranath, A.R., Pathak, J.L., Yu, S., Chung, C.Y. Bisphenol A (BPA) the mighty and the mutagenic. *Toxicol. Rep.* 2018; 5:76–84.
18. Zbucka-Krętowska, M., Łazarek, U., Milyk, W., Sidorkiewicz, I., Pierzyński, P., Milewski, R., Wołczyński, S., Czerniecki, J. Simultaneous analysis of bisphenol A fractions in maternal and fetal compartments in early second trimester of pregnancy. *J. Perinat. Med.* 2019; 47:765–770.

375 19. Corbel, T., Gayrard, V., Puel, S., Lacroix, M.Z., Berrebi, A., Gil, S., Viguié, C., Toutain, P.-
376 L., Picard-Hagen, N. Bidirectional placental transfer of Bisphenol A and its main metabolite,
377 Bisphenol A-Glucuronide, in the isolated perfused human placenta. *Reprod. Toxicol.* 2014; 47:51–
378 58.

379 20. Liao, C., Liu, F., Kannan, K. Bisphenol S, a New Bisphenol Analogue, in Paper Products and
380 Currency Bills and Its Association with Bisphenol A Residues. *Environ. Sci. Technol.* 2012b;
381 46:6515–6522.

382 21. Usman, A., Ahmad, M. From BPA to its analogues: Is it a safe journey? *Chemosphere* 2016;
383 158:131–142.

384 22. Wu, L.-H., Zhang, X.-M., Wang, F., Gao, C.-J., Chen, D., Palumbo, J.R., Guo, Y., Zeng, E.Y.
385 Occurrence of bisphenol S in the environment and implications for human exposure: A short
386 review. *Sci. Total Environ.* 2018; 615:87–98.

387 23. Liao, C., Liu, F., Alomirah, H., Loi, V.D., Mohd, M.A., Moon, H.-B., Nakata, H., Kannan, K.
388 Bisphenol S in Urine from the United States and Seven Asian Countries: Occurrence and Human
389 Exposures. *Environ. Sci. Technol.* 2012a; 46:6860–6866.

390 24. Siracusa, J.S., Yin, L., Measel, E., Liang, S., Yu, X. Effects of bisphenol A and its analogs on
391 reproductive health: A mini review. *Reprod. Toxicol.* 2018; 79:96–123.

392 25. Liu, J., Li, J., Wu, Y., Zhao, Y., Luo, F., Li, S., Yang, L., Moez, E.K., Dinu, I., Martin, J.W.
393 Bisphenol A Metabolites and Bisphenol S in Paired Maternal and Cord Serum. *Environ. Sci.*
394 *Technol.* 2017; 51:2456–2463.

395 26. Santoro, A., Chianese, R., Troisi, J., Richards, S., Nori, S.L., Fasano, S., Guida, M., Plunk, E.,
396 Viggiano, A., Pierantoni, R., Meccariello, R. Neuro-toxic and Reproductive Effects of BPA. *Curr.*
397 *Neuropharmacol.* 2019; 17:1109–1132.

398 27. Grandin, F.C., Lacroix, M.Z., Gayrard, V., Viguié, C., Mila, H., De Place, A., Vayssière, C.,
399 Morin, M., Corbett, J., Gayrard, C., Gely, C.A., Toutain, P.-L., et al. Is bisphenol S a safer
400 alternative to bisphenol A in terms of potential fetal exposure? Placental transfer across the
401 perfused human placenta. *Chemosphere* 2019; 221:471–478.

28. Lehmler, H.-J., Liu, B., Gadogbe, M., Bao, W. Exposure to Bisphenol A, Bisphenol F, and Bisphenol S in U.S. Adults and Children: The National Health and Nutrition Examination Survey 2013–2014. *ACS Omega* 2018; 3:6523–6532.
29. Huang, S., Li, J., Xu, S., Zhao, H., Li, Y., Zhou, Y., Fang, J., Liao, J., Cai, Z., Xia, W. Bisphenol A and bisphenol S exposures during pregnancy and gestational age – A longitudinal study in China. *Chemosphere* 2019; 237:124426.
30. Žalmanová, T., Hošková, K., Nevoral, J., Prokešová, Š., Zámotná, K., Kott, T., Petr, J. Bisphenol S instead of bisphenol A: a story of reproductive disruption by regrettable substitution – a review. *Czech J. Anim. Sci.* 2016; 18:433-449.
31. Rochester, J.R., Bolden, A.L. Bisphenol S and F: A Systematic Review and Comparison of the Hormonal Activity of Bisphenol A Substitutes. *Environ. Health Perspect.* 2015; 123:643–650.
32. Grignard, E., Lapenna, S., Bremer, S. Weak estrogenic transcriptional activities of Bisphenol A and Bisphenol S. *Toxicol. In Vitro* 2012; 26:727–731.
33. Molina-Molina, J.-M., Amaya, E., Grimaldi, M., Sáenz, J.-M., Real, M., Fernández, M.F., Balaguer, P., Olea, N. In vitro study on the agonistic and antagonistic activities of bisphenol-S and other bisphenol-A congeners and derivatives via nuclear receptors. *Toxicol. Appl. Pharmacol.* 2013; 272:127-138.
34. Carter, A.M., Enders, A.C., Pijnenborg, R. The role of invasive trophoblast in implantation and placentation of primates. *Phil. Trans. R. Soc. B.* 2015; 370:1663.
35. Chen, J.Z.-J., Sheehan, P.M., Brennecke, S.P., Keogh, R.J. Vessel remodelling, pregnancy hormones and extravillous trophoblast function. *Mol. Cell. Endocrinol.* 2012; 349:138–144.
36. Turco, M.Y., Moffett, A. Development of the human placenta. *Development* 2019; 146:22.
37. Hemberger, M., Hanna, C., Dean, W. Mechanisms of early placental development in mouse and humans. *Nat. Rev. Genet.* 2020; 17:27-43.
38. Mebratu, Y., Tesfaigzi, Y. How ERK1/2 activation controls cell proliferation and cell death: Is subcellular localization the answer? *Cell Cycle* 2009; 8:1168–1175.

428 39. Vaillancourt, C., Lanoix, D., Le Bellego, F., Daoud, G., Lafond, J. Involvement of MAPK
429 Signalling in Human Villous Trophoblast Differentiation. *Mini Rev. Med. Chem.* 2009; 9:962–
430 973.

431 40. García-Velasco, J.A., Arici, A. Chemokines and human reproduction. *Fertil. Steril.* 1999;
432 71:983–993.

433 41. Sokolov, D.I., Furaeva, K.N., Stepanova, O.I., Sel'kov, S.A. Proliferative and Migration
434 Activity of JEG-3 Trophoblast Cell Line in the Presence of Cytokines. *Bull. Exp. Biol. Med.* 2015;
435 159:550–556.

436 42. Msheik, H., Azar, J., El Sabeh, M., Abou-Kheir, W., Daoud, G. HTR-8/SVneo: A model for
437 epithelial to mesenchymal transition in the human placenta. *Placenta* 2020; 90:90–97.

438 43. Graham C. H., Hwley T. S., Hawlwy R. G., MacDougall J. R., Kerbel R. S., Khoo N., Lala P.
439 K. Establishment and characterization of first trimester human trophoblast cells with extended
440 lifespan. *Exp. Cell. Res.* 1993; 206:204-211.

441 44. Hannan, N.J., Paiva, P., Dimitriadis, E., Salamonsen, L.A. Models for Study of Human Embryo
442 Implantation: Choice of Cell Lines?. *Biol. Reprod.* 2010; 82:235–245.

443 45. Welshons, W.V., Wolf, M.F., Murphy, C.S., Jordan, V.C. Estrogenic activity of phenol red.
444 *Mol. Cell. Endocrinol.* 1988; 57:169–178.

445 46. Soto, A.M., Sonnenschein, C., Chung, K.L., Olea, N., Serrano, F.O. The E-SCREEN assay as
446 a tool to identify estrogens: an update on estrogenic environmental pollutants. *Environ. Health*
447 *Perspect.* 1995; 103:113-122.

448 47. Patel, S., Kilburn, B., Imudia, A., Armant, D.R., Skafar, D.F. Estradiol Elicits Proapoptotic
449 and Antiproliferative Effects in Human Trophoblast Cells. *Biol. Reprod.* 2015; 93:74.

450 48. Mosmann, T. Rapid colorimetric assay for cellular growth and survival: Application to
451 proliferation and cytotoxicity assays. *J. Immunol. Methods* 1983; 65:55–63.

452 49. Valbonesi, P., Ricci, L., Franzellitti, S., Biondi, C., Fabbri, E. Effects of Cadmium on MAPK
453 Signalling Pathways and HSP70 Expression in a Human Trophoblast Cell Line. *Placenta* 2008;
454 29:725–733.

50. Nevoral, J., Kolinko, Y., Moravec, J., Žalmanová, T., Hošková, K., Prokešová, Š., Klein, P., Ghaibour, K., Hošek, P., Štiavnická, M., Řimnáčová, H., Tonar, Z., Petr, J., Králíčková, M. Long-term exposure to very low doses of bisphenol S affects female reproduction. *Reproduction* 2018; 156:47–57.
51. Rochester, J.R. Bisphenol A and human health: A review of the literature. *Reprod. Toxicol.* 2013; 42:132–155.
52. Pergialiotis, V., Kotrogianni, P., Christopoulos-Timogiannakis, E., Koutaki, D., Daskalakis, G., Papantoniou, N. Bisphenol A and adverse pregnancy outcomes: a systematic review of the literature. *J. Matern. Fetal Neonatal Med.* 2018; 31:3320–3327.
53. Catenza, C.J., Farooq, A., Shubear, N.S., Donkor, K.K. A targeted review on fate, occurrence, risk and health implications of bisphenol analogues. *Chemosphere* 2021; 268:129273.
54. Tomza-Marciniak, A., Stępkowska, P., Kuba, J., Pilarczyk, B. Effect of bisphenol A on reproductive processes: A review of in vitro, in vivo and epidemiological studies. *J. App. Toxicol.* 2018; 38:51–80.
55. Mhaouty-Kodja, S., Belzunces, L.P., Canivenc, M.-C., Schroeder, H., Chevrier, C., Pasquier, E. Impairment of learning and memory performances induced by BPA: Evidences from the literature of a MoA mediated through an ED. *Mol. Cell. Endocrinol.* 2018; 475:54–73.
56. Le Magueresse-Battistoni, B., Multigner, L., Beausoleil, C., Rousselle, C. Effects of bisphenol A on metabolism and evidences of a mode of action mediated through endocrine disruption. *Mol. Cell. Endocrinol.* 2018; 475:74–91.
57. Ponniah, M., Billett, E.E., De Girolamo, L.A. Bisphenol A Increases BeWo Trophoblast Survival in Stress-Induced Paradigms through Regulation of Oxidative Stress and Apoptosis. *Chem. Res. Toxicol.* 2015; 28:1693–1703.
58. Wang, Z.-Y., Lu, J., Zhang, Y.-Z., Zhang, M., Liu, T., Qu, X.-L. Effect of Bisphenol A on invasion ability of human trophoblastic cell line BeWo. *Int. J. Clin. Exp. Pathol.* 2015; 8:14355-14364.

59. Mannelli, C., Szóstek, A.Z., Lukasik, K., Carotenuto, C., Ietta, F., Romagnoli, R., Ferretti, C.,
 Paulesu, L., Wołczynski, S., Skarzynski, D.J. Bisphenol A modulates receptivity and secretory
 function of human decidual cells: an in vitro study. *Reproduction* 2015; 150:115–125.

60. Ptak, A., Hoffmann, M., Gruca, I., Barć, J. Bisphenol A induce ovarian cancer cell migration
 via the MAPK and PI3K/Akt signalling pathways. *Toxicol. Lett.* 2014; 229:357–365.

61. Lavoie, H., Gagnon, J., Therrien, M. ERK signalling: a master regulator of cell behaviour, life
 and fate. *Nat. Rev. Mol. Cell. Biol.* 2020; 21:607–632.

62. Hatano, N., Mori, Y., Oh-hora, M., Kosugi, A., Fujikawa, T., Nakai, N., Niwa, H., Miyazaki,
 J., Hamaoka, T., Ogata, M. Essential role for ERK2 mitogen-activated protein kinase in placental
 development. *Genes. Cells* 2003; 8:847–856.

63. Daoud, G., Amyot, M., Rassart, É., Masse, A., Simoneau, L., Lafond, J. ERK1/2 and p38
 regulate trophoblasts differentiation in human term placenta. *J. Physiol.* 2005; 566:409–423.

64. Acconcia, F., Pallottini, V., Marino, M. Molecular Mechanisms of Action of BPA. *Dose-
 Response* 2015; 13:4.

65. Song, H., Zhang, T., Yang, P., Li, M., Yang, Y., Wang, Y., Du, J., Pan, K., Zhang, K. Low
 doses of bisphenol A stimulate the proliferation of breast cancer cells via ERK1/2/ERR γ signals.
Toxicol. in Vitro 2015; 30:521–528.

66. Chu, P.-W., Yang, Z.-J., Huang, H.-H., Chang, A.-A., Cheng, Y.-C., Wu, G.-J., Lan, H.-C.
 Low-dose bisphenol A activates the ERK signaling pathway and attenuates steroidogenic gene
 expression in human placental cells. *Biol. Reprod.* 2018; 98:250–258.

67. Li, X., Wang, Y., Wei, P., Shi, D., Wen, S., Wu, F., Liu, L., Ye, N., Zhou, H. Bisphenol A
 affects trophoblast invasion by inhibiting CXCL8 expression in decidual stromal cells. *Mol. Cell.
 Endocrinol.* 2018; 470:1–10.

68. Spagnoletti, A., Paulesu, L., Mannelli, C., Ermini, L., Romagnoli, R., Cintorino, M., Ietta, F.
 Low concentrations of Bisphenol A and para-Nonylphenol affect extravillous pathway of human
 trophoblast cells. *Mol. Cell. Endocrinol.* 2015; 412:56–64.

507 69. Lan, X., Fu, L.-J., Zhang, J., Liu, X.-Q., Zhang, H.-J., Zhang, X., Ma, M.-F., Chen, X.-M., He,
508 J.-L., Li, L.-B., Wang, Y.-X., Ding, Y.-B. Bisphenol A exposure promotes HTR-8/SVneo cell
509 migration and impairs mouse placentation involving upregulation of integrin- β 1 and MMP-9 and
510 stimulation of MAPK and PI3K signaling pathways. *Oncotarget* 2017; 8:51507–51521.

511 70. Basak, S., Srinivas, V., Duttaroy, A.K. Bisphenol-A impairs cellular function and alters DNA
512 methylation of stress pathway genes in first trimester trophoblast cells. *Reprod. Toxicol.* 2018;
513 82:72–79.

514 71. Filardi, T., Panimolle, F., Lenzi, A., Morano, S. Bisphenol A and Phthalates in Diet: An
515 Emerging Link with Pregnancy Complications. *Nutrients* 2020; 12:2-525.

516 72. Konieczna, A., Rutkowska, A., Rachoń, D. Health risk of exposure to Bisphenol A (BPA).
517 *Rocz Państ Zakł Hig* 2015; 66:5–11.

518 73. Tang, Z.-R., Xu, X.-L., Deng, S.-L., Lian, Z.-X., Yu, K. Oestrogenic Endocrine Disruptors in
519 the Placenta and the Fetus. *Int. J. Mol. Sci.* 2020; 21:1519.

520 74. Wei, P., Ru, D., Li, X., Shi, D., Zhang, M., Xu, Q., Zhou, H., Wen, S. Exposure to
521 environmental bisphenol A inhibits HTR-8/SVneo cell migration and invasion. *J. Biomed. Res.*
522 2020; 34:369–378.

523 75. Lunghi, L., Ferretti, M.E., Medici, S., Biondi, C., Vesce, F. Control of human trophoblast
524 function. *Reprod. Biol. Endocrinol.* 2007; 5:6.

525 76. Benincasa, L., Mandalà, M., Paulesu, L., Barberio, L., Ietta, F. Prenatal Nutrition Containing
526 Bisphenol A affects Placenta Glucose Transfer: Evidence in Rats and Human Trophoblast.
527 *Nutrients* 2020; 12:5.

528 77. Red-Horse, K., Drake, P.M., Fisher, S.J. Human pregnancy: the role of chemokine networks
529 at the fetal maternal interface. *Expert Rev. Mol. Med.* 2004; 6:1-14.

530 78. Zhang, H., Hou, L., Li, C.M., Zhang, W.Y. The chemokine CXCL6 restricts human trophoblast
531 cell migration and invasion by suppressing MMP-2 activity in the first trimester. *Hum. Reprod.*
532 2013; 28:2350–2362.

- 533 79. Du, M.-R., Wang, S.-C., Li, D.-J. The integrative roles of chemokines at the maternal–fetal
534 interface in early pregnancy. *Cell. Mol. Immunol.* 2014; 11:438–448.
- 535 80. Sharma, S., Godbole, G., Modi, D. Decidual Control of Trophoblast Invasion. *Am. J. Reprod.*
536 *Immunol.* 2016; 75:341–350.
- 537 81. Champion, H., Innes, B.A., Robson, S.C., Lash, G.E., Bulmer, J.N. Effects of interleukin-6 on
538 extravillous trophoblast invasion in early human pregnancy. *Mol. Hum. Reprod.* 2012; 18:391–
539 400.
- 540 82. Jovanović, M., Stefanoska, I., Radojčić, L., Vićovac, L. Interleukin-8 (CXCL8) stimulates
541 trophoblast cell migration and invasion by increasing levels of matrix metalloproteinase (MMP)2
542 and MMP9 and integrins $\alpha 5$ and $\beta 1$. *Reproduction* 2010; 139:789–798.
- 543 83. Jovanović, M., Vićovac, L. Interleukin-6 Stimulates Cell Migration, Invasion and Integrin
544 Expression in HTR-8/SVneo Cell Line. *Placenta* 2009; 30:320–328.

Figures




Figure 1: Effect of BPA or BPS on HTR-8/SVneo cell proliferation. Data are expressed as PE and reported as means \pm SE of 5 different experiments, each performed in quadruplicate. *p < 0.05, **p < 0.01 related to control (C, control cells exposed to vehicle alone, PE=1).





Figure 2: Effect of BPA or BPS on ERK1/2 phosphorylation in HTR-8/SVneo cells. [A] Representative immunoblots relative to phosphorylated and total ERK1/2. U0126= BPA or BPS 10^{-7} M treatment with the ERK1/2 inhibitor U0126. REF= external reference sample, loaded in each immunoblot. β -tubulin levels were also measured to assess equal loading of proteins. [B] Densitometric analysis of ERK1/2 phosphorylation, evaluated as the ratio between the phosphorylated and total forms, expressed as % of control (C) and reported as means \pm SE of 4 different experiments. * $p < 0.05$ related to C (control cells exposed to vehicle alone), # $p < 0.05$ related to 10^{-7} M BPA or BPS.




Figure 3: Effect of BPA or BPS on HTR-8/SVneo cell migration. [A] Densitometric data relative to cell migration, evaluated as the number of cells reaching the lower side of the membrane in Transwell inserts. [B] Representative light microscope images of Transwell migration assay (10 \times). Data are expressed as a percentage of the control (C) and reported as means \pm SE of 4 different experiments. * $p < 0.05$, ** $p < 0.01$ related to C (control cells exposed to vehicle alone).




Figure 4: Effect of Tamoxifen and U0126 on HTR-8/SVneo cells exposed to BPA or BPS. [A] Effect of 10^{-9} M BPA or BPS with or without TAM or U0126 on cell proliferation. [B] Effect of 10^{-7} M BPA or BPS with or without TAM or U0126 on cell migration. Data are reported as means \pm SE of 4 different experiments. * $p < 0.05$, ** $p < 0.01$ related to C (control cells exposed to vehicle alone).




Figure 5: Effect of BPA or BPS on IL-6, IL-8, and CCL2 secretion by HTR-8/SVneo cells. [A] IL-6 secretion (C=23.86 pg/ml); [B] IL-8 secretion (C=27.70 pg/ml); [C] CCL2 secretion (C=11.72 pg/ml). Data are expressed as % of control (C) and reported as means \pm SE of 3 different experiments. * $p < 0.05$ related to C (control cells exposed to vehicle alone).

