

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Natural distemper infection in stone martens (*Martes foina*): From infection to neutralizing antibodies

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Balboni A, S.F. (2021). Natural distemper infection in stone martens (*Martes foina*): From infection to neutralizing antibodies. RESEARCH IN VETERINARY SCIENCE, 138, 196-200 [10.1016/j.rvsc.2021.06.015].

Availability:

This version is available at: <https://hdl.handle.net/11585/857093> since: 2022-02-12

Published:

DOI: <http://doi.org/10.1016/j.rvsc.2021.06.015>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

This is the final peer-reviewed accepted manuscript of:

Balboni A, Savini F, Scagliarini A, Berti E, Naldi M, Urbani L, Fontana MC, Carra E, Gibelli LRM, Gobbo F, Bologna E, Zambelli D, Ceccherelli R, Battilani M. Natural distemper infection in stone martens (*Martes foina*): From infection to neutralizing antibodies. Research in Veterinary Science. 2021, 138: 196-200

The final published version is available online at:
<https://doi.org/10.1016/j.rvsc.2021.06.015>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

20 **Short Communication**

21

22 **Natural distemper infection in stone martens (*Martes foina*): from infection to neutralizing**
23 **antibodies**

24

25 Andrea Balboni ^{a*}, Federica Savini ^a, Alessandra Scagliarini ^b, Elisa Berti ^c, Marzia Naldi ^c,
26 Lorenza Urbani ^a, Maria Cristina Fontana ^d, Elena Carra ^d, Lucia Rita Maria Gibelli ^d, Federica
27 Gobbo ^e, Emanuela Bologna ^c, Donatella Zambelli ^c, Renato Ceccherelli ^c, Mara Battilani ^a.

28

29 ^a Department of Veterinary Medical Sciences, Alma Mater Studiorum-University of Bologna, Via
30 Tolara di Sopra 50, Ozzano Emilia (BO) 40064, Italy.

31 ^b Department of Experimental, Diagnostic and Specialty Medicine - Alma Mater Studiorum-
32 University of Bologna, Via Massarenti 9, Bologna 40138, Italy.

33 ^c Centro Tutela e Ricerca Fauna Esotica e Selvatica Monte Adone ODV, Via Brento 9, Sasso
34 Marconi (BO) 40037, Italy.

35 ^d Istituto Zooprofilattico Sperimentale della Lombardia ed Emilia Romagna “B. Ubertini”
36 (IZSLER), Via Antonio Bianchi 9, Brescia 25124, Italy.

37 ^e Istituto Zooprofilattico Sperimentale delle Venezie (IZSVe), viale Dell'Università 10, Legnaro
38 (PD) 35020, Italy.

39

40 Email addresses: Federica Savini (federica.savini3@unibo.it), Alessandra Scagliarini
41 (alessand.scagliarini@unibo.it), Elisa Berti (elisa@centrotutelafauna.org), Marzia Naldi
42 (marzia.naldi@virgilio.it), Lorenza Urbani (lorenza.urbani2@unibo.it), Maria Cristina Fontana
43 (mariacristina.fontana@izsler.it), Elena Carra (elena.carra@izsler.it), Lucia R.M. Gibelli
44 (luciarita.gibelli@izsler.it), Federica Gobbo (fgobbo@izsvenezie.it), Emanuela Bologna

45 (emanuelabo@gmx.com), Donatella Zambelli (zambellidonatella@icloud.com), Renato Ceccherelli
46 (apusvet.cruma@libero.it), Mara Battilani (mara.battilani@unibo.it).

47

48 * Corresponding author:

49 Andrea Balboni

50 Department of Veterinary Medical Sciences, Alma Mater Studiorum-University of Bologna,

51 Via Tolara di Sopra 50, Ozzano Emilia (BO) 40064, Italy.

52 Tel.: +39 051 2097083

53 E-mail address: a.balboni@unibo.it

54 ORCID: orcid.org/0000-0002-8049-6645

55

56 **Declaration of Competing Interest**

57 Declarations of interest: none.

58 **Abstract**

59 We report an outbreak of canine distemper virus (CDV) among stone martens (*Martes foina*) in
60 Italy. After being rescued in Northern Italy between April and June 2018, six subjects were kept in
61 a wildlife and exotic animal rescue center in Bologna province. Subjects have been monitored for
62 15 months in captivity. Within this time-lapse, two subjects died, while among the remaining four,
63 only one showed clinical symptoms referable to distemper. Surviving subjects have been regularly
64 tested for CDV by means of reverse transcriptase-PCR from conjunctival and oropharyngeal swabs
65 for eleven months. The identified viruses belonged to the Wildlife-Europe CDV genetic subgroup.
66 Neutralizing antibodies were detected at the end of the eleven months, when all subjects tested
67 reverse transcriptase-PCR negative. Our findings confirm the circulation of the Wildlife-Europe
68 CDV genetic subgroup (Europe 1/South America 1 lineage) within the Italian wildlife, and improve
69 knowledge on viral infection in stone martens.

70

71 **Keywords:** distemper; phylogenetic analysis; serology; stone marten; wildlife

Canine distemper is distributed worldwide and constitutes one of the most serious contagious diseases of carnivores, fluctuating in disease presentation (Greene and Vandeveld, 2012) and causing high lethality rate. Mustelids are a diverse group of carnivorous mammals known to be very susceptible to distemper (Philippa et al., 2008), displaying clinical signs similar to those in canids including anorexia, oculo-nasal mucopurulent discharge, multifocal dermal hyperkeratosis, emaciation, severe pruritus, intermittent diarrhoea, pneumonia, and rapid death (Mos et al., 2003; Williams et al., 1988). The stone marten (*Martes foina*) is widely distributed in Eurasia (IUCN, 2015), where is considered as possible canine distemper virus (CDV) wildlife reservoir (Myers et al., 1997; Philippa et al., 2008). Nevertheless, no information is available on spontaneous infection within this species. We report an outbreak of canine distemper in six stone martens held in Monte Adone wildlife rescue center in Bologna province.

This was a retrospective study; no stone martens were sampled exclusively for the purposes of this study. Only samples taken to diagnose the aetiology of the clinical disease and to monitor viral shedding and immunity before reintroduction into the wild, were used. Subjects were rescued in Northern Italy in 2018: two (#110A, #110B) at the end of April, three at the beginning of June (#246A, #246B, #246C) and one (#298) at the end of the same month (**Appendix A: Supplementary material**). All the stone martens were young, found in a state of apparent abandonment in anthropised areas (near human dwellings), and entered the structure in good general conditions, with the exception of #298, a male that showed mild sensory depression, which led to suspect to a car collision. Nonetheless, the subject fully recovered in five days. At the entrance of the subjects in the facility, an observation period (ranging from 2 to 30 days) was carried out in a separate isolation area before moving them to a fenced outdoor common area. In July 2018, on day 21 from rescue, subject #246A (female) showed hyperthermia and violent convulsions and died after 19 days (day 40 from rescue). One week later, subject #110A (female) was found dead, without showing any pathological sign in the previous days. On both subjects #246A and #110A, a post mortem CDV one-step real-time reverse transcription-PCR (RT-qPCR)

98 was performed on the RNA extracted from foot pads (**Appendix A: Supplementary material,**
99 Scagliarini et al., 2007) sampled in July 2018. Histological analysis was performed only on subject
100 #246A. In addition, qualitative two-step reverse transcription-PCR (RT-PCR) was performed to
101 amplify the hemagglutinin (H) gene of CDV identified in animals #246A and #110A (**Appendix A:**
102 **Supplementary material,** Demeter et al., 2007); these were subsequently sequenced, analysed by
103 BLAST (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) and aligned with 116 CDV reference strains
104 (**Appendix A: Supplementary material**) retrieved from GenBank
105 (<https://www.ncbi.nlm.nih.gov/genbank/>). Phylogeny was reconstructed by MEGA X version
106 10.1.7 (Kumar et al., 2018).

107 Ten days after death of subject #110A, at day 43 from its rescue, subject #298 showed visual
108 impairment, ocular nasal discharge, as well as lip ulcers and was quarantined. No symptoms were
109 reported during the following period for the remaining three subjects.

110 The four surviving animals were quarantined and viral RNA shedding monitored to identify when
111 they were no longer potentially infectious. Starting from the end of August 2018, all subjects were
112 tested seven times within nine months for CDV RNA by one-step RT-PCR from conjunctival and
113 oropharyngeal swabs (**Appendix A: Supplementary material,** Frisk et al., 1999). At the end of the
114 isolation period (June 2019), presence of neutralizing antibodies was also investigated to assess
115 whether animals were protected from reinfection before reintroduction into the wild (**Appendix A:**
116 **Supplementary material,** Confer et al., 1975).

117 Histology on entrails from subject #246A revealed the presence of both enteritis with de-
118 epithelialized, shortened and fused intestinal villi as well as interstitial pneumonia. Post-mortem
119 RT-qPCR for CDV from foot pads of subjects #246A and #110A tested positive with a quantity of
120 17 and 2.8×10^3 viral RNA copies/mg of tissue, respectively. Analyses performed on complete
121 CDV H gene nucleotide sequences (1824 bp) obtained from #246 and #110A (GenBank ID
122 MW015089 and MW015090) showed a unique non-synonymous nucleotide mutation in position
123 268 (corresponding to amino acid residue 90) that distinguished them with an overall nucleotide

identity >99,8% to the closest strain available in GenBank (ID MN044701). This strain was identified in a fox in North-Eastern Italy in 2018 (Bianco et al., 2020). Phylogeny revealed that the infection was caused by viruses belonging to the Wildlife-Europe CDV genetic subgroup, Europe 1/South America 1 lineage (Bianco et al., 2020; Panzera et al., 2015) (**Figure 1**), closely related to other CDV recently identified in wildlife and dog.

In **Table 1** the results of RT-PCR on oropharyngeal and conjunctival swabs sampled from survived animals are reported. Stone martens #298 and #110B tested positive until the third month (October 2018), while the remaining two subjects #246B and #246C tested positive until the fifth month (December 2018). Virus neutralization (VN) test revealed positive antibody titres of: 1:32 for subject #246C, 1:64 for both #110B and #298, and 1:128 for #246B.

Here we report a natural outbreak of canine distemper in six stone martens that showed different courses: two of them died, one showed mild signs while the remaining three displayed no symptoms. For all the animals the exact time of exposure is unknown. Nevertheless, it can be speculated that subject #246A might have been the source of infection, entering the structure during an incubation phase of the disease and developing convulsive crisis in July 2018, after 21 days, and dying 19 days later, in line with the data described in ferrets (Perpiñán et al., 2008). Furthermore, a concurrent subclinical infection of subject #298 cannot be excluded. Indeed, the mental dullness presented by the subject, initially ascribed to a car collision, might have been caused by the transient fever that occurs within the first days post CDV infection. Subsequent spread of the virus in the organism would have produced nasal and ocular discharge and lip ulcers 38 days later. As a matter of fact, asymptomatic carriers have been documented among other mustelids such as minks (Zhao et al., 2015). Also in dogs, it is estimated that at least 50% of CDV infections are subclinical (Peterson and Kutzler, 2011). The intermittent but protracted detection of the viral RNA (attributable to the sampling technique, the storing procedure between sampling and testing and the RT-PCR assay sensitivity) suggests that viral RNA excretion long lasted in all subjects. More specifically, in subjects #298 and #110B the viral excretion lasted at least three months while in the

150 remaining two subjects at least five months. This persistence recalls the pathogenesis described in
151 dogs, where shedding of virus can continue for as long as three to four months (Sykes, 2014).
152 Animals shed the virus in all secretions and since CDV is known for not persisting long in the
153 environment, mildly affected and recovering infected animals may play an important role in
154 maintaining transmission cycles in shelters (Miller and Hurley, 2009). This is particularly important
155 within rescue centres, housing animals of unknown medical histories and endangered wildlife. In
156 fact, CDV is considered of growing concern due to its propensity for host-switching and emergence
157 in new species, including endangered species (Feng et al., 2016; Sulikhan et al., 2018)..
158 Clinical manifestations and histological results confirmed that the gastrointestinal tract, the
159 respiratory and nervous systems were the most affected in stone martens, as already documented in
160 other mustelids (Perpiñán et al., 2008).

161 Morbilliviruses induce long-term immunity, in particular, immune response towards CDV infection
162 is considered lifelong (Schultz et al., 2010), thus detection of seropositive animals implies
163 protection against the virus. Sero-conversion of mink, in experimental conditions, has been
164 described to take place at 14 days post infection (dpi) reaching the highest titres at 21 dpi (Zhao et
165 al., 2015). Since we did not test subjects so early we cannot compare our results, nevertheless, it is
166 assumable that, as described in dogs, clinical signs resolve and the virus is cleared from most body
167 tissues when antibody titre increases (Greene and Vandeveld, 2012). Indeed, the neutralization test
168 is still considered the gold standard for measuring protection against infection, and antibody titres
169 correlate with the level of protection (Appel and Robson, 1973).

170 The identified viruses belonged to the Wildlife-Europe CDV genetic subgroup (Europe 1/South
171 America 1 lineage). This wildlife well-adapted CDV genetic group has been detected in Europe
172 (Monne et al., 2011). In Italy, it also caused the first spillover event, from wildlife to a non-
173 vaccinated domestic dog (Bianco et al., 2020). Coevolution and phylogenetic clustering of CDV
174 identified in both wildlife and domestic dogs have previously been reported (Piewbang et al., 2020),
175 describing an intricate epidemiological dynamics characterised by multiple host infections and

176 interspecies transmission (Duque-Valencia et al., 2019). In this light, the central role of wildlife in
177 the disease epidemiology should be exploited (Anis et al., 2020), enhancing passive surveillance of
178 the wide range of susceptible animal species to broaden the evolutionary analysis of CDV, a
179 pathogen that exhibits a high mutation rate, to undertake reliable wildlife conservation strategies.

180

181 **Funding**

182 This research did not receive any specific grant from funding agencies in the public,
183 commercial, or not-for-profit sectors.

184

185 **Declaration of Competing Interest**

186 Declarations of interest: none

187

188 **Appendix A: Supplementary material**

189 Supplementary data to this article can be found online at doi...

190

191 **Acknowledgements**

192 The Authors thank the wildlife rescue center “*Centro Tutela e Ricerca Fauna Esotica e*
193 *Selvatica Monte Adone ODV*” (<https://centrotutelafauna.org/>; Sasso Marconi, BO, Italy) where all
194 procedures, specimen collection and clinical management of the animals were carried out. The
195 Authors also thank Dr Offer Zeira and Dr Claudiaedda Giovannoni (Ospedale Veterinario San
196 Michele, Tavazzano con Villavesco, LO, Italy, <https://ospedalesanmichele.it/>) and the volunteers
197 working in the wildlife rescue center for medical support and the management of hospitalized
198 animals.

199 **References**

- 200 Anis, E., Needle, D.B., Stevens, B., Yan, L., Wilkes, R.P., 2020. Genetic characteristics of canine
201 distemper viruses circulating in wildlife in the United States. *J. Zoo. Wildl. Med.* 50, 790-
202 797.
- 203
- 204 Appel, M., Robson, D.S., 1973. A microneutralization test for canine distemper virus. *Am. J. Vet.*
205 *Res.* 34, 1459-1463.
- 206
- 207 Bianco, A., Zecchin, B., Fusaro, A., Schivo, A., Ormelli, S., Bregoli, M., Citterio, C.V., Obber, F.,
208 Dellamaria, D., Trevisiol, K., Lorenzetto, M., De Benedictis, P., Monne, I., 2020. Two
209 waves of canine distemper virus showing different spatio-temporal dynamics in Alpine
210 wildlife (2006–2018). *Infect. Genet. Evol.* 84, 104359.
- 211
- 212 Confer, A.W., Kahn, D.E., Koestner, A., Krakowka, S., 1975. Biological properties of a canine
213 distemper virus isolate associated with demyelinating encephalomyelitis. *Infect. Immun.* 11,
214 835-844.
- 215
- 216 Demeter, Z., Lakatos, B., Palade, E.A., Kozma, T., Forgách, P., Rusvai, M., 2007. Genetic diversity
217 of Hungarian canine distemper virus strains. *Vet. Mic.* 122, 258-269.
- 218
- 219 Duque-Valencia, J., Sarute, N., Olarte-Castillo, X.A., Ruíz-Sáenz, J., 2019. Evolution and
220 interspecies transmission of canine distemper virus-an outlook of the diverse evolutionary
221 landscapes of a multi-host virus. *Viruses.* 11, 582.
- 222
- 223 Feng, N., Yu, Y., Wang, T., Wilker, P., Wang, J., Li, Y., Sun, Z., Gao, Y., Xia, X., 2016. Fatal
224 canine distemper virus infection of giant pandas in China. *Sci. Rep.* 6, 27518.

225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249

Frisk, A.L., König, M., Moritz, A., Baumgärtner, W., 1999. Detection of canine distemper virus nucleoprotein RNA by reverse transcription-PCR using serum, whole blood, and cerebrospinal fluid from dogs with distemper. *J. Clin. Microbiol.* 37, 3634-3643.

Greene, C.E., Vandevelde, M., 2012. Canine distemper, in: Greene, C.E., (Eds), *Infectious diseases of the dog and cat*, fourth ed. Elsevier/Saunders, St. Louis, Mo, pp. 2542.

IUCN, 2015. *Martes foina*: Abramov, A.V., Kranz, A., Herrero, J., Choudhury, A., Maran, T.: The IUCN Red List of Threatened Species 2016: e.T29672A45202514.
<https://doi.org/10.2305/IUCN.UK.2016-1.RLTS.T29672A45202514.en>

Kumar, S., Stecher, G., Li, M., Knyaz, C., Tamura, K., 2018. MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Mol. Biol. Evol.* 35, 1547-1549.

Miller, L., Hurley, K., 2009. *Infectious disease management in animal shelters*. Wiley-Blackwell, Ames, Iowa.

Monne, I., Fusaro, A., Valastro, V., Citterio, C., Dalla Pozza, M., Obber, F., Trevisiol, K., Cova, M., De Benedictis, P., Bregoli, M., Capua, I., Cattoli, G., 2011. A distinct CDV genotype causing a major epidemic in Alpine wildlife. *Vet. Microbiol.* 150, 63-69.

Mos, L., Ross, P.S., McIntosh, D., Raverty, S., 2003. Canine distemper virus in river otters in British Columbia as an emergent risk for coastal pinnipeds. *Vet. Rec.* 152, 237-239.

250 Myers, D.L., Zurbriggen, A., Lutz, H., Pospischil, A., 1997. Distemper: not a new disease in lions
 251 and tigers. Clin. Diagn. Lab. Immunol. 4, 180-184.
 252

253 Panzera, Y., Sarute, N., Iraola, G., Hernández, M., Pérez, R., 2015. Molecular phylogeography of
 254 canine distemper virus: Geographic origin and global spreading. Mol. Phylogenet. Evol. 92,
 255 147-154.
 256

257 Perpiñán, D., Ramis, A., Tomás, A., Carpintero, E., Bargalló, F., 2008. Outbreak of canine
 258 distemper in domestic ferrets (*Mustela putorius furo*). Vet. Rec. 163, 246-250.
 259

260 Peterson, M.E., Kutzler, M.A., 2011. Small animal pediatrics: the first 12 months of life.
 261 Saunders/Elsevier, St. Louis, Mo.
 262

263 Philippa, J., Fournier-Chambrillon, C., Fournier, P., Schaftenaar, W., van de Bildt, M., van
 264 Herweijnen, R., Kuiken, T., Liabeuf, M., Ditcharry, S., Joubert, L., Bégner, M., Osterhaus,
 265 A., 2008. Serological survey for selected viral pathogens in free-ranging endangered
 266 European mink (*Mustela lutreola*) and other mustelids from South Western France. J. Wildl.
 267 Dis. 44, 791-801.
 268

269 Piewbang, C., Chansaenroj, J., Kongmakee, P., Banlunara, W., Poovorawan, Y., Techangamsuwan,
 270 S., 2020. Genetic adaptations, biases, and evolutionary analysis of canine distemper virus
 271 Asia-4 lineage in a fatal outbreak of wild-caught civets in Thailand. Viruses. 12, 361.
 272

273 Scagliarini, A., Dal Pozzo, F., Gallina, L., Vaccari, F., Morganti, L., 2007. TaqMan based real time
 274 PCR for the quantification of canine distemper virus. Vet. Res. Commun. 31, 261-263.
 275

276 Schultz, R.D., Thiel, B., Mukhtar, E., Sharp, P., Larson, L.J., 2010. Age and Long-term Protective
 277 Immunity in Dogs and Cats. J. Comp. Pathol. 142, S102-S108.
 278
 279 Sulikhan, N.S., Gilbert, M., Blidchenko, E.Y., Naidenko, S.V., Ivanchuk, G.V., Gorpenchenko,
 280 T.Y., Alshinetskiy, M.V., Shevtsova, E.I., Goodrich, J.M., Lewis, J.C.M., Goncharuk, M.S.,
 281 Uphyrkina, O.V., Rozhnov, V.V., Shedko, S.V., McAloose, D., Miquelle, D.G., Seimon,
 282 T.A., 2018. Canine distemper virus in a wild far eastern leopard (*Panthera pardus*
 283 *orientalis*). J. Wild. Dis. 54, 170-174.
 284
 285 Sykes, J.E., 2014. Canine Distemper Virus Infection, in: Sykes, J.E. (Eds.), Canine and Feline
 286 Infectious Diseases. Elsevier, pp. 152-165.
 287
 288 Williams E.S., Thome, E.T., Appel, M.J.G., Belitsky, D.W., 1988. Canine distemper in black footed
 289 ferrets (*Mustela nigripes*) from Wyoming. J. Wild. Dis. 24, 385-398.
 290
 291 Zhao, J., Shi, N., Sun, Y., Martella, V., Nikolin, V., Zhu, C., Zhang, H., Hu, B., Bai, X., Yan, X.,
 292 2015. Pathogenesis of canine distemper virus in experimentally infected raccoon dogs,
 293 foxes, and minks. Antiviral Res. 122, 1-11.

307 acronym of nation, host species, year of identification, lab numbers only for sequences generated in
308 this study, and the GenBank accession number. When the year of identification was not available,
309 the year of deposition of the nucleotide sequence in the GenBank database was indicated. Canine
310 morbillivirus genetic lineages: America 1 (NA1), America 2 (NA2), Europe 1/South America 1
311 (EU1/SA1), Europe 2/Europe-Wildlife (EU2), Europe 3/Arctic like (EU3), Asia 1 (AS1), Asia 2
312 (AS2), South Africa (ZA), South America 2 (SA2) and Rockborn like (RL). The genetic lineages
313 not containing CDV sequences obtained in this study are compressed. In grey: Wildlife Europe
314 genetic subgroup. Highlighted: sequences generated in this study.
315

316 **Table 1.** Results of reverse transcriptase-PCR (RT-PCR) on oropharyngeal and conjunctival swabs
 317 sampled from survived animals

Subject and samples	RT-PCR results at different sampling times						
	August 24, 2018	September 12, 2018	October 16, 2018	December 19, 2018	February 20, 2019	May 27, 2019	April 17, 2019
#110B							
OFS	N	N	N	N	N	N	N
CS	P	N	P	N	N	N	N
#246B							
OFS	P	N	N	N	N	N	N
CS	N	N	N	P	N	N	N
#246C							
OFS	N	N	N	N	N	N	N
CS	P	N	N	P	N	N	N
#298							
OFS	P	P	P	N	N	N	N
CS	P	P	P	N	N	N	N

318
 319 CS: conjunctival swab. N: negative. OFS: oropharyngeal swab. P: positive.