

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Straw competition and wheat root endophytism of *Trichoderma gamsii* T6085 as useful traits in the biological control of fusarium head blight

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Sarrocco S., Esteban P., Vicente I., Bernardi R., Plainchamp T., Domenichini S., et al. (2021). Straw competition and wheat root endophytism of *Trichoderma gamsii* T6085 as useful traits in the biological control of fusarium head blight. *PHYTOPATHOLOGY*, 111(7), 1129-1136 [10.1094/PHYTO-09-20-0441-R].

Availability:

This version is available at: <https://hdl.handle.net/11585/865295> since: 2022-02-23

Published:

DOI: <http://doi.org/10.1094/PHYTO-09-20-0441-R>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Phytopathology 2021 – 111:1129-1136

Straw Competition and Wheat Root Endophytism of Trichoderma gamsii T6085 as Useful Traits in the Biological Control of Fusarium Head Blight

By Sabrina Sarrocco, Pilar Esteban, Isabel Vicente, Rodolfo Bernardi, Tracy Plainchamp, Séverine Domenichini, Grazia Puntoni, Riccardo Baroncelli, Giovanni Vannacci, and Marie Dufresne

The final published version is available online at:
<https://doi.org/10.1094/PHYTO-09-20-0441-R>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

1 **Straw competition and wheat root endophytism of *Trichoderma gamsii* T6085 as**
2 **useful traits in the biocontrol of *Fusarium* head blight**

3

4 Sabrina Sarrocco^{1*}, Pilar Esteban¹, Isabel Vicente¹, Rodolfo Bernardi¹, Tracy
5 Plainchamp², Séverine Domenichini², Grazia Puntoni¹, Riccardo Baroncelli³,
6 Giovanni Vannacci¹ and Marie Dufresne²

7

8 ¹ Department of Agriculture, Food and Environment, University of Pisa, Via del
9 Borghetto 80, 56124 Pisa, Italy

10 ² Institute of Plant Sciences Paris-Saclay (IPS2), UMR9213/UMR1403, CNRS,
11 INRA, Université Paris-Sud, Université d’Evry, Université Paris-Diderot Sorbonne
12 Paris-Cité, Saclay Plant Sciences, 91405 Orsay, France

13 ³ Instituto Hispano-Luso de Investigaciones Agrarias (CIALE), Departamento de
14 Microbiología y Genética, Universidad de Salamanca, C/ Río Duero 12, Villamayor,
15 37185 Salamanca, Spain.

16

17 ***Corresponding author: S. Sarrocco**; E-mail address: sabrina.sarrocco@unipi.it

18

ABSTRACT

Sarrocco S., Esteban P., Vicente I., Bernardi R., Plainchamp T., Domenichini S.,
Puntoni G., Baroncelli R., Vannacci G. and Dufresne M. 2019. Straw competition and
wheat root endophytism of *Trichoderma gamsii* T6085 as useful traits in the
biocontrol of Fusarium head blight

Phytopathology

Trichoderma gamsii T6085 has been investigated for many years as a beneficial
isolate for use in the biocontrol of Fusarium head blight (FHB) of wheat caused
primarily by *Fusarium graminearum*. Previous work focused on application of T6085
to wheat spikes at anthesis, whereas application to soil before and/or at sowing has
received limited attention. In the present study, the competitive ability of T6085 on
plant residues against *F. graminearum* was investigated. Results showed a significant
reduction of wheat straw colonization by the pathogen and of the development of
perithecia, not only when T6085 was applied alone but also in the presence of a
Fusarium oxysporum isolate (7121), well known as a natural competitor on wheat
plant residues.

T6085 was able to endophytically colonize wheat roots, resulting in internal
colonization of the radical cortex area, without reaching the vascular system, as
confirmed by confocal microscopy. This intimate interaction with the plant resulted in
a significant increase of the expression of the plant defense-related genes *PAL1* and
PRI. Taken together, competitive ability, endophytic behavior, and host resistance
induction represent three important traits that can be of great use in the application of
T6085 against FHB, not only on spikes at anthesis but potentially also in soil before
and/or at sowing.

Keywords Competition_ *Fusarium graminearum* _ *Trichoderma gamsii* _ *Fusarium oxysporum* _ FHB_Endophytism

INTRODUCTION

Cereals are a dominant part of the diet of the world population and supply a large proportion of energy and nutrient needs. Wheat, together with rice and corn, are staple foods for over four billion people, corresponding to 26% of staple foods in Western Europe, (FAO, 2017). A reduction in the yield and quality of this crop threatens food security, particularly given the World Bank's prediction that global wheat production needs to increase by 60% by 2050. Diseases such as Fusarium head blight (FHB) are recognized as among the main causes of reduction in wheat production (Gilbert and Haber, 2013). FHB is mostly caused by a complex of *Fusarium* species, above all *F. graminearum* and *F. culmorum* which are the main pathogens responsible for significant yield reduction, reductions in kernel weight, size, germination rate, protein content and baking quality (McMullen et al., 2012). In addition, kernels can be frequently contaminated by mycotoxins such as trichothecenes, which are secondary metabolites that are extremely toxic for humans and animals (Willyerd et al., 2012). FHB is also economically damaging. Cultivar selection, assessment of disease risk by forecasting models, cultural practices, and fungicides can help control the disease, but no single strategy is fully effective against FHB (Gilbert and Tekauz, 2011; Wegulo et al., 2015; McMullen et al., 2012; Gilbert and Haber, 2013). Biological control could play a key role in protecting wheat against FHB, both in organic and integrated systems. From an epidemiological point of view, *F. graminearum* overwinters between successive cropping seasons on plant residues,

69 such as wheat straw, where it produces perithecia containing ascospores that are
70 released during spring and infect wheat spikes at flowering (Parry et al., 1995;
71 Champeil et al., 2004).

72 Crop residues left in the soil are recognized as the primary inoculum source of this
73 pathogen. Consequently, one or more beneficial fungi that can compete in soil and on
74 plant residue could be applied to reduce the pathogen's survival (Schoneberg et al.,
75 2015, Sarrocco and Vannacci, 2018; Sarrocco et al., 2019b).

76 *Trichoderma* spp., such as *T. atroviride*, *T. velutinum*, *T. harzianum* or *T. gamsii*, are
77 particularly good contenders as beneficial fungi due to their ability to compete for
78 different substrates (Inch and Gilbert, 2005; Gilbert and Habert, 2013; Schoneberg et
79 al., 2015). *T. gamsii* T6085 is a well-known beneficial isolate effective against FHB
80 causal agents, both in lab and field conditions, and is promising for future applications
81 in biological control (Matarese et al., 2012; Sarrocco et al., 2013; Sarrocco et al.,
82 2019a). In previous work (Sarrocco et al., 2019a), we performed an ecological study
83 on this isolate and demonstrated its ability to reduce *F. graminearum* growth on
84 different natural substrates, resulting in significantly lowered trichothecene
85 production and producing secondary metabolites that could affect the pathogen's
86 growth by interference competition.

87 In addition to being good competitors, some *Trichoderma* isolates act as root
88 endophytes of several plants where induction of resistance to later pathogen invasion
89 has been recorded frequently (Shoresh and Harman, 2008; Vinale et al., 2008; Fiorini
90 et al., 2016; Mendoza-Mendoza, 2018; Sarrocco et al., 2017). Endophytic
91 *Trichoderma* spp. are able to penetrate the first or second layers of plant roots, first
92 colonizing the epidermis and then the cortex, without entering the vascular system
93 (Chacòn et al., 2007). This intimate interaction with roots can stimulate defense

responses, with the activation of induced systemic resistance (ISR) involving jasmonic acid and ethylene (JA/ET) signaling, and/or of systemic acquired resistance (SAR), where the SA pathway and accumulation of PR (pathogenesis-related) proteins are activated (Shore et al., 2010; Pieterse et al., 2014).

The aim of the present study was to evaluate the possible application of *T. gamsii* T6085 in soil and/or crop residue in order to reduce the pathogen's initial inoculum and to establish a symbiotic interaction with the plant host. Important beneficial traits of *T. gamsii* T6085 were examined: competition against *F. graminearum* on crop residue, alone or in the presence of *F. oxysporum* (effective colonizer of wheat residues); the ability to endophytically colonize wheat roots; and the capacity to induce the expression of the plant defense-related genes *PAL1* and *PR1*. Results of this study will contribute to understanding the multiple mechanisms used by *T. gamsii* T6085 against FHB causal agents.

MATERIALS AND METHODS

Fungal and plant material

T. gamsii T6085, isolated from uncultivated soil in Crimea (Ukraine) (Matarese et al. 2012, Baroncelli et al., 2016; Vicente et al., 2020) and *F. oxysporum* 7121, isolated from wheat straw in a soil previously cultivated with wheat near Pisa, Italy (Sarrocco et al., 2012) are part of the Fungal Collection of the Plant Pathology & Mycology Lab (Dipartimento di Scienze Agrarie, Alimentari e Agro-Ambientali, University of Pisa). *F. graminearum* ITEM 124, from the fungal collection of CNR-ISPA (<http://www.ispa.cnr.it/Collection>), was kindly provided by Antonio Moretti, ISPA-CNR, Bari, Italy (Zapparata et al., 2017). All fungi were stored long-term on potato

dextrose agar (PDA) (BD, Difco) under mineral oil at 4°C, and actively grown on PDA (*T. gamsii* and *F. oxysporum*) or oat-meal agar (OA) (BD, Difco) (*F. graminearum*) at 24°C under a photoperiod of 12 h of light and 12 h of darkness.

Pathogenicity of *F. graminearum* was regularly tested by artificial inoculations on its host.

In the present work, seeds of the spring wheat cultivar Apogee, a very short life cycle cultivar, were used as plant material (Fu and Nelson, 1994; Bugbee et al., 1997).

Wheat straw colonization test

In order to evaluate the competitive ability of *T. gamsii* T6085, alone and in the presence of *F. oxysporum* 7121, against *F. graminearum* ITEM 124, wheat straw was inoculated following Schoneberg et al. (2015) with some modifications. The *F. oxysporum* isolate was included as potential competitor for wheat straw residue.

Wheat straw, collected from an experimental wheat field near Pisa (Italy), cut into 3-5 cm long pieces including one node, were maintained in deionized water for 24 h. The straw pieces were then autoclaved twice (24 h apart) for 20 min at 120°C, and then used in two different experiments (simultaneous inoculation and subsequent inoculation).

Simultaneous inoculation: In this first test, straw pieces were immersed for 1 h (in an orbital shaker at 100 rpm, room temperature) in 30 ml of an aqueous conidial suspension, made from 15-day old colonies, at a final concentration of 1×10^5 conidia ml⁻¹ of each isolate, both when isolates were used alone and when they were combined. The experiment consisted of: i) *F. graminearum* (FG); ii) *F. graminearum* + *F. oxysporum* (FG+FOX); iii) *F. graminearum* + *T. gamsii* (FG+T); iv) *F. graminearum* + *T. gamsii* + *F. oxysporum* (FG+T+FOX); and v) sterile water as

144 negative control (CONT). Inoculated straw pieces were transferred to empty Petri
145 plates and left for 48 h at room temperature, then moved to Petri dishes (90 mm
146 diameter) containing 25 g of sterilized vermiculite (autoclaved twice for 20 min at
147 120°C) moistened the day before with sterile deionized water. Each treatment
148 consisted of three plates, each containing three inoculated straw pieces. The entire
149 experiment was repeated three times.

150 Successive inoculation: In the second experiment, the pathogen was inoculated 48 h
151 before the antagonists. Straw pieces were initially immersed for 1 h in an aqueous
152 spore suspension of *F. graminearum* 124 (1×10^5 conidia ml⁻¹) then transferred to
153 empty Petri plates. After 48 h of incubation at room temperature, pieces were
154 immersed in 30 ml of a 1×10^5 conidia ml⁻¹ spore suspension of the two antagonists
155 (alone and in combination) for 1 h. Straw pieces were then incubated in empty plates
156 for 48 h, after which they were transferred to new Petri dishes (90 mm diameter)
157 containing 25 g vermiculite previously sterilized, as already described for the
158 simultaneous inoculation experiment. The same inoculum combinations were used as
159 listed for the simultaneous test.

160 Vermiculite plates from both experiments were incubated for 9 weeks at 19-25°C
161 under 12 h light and 12 h darkness with sterile water regularly added to maintain
162 humidity. This interval of incubation temperatures was chosen taking into account
163 that the optimum temperature for perithecium production of *F. graminearum* ranges
164 from 15 to 29.5°C, while for ascospore production it is from 25 to 28°C (Gilbert and
165 Tekauz, 2000).

166 The competitive effect of T6085 (alone and in the presence of *F. oxysporum*) was
167 evaluated on pathogen growth, quantified as fungal pathogen biomass by absolute
168 real-time PCR, together with their ability to reduce the number of perithecia

developed on plant residues. DNA extraction and quantification were performed following the same procedure as described in Sarrocco et al. (2019a) with some modifications: samples, consisting of three pieces of wheat straw for each replicate of each treatment, were ground with a mortar in liquid nitrogen and homogenized in 3 ml of extraction buffer. RNase treatment was not performed since the quality of material was sufficiently good to perform real-time PCR, and the RNase treatment would have led to a loss of DNA. As a standard control, the DNA of *F. graminearum* was extracted from 100 mg of mycelium collected from a PDA plate, using the DNeasy Plant Mini Kit (Qiagen, Germany) following the manufacturer's instructions. Absolute Real-Time PCR was performed in order to estimate fungal biomass. The absolute target quantity in samples was determined by the standard curve method according to Standard Curve Experiments (Applied Biosystems StepOne™ and StepOnePlus™ Real-Time PCR Systems PN 4376784F, Foster City, CA, USA). Fungal DNA in wheat straw was quantified by interpolation from a standard curve obtained with a standard DNA amplified in the same PCR run, as described in Sarrocco et al. (2019). Real-Time PCR reactions (20 µL) were carried out with DNA from fungal pure mycelium or from straw samples (20 ng), 250 nM primers (specific for *F. graminearum*, as described in Sarrocco et al., 2019) and 1x PowerUp™ SYBR™ Green Master Mix (Applied Biosystems) following the manufacturer's instructions. PCR was run under the following thermal cycling conditions: hold 95°C for 20s; 40 cycles at 95°C for 3s and 60°C for 30s.

At the end of the incubation time, the number of perithecia was counted by using a Leica MZ FLIII stereomicroscope (Manstretta and Rossi, 2016). In addition, some perithecia from each treatment were crushed and checked, by microscopic assessment, for the presence of mature ascospores.

The values of fungal growth resulting from Real-Time PCR (after log₁₀ transformation) and the number of perithecia were analyzed by the Tukey post hoc test using Systat (Systat Software, Inc., Chicago, IL) and assuming $P < 0.05$ as the significance level.

Wheat root endophytism

In order to evaluate the root endophytic ability of *T. gamsii*, the antagonist was applied to wheat seeds following Hubbard et al. (2012) with small modifications.

Apogee seeds were surface-sterilized for 3 min in 0.6% NaClO solution with gentle shaking followed by three rinses for 10 min in sterile distilled water and then incubated at 4°C in the dark for 4 days before use. Five sterilized seeds were placed on a PDA plate in a circle with seeds 3 cm apart from each other (corresponding to the diameter of a T6085 colony grown for 24 h in the same conditions). After 24 h, a 5-mm agar plug of T6085 (made from an actively growing colony on PDA) was placed at the center of the Petri dish. As the control, PDA dishes containing sterilized seeds were prepared with a sterile fresh PDA plug (5 mm diameter) in the center. Plates were incubated at 24°C in darkness for 5 days. The experiment was replicated twice, with each replicate consisting of three plates.

The internal colonization of wheat roots by T6085 was determined by re-isolating the antagonist from roots sterilized for 10 s in 95% ethanol, rinsed in sterile water for 10 s, submerged for 20 s in 2.5 % NaClO (Fiorini et al., 2016), then washed three times (2 for 20 s and the last one for 60 s) in sterile water, and finally placed on PDA with the addition of kanamycin (50 µg ml⁻¹) and ampicillin (100 µg ml⁻¹). Plates were incubated at 24°C in the dark for 4-6 days, until fungal development.

In addition, *T. gamsii* root colonization of wheat seedlings was examined by confocal microscopy (LSM880, Zeiss) of both intact and cross-sectioned stained fresh roots. After 5 days of incubation, roots were collected, cleaned gently to remove the excess external fungal colonization, cut into 1.5-2.0-cm lengths, and washed for 1 h in phosphate-buffered saline (PBS), pH 7.4. Roots to be observed whole were stored at 4°C in darkness in new PBS until use, while roots used for cross-sections were first embedded in 3% agarose, then stored at 4°C in new PBS (pH 7.4) until use. The 180- μ m cross-sections were made by a vibratome (VT1200S, Leica). Both intact and cross-sectioned roots were stained with a mix of wheat-germ agglutinin WGA-Alexa Fluor™ 488 and FM4-64 dyes (Thermo Fisher Scientific, Waltham, MA, USA). Samples were incubated for 2 min with the staining solution (10 μ g/mL WGA-Alexa fluor™ 488; 5 μ M FM4-64, 1X PBS in 0.02% Tween) before mounting in 50% glycerol for observation. Fungal material was stained with WGA-AlexaFluor™, while plant plasma membranes were stained with FM4-64 (Bolte et al., 2004).

Plant defense-related gene expression

Wheat roots (with and without *T. gamsii* from seedlings inoculation previously described) were ground in liquid N₂, and 100 mg was used for total RNA extraction using the RNeasy® Plant Mini Kit (Qiagen), followed by DNase I treatment (DNase I Amplification Grade, AMPD1 Sigma-Aldrich, St. Louis, MO), according to the manufacturer's instructions. cDNA synthesis was performed on 400 ng of RNA by using the Maxima First Strand cDNA synthesis kit (K1642 Applied Biosystems) according to the manufacturer's instructions. The expression of the defense-related genes phenylalanine ammonia lyase (*PAL1*) and pathogenesis-related protein 1 (*PR1*) (Shores and Harman, 2008a; 2008b; Bisen et al., 2016) was analyzed by quantitative

real-time PCR performed in a Rotor-Gene Q cycler (Qiagen) with QuantiNOVA SYBR® Green PCR Master MIX 2x (Qiagen) in 20-µl reactions containing 1 µl of cDNA (20 ng) and 0.7µM of each primer (Table 1). Amplification conditions consisted of an initial activation (95°C, 2 min) followed by 40 cycles of denaturation (95°C for 5 sec) and combined annealing/extension (60°C, 10 sec). All reactions were performed in triplicate. Threshold cycles (Ct) were calculated using the β-tubulin gene as housekeeping control. Actin and 18S genes were also tested, but β-tubulin was chosen due to its expression stability. Data were expressed as $2^{-\Delta\Delta C_t}$ to calculate fold differences (Livak and Schmittgen, 2001). Values obtained from the three biological replicates were consistent and submitted to ANOVA (Systat Software, Inc.), assuming $P < 0.05$ as the significance level, to compare gene expression between control (uninoculated) and wheat roots inoculated with T6085. Primers, listed in Table 1, were checked for efficiency and dimer formation.

RESULTS

Wheat straw colonization

In the first experiment, where *F. graminearum* ITEM 124 and the two antagonists were inoculated simultaneously, both *T. gamsii* T6085 and *F. oxysporum* 7121, alone and in combination, were able to significantly reduce the pathogen's growth, expressed as DNA concentration measured by absolute real-time PCR (Figure 1A) after 9 weeks of incubation. A trace amount of *F. graminearum* DNA was detected in uninoculated wheat straw.

A similar trend was observed for the number of perithecia developed by *F.*

graminearum on plant residues after 9 weeks of incubation, when all fungi were inoculated at the same time. When in presence of *T. gamsii* and *F. oxysporum*, alone and co-inoculated, a significantly lower number of perithecia was counted compared with that observed when the pathogen was alone (Figure 1B). No perithecia were detected on the uninoculated wheat straw.

At the end of the incubation time of the second experiment (pathogen inoculated before the antagonists), there was a significant reduction of the pathogen's growth in the presence of *T. gamsii* and *F. oxysporum*, with a greater reduction when the two antagonists were applied together (Figure 1C). A trace amount of DNA of *F. graminearum* was detected in the uninoculated wheat straw used as a control.

When perithecia developed by *F. graminearum* were counted on wheat straw incubated for 9 weeks after the consecutive inoculation of the pathogen and of the two antagonists, a significant reduction in the production of sexual structures was detected, as observed with the first experiment. As shown in Figure 1D, the addition, after 48 hours, of *T. gamsii* and *F. oxysporum*, alone and in combination, resulted in a significantly lower number of perithecia, which was not significant different from the number observed on the uninoculated control (no perithecia).

Wheat root endophytism

At the end of the incubation period (5 days in darkness) all roots developed from sterile seeds appeared to be colonized externally by *T. gamsii*. In some cases a diffuse sporulation began to occur all along the root length. When surface-sterilized roots were plated on PDA with added antibiotics, *T. gamsii* developed from all the

inoculated roots, thus showing a high endophytic ability. No fungal colonization was observed in roots from uninoculated control seeds.

Confocal microscopic observation of double-stained fresh roots, both intact and cross-sectioned, after 5 days of incubation in the presence of *T. gamsii* revealed a superficial fungal root colonization (Figure 2A).

When cross-sections were made, T6085 could be observed in the internal part of the roots. *T. gamsii* hyphae occupied the cortex layer almost reaching but not colonizing the vascular system (Figure 2B). The fungus was able to grow both intra-cellularly and inter-cellularly (Figure 2C).

Plant defense-related gene expression

To test whether wheat roots response to *T. gamsii* endophytic colonization involved a differential activation of defense-related genes, the expression of two selected defense genes, *PAL1* and *PR1*, was analyzed by qRT-PCR. At the end of the 5-day incubation time, the expression of both genes was significantly up-regulated when wheat roots were colonized by T6085 compared with the non-inoculated control exhibiting very weak expression (Figure 3). Specifically, in the presence of *T. gamsii* the expression of *PAL1* and *PR1* genes were induced 3.69-fold ($P=0.0001$) and 257.07-fold ($P=0.0004$), respectively, thus providing preliminary evidence of an induction of host defenses by T6085.

DISCUSSION

Fusarium graminearum, one of the main causal agents of FHB, overwinters as perithecia in crop residue. These resting structures are not only a mean of survival but

produce initial inoculum in the form of ascospores, which will infect the spikes at anthesis (Gilbert and Tekauz, 2000). This survival strategy renders soil conservation strategies such as sod seeding and minimum tillage more prone to the disease. Taking into account the pathogen's survival phase in the disease cycle, the application of one or more antagonists, such as *Trichoderma* isolates, which compete with the pathogen for the colonization of crop residues has the potential to reduce this primary inoculum source (Naef et al., 2006; Schoneberg et al., 2015; Sarrocco and Vannacci, 2018; Sarrocco et al., 2019b). Thus, inoculating *Trichoderma* in the soil may be a good strategy from a competitive exclusion point of view.

Following previous encouraging results showing its antibiotic, mycoparasitic and competitive ability for nutrients (Matarese et al., 2012; Sarrocco et al., 2019a), an evaluation of *T. gamsii* T6085's activity in a more natural substrate - wheat straw - was needed to reveal the real potential as a biocontrol agent of FHB. In the present study, T6085 was applied onto straw pieces in the presence of *F. graminearum* in order to simulate a more natural environment. These tests showed that *T. gamsii* significantly reduced the growth of *F. graminearum* and the development of perithecia. The positive effect of this antagonist was demonstrated not only when the two fungi were inoculated simultaneously, but also when the pathogen was already present on the wheat straw when T6085 was applied, either alone or in presence of the natural competitor *F. oxysporum*. This experimental design resembles what happens in nature where the pathogen has already colonized crop residue and when a biocontrol agent needs to gain access to areas or resources previously held by the pathogen (Holmer and Stenlid, 1993; Boddy, 2000; Jensen et al., 2016).

Our results place *T. gamsii* T6085 in the group of fungi that can outcompete *F. graminearum* such as *T. atroviride*, *T. harzianum* and *C. rosea*, as well as

340 *Microsphaeropsis*, which are all able to colonize wheat residues or to reduce *F.*
341 *graminearum* sporulation on crop debris (Bujold et al., 2001; Naef and Défago, 2006;
342 Gromadzka et al., 2012; Sarrocco and Vannacci, 2018). Our competition tests are not
343 fully representative of what really happens in nature where, in addition to the
344 pathogen and the antagonist, many other biotic and abiotic factors make it a very
345 complex system. However, *in vitro* tests can be used to investigate those mechanisms,
346 which in natural complex systems such as wheat straw in soil, would be very difficult
347 to study (Crowther et al., 2018; Sarrocco et al., 2019a). As a next step, field
348 experiments are needed to confirm the performance of T6085 in more natural
349 conditions, as previously done in Argentina by Palazzini et al. (2013) with two
350 beneficial isolates of *Clonostachys rosea* that were able to reduce *Fusarium* spp. on
351 wheat stubble in different climatic conditions.

352 As a result of wheat straw colonization, we observed a significant reduction in
353 peritheciium development in the presence of *T. gamsii* T6085. Such positive results
354 were also obtained for other beneficial fungi such as *T. harzianum* T-22 or *C. rosea*
355 when inoculated with *Gibberella zeae* on crop residue (Schoneberg et al., 2015;
356 Gilbert and Haber, 2013). What makes *T. gamsii* even more interesting in this context
357 is its ability to suppress *F. graminearum* growth and peritheciium development on
358 wheat straw also in the presence of an *F. oxysporum* isolate, one of the major natural
359 competitors of *F. culmorum* and *F. graminearum* for wheat residues (Pereyra and Dill
360 Macky, 2008; Sarrocco and Vannacci, 2018). This illustrates *T. gamsii* T6085's
361 competitiveness and fitness in more complex situations.

362 Once applied to crop residue on the soil surface, antagonists are able to come into
363 contact with plant roots. The soil is where the first phases of plant growth occur and
364 where rhizosphere-competent fungi as well as endophytic fungi interact with plant

365 roots. Endophytism, i.e., the ability to colonize the first layers of plant root systems, is
366 common among *Trichoderma* isolates (Chacón et al., 2007; Fiorini et al., 2016). This
367 intimate relationship between *Trichoderma* and plants starts with the attachment on
368 the root surface followed by the formation of structures resembling appressoria, which
369 help the beneficial fungus to further penetrate inside the tissues (Yedidia et al., 1999,
370 2000; Viterbo and Chet, 2006). As previously observed for *T. virens* within wheat
371 roots, (Nogueira-Lopez et al., 2018), *T. gamsii* T6085 is able to endophytically
372 colonize wheat roots 1 week after germination, as confirmed by confocal microscopy
373 observations.

374 *Trichoderma* recognition by plants is followed by a cascade of reactions that could
375 lead to the induction of resistance to biotic and abiotic stresses in the host (Waller et
376 al., 2005; Hubbard 2010; Sarrocco et al, 2017). In our wheat seedling samples,
377 defense-related gene expression analysis showed a significant increase of the
378 expression of both *PAL1* and *PR1* genes during the endophytic colonization of roots
379 by T6085, thus letting us to hypothesize that both SAR and ISR could be induced,
380 even if at two different levels of over-expression of the related genes. This response
381 has been previously described in maize treated with other *Trichoderma* isolates, such
382 as *T. virens* or *T. harzianum* (Djonović et al., 2007; Madhavi et al., 2018), and
383 confirmed by Galletti et al. (2020) for *T. gamsii* isolates, with *T. gamsii* IMO5 able to
384 enhance the expression of genes related to ISR, while *T. gamsii* B21 able to enhance
385 the expression of SAR-related genes, thus suggesting an isolate-specific response.

386 These new results confirm the favorable antagonistic potential of *T. gamsii* T6085
387 against one of the major FHB pathogens in a more natural system (wheat straw) even
388 in the presence of a naturally occurring competitor, such as *F. oxysporum*. We further
389 document the ability of this beneficial fungus to induce host defense responses after

endophytic colonization of wheat roots, thus adding another biocontrol mechanism to the repertoire of this beneficial isolate.

LITERATURE CITED

- Baroncelli, R., Zapparata, A., Piaggieschi, G., Sarrocco, S., Vannacci, G. 2016. Draft whole-genome sequence of *Trichoderma gamsii* T6085, a promising biocontrol agent of *Fusarium* head blight on wheat. *Gen. Ann.* 4(1): e01747-15.
- Bisen, K., Keswani, C., Patel, J.S., Sarma, B.K., Singh, H.B. 2016. *Trichoderma* spp.: Efficient Inducers of Systemic Resistance in Plants. *Microbial-Mediated Induced Systemic Resistance in Plants*, 185–195.
- Boddy, L. 2000. Interspecific combative interactions between wood-decaying basidiomycetes. *FEMS Microbiol. Ecol.* 31:185-194.
- Bolte, S., Brown, S., Satiat-Jeunemaitre, B. 2004. The N-myristoylated Rab-GTPase m-Rabmcis involved in post-Golgi trafficking events to the lytic vacuole in plant cells. *J Cell Sci.* 117:943-954.
- Bugbee, B., Koerner, G., Albrechtsen, R., Dewey, W., Clawson, S. 1997. Registration of cultivars. *Crop Sci.* 37:626.
- Bujold, I., Paulitz, T. C., Carisse, O. 2001. Effect of *Microsphaeropsis* sp. on the production of perithecia and ascospores of *Gibberella zeae*. *Plant Dis.* 85:977–984.
- Chacón, M., Rodríguez-Galán, O., Benítez, T., Sousa, S., Rey, M., Llobell, A., Delgado-Jarana, J., 2007. Microscopic and transcriptome analysis of early colonization of tomato roots by *Trichoderma harzianum*. *Int. Microbiol.* 10:19–27.
- Champeil, A., Dore, T., Fourbet, J.F. 2004. *Fusarium* head blight: epidemiological origin of the effects of cultural practices on head blight attacks and the production of mycotoxins by *Fusarium* in wheat grains. *Plant Sci.* 166:1389-1415.

- 415 Crowther, T.W., Boddy, L., Maynard, D.S. 2017. The use of artificial media in fungal
416 ecology. *Fungal Ecol.* 10.1016/j.funeco.2017.10.007.
- 417 Djonović, S., Vargas, W.A., Kolomiets, M.V., Horndeski, M., Wiest, A., Kenerley
418 C.M. 2007. A proteinaceous elicitor Sm1 from the beneficial fungus *Trichoderma*
419 *virens* is required for induced systemic resistance in maize. *Plant Physiol.* 145:
420 875-889.
- 421 FAO. Staple Foods: What do people eat? FAO: Rome, Italy, 2017. Available online:
422 <http://www.fao.org/3/u8480e/u8480e07.htm>.
- 423 Fiorini, L., Guglielminetti, L., Mariotti, L., Curadi, M., Picciarelli, P., Scartazza, A.,
424 Sarrocco, S., Vannacci, G. 2016. *Trichoderma harzianum* T6776 modulates a
425 complex metabolic network to stimulate tomato cv Micro-Tom growth. *Plant Soil*
426 400(1):351-366.
- 427 Fu, B., Nelson, P.E. 1994. Conditions and constraints of food processing in space.
428 *Food Technol.* 48:113–127.
- 429 Galletti, S., Paris, R., Cianchetta S. 2020. Selected isolates of *Trichoderma gamsii*
430 induce different pathways of systemic resistance in maize upon *Fusarium*
431 *verticillioides* challenge. *Microb. Res.* 233: 126406.
- 432 Gilbert, J., Haber, S. 2013 Overview of some recent research developments in
433 *Fusarium* Head Blight of wheat. *Can. J Plant Path.* 35:149-174.
- 434 Gilbert, J., Tekauz, A. 2011. Strategies for management of fusarium head blight
435 (FHB) in cereals. *PS&C Prairie Soils and Crops J* 4: 97-104.
- 436 Gilbert, J. Tekauz, A. 2000. Review: recent developments in research on fusarium
437 head blight of wheat in Canada. *Can. J. Plant Pathol.* 22:1-8.

- 438 Gromadzka, K., Lenc, L., Sadowski, C.Z., Baturo-Cieśniewska, A., Chełkowski, J.,
439 Goliński, P., Bocianowski, J. 2012. Effects of fungicidal protection programs on
440 the development of *Fusarium* head blight and the accumulation of mycotoxins in
441 winter wheat. *Cer. Res. Comm.* 40(4):518–531.
- 442 Holmer, L., Stenlid, J. 2016. The importance of inoculum size for the competitive
443 ability of wood decomposing fungi. *FEMS Microbiol. Ecol.* 12:169-176.
- 444 Hubbard, K.E., Nishimura, N., Hitomi, K., Getzoff, E.D., Schroeder J.I. 2010. Early
445 abscisic acid signal transduction mechanisms: newly discovered components and
446 newly emerging questions. *Genes and Develop* 24:1695–1708.
- 447 Hubbard, M., Germida, J., Vujanovic, V. 2012. Fungal endophytes improve wheat
448 seed germination under heat and drought stress. *Botany* 90:137–149.
- 449 Inch, S. Gilbert, J. 2007. Effect of *Trichoderma harzianum* on perithecial production
450 of *Gibberella zeae* on wheat straw. *Biocontrol Sc. Technol.* 17:635-646.
- 451 Jensen, D.F., Karlsson, M., Sarrocco, S., Vannacci, G. 2016. Biological control using
452 microorganisms as an alternative to disease resistance. Pages 341-363, in:
453 Collinge DB (2016) *Biotechnology for Plant Disease Control*, New York and
454 London: Wiley. ISBN 978-1-118-86776-1.
- 455 Livak, K.J., Schmittgen, T.D. 2001. Analysis of relative gene expression data using
456 real-time quantitative PCR and the $2\Delta\Delta C(T)$ method. *Methods* 25(4):402–408.
- 457 Madhavi, G.B., Devi, G.U., Kumar, K.V.K., Ramesh, T. 2018. Evaluation of
458 *Pseudomonas fluorescens* and *Trichoderma harzianum* isolates in inducing
459 systemic resistance (ISR) in maize against *Rhizoctonia solani* f. sp. *Sasakii*. *Int. J.*
460 *Chem. Stud.* 6:628-632.
- 461 Manstretta, V., Rossi, V. 2016. Ascospore discharge by *Fusarium graminearum* as
462 affected by temperature and relative humidity. *Eur. J Plant Pathol.* 146:191–197.

- 463 Matarese, F., Sarrocco, S., Gruber, S., Seidl-Seiboth, V., Vannacci, G. 2012.
 464 Biocontrol of Fusarium Head Blight: interactions between *Trichoderma* and
 465 mycotoxigenic *Fusarium*. Microbiol. 158:98-106.
- 466 McMullen, M., Bergstrom, G., De Wolf, E., Dill-Macky, R., Hershman, D., Shaner,
 467 G., Van Sanford, D. 2012. A unified effort to fight an enemy of wheat and barley:
 468 Fusarium head blight. Plant Dis. 96:1712–1728.
- 469 Mendoza-Mendoza, A., Zaid, R., Lawry, R., Hermosa, R., Monte, E., Horwitz, B. A.,
 470 Mukherjee, P. K. et al. 2018. Molecular dialogues between *Trichoderma* and roots:
 471 role of the fungal secretome. Fungal Biol. Rev. 32, 62–85.
- 472 Morán-Diez, E., Hermosa, R., Ambrosino, P., Cardoza, R.E., Gutiérrez, S., Lorito,
 473 M., Monte, E. 2009. The ThPG1 endopolygalacturonase is required for the
 474 *Trichoderma harzianum*-plant beneficial interaction. Mol. Plant Microbe Interact.
 475 22:1021–31.
- 476 Naef, A., Défago, G. 2006- Population structure of plant-pathogenic *Fusarium* species
 477 in overwintered stalk residues from Bt-transformed and non-transformed maize
 478 crops. Eur. J Plant Pathol. 116(2):129–143.
- 479 Naef, A., Senatore, M., Défago, G. 2006. A microsatellite based method for
 480 quantification of fungi in decomposing plant material elucidates the role of
 481 *Fusarium graminearum* DON production in the saprophytic competition with
 482 *Trichoderma atroviride* in maize tissue microcosms. FEMS Microbiol. Ecol.
 483 55:211–220
- 484 Nogueira-Lopez, G., Greenwood, D.R., Middleditch, M., Winefield, C., Eaton, C.,
 485 Steyaert, J.M., Mendoza-Mendoza, A. 2018. The apoplastic secretome of
 486 *Trichoderma virens* during interaction with maize roots shows an inhibition of

- 487 plant defence and scavenging oxidative stress secreted proteins. *Front Plant Sci.*
 488 9:409.
- 489 Palazzini, J.M., Groenenboom-de Haas, B.H., Torres, A.M., Köhl, J., Chulze, S.N.
 490 2012. Biocontrol and population dynamics of *Fusarium* spp. on wheat stubble in
 491 Argentina. *Plant Pathol.* 62:859-866.
- 492 Parry, D.W., Jenkinson, P., McLeod, L. 1995. *Fusarium* ear blight (scab) in small
 493 grains - a review. *Plant Pathol.* 44:207–238.
- 494 Pereyra, S.A., Dill-Macky, R. 2008. Colonization of the residues of diverse plant
 495 species by *Gibberella zeae* and their contribution to *Fusarium* head blight
 496 inoculum *Plant Dis.* 92:800-807.
- 497 Pieterse, C.M.J., Zamioudis, C., Berendson, R., Weller, D., van Wees, S.C.M.,
 498 Bakker, P.A.H.M. 2014. Induced systemic resistance by beneficial microbes. *Ann.*
 499 *Rev. Phytopath.* 52:347– 375.
- 500 Rodriguez, R.J., Henson, J., Van Volkenburgh, E., Hoy, M., Wright, L., Beckwith, F.
 501 Kim, Y-O, Redman, R.S. 2008. Stress tolerance in plants via habitat-adapted
 502 symbiosis. *The ISMA J.* 2:404e416.
- 503 Sarrocco S, Matarese F, Baroncelli, R, Seidl-Seiboth, V, Kubicek, CP, Vannacci, G,
 504 Vergara, M. 2017. The constitutive endopolygalacturonase TvPG2 regulates the
 505 induction of plant systemic resistance by *Trichoderma virens*. *Phytopathology.*
 506 107(5):537-544.
- 507 Sarrocco, S., Matarese, F., Moncini, L., Pachetti, G., Ritieni, A., Moretti et al. 2013.
 508 Biocontrol of *Fusarium* head blight by spike application of *Trichoderma gamsii*. *J*
 509 *Plant Pathol.* S1:19-27.
- 510 Sarrocco, S., Matarese, F., Moretti, A., Haidukowski, M., Vannacci, G. 2012. DON on
 511 wheat crop residues: effects on mycobiota as a source of potential antagonists of

- 512 *Fusarium culmorum*. *Phytopath. Med.* 51:225-235.
- 513 Sarrocco, S., Mauro, A., Battilani, P. 2019a. Use of Competitive Filamentous Fungi
514 as an Alternative Approach for Mycotoxin Risk Reduction in Staple Cereals:
515 State of Art and Future Perspectives. *Toxins* 11:701.
- 516 Sarrocco, S., Valenti, F., Manfredini, S., Esteban, P., Bernardi, R., Puntoni G.,
517 Baroncelli, R., Haidukowski, M., Moretti, A., Vannacci, G. 2019. Is exploitation
518 competition involved in a multitrophic strategy for the biocontrol of *Fusarium*
519 Head Blight? *Phytopathology* 109(4):560-570.
- 520 Sarrocco, S., Vannacci, G. 2018b. Preharvest application of beneficial fungi as a
521 strategy to prevent postharvest mycotoxin contamination: a review. *Crop Prot.*
522 110:160-170.
- 523 Schöneberg, A., Musa, T., Voegelé, R., Vogelgsang, S. 2015. The potential of
524 antagonistic fungi for control of *Fusarium graminearum* and *Fusarium*
525 *crookwellense* varies depending on the experimental approach. *J. Appl. Microbiol.*
526 118:1165–1179-
- 527 Shores, M., Harman, G.E. 2008a. The molecular basis of shoot responses of maize
528 seedlings to *Trichoderma harzianum* T22 inoculation of the root: a proteomic
529 approach. *Plant Physiol.* 147:2147–63.
- 530 Shores M, Harman GE. 2008b. The relationship between increased growth and
531 resistance induced in plants by root colonizing microbes. *Plant Sign. Behav.* 3:737–39
- 532 Shores, M., Mastouri, F., Harman, GE. 2010. Induced systemic resistance and plant
533 responses to fungal biocontrol agents. *Ann. Rev. Phytopath.* 48:21– 43.
- 534 Vicente, I., Baroncelli, R., Moran-Diez, M.E., Bernardi, R., Puntoni, G., Hermosa, R.,
535 Monte, E., Vannacci, G., Sarrocco, S. 2020. Combined comparative genomics and

- 536 gene expression analyses provide insights into the terpene synthases inventory in
537 *Trichoderma*. *Microorganisms*, 8(10):1603.
- 538 Vinale, F., Sivasithamparam, K., Ghisalberti, E.L., Marra, R., Berbetti, M.J., Li, H.,
539 Woo, S.L., Lorito, M. 2008. A novel role for *Trichoderma* secondary metabolites
540 in the interactions with plants. *Physiol. Mol. Plant Path.* 72(1-3):80-86.
- 541 Viterbo, A., Chet, I. 2006. *TasHyd1*, a new hydrophobin gene from the biocontrol
542 agent *Trichoderma asperellum*, is involved in plant root colonization. *Mol. Plant*
543 *Pathol.* 7:249–258.
- 544 Waller, F., Achatz, B., Baltruschat, H., Fodor, J., Becker, K., Fischer, M., Heier, T.,
545 Hu ckelhoven, R., Neumann C., von Wettstein, D., Franken, P., Koge K.H., 2005.
546 The endophytic fungus *Piriformospora indica* reprograms barley to salt-stress
547 tolerance, disease resistance, and higher yield. *Proc. Natl. Acad. Sci. USA*
548 102:13386–13391.
- 549 Wegulo, S.N., Baenziger, P.S., Nopsa, J.H., Bockus, W.W., Hallen-Adams, H. 2015.
550 Management of Fusarium head blight of wheat and barley. *Crop Prot.* 73:100 -
551 107.
- 552 Willyerd, K.T., Li, C., Madden, L.V., Bradley, C.A., Bergstrom, G.C., Sweets, L.E.,
553 McMullen, M., Ransom, J.K., Grybauskas, A., Osborne, L., Wegulo, S.N.,
554 Hershman, D.E., Wise, K., Bockus, W.W., Groth, D., Dill-Macky, R., Milus, E.,
555 Esker, P.D., Waxman, K.D., Adee, E.A., Ebelhar, S.E., Young, B.G., Paul, P.A.
556 2012. Efficacy and stability of integrating fungicide and cultivar resistance to
557 manage Fusarium head blight and deoxynivalenol in wheat. *Plant Dis.* 96:957-967.
- 558 Yedidia I., Benhamou N., Chet I. 1999. Induction of defense responses in cucumber
559 plants (*Cucumis sativus* L.) by the biocontrol agent *Trichoderma harzianum*. *Appl.*
560 *Environ. Microbiol.* 65:1061-70

- 561 Yedidia I., Benhamou N., Kapulnik Y., Chet I. 2000. Induction and accumulation of
562 PR proteins activity during early stages of root colonization by the mycoparasite
563 *Trichoderma harzianum* strain T-203. Plant Physiol. Biochem. 38:863-73
- 564 Zapparata, A., Da Lio, D., Somma, S., Vicente Muñoz, I., Malfatti, L., Vannacci, G.,
565 Moretti, A., Baroncelli, R., Sarrocco, S. 2017. Genome sequence of *Fusarium*
566 *graminearum* ITEM 124 (ATCC 56091), a mycotoxigenic plant pathogen. Gen.
567 Announc. 5(45):e01209-17.

Table 1. List of specific primers used for qRT-PCR.

Target gene	Primer	Sequence	Expected amplicon (bp)
Induction of Resistance			
<i>Triticum aestivum</i> pathogenesis-related protein 1 (<i>PR1</i>)	PR1A-F	CGCAGAACTCGCCTCAGGAC	87
	PR1A-R	GCTTCGTGCTCCAGGTCACC	
<i>Triticum aestivum</i> inducible phenylalanine ammonia-lyase (<i>PAL</i>)	PALB-F	ATCTCATCCAGGAAGACGCCG	176
	PALB-R	CCCATGTTGTTTCATGCTCAGGG	
Housekeeping			
<i>Triticum aestivum</i> β-tubulin 1 (<i>TUB1</i>)	TUBA-F	AAGTTCTGGGAGGTGGTGTGC	105
	TUBA-R	CTCGTTGTAGTAGACGTTGACGC	
<i>Triticum aestivum</i> actin (<i>ACT</i>)	ACTB-F	GCCGTTCTGTCCTTGTATGCC	158
	ACTB-R	ATTAGATTATCCGTGAGGTCCCG	
<i>Triticum aestivum</i> 18S rRNA gene	18S-F	GCTCGAAGACGATCAGATACCG	146
	18S-R	TTCAGCCTTGCGACCATACTC	

Primers were manually designed and *in silico* verified them by using bioinformatic tools (Netprimer and IDT oligo analyzer tool). Further, primers have been checked by ReaTime_PCR in order to evaluate their efficiency and to exclude dimmer formation.




Figure 1: Amount of DNA per sample (A), and number of perithecia per straw piece (B) of *Fusarium graminearum* ITEM 124 in wheat residue simultaneously inoculated with the pathogen and the two antagonists *T. gamsii* T6084 and *F. oxysporum* 7121 (alone and in combination), after 9 weeks of incubation. Amount of DNA per sample (C) and number of perithecia per straw piece (D) of *F. graminearum* on wheat straw consecutively inoculated with the pathogen (48 h before) and the two antagonists (alone and in combination), after 9 weeks of incubation. *F. graminearum* DNA was measured by absolute real-time polymerase chain reaction performed on 20 ng of total DNA extracted from straw samples and submitted to log₁₀ transformation. CONT = uninoculated wheat straw; FG = straw pieces inoculated with *F. graminearum*; FG+FO = straw pieces simultaneously inoculated with *F. graminearum* and *F. oxysporum*; FG+T = straw pieces simultaneously inoculated with *F. graminearum* and *Trichoderma gamsii*; and FG+FO+T = straw pieces simultaneously inoculated with *F. graminearum*, *T. gamsii*, and *F. oxysporum*. Values are means and standard deviations of 3 replicates. Different letters correspond to significantly different values (ANOVA, P<0.05).

176x164mm (300 x 300 DPI)




Figure 2. Colonization pattern of *Trichoderma gamsii* T6085 in wheat seedling roots 7 days after inoculation: A) close-up of *T. gamsii* hyphae inhabiting epidermal cells of wheat root; B) cross-section of wheat root showing internal colonization of epidermal and cortical layers near to vascular system; C) arrows indicate intracellular (dashed line) and intercellular (continuous line) colonization by *T. gamsii* hyphae. Fungal cells were labeled with WGA-Alexa Fluor 488 (green channel); the plant cell wall was labeled with FM4-64 (red channel).

153x185mm (300 x 300 DPI)




Figure 3. Induction of defense-related genes in wheat seedling roots colonized by *Trichoderma gamsii* T6085. A) relative expression of the PAL1 gene; B) relative expression of the PR1 gene. The relative quantity of transcripts was calculated using the comparative cycle threshold method ($2^{-\Delta\Delta Ct}$). The wheat TUB1 gene was used as an endogenous control to normalize for differences in input RNA between the different samples. Data represent mean values of 3 independent experiments. Within each graph at different letters correspond values significantly different (ANOVA).

187x73mm (300 x 300 DPI)