



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

ARCHIVIO ISTITUZIONALE  
DELLA RICERCA

## Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Soil organic carbon stock assessment in forest ecosystems through pedogenic horizons and fixed depth layers sampling: What's the best one?

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

De Feudis, M., Falsone, G., Vianello, G., Agnelli, A., Vittori Antisari, L. (2022). Soil organic carbon stock assessment in forest ecosystems through pedogenic horizons and fixed depth layers sampling: What's the best one?. *LAND DEGRADATION & DEVELOPMENT*, 33(9), 1446-1458 [10.1002/ldr.4253].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/886117> since: 2022-11-24

*Published:*

DOI: <http://doi.org/10.1002/ldr.4253>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

De Feudis, M., Falsone, G., Vianello, G., Agnelli, A., & Vittori Antisari, L. (2022). Soil organic carbon stock assessment in forest ecosystems through pedogenic horizons and fixed depth layers sampling: What's the best one? *Land Degradation & Development*, 33( 9), 1446– 1458.

The final published version is available online at: <https://doi.org/10.1002/ldr.4253>

#### Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

De Feudis Mauro (Orcid ID: 0000-0002-1899-0730)  
Falsone Gloria (Orcid ID: 0000-0002-0072-9139)  
Agnelli Alberto (Orcid ID: 0000-0002-2236-9103)

## Manuscript title

**Soil organic carbon stock assessment in forest ecosystems through pedogenic horizons and fixed depth layers sampling: what's the best one?**

Running title

Pedogenic horizons sampling for an accurate forest SOC assessment

Mauro De Feudis<sup>1</sup>, \*, Gloria Falsone<sup>1</sup>, Gilmo Vianello<sup>2</sup>, Aberto Agnelli<sup>3</sup>, Livia Vittori Antisari<sup>1</sup>

<sup>1</sup>Department of Agricultural and Food Sciences, Alma Mater Studiorum - University of Bologna, Via Fanin, 40, 40127 Bologna, Italy.

<sup>2</sup>Centro Sperimentale per lo Studio e l'Analisi del Suolo (CSSAS), Alma Mater Studiorum-University of Bologna, Bologna, Italy.

<sup>3</sup>Department of Agricultural, Food and Environmental Sciences, University of Perugia, Borgo XX giugno, 74, 06121 Perugia, Italy.

\* Corresponding author:

Mauro De Feudis

Department of Agricultural and Food Sciences

Alma Mater Studiorum – University of Bologna

Via Fanin, 40; 40127 Bologna, Italy

e-mail: mauro.defeudis2@unibo.it

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1002/ldr.4253](https://doi.org/10.1002/ldr.4253)

This article is protected by copyright. All rights reserved.

## Abstract

In soil organic carbon (SOC) survey the role of sampling approach is crucial and should not be underestimated. In this sense, the appropriateness of soil sampling by pedogenic horizons (PGH) and fixed-depth layers (FDL) in forest is still field of debate. The present work aimed to a) study the spatial variability of SOC concentrations and C stocks through PGH and FDL samplings in uneven-aged sweet chestnut, European beech and Norway spruce forests of Apennine chain (Italy); b) clarify the major advantages and drawbacks of sampling by PGH and FDL. In a representative area (18×18 m) of each forest, a soil profile was investigated and 8 additional sampling points were selected. The sampling was performed both by PGH and FDL (0–15 and 15–30 cm). For each forest, no SOC content and C stock differences in 0–30 cm soil depth were found between PGH (58.8±5.3 g kg<sup>-1</sup> and 103±7 g ha<sup>-1</sup>, respectively) and FDL (52.7±4.3 g kg<sup>-1</sup> and 117±9 Mg ha<sup>-1</sup>, respectively) sampling. However, under Norway spruce, PGH sampling pointed out that C was greatly stored in Oe and Oa horizons (51.1 vs 146 Mg ha<sup>-1</sup> in the 0-30 cm layer). The higher coefficient of variation obtained when soil horizons were considered (from 19.2 to 72.8% for PGH and from 16.5 to 25.7% for FDL) suggested that PGH sampling preserved the information about the spatial variability within an ecosystem. The information loss about SOC vertical and spatial distribution would indicate the inability of FDL sampling to support decision-making plans addressed for sustainable use of soil resource.

## Keywords

Mini-sampler; soil horizons; SOC stock; forest soil; cambisols

## 1. Introduction

The report published by the Intergovernmental Panel on Climate Change in 2018 (IPCC, 2018) highlighted the urgent need for actions on global scales to prevent the increase of the mean air temperature of 1.5°C compared to the pre-industrial levels. At the global scale, soils are the largest organic carbon (C) sink in terrestrial ecosystems, containing approximately 1500 Pg C in 1 m of depth which is 2–3 folds the amount of C found in plant biomass and atmosphere (Batjes, 2014; Vicente-Vicente et al., 2016). Therefore, while a slight decrease of the soil organic C (SOC) content can lead to significant changes in atmospheric CO<sub>2</sub> (IPCC, 2013), the increase of C sequestration in form of SOC can have an important role on mitigation of the current global warming (Minasny et al., 2017).

Globally, forest ecosystems cover approximately more than 4 billion hectares (FAOSTAT, 2020) and the most recent estimates reported that they store about 650 Pg of C, of which 44 % in the biomass, 11 % in dead wood and litter, and 45 % into the soil (FAO, 2010). In Italy, forested areas amount for about 10 million hectares (FAOSTAT, 2020) and their soils store from 80 to 130 Mg C ha<sup>-1</sup> to 1 m depth (Caddeo et al., 2019). Italian forests are mainly distributed in mountainous areas (Cesaro and Romano, 2017), where soils are often characterized to be poorly developed, shallow and skeletal due to limited pedogenic processes and high soil erosion rates (FAO, 2015; Guerra et al., 2020; Alewell et al., 2008). Further, mountain soils are recognized to have a high sensitivity to climate change (Hagedorn et al., 2010), which make them highly vulnerable to SOC decline, erosion and further soil degradation (Conforti et al., 2013; Bojko and Kabala, 2016), which, in turn, can affect the ecosystem functions supplied by forest soil such as biomass production, nutrient and water supply, C sequestration, and biodiversity conservation (Vicente-Vicente et al., 2019; Baveye et al., 2016).

SOC amount is the result of a combination of various environmental factors as well as vegetation composition and climate. To this regard, several studies pointed out that higher SOC concentrations occur under broadleaf than coniferous forests (e.g., Cha et al., 2019; Massaccesi et al., 2020) and

SOC stock is negatively correlated with temperature and positively correlated with precipitation (e.g., Choudhury et al., 2016; Jobbágy and Jackson, 2000; Massaccesi et al., 2020). Further, also parent material can affect the size of organic C pool stored in soil. According to Yang et al. (2018) and Mao et al. (2020), limestone-derived soils tend to promote C stock due to the presence of calcium and magnesium which link the organic compound to soil minerals. In acidic soils, SOC is stabilized in organic- mineral complexes through the binding action of Fe and Al ions (Su et al., 2021).

SOC plays a key role in the functioning of forest ecosystems by regulating nutrient cycling, water storage and availability, erosion and flooding, and supporting biota habitat (Brang et al., 2006; Adhikari and Hartemink, 2016; Minasny and McBratney, 2018; Rabbi et al., 2018). Because of the large number of environmental functions provided by SOC, its adequate monitoring is the first step towards preserving SOC and thus limiting soil degradation processes.

Several studies were performed in European forests to give a large-scale view of the current state and changes in forest soils (e.g., Wellbrock and Bolte, 2019; Hiederer et al., 2011). At local scale, however, the measurement of SOC concentration and, consequently, the determination of C stock pose several critical points in their assessment, and the soil sampling approaches are still field of debate. Indeed, with the exception of the organic layers, namely the O horizons, for which there is a general agreement on sampling by horizons (e.g., Koga et al., 2020; Lull et al., 2020), in forest ecosystems mineral soil samples can be collected both by horizons (e.g., De Feudis et al., 2021b; Massaccesi et al., 2020; Valtera and Šamonil, 2018) or by fixed-depth layers (Canedoli et al., 2020; Gross et al., 2018; Tesfaye et al., 2016). The ICP Forest (Cools and De Vos, 2020) designed a manual for the monitoring of the main soil properties based on both sampling types. The choice to sample by pedogenic horizons (PGH) or by fixed depth layers (FDL) is driven by the research aims and the advantages and disadvantages of the two approaches. Soil sampling by FDL is usually employed for modelling and mapping (e.g., De Feudis et al., 2021a; Orton et al., 2016; Sulieman et al., 2018). The major strength of soil sampling by FDL is related to its greater ease execution. It

Accepted Article

does not need experts for the delineation of the soil horizons, it is less time consuming compared to the sampling by PGH and is, therefore, cheaper. Sampling by FDL is statistically efficient because not all soil profiles have all the same morphology, although it does not give any information about type and thickness of the soil horizons. The sampling by PGH is useful when vertical and lateral changes of soil properties (e.g., nutrients and SOC) are studied (Grüneberg et al., 2010; Shahbazi et al., 2019; Sharififar et al., 2019), or when soil processes regulating properties and functions are investigated. Additionally, the soil sampling by PGH provide the same data that can be obtained collecting samples by fixed depth, because during the pedological survey the thickness and depth of each horizon are recorded.

In the present study we determined the soil C concentration and stocks in three widespread forest ecosystems, (chestnut, beech and spruce forests) sampling by both PGH and FDL and considering the upper 30 cm of soil. We choice to investigate the soil down to 30 cm of depth because it has been reported that the SOC in the topsoil is noticeably responsive to global climate change and human disturbances (Don et al., 2011; Grüneberg et al., 2014; Li et al., 2017) and such interval is worldwide used for the soil organic carbon monitoring (Makipaa et al., 2012; IPCC, 2003; Hiederer et al., 2011). In this context, it is important to highlight that, though the upper part of the soil (i.e., 0 – 10 cm) has higher SOC concentration compared to the subsoil (e.g., Massaccesi et al., 2020; De Feudis et al., 2021b), the latter can considerably contribute to the C stocks of the whole soil (Grüneberg et al., 2010). The main aims of this study were a) to study the spatial variability of SOC concentrations and C stocks through PGH and FDL samplings in the three forest ecosystems investigated; b) to clarify the major advantages and drawbacks of sampling by soil horizons and fixed depths. We hypothesized that: a) SOC concentration and C stock values do not change in 0–30 cm soil depth between the two sampling approaches; b) PGH sampling allows to keep information about horizontal and vertical SOC distribution.

## 2. Materials and methods

### 2.1. Study sites description

Three forest types that are widespread distributed in the Italian mountain areas (Cislaghi et al., 2021), uneven-aged sweet chestnut (*Castanea sativa* Mill.), European beech (*Fagus sylvatica* L.) and Norway spruce (*Picea abies* (L.) H. Karst.) forests, were selected as study sites (Figure 1) in the northern Apennine chain (Italy). The three study sites had similar lithology made of stratified arenaceous–pelitic turbidites (Bertolini et al., 2005) belonging to the Miocene period, with feldspars, micas and quartz as the main minerals (Vittori Antisari et al., 2013). In all sites, the soils were classified as Dystric Cambisols (FAO, 2014).

The sweet chestnut forest was located at about 700 m above sea level (a.s.l.), with a North–Northwest exposure (315°N) and a slope ranging from 12 to 18%. The mean annual air temperature and precipitation were 10.0 °C and 1403 mm, respectively. The European beech and Norway spruce forests were placed at about 1300 m a.s.l., with a Northeast exposure (47°N) and a slope ranging from 19 to 24% for the beech forest, and a South–Southwest exposition (225°N) and a slope ranging from 10 to 12% for the spruce one. The mean annual air temperature was 6.9 °C and the mean annual precipitation was 2098 mm for both forest type.

The chestnut forest had a density of 448 plants ha<sup>-1</sup> and the average tree diameter at breast height (DBH) was of 31 cm, 24 cm and 10 cm for 39, 30 and 31 % of total plants, respectively. The beech forest had a density of 1105 plants ha<sup>-1</sup> and the average tree DBH was of 34 cm, 22 cm and 12 cm for 25, 40 and 35 % of total plants, respectively. The spruce forest had a density of 608 plants ha<sup>-1</sup> and the average tree DBH was of 54 cm and 42 cm for 27 and 73 % of total plants, respectively.

### 2.2. Soil sampling design

In each site, an area of 324 m<sup>2</sup> (18 m × 18 m) was selected. In order to obtain comparable data among the forest types, we selected investigation areas with healthy plants, no herb coverage, no invasive plants intrusion, similar slope and no evident erosive soil processes. At the centre of the selected area, a representative soil profile was described down to BC or C horizon (Figure 2). The



morphological features (i.e., depth, thickness, structure, etc.) of organic and mineral horizons were described according to Schoeneberger et al. (2012) and are reported in Table S1 of the Supplementary Materials. In each area, the Oi horizon has been collected within one 100×100 cm sampling frame that was placed near the point where the soil profile was excavated. Soil samples were collected from the profile both by considering all the pedogenic horizons (PGH) and by fixed-depth layers at 0–15 cm (comprising the thin Oe horizon) and 15–30 cm (FDL) (Figure 2). Further, 8 additional sampling points were chosen at a distance of 6 m from the soil profile and 6 m from one to each other (Figure 3). In such sampling points, two soil cores from 0 to 30 cm of depth were collected through metallic mini-samplers made by two half-cylinders (height 30 cm, diameter 4 cm, weight 700 g) (Figure 3) that were hammered into the soil starting from Oe horizon. Once in the laboratory, one of the two soil cores collected in each sampling point was divided according the recognized pedogenic horizons, whereas the other was divided in two samples by depth (0–15 cm and 15–30 cm) (Figure 2).

### 2.3. Bulk density determination

#### 2.3.1 Bulk density for Oi and Oe horizons

In each area, one sample from Oi horizon and one sample from Oe horizon were collected and their thickness were measured. The collected samples were oven-dried at 105 °C and then their mass was weighted. The bulk density (BD) of Oi and Oe horizons was calculated taking into account the 105°C oven-dried collected mass (g), the sampling area (cm<sup>2</sup>) and the horizon thickness (cm).

Conversely to Oi and Oe, the Oa horizon was hard to identify in the field and it was classified as an organic horizon only after it was analysed for its C content (130 g kg<sup>-1</sup>, on average). Therefore, the BD of the Oa horizon was calculated by a pedotransfer function (see section 2.3.3).

The measured BD of the organic horizons under Sweet chestnut, Norway spruce and European beech forests were 0.028, 0.084 and 0.068 g cm<sup>-3</sup> for Oi and 0.062, 0.099 and 0.157 g cm<sup>-3</sup> for Oe,

respectively. The calculated mean BD for the Oa horizon, present only under Norway spruce, was  $0.73 \text{ g cm}^{-3}$ .

### 2.3.2 Bulk density for 0–15 and 15–30 soil layers of the representative soil profile

Undisturbed soil samples were collected from each soil profile by a steel core (diameter 5 cm; height 5 cm) at 0–5, 5–10, 10–15, 15–20, 20–25 and 25–30 cm depth starting from the upper limit of Oe horizon. The undisturbed soil samples collected by steel-core were oven-dried at  $105 \text{ }^{\circ}\text{C}$ , weighed, and the dry weight was divided by the volume of the cores. The obtained density values were adjusted by subtracting the mass and the volume of skeleton to determine the BD of the fine earth fraction. The mass and the volume occupied by roots inside the steel cores accounted less than 1% and were considered negligible. Then, to obtain the BD of the 0–15 cm depth layer, the BD values of the samples collected at 0–5, 5–10, 10–15 cm depths were averaged. Similarly, the BD of 15–30 cm depth layer was obtained by averaging the BD values of the samples collected at 15–20, 20–25 and 25–30 cm depths. The BD values of 0–15 cm and 15–30 cm depth layers were then used to develop the pedotransfer function (PTF) that was used for the calculation of the BD values of the soil samples collected both by PGH and FDL.

### 2.3.3 Pedotransfer function (PTF) development for estimating soil bulk density

The BD values of the soil samples collected both by PGH and FDL were calculated by the following PTF:

$$BD = a + b \times \log(\text{sand}) + c \times \log(\text{SOC}) \quad [1]$$

where  $a = 3.80429$ ,  $b = -0.42052$ ,  $c = -0.08518$ .

The PTF used for the present study was developed taking into account sand and SOC contents because of their major influence on BD (Chen et al., 2018). The obtained PTF had an adjusted  $R^2 = 0.74$  and  $P < 0.001$  both for  $\log(\text{sand})$  and  $\log(\text{SOC})$ .

The PTF was developed by merging two datasets: a set of data obtained from the samples collected by depth within the soil profile of the three study areas, as previously described, and a set of additional data obtained from a Dystric Cambisol under chestnut forest located close to the sites under study and with a similar lithology, which it is known to have relevant influence on soil BD (Patton et al., 2019), and soil type.

The BD values of both datasets (Table S2 of the Supplementary Materials) are in accordance with those reported in literature for similar pedological conditions (Heinze et al., 2018; De Feudis et al., 2020a; De Feudis et al., 2021b).

#### 2.4. Particle size distribution and organic carbon content and stock

The soil samples collected from the soil profiles and by mini-samplers were air-dried and sieved through a 2-mm mesh. The particle size distribution was determined by the pipette method (Gee and Bauder, 1986). SOC content was determined by loss-on-ignition method according to Schulte and Hopkins (1996) and Cambardella et al. (2001).

The C stock [ $\text{Mg ha}^{-1}$ ] of each horizon and fixed depth layers was calculated by the following equation (FAO, 2019):

$$C_{\text{stock}} = \text{SOC} \times \text{BD} \times (1 - V_g) \times th \times 0.1 \quad [2]$$

where  $\text{SOC}$  is the organic carbon content ( $\text{g kg}^{-1}$ ),  $\text{BD}$  is the mass of the fine earth per volume of fine earth ( $\text{g fine earth cm}^{-3}$  fine earth),  $V_g$  is the coarse fragment content ( $\text{Ø} > 2 \text{ mm}$ ) expressed as percent by volume evaluated on the soil profiles,  $th$  is the thickness of the considered soil horizon or fixed depth layer (cm),  $0.1$  is the conversion factor for converting  $\text{mg C cm}^{-2}$  to  $\text{Mg C ha}^{-1}$ .

To test the data transferability from soil horizon sampling procedure to soil fixed depth layers, the data of SOC content and stock from soil horizons were also expressed referring to soil fixed depth as weighted average based on soil horizon thickness.

### 2.5. Statistical analysis

The graphical analysis of residuals was used to verify the normality and homoscedasticity of the data (Schützenmeister et al., 2012). If the assumptions of normality and homoscedasticity were not satisfied, the data were transformed according to the Box and Cox (1964) procedure. Then, within each soil horizon, one-way analysis of variance (ANOVA) was carried out to assess the effect of forest type on soil horizons thickness, SOC content, C stock and amount of sand, silt and clay. On data from soil horizons and fixed depth layers, two-way ANOVA was applied to assess the effect of soil sampling approach (soil horizons vs fixed depth layers) and forest type on SOC content and C stock.

The comparison between the means was performed with Tukey's honest significant differences with a significance level of 0.05. In order to evaluate the short-scale variability of C stock in each study site, the coefficient of variation (CV) was calculated (Rosemary et al., 2017) for soil horizons and 0–15 and 15–30 cm soil depth layers. The statistical analyses were performed using the R 4.1.1 statistical software.

## 3. Results

### 3.1. Pedogenic horizons in 0–30 cm soil depth

Soils of the selected study sites showed a similar soil horizon sequence (Oe, A, AB and Bw) in the upper 30 cm of soil with the exception of Norway spruce forest where also an Oa horizon was detected (Table S1 of the Supplementary Materials and Figure 4).

All study sites had a sandy loam texture (Figure 5). The A and AB horizons of soils under chestnut and Norway spruce forests showed the lowest sand content and the highest silt content (Figure 5a,

b). The amount of clay did not differ among the three forest types (87 and 111 g kg<sup>-1</sup> on average, for A and AB horizons, respectively; Figure 5c). In Bw horizon, soil under chestnut forest showed the lowest sand content and the highest silt content (369 and 485 g kg<sup>-1</sup>, respectively; Figure 5a, b). Clay in Bw horizon showed some differences between chestnut and spruce forests (Figure 5c); in particular we found higher clay content under the former forest type than in the latter one (146 and 107 g kg<sup>-1</sup>, respectively).

SOC concentration always showed the lowest values under chestnut forest (Figure 6a). Norway spruce and European beech forests showed similar SOC content in Oe and Bw horizons, while higher SOC content was found in beech than in Norway spruce forest for A and AB horizons (Figure 6a).

Taking in consideration the total amount of organic carbon stored in each soil horizon, chestnut forest showed the lowest C stock in Oe (3.2 Mg ha<sup>-1</sup>), A (14.9 Mg ha<sup>-1</sup>) and AB (16.4 Mg ha<sup>-1</sup>) horizons. No differences occurred between Norway spruce and beech forests where the C stock in Oe, A and AB horizons were on average 9.0, 31.0 and 41.6 Mg ha<sup>-1</sup>, respectively (Figure 6b).

### 3.2. Soil organic carbon stock and content in 0–15 and 15–30 cm soil depths

In Figure 7, the values of SOC content and C stock observed in 0–15 and 15–30 cm soil depths sampled through the fixed depth interval approach and those calculated using the data of each identified soil horizon were reported. At both soil depths, soil under chestnut forest showed the lowest SOC content, while no SOC content differences occurred between Norway spruce and European beech forests (Figure 7a, b). For C stock in 0–15 cm soil depth some few differences between the two sampling approaches were observed (Figure 7c). Through the FDL sampling, the lowest C stock was found in soil under chestnut forest, while the highest one under Norway spruce forest. Through the calculated values from PGH sampling, the lowest C stock was observed under chestnut forest, but no differences occurred between European beech and Norway spruce forests.

For C stock in 15–30 cm soil depth, through both sampling approaches, the highest C stock was observed in soil under Norway spruce forest (Figure 7d).

### *3.3. Short-scale variability of soil organic carbon stock*

The coefficient of variation (CV) showed higher values when the soil sampling was performed considering the soil horizons compared to the values obtained through dividing soil samples in the two depth intervals (Table 1). The differences of the CV between the two approaches were pronounced for the surface layer. The CV tended to decrease from Oe to Bw horizon in chestnut and Norway spruce forests. Under beech forest the CV values did not change with exception of AB horizon where an increase was observed. Taking in consideration the fixed depth layers, the CV decreased in Norway spruce forest and negligible differences were found for chestnut and beech forests (Table 1).

Comparing the study sites, the Oe horizon showed the highest CV in chestnut forest, while the lowest one was observed in beech forest. The CV in A horizon showed the trend Norway spruce > chestnut > beech, but no differences occurred for the AB horizon. In Bw horizon the CV showed the highest value under beech forest while the lowest one under chestnut forest. Compared to the horizons, the CV values of 0–15 and 15–30 cm layers showed weaker differences among the study sites. For the 0–15 cm layer the highest CV was found in Norway spruce forest and the lowest one in chestnut forest. The trend beech > Norway spruce > chestnut trend was observed for the 15–30 cm soil depth (Table 1).

## **4. Discussion**

### *4.1. Quantity of soil organic carbon storage by horizon sampling*

As showed by the soil sampling by PGH, in the 0–30 cm soil depth interval below Oi horizon, the Oe horizon was present in all sites as well as A–AB–Bw sequence in mineral soil (Figure 4). Under Norway spruce, the Oa horizon underlie the Oe one. Under Norway spruce forest, the sampling by

PGH pointed out how the organic material tends to accumulate on soil surface as organic horizon likely due to the chemical recalcitrance to microbial attack of coniferous plant residues (Cha et al., 2019; Díaz-Pinés et al., 2011). To this regard, in the literature is reported a higher C:N ratio of Norway spruce litter layer compared to that of beech and chestnut forests (Cremer et al., 2016; Sariyildiz and Anderson, 2005) indicating a lower degradability of conifers compared to broadleaves (Massaccesi et al., 2020) and explaining the highest storage of organic C in the 0–30 cm of soil depth under Norway spruce, as observed also in a study conducted in similar ecozone by Vittori Antisari et al. (2015).

The lower C stock in the Oe, A and AB horizons under chestnut forest compared to Norway spruce and European beech forests could be also attributed to climatic factors (Figure 6b). In fact, compared to the chestnut forest, Norway spruce and European beech forests were located at higher altitude where a lower mean air temperature limited the microbial activity and promote SOC storage (De Feudis et al., 2019; Massaccesi et al., 2020). Although in chestnut forest we observed the lowest C stock (Figure 7c, d), PGH approach allowed to observe a similar thickness of A and AB horizons (Figure 4). This fact would indicate that chestnut forest promoted organic carbon deepening (De Feudis et al., 2020b; Gautam et al., 2017), independently from soil organic matter turnover, likely due to the high appetibility of chestnut residues for soil fauna and microbes (Anderson, 1973). The deepening of organic matter in soil can be considered a positive aspect because it could prevent soil degradation due to the presence of stable SOC forms in depth (e.g., Kramer et al., 2017; Soucémariadin et al., 2018).

Compared to Norway spruce forest, located at similar altitude, beech forest had a lower C stock in 0–30 cm soil depth (Figure 7c, d), although in A and AB horizons the concentration of SOC was higher in beech forest than in Norway spruce soils (Figure 6a). The lower soil C stock under beech has been mainly attributed to the high amount of skeleton found between 10 and 30 cm depth (see Table S1 of the Supplementary Materials) which lowered the C stock (Rytter, 2012).

Accepted Article

Sampling by PGH allowed us to detect that a large part of the soil organic pool of Norway spruce forest had a substantial proportion of weakly stabilised components, as observed by the soil horization. In fact, the presence of a Oa horizons, rich in SOC scarcely associated to mineral particles, made the spruce ecosystem more susceptible than broadleaves forest to SOC reduction following natural or anthropic disturbances (climatic change, wildfires, forest cutting and land use change) and, hence, soil resilience decline (Bonetti et al., 2017).

#### 4.2. Comparison between sampling methods

In the three forest types, the lack of differences observed for C stock in 0–30 soil depth between the sampling by PGH and by FDL (Figure 7c, d) demonstrated the reliability of both methodologies for SOC estimation. Our findings are in contrast to those observed by Francaviglia et al. (2017) in vineyards and by Parras-Alcántara et al. (2015) in Mediterranean natural areas. Both studies reported a higher C stock estimation through FDL sampling than through PGH sampling. The authors attributed the higher C stock by FDL to the mixing of horizons through such sampling approach which affects BD and gravel content. In this sense, because of the pivotal role of the amount of skeleton and BD on C stock estimation, in the present study the relative volume occupied by the skeleton was considered and the BD was calculated by PTF built-up through data of soils located in the same study area. It is noteworthy that to neglect the relative volume occupied by stones and gravels in C stock calculation cause the overestimation of the amount of organic C stored in soil (Rytter, 2012; Wang et al., 2018), although some authors reported skeleton itself contain organic C (Agnelli et al., 2002, 2008) and should be considered for C stock estimation (Corti et al., 2002). However, since one of the aims of the present study was to compare two soil sampling approaches for C stock estimation, and skeleton was found only under beech forest within the 0–30 soil depth, we evaluated the relative volume occupied by skeleton but not its organic C content.

The data on SOC storage, calculated by both PGH and FDL sampling, suggested the key combined effect of the organic residues' quality (Sariyildiz, 2003; 2008) and climate (Massaccesi et al., 2020;



Kumar et al., 2021) on C stock. The sampling by PGH pointed out that under Norway spruce forest the organic material tends to accumulate on soil surface as organic horizon. Such results are in accordance with previous studies (e.g., Labaz et al., 2014; Kuznetsova et al., 2019) which found a greater organic horizon thickness under coniferous than under broadleaved forests. The findings obtained through the sampling by FDL would indicate a positive effect of conifers on SOC storage (Figure 7c, d), but the “reading” of the soil horizonation, as occurred with PGH sampling, allowed to assess that SOC under spruce forest was greatly stored in the organic horizons (Oe and Oa), where it accounted for about one third of the 0-30 cm C stock (Figure 6b), because of the recalcitrant nature of the spruce litter. The noticeable storage of organic matter in the organic horizons rather than in the mineral ones, as occurred for beech and chestnut forests, makes the spruce forest susceptible to lose relevant amount of SOC in case of wildfires (Poirier et al., 2014), which have increased dramatically in frequency and extent in the European Mediterranean region (Oliveira et al., 2018). Further, the poorly degradable litter together with the siliceous substrate of our study sites might promote the formation of Mor humipedon (Zanella et al., 2018) which is more vulnerable to erosion compared to Mull, Moder and Amphi humipedon types (Pintaldi et al., 2018). Hence, in a view of SOC monitoring, our findings demonstrated that the sampling by PGH draws a better picture of SOC distribution along depth and its potential susceptibility to external factors leading to degradation.

Besides the large information about soil processes that the sampling by PGH can provide (Jiang et al., 2021), through this approach also the information about the spatial variability within each study site is preserved. In fact, we noted higher values of CV when soil horizons were considered compared to fixed depth intervals and this difference was marked for the 0–15 cm soil depth interval (Table 1). The loss of information on spatial variability would indicate the low reliability of FDL sampling to support decision-making plans addressed for sustainable use and preservation actions of soil and forest resource (Denton et al., 2017; Kirchen et al., 2017). Indeed, accurate

information about soil variability is important in environmental modelling and prediction (Zeraatpisheh et al., 2019; Rosemary et al., 2017; Taghizadeh-Mehrjardi et al., 2016).

## **5. Conclusion**

The overall data of the present study suggested that both the soil sampling by PGH and by FDL could be considered reliable for C stock examination in forest ecosystems. However, it is necessary to specify that carefulness should be paid for soil BD and skeleton amount when C stock is calculated. Our findings highlighted that, although both sampling approaches resulted appropriate to evaluate the effects of biotic and abiotic factors (e.g., vegetation and climate, respectively) on C stock, the sampling by PGH provides information about SOC processes. In fact, through the identification and characterization of soil horizons it is possible to predict how and how much the considered biotic and abiotic factors influence C stock and pools. The present study showed that the evaluation of soil properties (i.e., C stock) through soil depth layers provided a lower information about spatial variability inside the selected areas compared to the same evaluation through soil horizon sampling. Our results drawn a clear picture where sampling by PGH is appreciable in order to gain knowledge useful both for scientific purposes and a sustainable land use management of natural ecosystems.

## **Conflict of interest**

The authors declare no conflict of interest.

## **References**

- Adhikari, K., Hartemink, A.E., 2016. Linking soils to ecosystem services - A global review. *Geoderma* 262, 101–111. DOI: 10.1016/j.geoderma.2015.08.009.
- Agnelli, A., Celi, L., Degl’Innocenti, A., Corti, G., Ugolini F.C. 2002. The changes with depth of humic and fulvic acids extracted from fine earth and rock fragments of a forest soil. *Soil*

Science 167, 524-538. DOI: 10.1097/00010694-200208000-00004.

Agnelli, A., Celi, L., Corti, G., Condello, L. 2008. Organic matter stabilization in soil aggregates and rock fragments as revealed by low-temperature ashing (LTA) oxidation. *Soil Biol. Biochem.* 40, 1379-1389. DOI: 10.1016/j.soilbio.2007.12.008.

Alewell, C., Meusburger, K., Brodbeck, M., Bänninger, D., 2008. Methods to describe and predict soil erosion in mountain regions. *Landsc. Urban Plan.*, 88, 46-53. DOI: 10.1016/j.landurbplan.2008.08.007.

Anderson, J.M., 1973. The breakdown and decomposition of sweet chestnut (*Castanea sativa* Mill.) and beech (*Fagus sylvatica* L.) leaf litter in two deciduous woodland soils. *Oecologia* 12, 275-288. DOI: 10.1007/BF00347567.

Batjes, N.H., 2014. Total carbon and nitrogen in the soils of the world. *Eur. J. Soil Sci.* 65, 10–21. DOI: 10.1111/ejss.12114\_2.

Baveye, P.C., Baveye, J., Gowdy, J., 2016. Soil “ecosystem” services and natural capital: critical appraisal of research on uncertain ground. *Frontiers in Environmental Science* 4, 1-49. DOI: 10.3389/fenvs.2016.00041.

Bertolini, G., Guida, M., Pizziolo, M., 2005. Landslides in Emilia-Romagna region (Italy): Strategies for hazard assessment and risk management. *Landslides* 2, 302–312. DOI: 10.1007/s10346-005-0020-1.

Bojko, O., Kabala, C., 2016. Transformation of physicochemical soil properties along a mountain slope due to land management and climate changes — a case study from the Karkonosze Mountains, SW Poland. *Catena* 140, 43-54. DOI: 10.1016/j.catena.2016.01.015.

Bonetti, J. de A., Anghinoni, I., Moraes, M.T., Fink, J.R., 2017. Resilience of soils with different texture, mineralogy and organic matter under long-term conservation systems. *Soil Tillage Res.* 174, 104-112. DOI: 10.1016/j.still.2017.06.008.

Box, G.E.P., Cox, D.R., 1964. Analysis of transformations. *J. R. Stat. Soc. Ser. B Stat Methodol.* 26, 211–252. DOI: 10.1111/j.2517-6161.1964.tb00553.x.

- Accepted Article
- Brang, P., Schönenberger, W., Frehner, M., Schwitter, R., Thormann, J.J., Wasser, B., 2006. Management of protection forests in the European Alps: An overview. *For. Snow Landsc. Res.* 80, 23–44.
- Caddeo, A., Marras, S., Sallustio, L., Spano, D., Sirca, C., 2019. Soil organic carbon in Italian forests and agroecosystems: Estimating current stock and future changes with a spatial modelling approach. *Agric. For. Meteorol.* 278. DOI: 10.1016/j.agrformet.2019.107654.
- Cambardella, C.A., Gajda, A.M., Doran, J.W., Wienhold, B.J., Kettler, T.A., 2001. Estimation of particulate and total organic matter by weight loss-on-ignition. In: *Assessment methods for soil carbon (advances in soil science)* (pp. 349-359).
- Canedoli, C., Ferrè, C., Abu El Khair, D., Comolli, R., Liga, C., Mazzucchelli, F., Proietto, A., Rota, N., Colombo, G., Bassano, B., Viterbi, R., Padoa-Schioppa, E., 2020. Evaluation of ecosystem services in a protected mountain area: Soil organic carbon stock and biodiversity in alpine forests and grasslands. *Ecosyst. Serv.* 44. DOI: 10.1016/j.ecoser.2020.101135.
- Cesaro, L., Romano, R., 2017. Forestry in Italy: state of health and management. A challenge for the future.
- Cha, J.Y., Cha, Y.K., Oh, N.H., 2019. The effects of tree species on soil organic carbon content in South Korea. *J. Geophys. Res. Biogeosciences* 124, 708–716. DOI: 10.1029/2018JG004808.
- Chen, S., Richer-de-Forges, A.C., Saby, N.P.A., Martin, M.P., Walter, C., Arrouays, D., 2018. Building a pedotransfer function for soil bulk density on regional dataset and testing its validity over a larger area. *Geoderma* 312, 52–63. DOI: 10.1016/j.geoderma.2017.10.009.
- Choudhury, B.U., Fiyaz, A.R., Mohapatra, K.P., Ngachan, S., 2016. Impact of land uses, agrophysical variables and altitudinal gradient on soil organic carbon concentration of North-Eastern Himalayan Region of India. *L. Degrad. Dev.* 27, 1163–1174. DOI: 10.1002/ldr.2338.
- Cislaghi, A., Alterio, E., Fogliata, P., Rizzi, A., Lingua, E., Vacchiano, G., Bischetti, G.B., Sitzia, T., 2021. Effects of tree spacing and thinning on root reinforcement in mountain forests of the European Southern Alps. *For. Ecol. Manage.* 482. DOI: 10.1016/j.foreco.2020.118873.

- Conforti, M., Buttafuoco, G., Leone, A.P., Aucelli, P.P.C., Robustelli, G., Scarciglia, F., 2013. Studying the relationship between water-induced soil erosion and soil organic matter using Vis–NIR spectroscopy and geomorphological analysis: a case study in a southern Italy area. *Catena*, 110, 44-58. DOI: 10.1016/j.catena.2013.06.013.
- Cools, N., De Vos, B., 2020. Part X: Sampling and Analysis of Soil. In: UNECE ICP Forests Programme Co-ordinating Centre (ed.): Manual on methods and criteria for harmonized sampling, assessment, monitoring and analysis of the effects of air pollution on forests.
- Corti, G., Ugolini, F.C., Agnelli, A., Certini, G., Cuniglio, R., Berna, F., Fernández Sanjurjo, M.J., 2002. The soil skeleton, a forgotten pool of carbon and nitrogen in soil. *Eur. J. Soil Sci.* 53, 283–298. DOI: 10.1046/j.1365-2389.2002.00442.x.
- Cremer, M., Kern, N.V., Prietzel, J., 2016. Soil organic carbon and nitrogen stocks under pure and mixed stands of European beech, Douglas fir and Norway spruce. *For. Ecol. Manage.* 367, 30–40. DOI: 10.1016/j.foreco.2016.02.020.
- De Feudis, M., Cardelli, V., Massaccesi, L., Trumbore, S.E., Vittori Antisari, L., Cocco, S., Corti, G., Agnelli, A., 2019. Small altitudinal change and rhizosphere affect the SOM light fractions but not the heavy fraction in European beech forest soil. *Catena* 181, article 104091. DOI: 10.1016/j.catena.2019.104091.
- De Feudis, M., Falsone, G., Gherardi, M., Speranza, M., Vianello, G., Vittori Antisari, L., 2021a. GIS-based soil maps as tools to evaluate land capability and suitability in a coastal reclaimed area (Ravenna, northern Italy). *Int. Soil Water Conserv. Res.* 9, 167-179. DOI: 10.1016/j.iswcr.2020.11.007.
- De Feudis, M., Falsone, G., Vianello, G., Vittori Antisari, L., 2020a. The conversion of abandoned chestnut forests to managed ones does not affect the soil chemical properties and improves the soil microbial biomass activity. *Forests* 11(8), 786. DOI: 10.3390/f11080786.
- De Feudis, M., Falsone, G., Vianello, G., Vittori Antisari, L., 2020b. Stable organic carbon pool rises in soil under chestnut (*Castanea sativa* Mill.) forest for timber production after 15 years

since grafting onto satin-cut stumps. *EQA - Int. J. Environ. Qual.* 40, 1–10. DOI: 10.6092/issn.2281-4485/10731.

De Feudis, M., Falsone, G., Vittori Antisari, L., 2021b. Mid-term (30 years) changes of soil properties under chestnut stands due to organic residues management: An integrated study. *Catena* 198, article 105021. DOI: 10.1016/j.catena.2020.105021.

Denton, O.A., Aduramigba-Modupe, V.O., Ojo, A.O., Adeoyolanu, O.D., Are, K.S., Adelana, A.O., Oyedele, A.O., Adetayo, A.O., Oke, A.O., 2017. Assessment of spatial variability and mapping of soil properties for sustainable agricultural production using geographic information system techniques (GIS). *Cogent Food Agric.* 3, 1279366. DOI: 10.1080/23311932.2017.1279366.

Díaz-Pinés, E., Rubio, A., Van Miegroet, H., Montes, F., Benito, M., 2011. Does tree species composition control soil organic carbon pools in Mediterranean mountain forests? *For. Ecol. Manage.* 262, 1895–1904. DOI: 10.1016/j.foreco.2011.02.004.

Dixon, R.K., Solomon, A.M., Brown, S., Houghton, R.A., Trexler, M.C., Wisniewski, J., 1994. Carbon pools and flux of global forest ecosystems. *Science* 263, 185-190. DOI: 10.1126/science.263.5144.185.

Don, A., Schumacher, J., Freibauer, A., 2011. Impact of tropical land-use change on soil organic carbon stocks: a meta-analysis. *Glob. Chang. Biol.*, 17, 1658-1670. DOI: 10.1111/j.1365-2486.2010.02336.x.

FAO, 2010. Global forest resources assessment 2010, Main report, FAO Forestry Paper 163. Rome, Italy.

FAO, 2019. Measuring and modelling soil carbon stocks and stock changes in livestock production systems: Guidelines for assessment (Version 1). In *Livestock environmental assessment and performance (LEAP) partnership*. Rome, Italy: FAO. 170 pp. Licence: CC BY-NC-SA 3.0 IGO.

FAO, 2015. *Understanding Mountain Soils: A contribution from mountain areas to the International*

Year of Soils 2015, by Romeo, R., Vita, A., Manuelli, S., Zanini, E., Freppaz, M. & Stanchi, S. Rome, Italy. Available from: [https://www.researchgate.net/publication/282150416\\_Understanding\\_Mountain\\_Soils\\_A\\_contribution\\_from\\_mountain\\_areas\\_to\\_the\\_International\\_Year\\_of\\_Soils\\_2015](https://www.researchgate.net/publication/282150416_Understanding_Mountain_Soils_A_contribution_from_mountain_areas_to_the_International_Year_of_Soils_2015)[accessed Nov 10 2021].

FAO. World Reference base for Soil Resources 2014. In: International Soil Classification System for Naming Soils and Creating Legends for Soil Maps; FAO: Rome, Italy, 2014.

FAOSTAT, 2020. Food and Agriculture Organization of the United Nations, 2017. Food and Agriculture Organization of the United Nations – Statistic division.

Francaviglia, R., Renzi, G., Doro, L., Parras-Alcántara, L., Lozano-García, B., Ledda, L., 2017. Soil sampling approaches in Mediterranean agro-ecosystems. Influence on soil organic carbon stocks. *Catena* 158, 113–120. DOI: 10.1016/j.catena.2017.06.014.

Gautam, M.K., Lee, K.S., Song, B.Y., Bong, Y.S., 2017. Site related  $\delta^{13}\text{C}$  of vegetation and soil organic carbon in a cool temperate region. *Plant Soil* 418, 293–306. DOI: 10.1007/s11104-017-3284-z.

Gee, G.W., Bauder, J.W., 1986. *Methods of Soil Analysis: Part 1—Physical and Mineralogical Methods*; SSSA Book Series; Soil Science Society of America: Madison, WI, USA; American Society of Agronomy: Madison, WI, USA; Volume sssabookse, ISBN 978-0-89118-864-3.

Gross, C.D., James, J.N., Turnblom, E.C., Harrison, R.B., 2018. Thinning treatments reduce deep soil carbon and nitrogen stocks in a coastal pacific northwest forest. *Forests* 9 (5), 238. DOI: 10.3390/f9050238.

Grüneberg, E., Schöning, I., Kalko, E.K.V., Weisser, W.W., 2010. Regional organic carbon stock variability: A comparison between depth increments and soil horizons. *Geoderma* 155, 426–433. DOI: 10.1016/j.geoderma.2010.01.002.

Grüneberg, E., Ziche, D., Wellbrock, N., 2014. Organic carbon stocks and sequestration rates of forest soils in Germany. *Glob. Chang. Biol.* 20, 2644-2662. DOI: 10.1111/gcb.12558.

Guerra, C.A., Rosa, I.M., Valentini, E., Wolf, F., Filipponi, F., Karger, D.N., Eisenhauer, N., 2020.

Global vulnerability of soil ecosystems to erosion. *Landscape Ecol.*, 1–20. DOI: 10.1007/s10980-020-00984-z.

Hagedorn, F., Mulder, J., Jandl, R., 2010. Mountain soils under a changing climate and land-use.

*Biogeochemistry* 97, 1–5. DOI: 10.1007/s10533-009-9386-9.

Heinze, S., Ludwig, B., Piepho, H., Mikutta, R., Don, A., Wordell-dietrich, P., Helfrich, M., Hertel,

D., Leuschner, C., Kirfel, K., Kandeler, E., Preusser, S., Guggenberger, G., Leinemann, T.,

Marschner, B., 2018. Factors controlling the variability of organic matter in the top- and

subsoil of a sandy Dystric Cambisol under beech forest. *Geoderma* 311, 37-44. DOI:

10.1016/j.geoderma.2017.09.028.

Hiederer, R., Micheli, E., Durrant, T., 2011. Evaluation of the BioSoil demonstration project Soil

Data Analysis, JRC Scientific and Technical Reports, Publication Office of the European

Union

IPCC, 2003. Good Practice Guidance for Land Use, Land Use Change and Forestry. Penman, J.,

Gytarsky, M., Hiraishi, T., Krug, T., Kruger, D., Pipatti, R., Buendia, L., Miwa, K., Ngara, T.,

Tanabe, K., Wagner, F., (Eds). IPCC/OECD/IEA/IGES, Hayama, Japan

IPCC, 2013: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I

to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker,

T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex

and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New

York, NY, USA, pp 1535.

IPCC, 2018. Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming

of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in

the context of strengthening the global response to the threat of climate change, sustainable

development, and efforts to eradicate poverty [Masson-Delmotte, V., Zhai, P., Pörtner, H.-O.,

Roberts, D., Skea, J., Shukla, P.R., Pirani, A., Moufouma-Okia, W., Péan, C., Pidcock, R.,



Connors, S., Matthews, J.B.R., Chen, Y., Zhou, X., Gomis, M.I., Lonnoy, E., Maycock, T., Tignor, M., and Waterfield T. (eds.)].

Jiang, Z.-D., Owens, P.R., Zhang, C.-L., Brye, K.R., Weindorf, D.C., Adhikari, K., Sun, Z.-X., Sun, F.-J., Wang, Q.-B., 2021. Towards a dynamic soil survey: Identifying and delineating soil horizons in-situ using deep learning. *Geoderma* 401, article 115341. DOI: 10.1016/j.geoderma.2021.115341.

Jobbágy, E., Jackson, R., 2000. The vertical distribution of soil organic carbon and its relation to climate and vegetation. *Ecol. Appl.* 10, 423–436. DOI: 10.1890/1051-0761(2000)010[0423:TVDOSO]2.0.CO;2.

Koga, N., Shimoda, S., Shirato, Y., Kusaba, T., Shima, T., Niimi, H., Yamane, T., Wakabayashi, K., Niwa, K., Kohyama, K., Obara, H., Takata, Y., Kanda, T., Inoue, H., Ishizuka, S., Kaneko, S., Tsuruta, K., Hashimoto, S., Shinomiya, Y., Aizawa, S., Ito, E., Hashimoto, T., Morishita, T., Noguchi, K., Ono, K., Katayanagi, N., Atsumi, K., 2020. Assessing changes in soil carbon stocks after land use conversion from forest land to agricultural land in Japan. *Geoderma* 377, Article 114487. DOI: 10.1016/j.geoderma.2020.114487.

Kirchen, G., Calvaruso, C., Granier, A., Redon, P.O., Van der Heijden, G., Bréda, N., Turpault, M.P., 2017. Local soil type variability controls the water budget and stand productivity in a beech forest. *For. Ecol. Manage.* 390, 89–103. DOI: 10.1016/j.foreco.2016.12.024.

Kramer, M.G., Lajtha, K., Audfenkampe, A., 2017. Depth trends of soil organic matter C:N and <sup>15</sup>N natural abundance controlled by association with minerals. *Biogeochemistry* 136, 237–248. DOI: 10.1007/s10533-017-0378-x.

Kumar, A., Kumar, M., Pandey, R., Guo, Y.Z., Cabral-Pinto, M., 2021. Forest soil nutrient stocks along with an altitudinal range of Uttarakhand Himalayas: An aid to Nature based climate Solutions. *Catena* 207, article 105667. DOI: 10.1016/j.catena.2021.105667.

Kuznetsova, A.I., Lukina, N.V., Tikhonova, E.V., Gornov, A.V., Gornova, M.V., Smirnov, V.E., Geraskina, A.P., Shevchenko, N.E., Tebenkova, D.N., Chumachenko, S.I., 2019. Carbon stock

in sandy and loamy soils of coniferous–broadleaved forests at different succession stages. *Eurasian Soil Sci.* 52, 756–768. DOI: 10.1134/S1064229319070081.

Labaz, B., Galka, B., Bogacz, A., Waroszewski, J., Kabala, C., 2014. Factors influencing humus forms and forest litter properties in the mid-mountains under temperate climate of southwestern Poland. *Geoderma* 230–231, 265–273. DOI: 10.1016/j.geoderma.2014.04.021.

Li, Y.C., Li, Y.F., Chang, S.X., Liang, X., Qin, H., Chen, J.H., Xu, Q.F., 2017. Linking soil fungal community structure and function to soil organic carbon chemical composition in intensively managed subtropical bamboo forests. *Soil Biol. Biochem.* 107, 19–31. DOI: 10.1016/j.soilbio.2016.12.024.

Lull, C., Bautista, I., Lidón, A., del Campo, A.D., González-Sanchis, M., García-Prats, A., 2020. Temporal effects of thinning on soil organic carbon pools, basal respiration and enzyme activities in a Mediterranean Holm oak forest. *For. Ecol. Manage.* 464, article 118088. DOI: 10.1016/j.foreco.2020.118088.

Makipaa, R., Liski, J., Guendehou, S., Malimbwi, R., Kaaya, A., 2012. Soil carbon monitoring using surveys and modelling: General description and application in the United Republic of Tanzania. Food and Agriculture Organization of the United Nations, Rome.

Miao, X., Van Zwieten, L., Zhang, M., Qiu, Z., Yao, Y., Wang, H., 2020. Soil parent material controls organic matter stocks and retention patterns in subtropical China. *J. Soils Sediments* 20, 2426–2438. DOI:10.1007/s11368-020-02578-3.

Massaccesi, L., De Feudis, M., Leccese, A., Agnelli, A., 2020. Altitude and vegetation affect soil organic carbon, basal respiration and microbial biomass in apennine forest soils. *Forests* 11, 1–13. DOI: doi.org/10.3390/f11060710.

Minasny, B., Malone, B.P., McBratney, A.B., Angers, D.A., Arrouays, D., Chambers, A., Chaplot, V., Chen, Z.S., Cheng, K., Das, B.S., Field, D.J., Gimona, A., Hedley, C.B., Hong, S.Y., Mandal, B., Marchant, B.P., Martin, M., McConkey, B.G., Mulder, V.L., O'Rourke, S., Richer-de-Forges, A.C., Odeh, I., Padarian, J., Paustian, K., Pan, G., Poggio, L., Savin, I.,

Stolbovoy, V., Stockmann, U., Sulaeman, Y., Tsui, C.C., Vågen, T.G., van Wesemael, B., Winowiecki, L., 2017. Soil carbon 4 per mille. *Geoderma* 292, 59–86. DOI: 10.1016/j.geoderma.2017.01.002.

Minasny, B., McBratney, A.B., 2018. Limited effect of organic matter on soil available water capacity. *Eur. J. Soil Sci.* 69, 39–47. DOI: 10.1111/ejss.12475.

Oliveira, S., Félix, F., Nunes, A., Lourenço, L., Laneve, G., Sebastián-López, A., 2018. Mapping wildfire vulnerability in Mediterranean Europe. Testing a stepwise approach for operational purposes. *J. Environ. Manag.* 206, 158-169. DOI: 10.1016/j.jenvman.2017.10.003.

Orton, T.G., Pringle, M.J., Bishop, T.F.A., 2016. A one-step approach for modelling and mapping soil properties based on profile data sampled over varying depth intervals. *Geoderma* 262, 174–186. DOI: 10.1016/j.geoderma.2015.08.013.

Parras-Alcántara, L., Lozano-García, B., Brevik, E.C., Cerdá, A., 2015. Soil organic carbon stocks assessment in Mediterranean natural areas: A comparison of entire soil profiles and soil control sections. *J. Environ. Manage.* 155, 219–228. DOI: 10.1016/j.jenvman.2015.03.039.

Patton, N.R., Lohse, K.A., Seyfried, M., Will, R., Benner, S.G., 2019. Lithology and coarse fraction adjusted bulk density estimates for determining total organic carbon stocks in dryland soils. *Geoderma* 337, 844–852. DOI: 10.1016/j.geoderma.2018.10.036.

Pintaldi, E., D'Amico, M.E., Stanchi, S., Catoni, M., Freppaz, M., Bonifacio, E., 2018. Humus forms affect soil susceptibility to water erosion in the Western Italian Alps. *Appl. Soil Ecol.* 123, 478-483. DOI: 10.1016/j.apsoil.2017.04.007.

Poirier, V., Paré, D., Boiffin, J., Munson, A.D., 2014. Combined influence of fire and salvage logging on carbon and nitrogen storage in boreal forest soil profiles. *For. Ecol. Manag.* 326, 133-141. DOI: 10.1016/j.foreco.2014.04.021.

Rabbi, S.M.F., Tighe, M.K., Flavel, R.J., Kaiser, B.N., Guppy, C.N., Zhang, X., Young, I.M., 2018. Plant roots redesign the rhizosphere to alter the three-dimensional physical architecture and water dynamics. *New Phytol.* 219, 542–550. DOI: 10.1111/nph.15213.

- Rosemary, F., Vitharana, U.W.A., Indraratne, S.P., Weerasooriya, R., Mishra, U., 2017. Exploring the spatial variability of soil properties in an Alfisol soil catena. *Catena* 150, 53–61. DOI: 10.1016/j.catena.2016.10.017.
- Rytter, R.M., 2012. Stone and gravel contents of arable soils influence estimates of C and N stocks. *Catena* 95, 153–159. DOI: 10.1016/j.catena.2012.02.015.
- Sariyildiz, T., 2003. Litter decomposition of *Picea orientalis*, *Pinus sylvestris* and *Castanea sativa* trees grown in Artvin in relation to their initial litter quality variables. *Turk. J. Agric. For.* 27, 237-243.
- Sariyildiz, T., 2008. Effects of gap-size classes on long-term litter decomposition rates of beech, oak and chestnut species at high elevations in Northeast Turkey. *Ecosystems* 11, 841-853. DOI: 10.1007/s10021-008-9164-x.
- Sariyildiz, T., Anderson, J.M., 2005. Variation in the chemical composition of green leaves and leaf litters from three deciduous tree species growing on different soil types. *For. Ecol. Manage.* 210, 303–319. DOI: 10.1016/j.foreco.2005.02.043.
- Schoeneberger, P.J., Wysocki, D.A., Benham, E.C., 2012. *Soil Survey Staff Field Book for Describing and Sampling Soils; Version 3.0*; Natural Resources Conservation Service, National Soil Survey Center: Lincoln, NE, USA.
- Schulte, E.E., Hopkins, B.G., 1996. Estimation of organic matter by weight loss-on-ignition. *Soil Organic Matter: Analysis and Interpretation*, 46, 21e31. DOI: 10.2136/sssaspecpub46.c3.
- Schützenmeister, A., Jensen, U., Piepho, H.-P., 2012. Checking normality and homoscedasticity in the general linear model using diagnostic plots. *Commun. Stat. Simulat. Comput.* 41, 141-154. DOI: 10.1080/03610918.2011.582560.
- Shahbazi, F., Hughes, P., McBratney, A.B., Minasny, B., Malone, B.P., 2019. Evaluating the spatial and vertical distribution of agriculturally important nutrients — nitrogen, phosphorous and boron — in North West Iran. *Catena* 173, 71–82. DOI: 10.1016/j.catena.2018.10.005.
- Sharififar, A., Sarmadian, F., Alikhani, H., Keshavarzi, A., Asghari, O., Malone, B.P., 2019. Lateral

and vertical variations of soil organic and inorganic carbon content in Aridisols and Entisols of a rangeland. *Eurasian Soil Sci.* 52, 1051–1062. DOI: 10.1134/S1064229319090084.

Soucémariadin, L.N., Cécillon, L., Guenet, B., Chenu, C., Baudin, F., Nicolas, M., Girardin, C., Barré, P., 2018. Environmental factors controlling soil organic carbon stability in French forest soils. *Plant Soil* 426, 267–286. DOI: 10.1007/s11104-018-3613-x.

Su, F., Xu, S., Sayer, E.J., Chen, W., Du, Y., Lu, X., 2021. Distinct storage mechanisms of soil organic carbon in coniferous forest and evergreen broadleaf forest in tropical China. *J. Environ. Manage.* 295, article 113142. DOI: 10.1016/j.jenvman.2021.113142.

Suliman, M., Saeed, I., Hassaballa, A., Rodrigo-Comino, J., 2018. Modeling cation exchange capacity in multi geochronological-derived alluvium soils: An approach based on soil depth intervals. *Catena* 167, 327–339. DOI: 10.1016/j.catena.2018.05.001.

Taghizadeh-Mehrjardi, R., Nabiollahi, K., Kerry, R., 2016. Digital mapping of soil organic carbon at multiple depths using different data mining techniques in Baneh region, Iran. *Geoderma* 266, 98–110. DOI: 10.1016/j.geoderma.2015.12.003.

Tesfaye, M.A., Bravo, F., Ruiz-Peinado, R., Pando, V., Bravo-Oviedo, A., 2016. Impact of changes in land use, species and elevation on soil organic carbon and total nitrogen in Ethiopian Central Highlands. *Geoderma* 261, 70–79. DOI: 10.1016/j.geoderma.2015.06.022.

Valtera, M., Šamonil, P., 2018. Soil organic carbon stocks and related soil properties in a primary *Picea abies* (L.) Karst. volcanic-mountain forest. *Catena* 165, 217–227. DOI: 10.1016/j.catena.2018.01.034.

Vicente-Vicente, J.L., García-Ruiz, R., Francaviglia, R., Aguilera, E., Smith, P., 2016. Soil carbon sequestration rates under Mediterranean woody crops using recommended management practices: A meta-analysis. *Agric. Ecosyst. Environ.* 235, 204–214. DOI: 10.1016/j.agee.2016.10.024.

Vittori Antisari, L., Falsone, G., Carbone, S., Marinari, S., Vianello, G., 2015. Douglas-fir reforestation in North Apennine (Italy): performance on soil carbon sequestration, nutrients

stock and microbial activity. *Appl. Soil Ecol.* 86, 82–90. DOI: 10.1016/j.apsoil.2014.09.009.

Vittori Antisari, L., Falsone, G., Carbone, S., Vianello, G., 2013. Short-term effects of forest recovery on soil carbon and nutrient availability in an experimental chestnut stand. *Biol. Fertil. Soils* 49, 165–173. DOI: 10.1007/s00374-012-0708-z.

Wang, X., Yoo, K., Wackett, A., Gutknecht, J., Amundson, R., Heimsath, A., 2018. Soil organic carbon and mineral interactions on climatically different hillslopes. *Geoderma* 322, 71-80. DOI: 10.1016/j.geoderma.2018.02.021.

Wellbrock, N., Bolte, A., 2019. Status and dynamic of forest in Germany. Springer International Publishing

Yang, S., Cammeraat, E., Jansen, B., den Haan, M., van Loon, E., Recharte, J., 2018. Soil organic carbon stocks controlled by lithology and soil depth in a Peruvian alpine grassland of the Andes. *Catena* 171, 11–21. DOI: 10.1016/j.catena.2018.06.038.

Zanella, A., Ponge, J.F., Jabiol, B., Sartori, G., Kolb, E., Gobat, J.M., Bayon, R.C. Le, Aubert, M., Waal, R. De, Delft, B. Van, Vacca, A., Serra, G., Chersich, S., Andretta, A., Cools, N., Englisch, M., Hager, H., Katzensteiner, K., Brêthes, A., Nicola, C. De, Testi, A., Bernier, N., Graefe, U., Juilleret, J., Banas, D., Garlato, A., Obber, S., Galvan, P., Zampedri, R., Frizzera, L., Tomasi, M., Menardi, R., Fontanella, F., Filoso, C., Dibona, R., Bolzonella, C., Pizzeghello, D., Carletti, P., Langohr, R., Cattaneo, D., Nardi, S., Nicolini, G., Viola, F., 2018. Humusica 1, article 4: Terrestrial humus systems and forms — Specific terms and diagnostic horizons. *Appl. Soil Ecol.* 122, 56–74. DOI: 10.1016/j.apsoil.2017.07.005.

Zeraatpisheh, M., Ayoubi, S., Jafari, A., Tajik, S., Finke, P., 2019. Digital mapping of soil properties using multiple machine learning in a semi-arid region, central Iran. *Geoderma* 338, 445-452. DOI: 10.1016/j.geoderma.2018.09.006.

**Table 1.** Coefficient of variation to describe the spatial variability of C stock calculated both for each identified soil horizon (Oe, Oa, A, AB and Bw) and soil depth layer (0–15 and 15–30 cm) of the plots selected inside chestnut, Norway spruce and European beech forests.

Forest type	Oe	Oa	A	AB	Bw	0–15	15–30
	%						
Chestnut	72.8		36.2	49.2	19.2	17.7	16.5
Norway spruce	65.3	50.3	60.4	49.8	26.0	25.7	16.9
European Beech	35.5		27.7	46.1	35.1	21.0	19.5

## Figure captions

**Figure 1.** Map of the study area showing the sampling locations (Sweet chestnut, Norway spruce and European beech forests).

**Figure 2.** Collection of samples from 0–30 cm of soil depth for soil profile and for the 8 additional sampling points by horizon (on upper left) and fixed 0–15 and 15–30 cm (on upper right), and pictures of the representative soil profiles (on the bottom) dug under Sweet chestnut, Norway spruce and European beech forests.

**Figure 3.** Sampling design scheme carried out in each study site (left side) and the engaged mini-sampler (right side). On left, the full circle indicates the representative soil pit, the empty circles are sampling points where the mini samplers were engaged.

**Figure 4.** Mean thickness of soil horizons identified in 0–30 cm depth interval under chestnut (dark grey bars), Norway spruce (white bars) and European beech (light grey bars) forests. Error bars indicate standard errors (n=9). Different letters, where included, indicate significant differences among forest types for each soil horizon (One-way ANOVA and Tukey HSD test,  $p \leq 0.05$ ).

**Figure 5.** Sand (a) silt (b) and clay (c) contents in mineral horizons identified in 0–30 cm soil depth interval under chestnut (dark grey bars), Norway spruce (white bars) and European beech (light grey bars) forests. Error bars indicate standard errors (n=9). Different letters, where included, indicate significant differences among forest types for each soil horizon (One-way ANOVA and Tukey HSD test,  $p \leq 0.05$ ).




**Figure 6.** (a) Soil organic carbon (SOC) content and (b) C stock in soil horizons identified in 0–30 cm depth interval under chestnut (dark grey bars), Norway spruce (white bars) and European beech (light grey bars) forests. Error bars indicate standard errors (n=9). Different letters, where included, indicate significant differences among forest types for each soil horizon (One-way ANOVA and Tukey HSD test,  $p \leq 0.05$ ).


**Figure 7.** Soil organic carbon (SOC) content in (a) 0–15 and (b) 15–30 cm soil depth layers, and C stock in (c) 0–15 and (d) 15–30 cm soil depth layers, under chestnut, Norway spruce (Spruce) and European beech (Beech) forests obtained through fixed depth layers (grey bars) and soil horizons (white bars) sampling approaches. Error bars indicate standard errors (n=9). Different letters, where included, indicate significant differences among the means (Two-way ANOVA and Tukey HSD test,  $p \leq 0.05$ ).  $p$  values of forest type  $\times$  sampling approach: 0.5829, 0.4810, 0.7279 and 0.6897 for barplot a, b, c and d, respectively.




Sampling by horizon




Sampling by fixed depth intervals




Sweet chestnut





Norway spruce






European beech











■ Sweet chestnut   □ Norway spruce   ■ European beech



■ Sweet chestnut    □ Norway spruce    ■ European beech

