

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

On the trail of medieval wolves: ancient DNA, CT-based analyses and palaeopathology of a 1000-year-old wolf cranium from the Po Valley (northern Italy)

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Iurino D.A., Cilli E., Caniglia R., Fabbri E., Mecozzi B., Ciucani M.M., et al. (2023). On the trail of medieval wolves: ancient DNA, CT-based analyses and palaeopathology of a 1000-year-old wolf cranium from the Po Valley (northern Italy). *HISTORICAL BIOLOGY*, 35(6), 976-987 [10.1080/08912963.2022.2071710].

Availability:

This version is available at: <https://hdl.handle.net/11585/892344> since: 2023-06-27

Published:

DOI: <http://doi.org/10.1080/08912963.2022.2071710>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Iurino, Dawid A., Elisabetta Cilli, Romolo Caniglia, Elena Fabbri, Beniamino Mecozzi, Marta Maria Ciucani, Raffaele Sardella, and Davide Persico. 2023. "On the Trail of Medieval Wolves: Ancient DNA, CT-Based Analyses and Palaeopathology of a 1000-Year-Old Wolf Cranium from the Po Valley (Northern Italy)." *Historical Biology* 35 (6): 976–87.

The final published version is available online at:

<https://doi.org/10.1080/08912963.2022.2071710>.

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

On the trail of medieval wolves: Ancient DNA, CT-based analyses and paleopathology of a 1000-year-old wolf cranium from the Po Valley (northern Italy)

Dawid A. Iurino^{a*}, Elisabetta Cilli^{b*}, Romolo Caniglia^c, Elena Fabbri^c, Beniamino Mecozzi^a,
Marta Maria Ciucani^d, Raffaele Sardella^a, Davide Persico^e

^aDipartimento di Scienze della Terra, PaleoFactory, Sapienza Università di Roma, Roma, Italy;

^bLaboratorio del DNA antico, Dipartimento di Beni Culturali, Università di Bologna, Campus di Ravenna, Italy;

^cArea per la Genetica della Conservazione BIO-CGE, Istituto Superiore per la Protezione e la Ricerca Ambientale (ISPRA), Bologna, Italy;

^dSection for Evolutionary Genomics, The GLOBE Institute, University of Copenhagen, Copenhagen, Denmark;

^eDipartimento di Scienze chimiche della vita e della sostenibilità ambientale (SCVSA), Università degli Studi di Parma, Parma, Italy;

*Corresponding authors: dawid.iurino@uniroma1.it, elisabetta.cilli@unibo.it

ORCiDs:

Dawid Adam Iurino: <http://orcid.org/0000-0002-3702-9726>

Elisabetta Cilli: <https://orcid.org/0000-0003-0407-267X>

Romolo Caniglia: <https://orcid.org/0000-0002-3641-9490>

Elena Fabbri: <https://orcid.org/0000-0002-9957-8202>

Beniamino Mecozzi: <http://orcid.org/0000-0002-7097-3217>

26 Marta Maria Ciucani: <https://orcid.org/0000-0002-8998-4899>

27 Raffaele Sardella: <http://orcid.org/0000-0002-9752-6281>

28 Davide Persico: <https://orcid.org/0000-0001-5194-9724>

29

30 **Abstract**

31 The Middle Ages represented a crucial period for the evolutionary history of wolves (*Canis*
32 *lupus*), marked by both significant ecosystem changes, especially through the degradation
33 of wooded landscapes and heavy persecution, that drove this species to a dramatic
34 demographic decline. In Europe, informative and well-documented wolf remains from the
35 Medieval Ages are exceptionally rare and are mostly represented by teeth and postcranial
36 elements. In this study, we describe a well-preserved wolf cranium dated to ca. 967 - 1157
37 AD from the Po Valley (northern Italy). The specimen was analysed through a
38 multidisciplinary approach including CT-based, ancient DNA, and palaeopathological
39 analyses. Morphological and genetic data supported the assignment of this sample to
40 *Canis lupus* species. CT-based analyses indicated a typical wolf-like morphology falling
41 into the extant variability of the medium-sized subspecies *C. lupus italicus*, whereas
42 paleopathological analyses indicated a severe periodontitis. Phylogenetic analyses
43 showed that the Po valley wolf had a unique and never described mtDNA control region
44 haplotype, testifying variability in the ancient Italian wolf, which has now been lost. This
45 study provides the first comprehensive description of a wolf from the Middle Ages, adding
46 useful information for a deeper knowledge about population dynamics, variability, and
47 diseases of this species.

48

49 **Keywords:** *Canis lupus*; Medieval period; aDNA; Genetics; Morphology; Palaeopathology.

50

51 **Introduction**

52 The wolf (*Canis lupus* Linnaeus, 1758) represents one of the most iconic predators
53 throughout human history. Its complex social behaviour, ecological adaptability and pack
54 hunting abilities acquired over ~350 Ka of evolution ([Brugal and Boudadi-Maligne 2011](#);
55 [Sardella et al. 2014](#)), make the wolf one of the most successful and widespread species of
56 the genus *Canis*. The species occurs in Eurasia and North America where it inhabits both
57 natural (e.g., forests, grasslands, Mediterranean scrub) and anthropised (e.g., agricultural
58 fields, urban peripheries) environments. In Europe, after centuries of persecution,
59 abundance declines, and spatial contractions ([Breitenmoser 1998](#)), the wolf has
60 recolonised most of its original distribution range in the Iberian, Italian and Balkan
61 Peninsulas, as well as in several areas of central-eastern and north-western countries
62 ([Chapron et al. 2014](#)). Paleontological studies attest a much wider wolf distribution during
63 the Late Pleistocene ([Brugal and Boudadi-Maligne 2011](#); [Sardella et al. 2014](#)) and
64 Holocene ([Sommer and Benecke 2005](#)) in almost all the European continent, with a
65 discrete presence even on large islands such as the Great Britain and Sicily ([Angelici and](#)
66 [Rossi 2018](#)). In particular, paleontological investigations on the genus *Canis* have
67 significantly contributed also to define several morphological ([Sardella et al. 2014](#); [Mecozzi](#)
68 [and Bartolini Lucenti 2018](#); [Mecozzi et al. 2017](#); [Mecozzi et al. 2020](#)) and chronological
69 ([Brugal and Boudadi-Maligne 2011](#)) aspects concerning the evolution of Quaternary
70 European wolves.

71 For example, Italian fossil records of *C. lupus*, consisting of thousands of cranial and
72 postcranial remains spanning from the Middle Pleistocene to the Holocene, confirmed the
73 key role played by the Italian Peninsula, due to its geographical position, in affecting Italian
74 wolf dispersal routes and diversification dynamics during the Quaternary.

75 Recently, ancient DNA analyses performed on Late Pleistocene and Holocene remains of
76 *C. lupus* and *C. lupus familiaris* provided new insights into ancient Italian and European
77 wolf population dynamics and allowed to detect the oldest occurrence of domestic dogs in
78 Italy, suggesting the presence of a possible domestication site also in the Peninsula
79 ([Ciucani et al. 2019](#); [Angelici et al. 2019](#); [Boschin et al. 2020](#); [Koupadi et al. 2020](#)).
80 Additionally, ancient DNA analyses on European canid samples from the nineteenth to the
81 early twentieth centuries showed that most western European wolf populations
82 experienced a drastic spatio-temporal reduction of their genetic diversity and that among
83 them the Italian wolf population suffered the most severe and protracted bottleneck
84 ([Dufresnes et al. 2018](#)).

85 Conservation genetic studies analysing extant canid samples revealed that such numerical
86 contractions and genetic variability loss for the Italian wolf population have continued until
87 the early 1970s ([Montana et al. 2017a,b](#)), when it reached the verge of extinction counting
88 only approximately 100 individuals surviving in the central and southern Apennines ([Zimen](#)
89 [and Boitani 1975](#)).

90 Since then, thanks to legal protection and socio-ecological changes ([Chapron et al. 2014](#)),
91 Italian wolves significantly increased with 1.600 – 1.900 individuals estimated between
92 2009 and 2013 ([Galaverni et al. 2016](#)), and more than 2.000 wolves are likely present in
93 Italy today.

94 Interestingly, despite this wide availability of paleontological and genetic data on ancient
95 and recent European wolf populations, morphological and genetic studies of wolves living
96 during intermediate historical periods such as the Middle Ages are underrepresented. The
97 few available documents from these periods are solely reports of hunting bounties and
98 commercial exploitations, attesting the impact of human activities on wolf survival.

99 However, Middle Ages certainly represented a significant period in the evolutionary history
100 of the wolf in Europe.

101 Though planned and rewarded hunting activities against the wolf dates back to the sixth
102 century B.C., when Solone of Athens introduced a 1 to 5 drachma bounty for every wolf
103 killed ([Boitani 2003](#)), persecutions on wolf populations became deeply impactful only from
104 the beginning of the Medieval period. Since then, wolves were systematically killed
105 throughout Europe until the end of the 1800s ([Mallinson 1978](#); [Zimen 1978](#)) for multiple
106 reasons, from personal and livestock protection to completely irrational motivations linked
107 to cults and popular beliefs ([Boitani 2003](#); [Pluskowsky 2006](#)).

108 Hunting pressures in Europe were as extensive as that the wolf was completely extirpated
109 in the British Isles ([Harmer and Shipley 1902](#)) and Denmark ([Zimen 1978](#); [Boitani 1986](#))
110 around the mid-1700s, in Switzerland and Germany before the end of the nineteenth
111 century ([Zimen 1978](#); [Boitani 1986](#)), whereas the last individuals were slain in Sweden and
112 Norway in 1966 and 1973, respectively ([Turi 1931](#); [Promberger and Schröder 1993](#)). In
113 France, persecutions started with Charlemagne who issued two special laws between 800
114 and 813 AD (*Capitolare de Villis* and *Capitolare Aquisgranensis*) creating specialised wolf
115 hunters called “*louveterie*” dismissed during the French Revolution and revived in 1814
116 ([Hainard 1961](#)). In 1883 French wolves killed reached the impressive number of 1.386 and
117 the last individual was spotted in 1934 ([Beaufort 1987](#)). Similarly, extensive killings were
118 perpetuated also in Spain by Carlo VI in 1404 until a few decades ago ([Grande del Brio](#)
119 [1984](#); [Blanco et al. 1990](#)), while in Italy wolf bounties are reported from the twelfth century
120 to the 1950s ([Boitani 2003](#)). However, both in Spain and in Italy wolves never
121 disappeared.

122 Despite such hunting pressures protracted for centuries, in Europe osteological remains of
123 Medieval wolves are extremely limited ([Pluskowsky 2006](#) and literature therein), leading to

124 a sort of paradox, because in the scientific literature more data are available for European
125 Pleistocene wolves than for those from a few centuries ago. Among Medieval
126 archaeozoological remains, the proportion of wild to domestic species varies according to
127 the geographical and cultural context, but the presence of *C. lupus* is always scarce. This
128 shortage might be due to 1) the poorly preserved nature of the samples/specimens, which
129 often makes the distinction between wolves and dogs morphologically unfeasible; 2)
130 Medieval remains of wild animals mainly represented by game species as ungulates,
131 lagomorphs, and birds; 3) wolf carcasses generally ignored without gathering their bones
132 since wolves were usually killed for defence, bounties or their pelts, but rarely exploited as
133 a food resource.

134 This scarcity of European wolf remains in Medieval archaeological contexts is not only a
135 question concerning the human-wolf relationship, but also results in a serious lack of
136 biological and ecological information for a more exhaustive comprehension of the
137 dynamics and phenomena affecting the evolution of past populations of *C. lupus*.

138 In this scenario, we morphologically and genetically analysed a perfectly preserved
139 cranium of *C. lupus* recovered from the Po Valley (northern Italy) dated to ca. 967 – 1.157
140 AD. which might be crucial to improve our knowledge on Middle Age wolves and their
141 relation with the extant and fossil Italian forms. This cranium was only briefly reported in
142 [Ciucani et al. \(2019\)](#), but in this study, we provide the first comprehensive description of
143 the specimen based on a multidisciplinary approach, including palaeopatological, aDNA,
144 and tomographic analyses, to advance more detailed inferences on wolf dispersal
145 dynamics and genetic variability during the Middle Ages.

146

147 **Geographical and geological background**

148 The Po Valley ([Figure 1](#)) represents an area consisting of a layer of sediments of fluvial
149 and glacial origin, from the Pleistocene (2,6 – 0,01 Ma) to the Holocene (from 12.000
150 years ago to today) overlying a marine substrate that rarely emerges (Plio-Pleistocene)
151 ([Persico et al. 2006a](#); [Bona and Corbetta 2009](#)).

152 Since the beginning of the Quaternary, glacial and interglacial phases have alternated
153 throughout the Pleistocene characterized by general variations in the global average
154 temperature with the consequent development or regression of polar and alpine glaciers
155 on the Earth's surface. Between one glacial period and the next, interglacial phases
156 alternated, during which the rise in the average temperature of the Earth caused a
157 progressive retreat of the ice and a consequent rise in sea level. In this context, the action
158 of the Alpine and Apennine rivers filled the Po basin, generating the plain. In particular, the
159 more superficial layers of the plain derive from the material transported by the rivers at the
160 end of the last glacial period (110.000 – 12.000 years ago) following the erosion of the
161 morainic bodies previously generated by the advancement of the glacial front during the
162 cold phase ([Persico 2021](#)).

163 The end of the Pleistocene sees a morphological arrangement of the plain quite similar to
164 the current one, with a hydrographic network in which the main rivers are still recognizable
165 today, with the Po as the main collector. The main traceable geomorphologies also date
166 back to the last glacial phase. In this stretch of the plain, the Po River has a tortuous
167 course. This morphology is due to the low slope of the plain, the intensity of the erosive
168 and transport processes, to the geomorphology of the territory.

169 The low slope of the plain forces the river to erode the concave (external) bank and to
170 deposit material on the convex (internal) bank. The deposited sediments make up the
171 meander bar.

172 Following the floods, coarse sediments are deposited, which often contain fossil remains.
173 Under normal riverbed conditions, the reduced energy of the water involves the transport
174 of medium, fine, or very fine sediments such as gravels, sands, and silts. On the other
175 hand, when the flow of the river increases, coinciding with seasonal flood events, erosion
176 and transport increase, causing the deposition of larger clasts on the bars and with them
177 also the fossil bone remains. The ease with which fossil bones are transported by river
178 water varies according to their shape, size, and weight, and their frequency of discovery is
179 generally related to the presumed proximity of the primary reservoir sites, never directly
180 identified.

181 The river bar where the studied fossil was found is located downstream of the “Boschi
182 Maria Luigia” Park, near the Coltaro di Sissa Trecasali (PR) (Figure 1). The bar has a
183 crescent shape and is located on the right bank of the Po River, inside the meander.
184 Sedimentologically, this bar is characterized by gravel, sand, and silt, with the second
185 component in clear prevalence. On this beach, there are notoriously few finds, for the most
186 ancient fossil remains, it is necessary to shift attention to the river bars upstream of the Po
187 until reaching Cremona.

188 The ongoing research by one of the authors (D.P.) testifies to a sedimentary classification
189 from Cremona to Casalmaggiore (CR), with a drastic decrease in the size of the clasts
190 towards the second city, located a few kilometers downstream of the site where the wolf
191 was found. This sedimentary characterization is linked to the distribution of paleontological
192 finds along the river that show ancient, large, and well-preserved remains, where the
193 granulometry of the sediments is greater, with more modern, smaller, or fragmented
194 remains, where the sedimentology shows a reduced granulometry. This correlation can be
195 easily explained by correlating the size of the clasts, the transport energy of the river

196 water, and the lesser or greater depth of the fossil formation under the river: superficial
197 near Cremona and deep in nearby Casalmaggiore (CR).

198

199 **Materials and methods**

200 ***Sample collection***

201 The wolf (MSDP 348) specimen ([Figure 2](#)) analysed in the present study was recovered in
202 September 2016 on the alluvial bar of the Po River, right bank near Coltaro (PR) but in
203 Gussola (CR) territory, Lombardy Region (North Italy) (45° 58' 53" N; 10° 19' 49" E;
204 altitude: 21 m a.s.l.). The area, which is well-known for its numerous paleontological
205 Quaternary records ([Persico et al. 2006](#)), consists of a crescent-shaped meander bar
206 (about 800 meters), located in the meander of Isola Maria Luigia, a few kilometres
207 upstream to Casalmaggiore (CR) ([Figure 1](#)). The fossil was found in the typical
208 allochthonous position. For this reason, there are no stratigraphic information, and the
209 dating was possible only with a radiometric method (see below). Sample MSDP 348 is
210 currently housed in the Carnivora fossil section of the Po River at the Paleoanthropological
211 Museum of San Daniele Po (Cremona, Northern Italy).

212 Craniodental measurements were taken with a manual calliper to the nearest 0.1 mm.
213 Comparative samples, used for both morphological and biometric analyses, are largely
214 based on our personal dataset (complete list in [Tables S1, S2, Figures S1, S2](#)) which
215 includes extant *Canis lupus italicus* and Late Pleistocene *Canis lupus* from several Italian
216 paleontological sites. For the biometric comparison we considered the cranial
217 measurements following [von den Driesch \(1976\)](#): the total length (TL), condylobasal length
218 (CBL), palatal length (PL), greatest breadth of the mastoid processes (GMB), greatest
219 neurocranial breadth (GNB), greatest breadth of the palate at the level of P⁴ (GBP), lower

220 palatal breadth at level of C¹ (LPB). Dental traits were also considered: length (L) and
221 breadth (B) of I³, P², P³, P⁴, M¹ and M². Angular measurements were obtained using the
222 digital tool available on Materialise Mimics 20. The age estimation was performed following
223 [Gibson et al. \(2000\)](#) while gender determination was estimated following ([Siracusa and Lo](#)
224 [Valvo 2004](#)).

225

226 ***Tomography***

227 Sample MSDP 348 and a selection of extant *Canis lupus italicus* ([Tables S3, S4](#)) were
228 acquired using a Philips Brilliance CT 64-channel scanner at M.G. Vannini Hospital
229 (Rome). The scanning resulted in 659 slices with dimensions of 768 × 768 pixels. The
230 slice thickness is 0,8 mm with an interslice space of 0,4 mm. The CT data were processed
231 using Materialise Mimics 21.0 while the final 3D model was rendered with ZBrush 4R6
232 ([Boscaini et al. 2020](#)).

233

234 ***Radiometric dating***

235 Sampling for radiometric dating was performed in the Ancient DNA Lab of the Cultural
236 Heritage Department of University of Bologna. Subsequently the material was analysed by
237 the Dating and Diagnostics Center (CEDAD) of the University of Salento. Radiometric
238 dating was performed by conventional radiocarbon method using the high-resolution mass
239 spectrometry technique (AMS). The dating results have been corrected for the effects of
240 isotopic fractioning both through the measurement of the $\delta^{13}\text{C}$ term carried out directly
241 with the accelerator and for the bottom of the measurement. Samples of known
242 concentration of Oxalic Acid provided by NIST (National Institute of Standards and
243 Technology) have been used for quality control of the results. For the determination of the

244 experimental error on the radiocarbon date, both the scattering of data around the average
245 value and the statistical error deriving from the $\delta^{14}\text{C}$ count were taken into account.
246 Radiocarbon dating for the samples was then calibrated at the calendar age using the
247 software OxCal Ver. 3.10 based on atmospheric data ([Reimer et al. 2013](#)).

248

249 ***Ancient DNA standards and analyses***

250 Molecular analyses were conducted at the ancient DNA laboratory of the University of
251 Bologna (Ravenna Campus) following strict criteria (see [Supplemental Material](#)) in order to
252 support the authenticity of the results and prevent possible contamination from exogenous
253 DNA ([Cooper and Poinar 2000](#); [Gilbert et al. 2005](#); [Knapp et al. 2012](#); [Llamas et al. 2017](#)).

254 A sub-sampling for molecular analyses was performed on a tooth encased in the alveolus
255 of the maxilla, drilling directly in the root from the tooth neck. The surface was superficially
256 decontaminated by slight abrasion with a sterile diamond drill and then exposed to UV
257 radiation for 30 minutes. Then, DNA was extracted from the root powder with a silica-
258 based protocol ([Cilli et al. 2020](#)) modified from [Dabney et al. \(2013\)](#). Amplicons of 361 bp
259 of the mitochondrial DNA were obtained by means of three overlapping fragments
260 amplified by Polymerase Chain Reaction (PCR) using three couples of primers (See
261 [Supplemental Material](#)).

262 Excluding primers, the three overlapping fragments covered the following nucleotide
263 positions 15431 - 15792 of the Italian wolf mitochondrial genome (Genbank accession
264 number KU644662). PCR reactions were conducted in triplicate, to ascertain the presence
265 of possible miscoding lesions.

266 The purified amplicons were sequenced at the Unit for Conservation Genetics (Ozzano
267 dell'Emilia, Bologna, Italy), Department for the Monitoring and Protection of the

268 Environment and for Biodiversity Conservation of the Italian Institute for Environmental
269 Protection and Research (ISPRA).

270 Electropherograms were manually analysed and visually edited using the ABI software
271 SEQSCAPE v.2.5. Consensus sequences were reconstructed by using the three
272 replicates for each fragment, using the Italian wolf mitochondrial genome (Genbank
273 accession number KU644662) as a reference.

274 Blastn ([Altschul et al. 1990](#)) was checked in order to analyze the sequence in the context
275 of the ancient and modern genetic variability available in GenBank database.

276 A database of published homologous mtDNA sequences ([Table S5](#)) was constructed to
277 analyse our data in the wider context of ancient and modern European canids. In
278 particular, a 331-bp alignment containing ancient and modern wolf and dog sequences
279 was used for phylogenetic reconstructions. Instead, a 278-bp alignment, containing only
280 ancient and modern wolf sequences was used for within-species network analyses. The
281 dimension of the two alignments was determined by the maximum length of the sequences
282 included in the reference selected (See [Table S5](#)).

283 The software DnaSP v.5.10.01 ([Librado and Rozas 2009](#)) was used to identify identical
284 sequences and to collapse them into unique haplotypes.

285 The 331-bp alignment was run in JModeltest2 ([Darriba et al. 2012](#)) to estimate the best
286 nucleotide substitution model through the Akaike information criterion (AIC) and to
287 construct in MEGA-X v.10.2.5 ([Kumar et al. 2018](#)) a Neighbor-Joining (NJ; [Saitou and Nei
288 1987](#)) phylogenetic tree, which was rooted using a Coyote sequence (*Canis latrans*,
289 GenBank access number FJ213927.2; [Fain et al. 2010](#)) as an outgroup and whose
290 internode supports were obtained by 1,000 bootstrap replicates ([Felsenstein 1985](#)).
291 Finally, the obtained tree was visualized and edited in FigTree V.1.4.4 (Rambaut 2010).

292 A Median-Joining network (MJ), based on the 278-bp alignment, was constructed by using
293 Network v.10.2 (<https://www.fluxus-engineering.com/sharenet.htm>) with no pre- or post-
294 processing steps.

295

296 **Results**

297 ***Radiometric dating***

298 Radiometric dating performed on a bone sample taken from the studied cranium is
299 supported by a high probability of correctness (94%) and highlights a young age of the
300 fossil (967 – 1.157 AD, [Figure S3](#)). The determined age is in agreement with the initial
301 fossilization state identified.

302

303 ***Outer cranial anatomy***

304 Sample MSDP 348 ([Figure 2](#)) consists of an almost complete and well preserved cranium
305 missing of both the zygomatic arches, the left I¹, I², C¹, P¹ and the right I¹-I³, C¹, P¹. The
306 bone tissue appears beige in colour, compact and quite mineralised but light. The dental
307 enamel is white-yellowish with orange traces of iron oxides. The cranial sutures appear
308 completely fused with the exception of some portions of the fronto-parietal and
309 presphenoidal sutures, where the welding is weaker. Both the canines are missing and
310 their alveolar cavities are partially exposed due to the loss of bone tissue. The bone
311 surrounding the right alveolus appears compact with sharp edges, while a marked swelling
312 with highly vascularised newly formed bone tissue with smooth edges characterising the
313 left alveolus. The abnormal morphologies of the latter, suggest a lesion of pathological
314 origin rather than the result of taphonomic processes (see Paleopathology). In contrast,
315 the fractures with smooth edges of the zygomatic arches, the strong abrasion of both the

zygomatic processes of the frontals, which partially expose the cavity of the frontal sinuses, as well as the partial exposure of the tooth roots of the cheek teeth and the cavity affecting the right lacrimal and palatine bones, are of taphonomic origin and probably due by a low-energy river transport (rafting). On both sides, the orbital fissure, the anterior alar foramen and the oval foramen are obliterated with very fine gravels (2-4 mm) which support the hypothesis of a long permanence in river waters.

The cranium MSDP 348 shows a typical wolf-like morphology that falls into the variability of extant specimens of *C. lupus italicus* (Table S1): the braincase is globose with a prominent sagittal crest distally protruding over the occipital condyles; the rostrum is robust and elongated, the frontals are quite narrow and elevated with a moderate step; the nasal bones end in correspondence of the maxillofrontal suture; the tympanic bullae are large and oval in shape in ventral view; the orbital angle is approximately 41.5°. On the distal margin of the anterior palate, the incisive foramina extend from the anterior margin of the canine alveoli to almost reach the mesial margin of the P¹. The arrangement of the upper tooth row in the studied specimen is overlapping with those of the extant wolves, indeed, the C¹-P² are arranged on the same axis, which changes in P³-P⁴ and is different again in M¹-M². However, some minor differences with the extant form are noticeable. In MSDP 348 the distal portion of the rostrum is slightly more flattened, the palate is larger at the level of P²-P³ and the frontal bones are less wider with a more pronounced postorbital constriction (Figure 3). Taking into account the Italian crania of *C. lupus* from Middle to Late Pleistocene, is important to remark that recent studies (Sardella et al. 2014; Mecozzi et al. 2017; Mecozzi and Bartolini Lucenti 2018; Mecozzi et al. 2020) attested the presence of two distinct chronological morphotypes, consisting of “slender forms” spanning from the late Middle to early Late Pleistocene and “robust forms” from the beginning of MIS 3. According to the authors, suggesting a dispersal wave of canids from continental Europe probably adapted to cold climatic conditions. In this context, the specimen MSDP 348

342 resembles in overall morphology the robust forms from Vignone, Grotta Ladrenizza and
343 Grotta Grimaldi with which it shares a dorsoventrally flattened rostrum and a wide and
344 straight anterior portion of the palate without the constriction at the P²-P³ observed in
345 extant Italian wolves.

346 Also, the teeth are morphologically almost indistinguishable from those of fossil and extant
347 specimens. In MSDP 348, the left I³, the P⁴ and M¹ of both the hemiarches are quite worn,
348 whereas the P², P³ and M² only show a slight wear on the cusp tips. The I³ is separated
349 from the I² alveolus by a short diastema, but the development of its basal cingula cannot
350 be observed due to the advanced wear of the crown. The P¹ is monocusped and meso-
351 distally elongated. The P² is meso-distally elongated with an asymmetric protocone and a
352 weak distal cingulum. Also the P³ is meso-distally elongated, but possesses a symmetric
353 protocone, a small distal accessory cusp and a weak distal cingulum. The P⁴ shows a
354 weak cusp on the protocone, located slightly anteriorly than the mesial margin of the tooth
355 and in occlusal view it appears lingually expanded. Both the paracone and metastyle are
356 stout but not very high due to wear. The mesial cingulum is marked, whereas those labial
357 and lingual are less developed. The M¹ possesses a nearly squared profile in occlusal
358 view. It has a large paracone, considerably more developed compared to the metacone.
359 The trigon basin is wider and deeper than the talon basin. The protocone is quite large,
360 whereas the metaconule and the hypocone are weakly developed. The mesial and buccal
361 cingula are pronounced. Also the M² is quite squared in shape in occlusal view, with a
362 paracone slightly larger than the metacone. The protocone is large and rounded in profile,
363 the basin is less deep and the hypocone cingulum is well developed. Distal and buccal
364 cingula are marked.

365 Considering the morphology of the cheek teeth, [Mecozzi and Bartolini Lucenti \(2018\)](#)
366 reported the presence of marked cusps and marked cingula, together to the occurrence of

367 mesial and distal accessory cusps, in the robust specimens from the Late Pleistocene of
368 Italy. The cranium MSDP 348 shows an intermediate morphology between the slender and
369 robust morphotypes, as the main cusps of the teeth are stout and well developed, whereas
370 the accessory ones and cingula are less pronounced.

371

372 ***Inner cranial anatomy***

373 The brain and frontal sinuses of MSDP 348 show an overlapping size and morphology with
374 those of the extant sample (Figure 3, Tables S3, S4). The frontal sinuses are large and
375 well-developed extending from the antorbital constriction up to the fronto-parietal suture. In
376 dorsal view, they appear slightly narrower compared to those of the extant sample. The
377 edges of the sinuses are lobed and the surface is smooth. Laterally, they appear
378 prominent with the typical triangular profile (Figures 3a-d).

379 The brain endocast of MSDP 348 lacks both olfactory bulbs, the telencephalon is globular
380 in shape with marked convolutions. In dorsal view, a marked longitudinal fissure divides
381 the cerebrum into two roughly symmetrical hemispheres characterised by the following
382 gyri, from front to back: orbital, prorean, lateral, sigmoid, coronal, endolateral, ectolateral,
383 suprasylvian and ectosylvian, including the respective sulci. The frontal pole cortex is
384 antero-posteriorly elongated and the orbital gyrus is laterally expanded forming a “bump”
385 bounded by the intraorbital sulcus (Figures 3e-h). The orbital region is characterized by the
386 presence of three main sulci: the prorean sulcus, the intraorbital sulcus and a third sulcus
387 considered by [Lyras and Van der Geer \(2003\)](#) as typical of *C. lupus*, *C. simensis* and *C.*
388 *rufus*. The overall brain morphology of MSDP 348 corresponds to that of the extant wolf,
389 except for the wider and more marked pseudosylvian fissure.

390

391 **Biometric analysis**

392 Figures 4a, S1 and Table S1 show that the specimen MSDP 348 fall within the cranial
393 variability of the extant Italian wolf occupying the lowest dimensional ranges. The average
394 size of MSDP 348 is between that of the Late Pleistocene specimen from Grotta Romanelli
395 – very small and thus referred to the “slender form” – and those of larger and more robust
396 specimens from the Late Pleistocene sites of Grotta Ladrenizza, Covoli di Velo,
397 Romagnano and Broion (Figure 4a). Indeed, the latter occupies the highest values
398 reached by the extant forms and are considerably larger than MSDP 348. Compared to the
399 sample, MSDP 348 would appear to have a relatively narrow braincase (TL-GNB) and a
400 fairly wide palate (GBP-LPB) (Figure 3, Table S1).

401 The teeth measurements indicate that MSDP 348 has a smaller I³ and a larger breadth of
402 the P⁴ and M¹ compared to the extant *C. lupus italicus* (Figures 4b, S2, Table S2). All other
403 measures fall comfortably within the tooth variability of the extant form. Biometrically the
404 teeth of MSDP 348 resemble those of the Late Pleistocene wolves, except for the larger
405 breadth of the M¹ and M² and the shorter P².

406 The teeth of the specimen from Grotta Romanelli are the smallest of the sample, except
407 for the length and breadth of the P² which are similar to those of the other specimens,
408 including MSDP 348.

409

410 **Age and gender determination**

411 At first glance, the overall morphology and size of MSDP 348 clearly indicates an adult
412 individual, but to better define the age in terms of years, we considered the welding degree
413 of the cranial sutures and the tooth wear. According to Barone (2006), in domestic dogs
414 the complete welding of all the cranial sutures occurs at the age of 3-4 years. In MSDP

348 the cranial sutures are completely welded except the fronto-parietal and
presphenoidal ones where the welding is weaker suggesting an age of approximately 3
years. In contrast, the I³, P⁴ and M¹ are quite worn to the point that a portion of the pulp
cavity is exposed on the I³ and the main cusps of the P⁴ and M¹ are completely blunted.
Following the age estimations proposed by [Gipson \(2000\)](#) based on the tooth wear of the
wolf, the upper carnassial of the Po Valley specimen corresponds to an individual between
6 and 8 years of age. The discrepancy between the estimated ages from sutures and teeth
is considerable and it is likely that some sutures have loosened as a result of taphonomic
processes such as a long presence in water. Teeth are generally less affected by the
environmental processes as the tooth enamel is harder and more resistant compared to
the bones tissue, moreover in the case of MSDP 348 the wear surfaces are all located on
the occlusal plane of the teeth, indicating that their origin is due to chewing activity and not
to taphonomic processes. Therefore, we consider the age estimated from teeth more
reliable than that obtained from the cranial sutures.

In the wolf skulls there is a certain sexual dimorphism that can be evaluated through the
Storer index. Its formula is as follows:

$$x_m - x_f / (x_m + x_f) : 2 \times 100$$

Where:

x_m = average value for a character for males

x_f = mean value of the same character for females

The index value is negative when the female is larger than the male, it is equal to zero
when there are no differences between the sexes and it is positive when the male is larger
than the female ([Siracusa and Lo Valvo 2004](#)). In the studied sample the incompleteness
of the fossil did not allow the survey of all the parameters for the calculation of the index so
that only some measures (condylobasal length and total length of the cranium) were
compared with the values present in the literature ([Syracuse and Lo Valvo 2014](#)) ([Table 1](#)).

441 From the comparison in Table 1 it can however be estimated that MSDP 348 sample might
442 belong to a female wolf.

443

444 ***Ancient DNA analyses***

445 Molecular analyses allowed us to obtain reliable data, in accordance with the guidelines
446 and standards applied, for all the three replicated amplifications of the three different
447 fragments, resulting in a 361-bp mtDNA consensus sequence. Moreover, the sequence
448 was confirmed also by the comparison with the previously published 57-bp fragment
449 (Ciucani et al., 2019). The sequence of MSDP 348 here obtained is available on GenBank
450 (acc. n. OL906402).

451 After blasting, the Medieval sequence returned to be unique and showed no homology
452 with the data stored in the GenBank database. It presents one mutation of difference from
453 the wolf haplotype named W15 by [Randi et al. \(2000\)](#) and by [Montana et al. \(2017b\)](#) and 2
454 mutations of difference from other 4 modern wolf sequences: 3 belonging to samples
455 collected in Italy (KU696389 and KU6446621, [Koblmüller et al. 2016](#), and KF661048,
456 [Thalmann et al. 2013](#)) and all sharing the typical Italian haplotype W14 ([Randi et al. 2000](#);
457 [Montana et al. 2017b](#)) and 1 belonging to a Belarusian sample (KU696390, [Koblmüller et](#)
458 [al. 2016](#)). The subsequent haplotypes, showing more than 2 mutations of difference from
459 the Medieval sequence we obtained, are from Bulgaria (KU696388) and Poland
460 (KF661045).

461 The MSDP 348 sequence was also analysed against the shorter (278 bp) haplotypes
462 retrieved in the historical (19th and 20th centuries) published dataset by [Dufresnes et al.](#)
463 [\(2018\)](#), showing again no matches with them.

464 The best evolutionary model for the alignment of 141 canid mtDNA sequences we built
465 was the Kimura 2-parameters model (Kimura 1980) with the rate variation among sites to
466 Gamma distribution (G shape parameter = 1). The reconstructed NJ phylogenetic tree

467 (Figure 5), despite low supports for most nodes due to the limited length of the analysed
468 fragment, well confirmed the distinction between the two main wolf mitochondrial
469 haplogroups Hg1 and Hg2 proposed by Pilot et al. (2010) using partial mtDNA genomes
470 and clearly highlighted also the distinction between the three main dog clades (A, B, C)
471 observed by Thalmann et al. (2013) using complete mtDNA genomes. Among
472 haplogroups, four macro-groups were also identifiable in the tree (Figure 5). Macro-groups
473 4 and 5 include both wolf and dog (belong to canine clades A and B) haplotypes obtained
474 from both modern and ancient samples. Macro-group 3 predominately includes modern
475 dog haplotypes (clade A), except for four branches represented by ancient wolf
476 haplotypes. Macro-group 1 hosts almost exclusively dog haplotypes belonging to clade C,
477 whereas macro-group 2, which includes the sequence obtained from sample MSDP 348,
478 is exclusively composed of ancient and modern wolf haplotypes.

479 In particular, sample MSDP 348 is clearly located in a specific wolf clade, which includes
480 only mtDNA sequences from the Mediterranean Basin belonging to haplogroup 2 (Pilot et
481 al. 2010). In fact, MSDP 348 haplotype is clearly connected with W15 and W14
482 haplotypes, differing only one and two mutations from them, respectively.

483 Network analyses (Figure 6) confirmed the placement of sample MSDP 348 in the wolf
484 haplogroup 2, and in this shorter alignment, it matches with the W15 wolf haplotype (Randi
485 et al. 2000). Samples included in the shorter alignment showed an almost complete
486 turnover in the wolf haplotypic composition in the European area, moving from Late
487 Pleistocene to the current era. Indeed, all but one haplotype resulted to be different
488 between the two considered periods: Late Pleistocene – Holocene (>2.000 yrs BP) and
489 Holocene (<2.000 yrs BP) – present.

490

491 ***Paleopathology***

492 Among the various diseases affecting the oral cavity of canids, periodontitis is one of the
493 most common (Strillou et al. 2010; Iurino et al. 2015). Caused by specific microorganisms
494 which lead to the formation of periodontal pockets and the loss of alveolar bone tissue, it is
495 often the result of pre-existing gingivitis, gingival lesions or accumulation of supragingival
496 plaque and calculus (Brown et al. 1989; Pejčić et al. 2007; Strillou et al. 2010). In canids, a
497 chronic inflammation of the oral cavity provokes an increase of the gingival blood flow from
498 200 to 400% with a consequent enlargement of the vascular pores in the maxillary and
499 dentary bones (Kaplan et al. 1982; Hock and Kim 1987; Zoellner 1990), as can be
500 observed on the left canine alveolus of MSDP 348 (Figure 5). The protrusion of bacterial
501 infection may have triggered the periodontitis in MSDP 348 by forming gingival pockets
502 and causing the total loss of the canine and the retraction of the bone tissue. The most
503 serious lesions can be noticed on the outer edge of the alveolus and within the alveolar
504 cavity. Here the bone tissue appears swollen and highly vascularized with a consistent
505 missing portion of the maxillary bone forming a large hole (max diameter of 12.2 mm) that
506 connect the alveolar cavity with the nasal one (Figure 7). The smooth edges of the cavity
507 and its sheltered position exclude a taphonomic origin of the lesion. Additional small
508 cavities connect the alveolus of the left canine with that of P¹, reaching in part the anterior
509 root of the P².

510 The prevalence and severity of periodontitis in canids increase mainly with age, although
511 multiple factors such as the nature of the infection, previous lesions, genetics and diet may
512 favour the onset of this disease and influence its pathological course (Strillou et al. 2010).
513 Measurements carried out on spontaneous or experimental cases demonstrated that 4–6
514 months are needed to provoke a root exposure of 5 mm (Strillou et al. 2010; Kinoshita et
515 al. 1997; Pallua and Suschek 2011). Iurino et al (2013) documented a case of severe
516 periodontitis in a Pleistocene *Cuon alpinus* from the San Sidero locality (southern Italy)
517 inferring a pathological course of about 6 months. In MSDP 348, the loss of the alveolar

bone has exposed at least one half of the canine root (12.5 mm), thus indicating a protracted periodontitis for a period of about 12 months. Although the severity of the lesions has contributed to debilitating the individual, probably compromising its chewing and predatory abilities, however, on the basis of the available data it is not possible to establish whether its death was directly related to this pathology.

Discussion and conclusions

The nearly complete cranium MSDP 348 recovered from the alluvial bar of the Po River revealed an age of ca. 967-1157 AD by radiometric dating. This chronology makes the sample of particular relevance for the study of past wolf populations as the archaeozoological record of well-preserved remains of *C. lupus* are extremely rare, especially those of the Middle Ages.

Biometric and CT-based analyses indicate that the specimen MSDP 348 falls within the cranial variability of the extant *Canis lupus italicus* occupying the lowest dimensional ranges. The tooth wear allows to refer the cranium to an adult individual between 6 and 8 years, while gender determination, performed following [Syracuse and Lo Valvo \(2014\)](#), suggests that it could be a female individual. This specimen shows clear evidence of a severe periodontitis which caused the complete loss of the left canine producing a large hole connecting the alveolus with the nasal cavity. This condition probably severely debilitated the subject, although it is not possible to establish whether the death was a consequence of this disease.

Biometric and CT-based findings were clearly supported also by molecular investigations, which were performed to study this sample through a multidisciplinary approach especially to analyse its gene pool into the context of modern and ancient Eurasian wolf genetic variability. Ancient DNA methodologies and strict reliability criteria here applied clearly supported the authenticity of the data obtained. The strategy of analysis used in this study

544 based on the amplification of three short and adjacent fragments, successively
545 concatenated during bioinformatic analyses, allowed us to obtain a longer consensus
546 mtDNA sequence compared to the fragments analysed in similar aDNA studies based on
547 samples collected from the same latitudes ([Dufresnes et al. 2018](#); [Ciucani et al. 2019](#)).
548 Notably, the sequence we obtained in this study resulted to be even ~300 bp longer
549 respect also to the sequence previously obtained for the same sample by [Ciucani et al.](#)
550 [\(2019\)](#) and that allowed us to compare the results to a much wide database of ancient and
551 modern wolves and dogs.

552 Our phylogenetic analyses unambiguously agree with previous molecular modern and
553 ancient studies, resulting in similar topologies for both networks and trees ([Montana et al.](#)
554 [2017a](#); [Ciucani et al. 2019](#)).

555 Molecular comparisons showed that the MSDP 348 sequence has a unique haplotype, not
556 recorded in the ancient (for the data available so far) nor in the modern wolf variability.

557 Interestingly, phylogenetic analyses placed mtDNA gene pool of sample MSDP 348 within
558 the genetic variability of modern wolves, clearly distinct from that of dogs. In particular, it
559 falls in the wolf haplogroup 2, the oldest one, which today has been largely replaced by
560 haplogroup 1 that arrived in Europe at least 2.700 – 1.200 years ago ([Pilot et al. 2010](#)).

561 Phylogenetic analyses also showed that sample MSDP 348 haplotype is placed near the
562 Greek haplotype named W15 ([Randi et al. 2000](#)), from which it displays only one mutation
563 of difference. However, this difference did not emerge in the comparison based on the
564 shorter fragment previously obtained by [Ciucani et al. \(2019\)](#), because the distinctive
565 mutation is placed in the portion of the mitochondrial DNA that was not previously
566 investigated, highlighting the importance of considering wide amplicons even in aDNA
567 analyses and the success of the strategy here adopted in obtaining a longer mtDNA
568 sequence compared to previous aDNA studies.

569 This sequence represents part of the ancient wolf variability today lost, as a result of the
570 negative impact of overkilling and anthropic persecutions perpetrated in Medieval times
571 and, for western Europe, in particular in the last 150 years. The study of modern and
572 ancient samples allows us to better comprehend the dynamics of wolf populations and the
573 alternation of haplogroups in the European areas. This didn't occur in Italy, where only
574 haplogroup 2 is still present.

575 This study highlights the fragmentation and the few availabilities of archaeozoological and
576 paleogenomics data from the Medieval period, a period of dark centuries for the wildlife
577 and its management.

578 Despite *Canis lupus* is undoubtedly one of the most iconic and extensively studied
579 predators of all times, in Europe informative osteological remains of Medieval wolves are
580 extremely rare, limiting our comprehension of the dynamics and phenomena affecting the
581 evolution of past populations of *C. lupus*. The perfectly preserved cranium (MSDP 348)
582 dated to ca. 967-1.157 AD from the Po Valley (northern Italy) is here assigned to an adult
583 *C. lupus*. Tomographic and biometric analyses of both outer and inner cranial anatomy,
584 indicate a 6 – 8 years old female affected by a severe periodontitis, with size and
585 morphology similar to those of the smaller specimens of extant *C. lupus italicus*. Genetic
586 analysis reveals that the haplotype of MSDP 348 is related to the Greek haplotype W15,
587 but was never recorded before, representing a portion of ancient wolf variability today lost.
588 This study provides the most comprehensive description of a *C. lupus* from the Middle
589 Ages in Italy, and underlines how archaeozoological samples represent an essential
590 source of information to understand the dynamics, diversity and distribution of past and
591 present-day wolves.

592

593 **Acknowledgments**

594 We would like to thank M. Danti of the 'M.G. Vannini' Hospital of Rome for making the CT
595 scanning. We also thank D. Luiselli, F. Fontani and A. Latorre of the Ancient DNA
596 laboratory of the University of Bologna for their support and advice, N. Mucci (ISPRA) for
597 logistic support in the genetic analysis and E. Velli (ISPRA) for laboratory assistance. We
598 are grateful to Adam J. Andrews for English revision and useful comments. This work has
599 benefited from the equipment and the framework of the COMP-HUB Initiative, funded by
600 the 'Departments of Excellence' program of the Italian Ministry for Education, University
601 and Research (MIUR, 2018-2022).

602

603 **Disclosure statement**

604 No potential competing interest was reported by the authors.

605

606 **References**

- 607 Angelici FM, Ciucani MM, Angelini S, Annesi F, Caniglia R, Castiglia R, Fabbri E,
608 Galaverni M, Palumbo D, Ravegnini G, Rossi L., et al 2019. The Sicilian wolf: genetic
609 identity of a recently extinct insular population. *Zoological science*. 36(3):189-197.
- 610 Angelici FM, Rossi L. 2018. A new subspecies of grey wolf (Carnivora, Canidae), recently
611 extinct, from Sicily, Italy. *Bollettino del Museo Civico di Storia Naturale di Verona*
612 *(Botanica Zoologia)*. 42:3-15.
- 613 Altschul F, Gish G, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search
614 tool. *Journal of Molecular Biology*. 215:403–410.
- 615 Barone R. 2006. Comparative anatomy of domestic mammals. Tome 1. Osteology.
616 Edagricole-New Business Media. 690 pp.

- 617 Beaufort FG. 1987. Ecologie Historique du Loup en France. These de Doctorate, Universit
618 de Rennes I. Rennes (France). 4:1104 pp.
- 619 Blanco JC, Cuesta C, Reig S. 1990. El lobo en España: Una vision global. Pp. 69-93 in
620 Blanco JC, Cuesta C, Reig S, eds., *El lobo (Canis lupus) en España: Situation,*
621 *problematica y apuntes sobre su ecología.* ICONA, Madrid. 118 pp.
- 622 Boitani L. 1986. Dalla parte del lupo. Milano: L'airone di G. Mondatore Associati Spa.
- 623 Boitani L. 2003. Wolf Conservation and Recovery. Pp. 317-320 in Mech D, Boitani L, eds.,
624 *Wolves Behavior, Ecology and Conservation.* The University of Chicago Press. 448 pp.
- 625 Bona F, Corbetta C, 2009. Mammalofaune quaternarie delle alluvioni del Po (province di
626 Cremona, Lodi, Piacenza e Parma): analisi paleontologica e ricostruzione
627 paleoambientale, "Monografie di Pianura", n. 9, Provincia di Cremona, Cremona.
- 628 Boscaini A, Iurino DA, Mamani Quispe B, Andrade Flores R, Sardella R, Pujos F, Gaudin
629 TJ. 2020. Cranial anatomy and paleoneurology of the extinct sloth *Catonyx tarijensis*
630 (Xenarthra, Mylodontidae) from the late Pleistocene of Oruro, Southwestern Bolivia.
631 *Frontiers in Ecology and Evolution.* 8:69. doi: 10.3389/fevo.2020.00069
- 632 Boschini F, Bernardini F, Pilli E, Vai S, Zanolli C, Tagliacozzo A, Fico R, Fedi M, Corny J,
633 Dreossi D, Lari M, et al. 2020. The first evidence for Late Pleistocene dogs in Italy.
634 *Scientific reports.* 10(1):1-14.
- 635 Breitenmoser U. 1998. Large predators in the Alps: The fall and rise of man's competitors.
636 *Biol. Conserv.* 83(3):279-289.
- 637 Brown LJ, Oliver RC, Loe H. 1989. Periodontal disease in U.S. in 1981: prevalence,
638 severity, extent, and role in tooth mortality. *J. Period.* 60(7):363–370.

639 Brugal J P, Boudadi-Maligne M. 2011. Quaternary small to large canids in Europe:
640 taxonomic status and biochronological contribution. *Quat. Int.* 243:171–182.

641 Chapron G, Kaczensky P, Linnell JD, Von Arx M, Huber D, Andrén H, López-Bao JV,
642 Adamec M, Álvares F, Anders O, Balčiauskas L. 2014. Recovery of large carnivores in
643 Europe's modern human-dominated landscapes. *Science*. 346(6216):1517-1519.

644 Cilli E, Gabanini G, Ciucani MM, De Fanti S, Serventi P, Bazaj A, Sarno S, Ferri G,
645 Fregnani A, Cornaglia G, Gruppioni G, et al. 2020. A multifaceted approach towards
646 investigating childbirth deaths in double burials: Anthropology, paleopathology and
647 ancient DNA. *J. Archaeol. Sci.* 122:105219.

648 Poinar HN, Cooper A. 2000. Ancient DNA: do it right or not at all. *Science*.
649 5482(1139):416.

650 Ciucani MM, Palumbo D, Galaverni M, Serventi P, Fabbri E, Ravegnini G, Angelini S,
651 Maini E, Persico D, Caniglia R, Cilli E. 2019. Old wild wolves: ancient DNA survey
652 unveils population dynamics in Late Pleistocene and Holocene Italian remains. *PeerJ*.
653 7:e6424.

654 Dabney J, Knapp M, Glocke I, Gansauge MT, Weihmann A, Nickel B, Valdiosera C,
655 García N, Pääbo S, Arsuaga JL, Meyer M. 2013. Complete mitochondrial genome
656 sequence of a Middle Pleistocene cave bear reconstructed from ultrashort DNA
657 fragments. *Proceedings of the National Academy of Sciences*. 110(39):15758-15763.

658 Darriba D, Taboada GL, Doallo R, Posada D. 2012. jModelTest 2: more models, new
659 heuristics and parallel computing. *Nature methods*. 9(8):772-772.

660 Dufresnes C, Miquel C, Remollino N, Biollaz F, Salamin N, Taberlet P, Fumagalli L. 2018.
661 Howling from the past: historical phylogeography and diversity losses in European grey
662 wolves. *Proc. Royal Soc. B.* 285(1884):20181148.

663 Fain SR, Straughan DJ, Taylor BF. 2010. Genetic outcomes of wolf recovery in the
664 western Great Lakes states. *Conservat. Genet.* 11(5):1747-1765.

665 Felsenstein J. 1985. Confidence limits on phylogenies: an approach using the bootstrap.
666 *Evolution.* 39:783-791.

667 Galaverni M, Caniglia R, Fabbri E, Milanesi P, Randi E. 2016. One, no one, or one
668 hundred thousand: how many wolves are there currently in Italy? *Mammal Research.*
669 61(1):13-24.

670 Gilbert MT, Bandelt HJ, Hofreiter M, Barnes I. 2005. Assessing ancient DNA studies.
671 *Trends in ecology & evolution.* 20(10):541-544.

672 Gibson PS, Ballard WB, Nowak RM, Mech LD. 2000. Accuracy and precision of estimating
673 age of gray wolves by tooth wear. *Journal of Wildlife Management.* 64:752–758.

674 Grande del Brio R. 1984. El lobo iberico, biologia y mitologia. H. Blume, Madrid.

675 Harmer SF, Shipley SE. 1902. Cambridge natural history: Mammalia. (Beddard FE)
676 MacMillan, London.

677 Hainard R. 1961. Mammifères sauvage d'Europe. I. Delachaux et Niestle, Neuchatel
678 (Switzerland). 350 pp.

679 Hock JM, Kim S. 1987. Blood flow in healed and inflamed periodontal tissues of dogs. *J.*
680 *Periodont. Res.* 22(1):1–5.

681 Iurino DA, Danti M, Della Sala SW, Sardella R. 2013. Modern techniques for ancient
682 bones: vertebrate palaeontology and medical CT analysis. *Boll. Soc. Paleontol. Ital.*
683 52(3):145-155.

684 Iurino DA, Sardella R. 2015. Medical CT scanning and the study of hidden oral pathologies
685 in fossil carnivores. *Paläontologische Zeitschrift.* 89(2):251-259.

686 Kinoshita AOS, Takahashi KYS, Ishikawa I. 1997. Periodontal regeneration by application
687 of recombinant human bone morphogenic protein-2 to horizontal circumferential defects
688 created by experimental periodontitis in beagle dogs. *J. Periodontol.* 68(2):103–109.

689 Kaplan ML, Jeffcoat MK, Goldhaber P. 1982. Blood flow in gingiva and alveolar bone in
690 beagles with periodontal disease. *J. Periodont. Res.* 17(4):384–389.

691 Knapp M, Clarke AC, Horsburgh KA, Matisoo-Smith EA. 2012. Setting the stage - building
692 and working in an ancient DNA laboratory. *Annals of Anatomy - Anatomischer Anzeiger.*
693 194(1):3-6.

694 Koblmüller S, Vilà C, Lorente-Galdos B, Dabad M, Ramirez O, Marques-Bonet T, Wayne
695 RK, Leonard JA. 2016. Whole mitochondrial genomes illuminate ancient intercontinental
696 dispersals of grey wolves (*Canis lupus*). *J. Biogeogr.* 43(9):1728-1738.

697 Koupadi K, Fontani F, Ciucani MM, Maini E, De Fanti S, Cattani M, Curci A, Nenzioni G,
698 Reggiani P, Andrews AJ, Sarno S, et al. 2020. Population Dynamics in Italian Canids
699 between the Late Pleistocene and Bronze Age. *Genes.* 11(12):1409.

700 Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: Molecular Evolutionary
701 Genetics Analysis across computing platforms. *Molecular Biology and Evolution* 35:1547-
702 1549.

703 Librado P, Rozas J. 2009. DnaSP v5: a software for comprehensive analysis of DNA
704 polymorphism data. *Bioinformatics.* 25:1451–1452.

705 Llamas B, Valverde G, Fehren-Schmitz L, Weyrich LS, Cooper A, Haak W. 2017. From the
706 field to the laboratory: Controlling DNA contamination in human ancient DNA research
707 in the high-throughput sequencing era. *STAR.* 3(1):1-4.

708 Lyras GA, Van Der Geer AAE. 2003. External brain anatomy in relation to the phylogeny of
709 Caninae (Carnivora: Canidae). *Zool. J. Linnean Soc.* 138:505–522.

710 Mallinson J. 1978. The shadow of extinction: Europe's threatened wild mammals.
711 Macmillian, London.

712 Mecozzi B, Bartolini Lucenti B. 2018. The Late Pleistocene *Canis lupus* (Canidae,
713 Mammalia) from Avetrana (Apulia, Italy): reappraisal and new insights on the European
714 glacial wolves. *Ital. J. Geosci.* 137:138–150.

715 Mecozzi B, Iurino DA, Bertè DF, Sardella R. 2017. *Canis mosbachensis* (Canidae,
716 Mammalia) from the Middle Pleistocene of Contrada Monticelli (Putignano, Apulia,
717 southern Italy). *Boll. Soc. Paleontol. Ital.* 56(1):71-78.

718 Mecozzi B, Iurino DA, Profico A, Rosa C, Sardella R. 2020. The wolf from the Middle
719 Pleistocene site of Ostiense (Rome): the last occurrence of *Canis mosbachensis*
720 (Canidae, Mammalia) in Italy. *Hist. Biol.* 33(10):2031-2042.

721 Montana L, Caniglia R, Galaverni M, Fabbri E, Ahmed A, Bolfíková BČ, Czarnomska SD,
722 Galov A, Godinho R, Hindrikson M, Hulva P. 2017a. Combining phylogenetic and
723 demographic inferences to assess the origin of the genetic diversity in an isolated wolf
724 population. *PloS one.* 12(5):e0176560.

725 Montana L, Caniglia R, Galaverni M, Fabbri E, Randi E. 2017b. A new mitochondrial
726 haplotype confirms the distinctiveness of the Italian wolf (*Canis lupus*) population.
727 *Mammal. Biol.* 84:30-4.

728 Pallua N, Suschek CV. 2011. Tissue engineering: from Lab to Clinic. Springer Science &
729 Business Media.

730 Pejčić A, Obradović R, Kesić L, Kojović D. 2007. Smoking and periodontal disease a
731 review. *Facta Univ. Ser. Med. Biol.* 14(2):53–59.

732 Persico D, 2021. I Fossili delle alluvioni - ricerca, studio e descrizione della paleofauna
733 padana. Edizioni Delmiglio - Persico Dosimo (CR), pp. 224, ISBN 978-88-944250-9-3

734 Persico D, Cigala Fulgosi F, Ferrari V, Ravara S, Carubelli P, 2006a. I fossili del fiume Po
735 – Catalogo dei mammiferi delle alluvioni quaternarie del museo naturalistico
736 paleontologico di San Daniele Po. Edizioni Delmiglio, Persico Dosimo (CR).

737 Persico D, Guareschi E, Carubelli P, 2006b. Resti umani fossili delle alluvioni recenti
738 rinvenuti presso San Daniele Po (CR): indagine antropologica e geopaleontologica.
739 Pianura - Scienze e storia dell'ambiente padano, 20: 147-171.

740 Pilot M, Branicki W, Jędrzejewski W, Goszczyński J, Jędrzejewska B, Dykyy I, Shkvyrya
741 M, Tsingarska E. 2010. Phylogeographic history of grey wolves in Europe. *BMC Evol.*
742 *Biol.* 10(1):1-11.

743 Pluskowski A. 2006. Where are the wolves? Investigating the scarcity of European grey
744 wolf (*Canis lupus lupus*) remains in medieval archaeological contexts and its
745 implications. *International Journal of Osteoarchaeology*. 16(4):279-295.

746 Promberger C, Schröder W. 1993. Wolves in Europe: Status and perspectives. Munich
747 Wildlife Society, Ettal, Germany.

748 Randi E, Lucchini V, Christensen MF, Mucci N, Funk SM, Dolf G, Loeschcke V. 2000.
749 Mitochondrial DNA variability in Italian and East European wolves: detecting the
750 consequences of small population size and hybridization. *Conservation Biology*.
751 14(2):464-73.

752 Reimer PJ, Bard E, Bayliss A, Beck JW, Blackwell PG, Ramsey CB, Buck CE, Cheng H,
753 Edwards RL, Friedrich M, Grootes PM. 2013. IntCal13 and Marine13 radiocarbon age
754 calibration curves 0–50,000 years cal BP. *Radiocarbon*. 55(4):1869-1887.

755 Rambaut A (2010) FigTree v1.3.1. Institute of Evolutionary Biology, University of
756 Edinburgh, Edinburgh. <http://tree.bio.ed.ac.uk/software/figtree/> Saitou N, Nei M. 1987.

- 757 The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol.*
758 *Biol. Evol.* 4(4):406–425.
- 759 Sardella R, Bertè D, Iurino DA, Cherin M, Tagliacozzo A. 2014. The wolf from Grotta
760 Romanelli (Apulia, Italy) and its implications in the evolutionary history of *Canis lupus* in
761 the Late Pleistocene of Southern Italy. *Quat. Int.* 328:179–195.
- 762 Siracusa AM, Lo Valvo M. 2005. Confronti craniometrici tra lupi (*Canis lupus*) dell'Italia
763 continentale e della Spagna: primi dati. *Hystrix.* 15(2):31-38.
- 764 Sommer R, Benecke N. 2005. Late-Pleistocene and early Holocene history of the canid
765 fauna of Europe (Canidae). *Mammalian Biology.* 70(4):227-41.
- 766 Strillou X, Boutigny H, Soueidan A, Layrolle P. 2010. Experimental animal models in
767 periodontology: a review. *Open Dent. J.* 4:37–47.
- 768 Thalmann O, Shapiro B, Cui P, Schuenemann VJ, Sawyer SK, Greenfield DL, Germonpré
769 MB, Sablin MV, López-Giráldez F, Domingo-Roura X, Napierala H. 2013. Complete
770 mitochondrial genomes of ancient canids suggest a European origin of domestic dogs.
771 *Science.* 342(6160):871-874.
- 772 Turi JO. 1931. Turi's book of Lapland. London: Jonathan Cape.
- 773 Zimen E. 1978. Der Wolf: Mythos und Verhalten. Meyster Verlag, Muenchen. 373 pp.
- 774 Zimen E, Boitani L. 1975. Number and distribution of wolves in Italy. *Z. Saugetierk.*
775 40:102-112.
- 776 Zoellner H. 1990. The vascular response in chronic periodontitis. PhD Thesis. Department
777 of Preventive Dentistry. University of Sidney.

778

779 **Figure captions**

780 **Figure 1.** Location of the site and contextual photo of the specimen at the moment of the
781 discovery.

782 **Figure 2.** *Canis lupus* MSDP 348 from Po Valley (northern Italy), in left lateral (a), right
783 lateral (b), dorsal (c), ventral (d), frontal (e) and caudal (f) views. Scale bar: 50 mm.

784 **Figure 3.** Comparison of the inner cranial anatomy. *Canis lupus* MSDP 348 (a-d), in lateral
785 (a) and dorsal (b) views, brain endocast in dorsal (c) and lateral (d) views. Extant *Canis*
786 *lupus italicus* (e-h), in lateral (e) and dorsal (f) views, brain endocast in dorsal (g) and
787 lateral (h) views. Black arrows indicate the breadth of the palate and frontal sinuses. Scale
788 bars: 50 mm.

789 **Figure 4.** Biometric comparison between *Canis lupus* MSDP 348 and a selected sample of
790 fossil and extant *Canis lupus* (for the complete list of measurements see [Figure S1](#), [Tables](#)
791 [S1](#), [S2](#)). Cranium (a), condylobasal length (**CBL**), greatest neurocranial breadth (**GNB**),
792 greatest palatal breadth (**GBP**) and lower palatal breadth (**LPB**). Upper fourth premolar
793 (b), length (**L**) and breadth (**B**). Measurements in mm.

794 **Figure 5.** Neighbour-joining (NJ) phylogenetic tree computed in MEGA using a Kimura 2-
795 parameters model. The NJ tree was obtained aligning 331 base pairs of 141 canid mtDNA
796 sequences (wolf haplotypes are represented by orange dots and dog haplotypes by green
797 dots) and was rooted using a coyote (*Canis latrans*) homologue sequence. Bootstrap
798 percentages $\geq 50\%$ after 1.000 replicates are shown by asterisks. The two main wolf
799 haplogroups (HG1 and HG2) are indicated by yellow and light blue colors. Numbers from 1
800 to 4 show the main clades of the tree. Full and empty circles distinguish extant and extinct
801 haplotypes. In the upper right box, a zoom on tree's branches close to MSDP384 is
802 represented.

803 **Figure 6.** Median joining network. (a) Late Pleistocene and Holocene (>2.000 yrs BP) wolf
804 haplotypes. (b) Holocene (<2.000 yrs BP) and current wolf haplotypes. Haplogroups (HG)

805 1 and 2 are highlighted by the two shaded areas. Black dots represent median vectors.
806 Samples included in the network refer to the 278-bp alignment and are listed in [Table S5](#).
807 **Figure 7.** Periodontitis affecting the left canine alveolus of *Canis lupus* MSDP 348. Left
808 lateral photo (a) and inner view of the alveolus from the 3D model. The white asterisks
809 indicate the hole connecting the base of the alveolus with the nasal cavity. Scale bar: 2
810 cm.

811

812 **Table caption**

813 **Table 1.** The values for male and female wolves referred to *Canis lupus italicus* ([Siracusa](#)
814 [and Lo Valvo 2014](#)).