

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Characterization of the microbial community in ripened Pecorino Toscano cheese affected by pink discoloration

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Daghio, M., Pini, F., Espinoza-Tofalos, A., Conte, G., Mari, E., Giannerini, F., et al. (2022). Characterization of the microbial community in ripened Pecorino Toscano cheese affected by pink discoloration. *FOOD MICROBIOLOGY*, 104, 1-5 [10.1016/j.fm.2022.104006].

Availability:

This version is available at: <https://hdl.handle.net/11585/894505> since: 2022-09-22

Published:

DOI: <http://doi.org/10.1016/j.fm.2022.104006>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Matteo Daglio, Francesco Pini, Anna Espinoza-Tofalos, Giuseppe Conte, Eleonora Mari, Fabiola Giannerini, Luciana Giovannetti, Arianna Buccioni, Andrea Franzetti, Lisa Granchi, Marcello Mele, Giulia Rampazzo, Teresa Gazzotti, Elisa Zironi, Carlo Viti,

CHARACTERIZATION OF THE MICROBIAL COMMUNITY IN RIPENED PECORINO TOSCANO CHEESE AFFECTED BY PINK DISCOLORATION

Food Microbiology

Volume 104,

2022,

104006,

ISSN 0740-0020,

The final published version is available online at: <https://doi.org/10.1016/j.fm.2022.104006>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

1 Characterization of the microbial community in ripened Pecorino Toscano
2 cheese affected by pink discoloration
3

4 Matteo Daghio,^{1,*} Francesco Pini,¹ Anna Espinoza-Tofalos,² Giuseppe Conte,³ Eleonora Mari,¹

5 Fabiola Giannerini,⁴ Luciana Giovannetti,¹ Arianna Buccioni,¹ Andrea Franzetti,² Lisa Granchi,¹

6 Marcello Mele,³ Giulia Rampazzo,⁵ Teresa Gazzotti,^{5,6} Elisa Zironi,⁵ Carlo Viti¹

7
8 ¹*Dipartimento di Scienze e Tecnologie Agrarie Alimentari e Forestali, University of Florence,*

9 *Piazzale delle Cascine 18, 50144, Firenze, Italy*

10
11 ²*Dipartimento di Scienze dell'Ambiente e della Terra, University of Milano-Bicocca, Piazza della*

12 *Scienza 1, 20126, Milano, Italy*

13
14 ³*Dipartimento di Scienze Agrarie, Alimentari e Agro-ambientali, University of Pisa, Via del*

15 *Borghetto 80, 56124, Pisa, Italy*

16
17 ⁴*Caseificio Sociale Manciano, Loc. Piano di Cirignano, 58014, Manciano (GR), Italy*

18
19 ⁵*Department of Veterinary Medical Sciences - University of Bologna, via Tolara di Sopra 50,*

20 *40064, Ozzano Emilia, Italy*

21
22 ⁶*Health Sciences and Technologies-Interdepartmental Centre for Industrial Research (CIRI-SDV),*

23 *University of Bologna, via Tolara di Sopra 41/E, 40064 Ozzano Emilia, Italy*

24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

* Corresponding author: matteo.daghio@unifi.it

22
1
23
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Abstract

Pink discoloration defect can cause economic losses for cheese producers due to the impossibility to sell the defected cheese, but few knowledge is currently available on the causes of this defect. To gain more insight on the causes that lead to the formation of pink discoloration in Pecorino Toscano cheese with the Protected Designation of Origin (PDO) status, the bacterial community in defected and not defected cheese was characterized by high-throughput sequencing of bacterial 16S rRNA gene. The bacterial community in the defected cheese significantly differed compared to the control. The relative abundance of the genera *Acidipropionibacterium*, *Enterococcus*, *Escherichia/Shigella*, *Lactobacillus*, *Lentilactobacillus* and *Propionibacterium* was higher in the cheese with pink discoloration defect. The concentration of short chain fatty acids and of lactic acid in cheese was measured and a shift towards the production of propionate in the cheese with pink discoloration defect was observed. Furthermore, the possible involvement of microbially produced vitamin B₁₂ in the formation of pink discoloration was not supported by the data, since a tendency to a lower concentration of vitamin B₁₂ was measured in the defected cheese compared to the control.

Keywords: microbiota; pink discoloration; propionic bacteria; sheep cheese; vitamin B₁₂

1. Introduction

Sheep cheese includes several varieties of dairy products widely produced in Italy. Several of them, as Pecorino Toscano, have the Protected Designation of Origin (PDO, EC regulation 306/2010) status (Buccioni et al., 2012). Pecorino Toscano PDO cheese is ripened from a minimum of 20 days (for fresh cheese) to over 4 months (for ripened cheese). The ageing is carried out in conditioned chambers in which temperature and humidity favorite the proliferation of the bacteria responsible for aromas. Sometime, undesirable fermentation, due to a microbial contamination, may occur. Pink

48 discoloration (PD) of the loaf rind or of the cheese inner part is a very common defect. Pink
1
249 discoloration leads to economic losses for the producers due to the impossibility to sell the defected
3
4
50 cheese (Daly et al., 2012). Despite the negative consequences potentially linked to this defect, few
6
751 knowledge is available on the causes.

9
52 In cheese with added colorants (e.g., Cheddar) the PD defect has been often associated to the
10
11
153 degradation of the colorant itself (Daly et al., 2012). In cheese without colorant the defect has been
13
14
54 associated to Maillard browning or to the microbial activity, such as the activity of some lactobacilli
15
16
155 or propionic acid bacteria used as starter cultures (Daly et al., 2012).

18
19
56 *Thermus thermophilus* has been identified as a possible responsible of PD defect in Continental-
20
21
257 type cheese (Quigley et al., 2016). *T. thermophilus* is a Gram-negative, extremely thermophilic,
23
2458 aerobic, nonpathogenic microorganism, able to produce carotenoids (Tian and Hua, 2010), whose
25
26
59 occurrence can lead to the presence of a PD in cheese (Quigley et al., 2016). In a recent work, the
27
28
290 formation of PD close the rind of Pecorino Toscano cheese, has been attributed to the presence of
30
31
61 *Serratia liquefaciens*, a psychrotrophic and motile organism (Martelli et al., 2020). The authors
32
33
342 have hypothesized that the presence of *S. liquefaciens* on Pecorino Toscano cheese with PD defect
35
36
63 was due to an environmental contamination (Martelli et al., 2020). Despite the papers reported
37
38
394 above, which attempted to understand the mechanisms that lead to the PD defect in cheese, several
40
4165 aspects of this phenomenon remain unclear. Further understanding of the factors involved in the
42
43
66 development of PD in cheese can give the opportunity to develop strategies to avoid the formation
44
45
467 of this defect, with potential economic benefits for the producers.

48
4968 The aim of this study was to describe the microbial community associated to PD defect in Pecorino
50
5169 Toscano PDO cheese and to provide new hypotheses on the possible role of the bacterial
52
53
70 communities on the development of PD defect.

57
5872 **2. Materials and methods**

59
6073 *2.1. Samples*

74 Cheese making was realized in an industrial dairy processing plant located in Tuscany (Caseificio
1
275 Sociale Manciano, Manciano, Grosseto, Italy) and it was carried out in accordance with the
3
4
576 Protected Designation of Origin (PDO) disciplinary of Pecorino Toscano cheese: raw milk was
6
77 pasteurized and inoculated with milk cultures *Streptococcus salivarius* subsp. *thermophilus*,
8
9
78 *Lactobacillus delbrueckii* subsp. *bulgaricus*, *Lactococcus lactis* subsp. *lactis*, *Lactococcus lactis*
10
11
1279 subsp. *cremoris*. Veal rennet was added to the milk and the coagulation occurred at a temperature of
13
14
1580 35 °C within 30 minutes. The curds were broken at the size of a corn kernel. After, the curd was
16
1781 placed into 2 kg molds for the whey drain by manual pressing, located in a thermostatic chamber at
18
19
2082 35 °C for 2 h and then plunged in a salt solution (NaCl, 19% w/v) at 12 °C for 24 h. Cheese was
21
2283 ripened for 270 days at 8-10 °C.

2484 Cheese samples were collected from two lots of Pecorino cheese: the first lot was composed of non-
25
26
2785 defected cheese (4 samples were collected from 2 cheese units) and was used as control; the second
28
2986 lot was composed of non-defected cheeses (4 samples were collected from 2 cheese units), used as
30
31
3287 control, and defected cheese (8 samples were collected from 2 cheese units). Since the defected
33
3488 cheeses showed a PD shading from one flat side to the other, the slices were cut in half, the two
35
3689 samples were processed independently and considered in the defected group (Figure S1). The 8
37
38
3990 samples collected from the non-defected cheese were considered as the control group (4 cheese
40
4191 units, 2 samples for each cheese units) while the 8 samples collected from the cheese with PD (2
42
43
4492 cheese units, 4 samples each) were considered as the defected group.

48 4994 2.2. DNA extraction and 16S rRNA gene sequencing

50
5195 Around 25 g of each cheese sample were homogenized in a mixer and fat was removed. DNA was
52
53
5496 extracted from 200 mg of each homogenized sample with the DNeasy *mericon* Food Kit (Qiagen)
55
5697 according to the manufacturer's instructions.

5898 The V5-V6 hypervariable regions of the 16S rRNA gene were PCR-amplified using the 783F and
59
60
6199 1046R primers (Huber et al., 2007; Wang and Qian, 2009) as previously described (Daghio et al.,
62
63
64
65

100 2018). Briefly, the PCR was performed in $2 \times 50 \mu\text{L}$ reactions with GoTaq®Green Master Mix
1
201 (Promega Corporation, Madison, WI, USA) and $1 \mu\text{M}$ of each primer. Amplification conditions
3
4
102 were: 94°C for 5 min, 29 cycles with 94°C for 50 s, 47°C for 30 s, 72°C for 30 s and a final
6
103 elongation step of 72°C for 5 min. The amplicons were then purified with the Wizard®SV Gel and
8
104 PCR Clean-up System (Promega Corporation, Madison, WI, USA) according to the manufacturer's
10
11
105 instructions and quantified using Qubit® (Life Technologies, Carlsbad, CA, USA). All DNA
13
14
106 samples were tested for amplification inhibition by sample dilution. The libraries were sequenced
15
16
107 by MiSeq Illumina (Illumina, Inc., San Diego, CA, USA) using a $300 \text{ bp} \times 2$ paired-end protocol.
18
19
108 The sequencing produced a total of 1,656,884 reads with an average of $103,555 \pm 10,525$ reads per
20
21
109 sample (average \pm standard error).
22
23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

2.3. Bioinformatic elaboration

Bioinformatic elaborations were performed in R 4.0.3 (R Core Team, 2020) using DADA2 package
(Callahan et al., 2016), version 1.16.0. According to the quality profiles, forward reads were
truncated at 200 bases, and reverse reads were truncated at 180 bases. The first 20 bases were
removed from both the forward and the reverse reads. Low quality reads (i.e., reads with expected
errors higher than 2 and with Ns) were discarded. Specific error rates were estimated for the
forward reads and for the reverse reads. Filtered reads were dereplicated, the estimated error rates
were used to infer the Amplicon Sequence Variants (ASVs) (Callahan et al., 2017) and the reads
pairs were merged with default parameters. Chimeric sequences were removed and taxonomic
assignment (confidence 80%) for each ASV was performed against the RDP database (Cole et al.,
2014) using the assign Taxonomy function. Species assignment was performed by add Species
function (100% identity). Only the ASVs with a relative abundance of 0.01% (or higher) in at least
one sample were considered for further processing. After filtering, merging, removal of chimeric
sequences and removal of low abundance ASVs a total of 1,190,658 high-quality sequences were
obtained with an average of $74,416 \pm 7,958$ sequences per sample (average \pm standard error).

126
1
127
3
4
128
6
129
8
130
10
11
131
13
14
132
15
16
133
18
19
134
20
21
135
22
23
136
25
26
137
27
28
138
30
31
139
32
33
140
35
36
141
37
38
142
39
40
143
42
43
144
44
45
145
47
48
146
49
50
147
52
53
148
54
55
149
56
57
150
59
60
151
62
63
64
65

2.4. Chemical analyses

Short chain volatile fatty acids (SCVFAs) (C2:0, acetic; C3:0, propionic; C4:0, butyric; iso C4:0, isobutyric; C5:0, valeric; iso C5:0, isovaleric) and lactic acid were extracted as follows: 10 g of each sample were added to 50 mL of 0.1 N H₂SO₄ aqueous solution and homogenized for 2 min by UltraTurrax (IKA®-Werke GmbH & Co. KG, Staufen, Germany). After the extraction 15 mL of supernatant were centrifuged at 2500 rpm for 15 min. Five mL of the supernatant were microfiltered (0.22-µm) to remove solid particles. The resulting sample was directly injected in the HPLC apparatus using an Aminex 85 HPX-87 H ion exclusion column (300 mm × 7.8 mm; 9-µm particle size; Bio-Rad, Milan, Italy); the detection wavelength was 220 nm. The analyses were carried out applying an isocratic elution (flux 0.6 mL/min) with a 0.008 N H₂SO₄ solution as mobile phase; the injection loop was 20 µL. Individual SCVFAs and lactic acid were identified and quantified by means of an external calibration curve using a standard solution of 4.50 mg/mL of lactic acid, 5.40 mg/mL of acetic acid, 5.76 mg/mL of propionic acid, 7.02 mg/mL of butyric acid and isobutyric acid, 8.28 mg/mL of valeric acid and isovaleric acid in 0.1 N H₂SO₄ (69775, 338826, 402907, B103500, 58360, 75054, 129542, respectively; Sigma- Aldrich, Milano Italy). The molar concentration of each SCVFA and of lactic acid were estimated and the ratios lactic acid/acetic acid, lactic acid/propionic acid and propionic acid/acetic acid were calculated.

The samples of pecorino cheese were analyzed for the determination of the concentration of vitamin B₁₂. The main natural forms of cobalamin present in food are hydroxocobalamin, 5'-deoxyadenosylcobalamin, methylcobalamin and cyanocobalamin (vitamin B₁₂). All the different forms of cobalamin were converted to cyanocobalamin before analysis, because it is more stable than the others. Ultra-performance liquid chromatography coupled to triple-quadrupole mass spectrometry (UPLC-MS/MS) was used for the quantification of cyanocobalamin in Pecorino cheese. Methotrexate was used as internal standard (IS). The equipment employed consisted of a Waters Acquity UPLC® binary pump, coupled with a Waters Xevo TQ-S Micro triple quadrupole

152 mass spectrometer equipped with an ESCi™ Multi-Mode Ionization Source (Waters Corporation,
1 Milford MA, USA). Mass spectrometer operated in positive electrospray ionization (ESI+) mode
153 and analysis were performed in MRM (multiple reaction monitoring) mode, following two specific
2 transitions for the target analytes: 678.43 > 147.15, 678.43 > 359.15 for vitamin B₁₂ and 455.25 >
154 308.17, 455.25 > 134.17 for methotrexate (IS). The chromatographic separation was achieved on a
3 Waters Acquity BEH C18 UPLC® column (Waters Corporation, Milford MA, USA). The
4 chromatographic conditions were set as follow: constant flow of 0.350 mL/min; the mobile phase
155 was 5 mM ammonium formate in water acidified with 0.05% of formic acid (A) and acetonitrile
5 with 0.3% of formic acid (B). The extraction procedure was performed on 1 g of Pecorino cheese in
6 accordance with the protocol described by Zironi et al., 2013, with some modifications due to the
7 high lipid component that characterizes the matrix.
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

164 2.5. Statistical analysis

165 Data from 16S rRNA gene amplicons sequencing were further processed using the vegan package,
166 version 2.5.7 (Oksanen et al., 2020) in R 4.0.3 (R Core Team, 2020). To estimate the alpha-
167 diversity within each samples group a randomly rarefied dataset (28,000 sequences) was generated,
168 then, the Chao1 index and the Shannon diversity index were calculated. A non-metric
169 multidimensional scaling (NMDS) and a permutational multivariate analysis of variance
170 (PERMANOVA) based on Hellinger transformed ASV abundance data were performed using the
171 metaMDS and the adonis2 functions, respectively. Both the NMDS and the PERMANOVA were
172 performed on the Bray-Curtis dissimilarity index. A Kruskal-Wallis test (function `kruskal.test`) was
173 performed to identify the ASVs and the genera with a different relative abundance between the
174 defected cheese and the control cheese.

175 A Kruskal-Wallis test was performed in R 4.0.3 (R Core Team, 2020) to identify the significant
176 differences in the content of SCVFAs and of lactic acid, and in the content of vitamin B₁₂.

178 **3. Results and discussion**

179 The taxonomic composition of the microbial community of Pecorino Toscano PDO cheese was
180 investigated by high-throughput sequencing of 16S rRNA gene amplicons to identify bacteria
181 potentially associated to PD defect. The rarefaction analysis, performed on the detected ASVs,
182 indicated that the sequencing depth was enough to describe the biodiversity within the samples
183 (Figure S2). The presence of 74 ASVs was observed within the whole dataset: 18 ASVs were
184 detected only in the control cheese, 19 ASVs were detected only in the defected cheese and 37
185 ASVs were shared between the two groups (Figure S3). No differences in Chao1 index and in the
186 Shannon diversity index were observed between the defected cheese and the control cheese (Figure
187 1).

188 The NMDS plot clearly showed a separation between the bacterial community enriched in the
189 defected cheese and the bacterial community enriched in the control cheese (Figure 2). The
190 difference between the bacterial communities enriched in the two conditions was further confirmed
191 by the PERMANOVA ($R^2 = 0.37$, $p = 0.002$).

192 The starter culture was composed by microorganisms belonging to the genera *Streptococcus*,
193 *Lactobacillus* and *Lactococcus* which, in total, accounted for ~91 % of the sequences in the control
194 cheese and for ~88% of the sequences in the defected cheeses (Table 1).

195 To date the main genera that have been clearly associated with the cheese PD defect were *Serratia*
196 in Pecorino Toscano cheese (Martelli et al., 2020) and *Thermus* in continental cheese (Quigley et
197 al., 2016). Sequences classified within the genera *Serratia* and *Thermus* were not detected in the
198 whole dataset (Table S1). Furthermore, the presence of *Serratia* was linked to PD close to the rind
199 of cheese (Martelli et al., 2020), while in this study the discoloration defect was observed in the
200 inner part of the cheese (Figure S1). It is therefore possible to exclude the involvement of these
201 genera in the formation of PD defect in the present study. Quigley and colleagues also observed that
202 the pink color in presence of *T. thermophilus* was more intense raising the levels of *Lactobacillus*
203 *helveticus* in the starter culture and maintaining *Streptococcus thermophilus* at the same level, but

204 the biological reason of this difference was not determined (Quigley et al., 2016). A similar pattern
1
205 in the relative abundance of these two genera was observed also in this work with a higher ($p <$
3
4
206 0.05) ratio *Lactobacillus/Streptococcus* in the cheese with PD (0.024 ± 0.002) compared to the
6
207 control cheese (0.017 ± 0.002). Furthermore, Martelli and colleagues isolated different strains of
8
208 *Enterobacter* spp. together with *S. liquefaciens* from the pink spots on the rind of Pecorino Toscano
10
1209 (Martelli et al., 2020) and despite the PD was not attributed to the genus *Enterobacter* it is
13
14
210 interesting to observe that, in our study, the relative abundance of this genus was higher ($p < 0.01$)
15
16
211 in the samples collected by the defected cheese (Table 1).

18
19
212 Another difference possibly linked to PD defect is related to *Propionibacteriaceae* family. Three
20
21
213 ASVs were classified within this family, one ASV (ASV_21) was close to *Propionibacterium*
23
24
214 *freudenreichii* (Table S2), and the other two ASVs (ASV_27 and ASV_64) were close to
25
26
215 *Acidipropionibacterium olivae* and *Acidipropionibacterium damnosum* (Table S2), previously
27
28
216 classified as *Propionibacterium olivae* and *Propionibacterium damnosum* (Turgay et al., 2020),
30
31
217 both isolated from spoiled packaged green olives (Lucena-Padrós et al., 2014). However, ASV_27
32
33
218 and ASV_64 show a high similarity also to *Acidipropionibacterium jensenii* and to
35
36
219 *Acidipropionibacterium thoenii* (Table S2), two propionic bacteria associated to dairy products
37
38
220 (Turgay et al., 2020).

40
41
221 In our study, *P. freudenreichii* was more abundant in the defected cheese (Table 1, $p < 0.01$). *P.*
42
43
222 *freudenreichii* subsp. *shermanii* was previously described as a possible agent of PD in Swiss
45
46
223 cheese, but no mechanism was suggested (Daly et al., 2012; Park et al., 1967). Regarding the genus
47
48
224 *Acidipropionibacterium*, the other member of the family *Propionibacteriaceae*, detected in the
49
50
225 dataset (Table S1), its presence was observed only in the defected cheese (Table 1, $p < 0.01$).
52
53
226 Members of the genus *Acidipropionibacterium* can produce red pigments that can cause
54
55
227 discoloration defect in cheese (Turgay et al., 2020), therefore it is possible to hypothesize that
57
58
228 members of the family *Propionibacteriaceae* were the main actors involved in the formation of PD
59
60
229 defect in the Pecorino Toscano PDO cheese. The higher relative presence of propionic bacteria in
62
63
64
65

230 the defected cheese has been evaluated by sequencing of the 16S rRNA gene, therefore there are no
1
231 clues on their metabolic activity within the cheese. Propionic bacteria are characterized by
3
4
232 propionic fermentation, which leads to the production of acetate and propionate using lactate as the
6
233 substrate (Turgay et al., 2020). Therefore, to obtain more information on their activity the SCVFAs
8
9
234 and the lactic acid molar concentrations (Figure S4) and percentages were measured (Figure 3A).
10
11
235 Furthermore, the lactic acid/acetic acid, the lactic acid/propionic acid and the propionic acid/acetic
13
14
236 acid ratios were calculated (Figure 3B). The percentage of lactic acid was higher in the control
15
16
237 cheese compared to the defected cheese ($p < 0.01$), while the percentage of propionic acid ($p <$
18
19
238 0.01) and butyric acid ($p < 0.05$) was higher in the cheese with PD defect (Figure 3A). The ratios
20
21
239 lactic acid/acetic acid and lactic acid/propionic acid were considerably higher ($p < 0.01$) in the
23
24
240 control cheese compared to the defected cheese. These data suggest a higher consumption of lactate
25
26
241 for the propionic fermentation in the cheese in the presence of PD (Figure 3B). This observation is
27
28
242 in accordance with the highest abundance of propionic bacteria in the samples collected from
30
31
243 cheese with PD. Furthermore, the higher ratio propionic acid/acetic acid in the defected cheese ($p <$
32
33
244 0.01) suggested the presence of different pathways of propionic fermentation in the two groups of
35
36
245 cheese. Indeed, the fermentation seemed to shift towards the production of propionate in the cheese
37
38
246 with PD defect. Therefore, a change in the activity of propionic bacteria could be involved in the
40
41
247 production of PD defect in this study.

42
43
248 A possible metabolite responsible for PD could be vitamin B₁₂: vitamin B₁₂ is pink and it could be
45
46
249 produced by *P. freudenreichii* when it is growing in food-like conditions (Deptula et al., 2017).
47
48
250 Furthermore, microorganisms within the genera *Escherichia/Shigella* and *Lactobacillus*, which are
49
50
251 more abundant in the defected cheese (Table 1), have been shown to possess the genes encoding for
52
53
252 the enzymes required in vitamin B₁₂ biosynthetic pathway (Balabanova et al., 2021). There are four
54
55
253 forms of vitamin B₁₂: cyanocobalamin, hydroxocobalamin, methylcobalamin and
57
58
254 adenosylcobalamin (Prentice et al., 2013). Hydroxycobalamin, adenosylcobalamin, and
59
60
255 methylcobalamin are the major forms of vitamin B₁₂ in bovine milk and hard cheese (Gille and

256 Schmid, 2015). The involvement of vitamin B₁₂ in the PD defect in this study was excluded since
1
257 the content of vitamin B₁₂ did not differ in the defected cheese (24 ± 1 ng/g) compared to the
2
3
4
258 control cheese (29 ± 2 ng/g), though a tendency (p < 0.1) to a lower concentration in the defected
5
6
259 cheese was observed (Figure S5). These values are in accordance to the values reported in literature
7
8
260 for other dairy products (Gille and Schmid, 2015; Souci et al., 2008) since a vitamin B₁₂
9
10
11
261 concentration of 3.8 ng/g was reported for curd, but a concentration ranging from 10 ng/g to 31 ng/g
12
13
14
262 was reported for other cheeses (Gille and Schmid, 2015). The work of Prentice et al. (2013), on
15
16
263 formation of pink color in therapeutic proteins, suggests a threshold between 300 and 500 ng/g
17
18
19
264 hydroxycobalamin for visible pink. This threshold, even if referred to a different matrix, is well
20
21
265 above the average values found in this work.
22
23

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

4. Conclusions

Pink discoloration is an important issue in cheese manufacturing but its causes are still not completely understood. In this study, bacteria belonging to the genera *Thermus* and *Serratia*, which have been associated to PD defect were excluded as their presence was not observed in the samples analyzed. As well, data do not support the involvement of vitamin B₁₂ in the formation of PD. Considering the differences observed in the bacterial community of Pecorino Toscano PDO cheese with PD defect, the involvement of microorganisms belonging to the genera *Propionibacterium* and *Lactobacillus*, has been hypothesized. These genera were more abundant in the defected cheese compared to the control and a different propionic fermentation was observed in the two groups of cheese. Anyway, further studies could be helpful to elucidate the mechanisms that causes PD in Pecorino cheese.

References

Balabanova, L., Averianova, L., Marchenok, M., Son, O., Tekutyeva, L., 2021. Microbial and genetic resources for cobalamin (vitamin B₁₂) biosynthesis: from ecosystems to industrial

- 282 biotechnology. Int. J. Mol. Sci. 22, 4522. <https://doi.org/10.3390/ijms22094522>
- 283 Buccioni, A., Minieri, S., Conte, G., Benvenuti, D., Pezzati, A., Antongiovanni, M., Rapaccini, S.,
284 Mele, M., 2012. Changes in conjugated linoleic acid and C18:1 isomers profile during the
285 ripening of Pecorino Toscano cheese produced with raw milk. Ital. J. Anim. Sci. 11, e75.
286 <https://doi.org/10.4081/ijas.2012.e75>
- 287 Callahan, B.J., McMurdie, P.J., Holmes, S.P., 2017. Exact sequence variants should replace
288 operational taxonomic units in marker-gene data analysis. ISME J. 11, 2639–2643.
289 <https://doi.org/10.1038/ismej.2017.119>
- 290 Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P., 2016.
291 DADA2: high-resolution sample inference from Illumina amplicon data. Nat. Methods 13,
292 581–583. <https://doi.org/10.1038/nmeth.3869>
- 293 Cole, J.R., Wang, Q., Fish, J.A., Chai, B., McGarrell, D.M., Sun, Y., Brown, C.T., Porras-Alfaro,
294 A., Kuske, C.R., Tiedje, J.M., 2014. Ribosomal Database Project: data and tools for high
295 throughput rRNA analysis. Nucleic Acids Res. 42, D633–D642.
296 <https://doi.org/10.1093/nar/gkt1244>
- 297 Daghighi, M., Espinoza Tofalos, A., Leoni, B., Cristiani, P., Papacchini, M., Jalilnejad, E., Bestetti,
298 G., Franzetti, A., 2018. Bioelectrochemical BTEX removal at different voltages: assessment of
299 the degradation and characterization of the microbial communities. J. Hazard. Mater. 341,
300 120–127. <https://doi.org/10.1016/j.jhazmat.2017.07.054>
- 301 Daly, D.F.M., McSweeney, P.L.H., Sheehan, J.J., 2012. Pink discoloration defect in commercial
302 cheese: a review. Dairy Sci. Technol. 92, 439–453. <https://doi.org/10.1007/s13594-012-0079-0>
- 303 Deptula, P., Chamlagain, B., Edelmann, M., Sangsuwan, P., Nyman, T.A., Savijoki, K., Piironen,
304 V., Varmanen, P., 2017. Food-like growth conditions support production of active vitamin B12
305 by *Propionibacterium freudenreichii* 2067 without DMBI, the lower ligand base, or cobalt
306 supplementation. Front. Microbiol. 8, 368. <https://doi.org/10.3389/fmicb.2017.00368>
- 307 Gille, D., Schmid, A., 2015. Vitamin B12 in meat and dairy products. Nutr. Rev. 73, 106–115.

- 308 <https://doi.org/10.1093/nutrit/nuu011>
1
- 309 Huber, J.A., Mark Welch, D.B., Morrison, H.G., Huse, S.M., Neal, P.R., Butterfield, D.A., Sogin,
3
4
310 M.L., 2007. Microbial population structures in the deep marine biosphere. *Science*. 318, 97–
5
6
311 100. <https://doi.org/10.1126/science.1146689>
8
- 312 Lucena-Padrós, H., González, J.M., Caballero-Guerrero, B., Ruiz-Barba, J.L., Maldonado-
10
11
313 Barragán, A., 2014. *Propionibacterium olivae* sp. nov. and *Propionibacterium damnosum* sp.
13
14
314 nov., isolated from spoiled packaged Spanish-style green olives. *Int. J. Syst. Evol. Microbiol.*
15
16
315 64, 2980–2985. <https://doi.org/10.1099/ijs.0.063032-0>
18
- 316 Martelli, F., Bancalari, E., Neviani, E., Bottari, B., 2020. Novel insights on pink discoloration in
20
21
317 cheese: the case of Pecorino Toscano. *Int. Dairy J.* 111, 104829.
23
318 <https://doi.org/10.1016/j.idairyj.2020.104829>
25
- 319 Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R.,
27
28
320 O’Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., 2020.
30
31
321 vegan: Community Ecology Package. R package version 2.5-7. [https://CRAN.R-](https://CRAN.R-project.org/package=vegan)
32
33
322 [project.org/package=vegan](https://CRAN.R-project.org/package=vegan).
35
- 323 Park, H.S., Reinbold, G.W., Hammond, E.G., 1967. Role of Propionibacteria in split defect of
37
38
324 Swiss cheese. *J. Dairy Sci.* 50, 820–823. [https://doi.org/10.3168/jds.S0022-0302\(67\)87528-3](https://doi.org/10.3168/jds.S0022-0302(67)87528-3)
40
- 325 Prentice, K.M., Gillespie, R., Lewis, N., Fujimori, K., McCoy, R., Bach, J., Connell-Crowley, L.,
42
43
326 Eakin, C.M., 2013. Hydroxocobalamin association during cell culture results in pink
44
45
327 therapeutic proteins. *MAbs* 5, 974–981. <https://doi.org/10.4161/mabs.25921>
47
- 328 Quigley, L., O’Sullivan, D.J., Daly, D., O’Sullivan, O., Burdikova, Z., Vana, R., Beresford, T.P.,
49
50
329 Ross, R.P., Fitzgerald, G.F., McSweeney, P.L.H., Giblin, L., Sheehan, J.J., Cotter, P.D., 2016.
52
53
330 *Thermus* and the pink discoloration defect in cheese. *mSystems* 1, e00023-16.
54
55
331 <https://doi.org/10.1128/msystems.00023-16>
57
- 332 R Core Team, 2020. R: A language and Environment for Statistical Computing. R Foundation for
59
60
333 Statistical Computing, Vienna, Austria. <https://www.r-project.org/>.
61
62
63
64
65

334 Souci, S., Fachmann, W., Kraut, H., 2008. Food Composition and Nutrition Tables, 7th ed. St. ed.
 1
 335 Medpharm Scientific Publishers, CRC Press, New York, Washington D.C.
 3
 4
 336 Tian, B., Hua, Y., 2010. Carotenoid biosynthesis in extremophilic *Deinococcus-Thermus* bacteria.
 5
 6
 337 Trends Microbiol. 18, 512–520. <https://doi.org/10.1016/j.tim.2010.07.007>
 7
 8
 338 Turgay, M., Bachmann, H.-P., Irmeler, S., von Ah, U., Frö Hlich-Wyder, M.-T., Falentin, H.,
 9
 10
 11
 1339 Deutsch, S.-M., Jan, G., Thierry, A., 2020. *Propionibacterium* spp. and
 12
 13
 14
 340 *Acidipropionibacterium* spp., in: Reference Module in Food Sciences. Elsevier.
 15
 16
 1341 <https://doi.org/10.1016/B978-0-08-100596-5.23016-3>
 17
 18
 19
 342 Wang, Y., Qian, P.-Y., 2009. Conservative fragments in bacterial 16S rRNA genes and primer
 20
 21
 2243 design for 16S ribosomal DNA amplicons in metagenomic studies. PLoS One 4, e7401.
 23
 24
 244 <https://doi.org/10.1371/journal.pone.0007401>
 25
 26
 345 Zironi, E., Gazzotti, T., Barbarossa, A., Devicienti, C., Scardilli, M., Pagliuca, G., 2013. Technical
 27
 28
 2946 note: Development and validation of a method using ultra performance liquid chromatography
 30
 31
 347 coupled with tandem mass spectrometry for determination of vitamin B12 concentrations in
 32
 33
 348 milk and dairy products. J. Dairy Sci. 96, 2832–2836. <https://doi.org/10.3168/jds.2012-6451>
 35
 36
 349

351
1
352
3
353
5
6
354
8
355
10
11
356
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
357
39
40
358
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Tables

Table 1 – Average relative abundance of the microorganisms detected in the defected cheese and in the control cheese. Only the genera with a relative abundance of 0.1%, or higher, in at least one sample are reported. Values are reported average \pm standard error. N.D. = not detected (i.e., relative abundance = 0). Significance codes: * $p < 0.05$; ** $p < 0.01$.

Genus	Control (%)		Defect (%)		p-value
	Average	Std. Error	Average	Std. Error	
<i>Acidipropionibacterium</i>	N.D.		0.06	0.02	0.001 **
<i>Corynebacterium</i>	0.01	0.01	N.D.		0.317
<i>Enterococcus</i>	0.01	0.01	0.06	0.02	0.008 **
<i>Escherichia/Shigella</i>	0.07	0.02	0.19	0.04	0.046 *
<i>Lacticaseibacillus</i>	1.78	0.64	0.75	0.18	0.142
<i>Lactiplantibacillus</i>	0.41	0.13	0.49	0.08	0.401
<i>Lactobacillus</i>	1.44	0.18	1.96	0.18	0.046 *
<i>Lactococcus</i>	4.49	0.76	4.07	0.42	0.674
<i>Lentilactobacillus</i>	1.37	0.38	3.58	0.49	0.006 **
<i>Leuconostoc</i>	0.67	0.15	0.30	0.19	0.027 **
<i>Pediococcus</i>	0.24	0.07	0.04	0.02	0.004 **
<i>Propionibacterium</i>	0.06	0.06	0.09	0.02	0.016 **
<i>Streptococcus</i>	84.94	1.90	81.87	0.92	0.208
<i>Weissella</i>	0.00	0.00	0.04	0.02	0.371
Other genera	0.06	0.01	0.07	0.01	
Unclassified	4.44	0.20	6.44	0.45	

359
1
360
3
361
5
6
362
8
363
10
11
364
12
13
365
15
16
366
17
18
367
20
21
368
22
23
369
25
26
370
27
28
371
29
30
372
32
33
373
34
35
374
37
38
375
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure captions

Figure 1 – Alpha diversity indexes calculated for ASV abundance. The alpha diversity is not different in the defected cheese compared to the control cheese.

Figure 2 - Non Metric Mutidimensional Scaling (NMDS) based on the Bray-Curtis distance, calculated on the Hellinger transformed ASV relative abundance data. Stress = 0.061. The bacterial community enriched in the defected cheese and the bacterial community in the control cheese were different.

Figure 3 – Relative abundance of the detected SCVFAs and lactate in the control cheese and in the defected cheese (A). The proportion of lactic acid was lower in the defected cheese, and the proportion of propionic acid was lower in the control cheese. Ratios of fatty acids and of lactic acid in the control cheese and in the defected cheese (B). Lactic acid/acetic acid and lactic acid/propionic acid ratios were higher in the control cheese, while propionic acid/acetic acid ratio was higher in the defected cheese. Significance codes: * $p < 0.05$; ** $p < 0.01$.






Figure 1






Figure 3