

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

*Helicrysum italicum* (roth) G. Don, a promising species for the phytostabilization of polluted mine sites: A case study in the Montevecchio mine (Sardinia, Italy)

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Greggio, N., Buscaroli, A., Zannoni, D., Sighinolfi, S., Dinelli, E. (2022). *Helicrysum italicum* (roth) G. Don, a promising species for the phytostabilization of polluted mine sites: A case study in the Montevecchio mine (Sardinia, Italy). JOURNAL OF GEOCHEMICAL EXPLORATION, 242, 1-15 [10.1016/j.gexplo.2022.107088].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/895129> since: 2024-05-22

*Published:*

DOI: <http://doi.org/10.1016/j.gexplo.2022.107088>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

# ***Helicrysum italicum* (roth) G. Don, a promising species for the phytostabilization of polluted mine sites: a case study in the Montevecchio mine (Sardinia, Italy).**

Greggio Nicolas\*, Buscaroli Alessandro, Zannoni Denis, Sighinolfi Silvia, Dinelli Enrico.

Biological, Geological, and Environmental Sciences Department (BiGeA) - Alma Mater Studiorum – Università di Bologna, Operative Unit of Ravenna, Via S. Alberto, 163 - 48123 Ravenna, Italia.

Tel.: +39 0544 937311; Fax.: +39 0544 937303;

\*Corresponding author: Nicolas Greggio, e-mail address: nicolas.greggio2@unibo.it

## **Abstract**

Mine exploitations worldwide have generated a great amount of tailings, which still contain large quantities of Potentially Harmful Elements (PHEs) able to contaminate soil, water, air, wildlife, and the food chain. Phytoremediation is an option to immobilize and/or extract PHEs from polluted mining areas. This study aims at assessing the phytoremediation properties of *Helicrysum italicum* (roth) G. Don, and in particular the capacity to absorb, transfer and accumulate some PHEs, such as Cd, Cu, Ni, Pb and Zn, in the plant tissues. A restricted literature review (7 papers) is also proposed in order to outline the *H. italicum*'s behaviour and verify its possible use in phytoremediation strategy of polluted mine soils. A number of 22 contaminated sites from Montevecchio mine area (Sardinia, Italy) were sampled and the results compared with 6 uncontaminated sites. In each site both rhizospheric soil materials and *H. italicum* plants were sampled. Total composition and bioavailable fractions were analysed on soil samples. *Helicrysum italicum* roots, stems and leaves were separately analysed to extract PHEs; root/soil and leaf/root ratios were calculated to elucidate plant behaviour. Results show that Cd and Zn are the most bioavailable PHEs in contaminated sites compared to uncontaminated ones (300 and 500 folds, respectively), while Cd, Cu, Pb and Zn exceed the thresholds of the Italian environmental legislation when aqua regia extraction is executed. *Helicrysum italicum* plants growing on contaminated soils accumulate significantly more Cd, Ni, Pb and Zn than plants growing on uncontaminated soils, while no significant differences are found for Cu. For all considered PHEs the root/soil ratios are > 1 in both contaminated/uncontaminated sites meaning that *H. italicum* can be defined as a tolerant species for remediation of metal-polluted soils. The leaf/root ratios weakly > 1 of Zn, Pb and Cu and < 1 of Ni and Cd indicate *H. italicum* not suitable for phytoextraction. Our results are aligned with the available literature indicating *H. italicum* a tolerant species, especially for Cd,

Pb and Zn. The low leaf/root ratios, along with its being a spontaneous and perennial species able to propagate seeds directly on contaminated soils, recommended the use of *H. italicum* as pioneering strategy for the phytostabilization.

## Keywords

*Helicrysum italicum* (roth) G. Don, Potentially Harmful Elements, phytoremediation, mine soils, literature review

## 1. Introduction

Every historical civilization' advancement has required mined resources and also future human development and green technologies will depend on extensive Earth-extracted resources (Mills, 2020). Mining activities are widespread in the world, locally modifying the original environment and impacting biota even many years after their dismissal (Camizuli et al., 2018). Estimations for Europe, China and the USA reveal that about 0.1% of the land is represented by abandoned mining sites (Arbogast et al., 2000; EUROSTAT, 2012; Lin and Ho, 2003; Perez, 2012). Mine exploitation generates large amounts of tailings, which can still contain high quantities of Potentially Harmful Elements (PHEs). Tailings are poorly colonized by vegetation because of unfavourable chemical-physical conditions for plant growth (Martínez-Sánchez et al., 2012), thus PHEs could easily spread out in the environment contaminating soil, water, air, and wildlife reaching also the food chain (Dore et al., 2020). In many cases PHEs are essential elements at a low concentration (e.g., Cu, Zn, Mn, Fe) that turn to be toxic to one or more species when reaching higher concentrations (Bini and Wahsha, 2014; Vamerali et al., 2010). Other elements (e.g., As, Cd, Cr, Ni, and Pb) show toxic impacts on plants and animals even at low concentrations (Singh et al., 2011).

Remediation is the only intervention to reduce or definitively solve the environmental contamination problem. Conventional remediation technologies are often expensive, labour-intensive, destructive, and not eco-friendly (Meuser, 2013; Yao et al., 2012). On the other hand, phytoremediation, defined as the use of plants for degradation of xenobiotics or extraction/immobilization of PHEs from water or soil substrates (USEPA, 2000), is cost-effective, widely acceptable, sustainable, applicable in large areas, and economically exploitable, particularly when native plants are used (Pandey et al., 2015; 2016).

Plants for phytoremediation must be resistant to both contaminants and unfavourable climatic conditions like drought and heat, especially in Mediterranean areas (Poschenrieder et al., 2012), and should display a high growth rate to absorb considerable quantities of toxic

elements in their tissues (Mendez and Maier, 2008). For these reasons, plants are often native of the environment in which they will be used (Yoon et al., 2006).

Phytoremediation includes five types of strategies adopted by plants: phytoextraction/phytoaccumulation, phytostabilization, phytodegradation, rhizofiltration and phytovolatilization; only the first two are considered for PHEs (Mahar et al., 2016; Pandey and Bajpai, 2019). As reported by Mendez and Maier (2008), plants for phytostabilization should accumulate PHEs in roots and not transfer them to shoots, to avoid further transfer into the food chain. Instead, plants eligible for phytoextraction should be tolerant to PHEs, absorb and accumulate them in the aboveground plant parts, grow fast and be easy to harvest (Mendez and Maier, 2008).

A plant's phytoremediation capacity is generally assessed by means of a large number of different quantitative indicators (Buscaroli, 2017). These are calculated as ratios between element contents in aerial parts and roots, or as ratios between element contents in plant parts and soil. In literature, the element concentration in soil is assessed by adopting different analytical procedures such as X-ray fluorescence or by several wet extraction methods (e.g., Aqua regia, EDTA, DTPA, etc.) and thus, resulting ratios between plant parts could significantly differ. Abreu et al. (2008) defined the Bioconcentration Coefficient (BC) the ratio between the element content in leaves and available fraction of the corresponding soil element, extracted with DTPA aqueous solution. Plants are considered tolerant when the BC value is greater than 1 ( $BC > 1$ ). Similarly, Sidhu et al. (2017) named the same abovementioned ratio Bioconcentration Factor (BCF) and stated that the BCF values  $>1$  indicate the potential of a plant species for remediation of metal polluted soils. Regardless of the name, when the root/soil ratio is  $> 1$ , the plant is considered a tolerant species (Abreu et al., 2008) useful for remediation of metal polluted soils (Sidhu et al., 2017). Moreover, when the leaf/root ratio is  $> 1$  the element is efficiently transferred from roots to shoots proving that the plant is a phytoextractor while, if the leaf/root ratio is  $< 1$  no element translocation occurs, and the plant is suitable for phytostabilization (Bolan et al., 2011).

*Helicrysum italicum* (roth) G. Don is a perennial subshrub of the genus *Helicrysum* of the family Asteraceae, characteristic of the Mediterranean area, and it grows on barren, dry, sandy and poorly developed soils in a wide altitudinal range from the sea level up to 2200 m a.s.l. (Galbany-Casals et al., 2011; Ninčević et al., 2019). The scientific and industrial interest for *H. italicum* is increasing due to its rusticity, versatile biological activities, cosmetic and pharmaceutical applications, and ornamental uses (Bianchini et al., 2009; Melito et al., 2015; Ninčević et al., 2019). Moreover, for the utilization and commercialization of derivatives from *H. italicum* the European Union requires certified absence of chemical impurities and heavy elements (Bullitta et al., 2010).

106 *Helicrysum* spp., are indicated as metallophyte, metal tolerant plants growing on soils enriched  
107 or contaminated by several elements (Nkoane et al., 2003; 2007; Koosaletse-Mswela, 2015).  
108 *Helicrysum* spp. have already been considered for bio-remediation purposes in contaminated  
109 mine tailings (Bacchetta et al., 2017; 2018; Bini et al., 2017; Barbafieri et al., 2011; Cao et al.,  
110 2004; Leita et al., 1989, studied *H. italicum*) and soils (Brunetti et al., 2018, studied *H. italicum*)  
111 in Italy and in many other regions of the world (Conesa et al., 2006; 2011; García et al., 2002,  
112 2005, studied *H. decumbens*; Fitamo and Leta, 2010, studied *H. odoratissimum*; Hesami et  
113 al., 2018, studied *Helicrysum* Spp.). However, only the recent research paper by Brunetti et  
114 al. (2018) considered as many PHEs as this study, even if Brunetti's work was conducted as a  
115 pot experiment, while this work is an in situ experiment. The other papers alternatively  
116 investigated Pb, Zn, Cd and sometimes Cu. For these reasons there is still a lack of knowledge  
117 upon PHEs uptake and translocation in *Helicrysum* spp. in different environments, as well as  
118 their interaction mechanisms.

119 A previous study, conducted with the same criteria, was performed by Buscaroli et al. (2017)  
120 on *Dittrichia viscosa*, another rustic plant growing on the Montevecchio mine tailings. The ability  
121 of phytostabilization and translocation shown by *D. viscosa* in this environment, justifies also  
122 the interest for *H. italicum*.

123 The aim of this study is to evaluate *H. italicum* for phytoremediation applications in metal-  
124 contaminated sites. Major and trace elements total concentrations in soil samples were  
125 measured, while Cu, Cd, Fe, Ni, Pb and Zn were also quantified as bioavailable soil fractions  
126 to be compared with total amounts extracted from plants. Detailed objectives of this study are:  
127 i) to assess elements accumulation potential and interaction mechanisms in different parts of  
128 *H. italicum*; ii) to study elements uptake capability in the roots and the translocation to aerals  
129 plant parts; iii) to evaluate differences in plant behaviour in contaminated and uncontaminated  
130 sites; iv) to compare elements concentration and phytoremediation properties with the existing  
131 literature for *H. italicum* subspp.

132

## 133 **2. Materials and methods**

134

### 135 **2.1. Site description and sample collection**

136 Mining activities related to Pb and Zn extraction have been representing the main economic  
137 activity for centuries in the South-West Sardinian mining districts (Italy), largely impacting the  
138 environment (particularly soil and water) and landscape (Boni et al., 1999; Dore et al., 2020).  
139 In this area the mining activities ended in 1991 leaving many abandoned heaps of waste  
140 materials now exposed to gravity movements and water and wind weathering. It is estimated

141 that about 297 hectares are occupied by landfills and about 4.9 million m<sup>3</sup> is the volume of  
142 abandoned heaps. The cost for reclaiming activities is estimated at more than 485 million Euros  
143 (Italian Government, 2001). The mining area has been included in the Italian list of polluted  
144 sites since 2001 but, until today, no remediation activities have occurred.

145 In this work two broad areas were selected for the sample collection: one including the  
146 contaminated sites in the Montevecchio area (CS) and the other including the uncontaminated  
147 reference sites either close to the mining area or in the Emilia-Romagna area (US) (Fig. 1).  
148 The CS were entirely located in the Montevecchio mining district, in the Southwestern Sardinia,  
149 close to Montevecchio and Ingurtosu villages (Fig. 1A). The bedrock consists of low-grade  
150 meta-sedimentary and meta-volcanic Cambrian-Ordovician rocks, with intrusions of Arburese  
151 igneous complex occurred at the end of the Hercynian orogeny (Cuccurru et al., 2016; Moroni  
152 et al., 2019 and references within). This complex is constituted by granodiorite and leucogranite  
153 with radial fractures filled with acid and basic magmatic dykes, and with quartz and  
154 metalliferous hydrothermal deposits exploited by the Montevecchio-Ingurtosu mines (Moroni  
155 et al., 2019). The ore-veins are composed of galena, sphalerite and quartz with local intrusions  
156 of carbonates (Moroni et al., 2019). During the mines' activity (1848 - 1991), approximately 3  
157 Mt of Pb and Zn were extracted from the Montevecchio district. Nowadays several uncontrolled  
158 waste rock piles generate relevant sources of contamination (Caboi et al., 1993; Concas et al.,  
159 2006) due to the scarce vegetation cover and intense erosion.

160 The US include 2 sampling sites near but outside the Montevecchio mining district and 4 in the  
161 Appennine chain between eastern Emilia-Romagna and Tuscany (Fig. 1B). These sites are  
162 developed on different types of sedimentary materials. The two US near Montevecchio area  
163 were sampled on aeolian sandy deposits. In the Appennine, the bedrock is made up of  
164 alternations of sandstones and marls (Marnoso-Arenacea Formation) followed by a thin band  
165 of evaporitic gypsum (Gypsum Vein), formed during the Messinian salinity crisis. Close to the  
166 plain there are Pliocene clays and Pleistocene yellow sands (Lancianese and Dinelli, 2015).  
167 Among the four sampling sites, two were chosen on evaporitic gypsum (1 and 2 in Fig. 1B) and  
168 the other two on the Marnoso-Arenacea Formation (3 and 4 in Fig. 1B).

169 In these two broad areas, 28 sample sites were selected and sampled. At each site the entire  
170 *H. italicum* plants and a composite rhizospheric soil, sampled at a depth of 5-30 cm, were  
171 collected, stored in plastic bags and brought to the laboratory for analysis. Overall, 22 soil  
172 samples and 22 plants were collected in CS corresponding to the major mine tailings deposits  
173 (Fig. 1A), while 6 soil samples and 6 plants were collected in US (Fig. 1A and 1B).

174




Figure 1. Location of sampling sites in the two study areas (a and b) with their main lithologies and related legend. Modified from ISPRA (<http://sgi2.isprambiente.it/arcgis/rest/services/servizi/cartageologica500k/MapServer>).

## 2.2. Chemical analysis of solid material

Soil material was air-dried at room temperature for two weeks, crushed and sieved through a 2 mm screen, and this fraction was retained for analysis. Soil sand, silt and clay determination was performed according to the hydrometer method devised by Day (1965). Soil reaction (pH)

180 and electrical conductivity (EC) were determined in a 1:2.5 (m/V) soil-water suspension. The  
181 total limestone was determined by volumetric method according to ISO 10693 method (ISO,  
182 1995a). Total Organic Carbon (TOC) and Total Nitrogen (TN) were determined by elemental  
183 analyser Thermo Fisher EA Flash 2000 CHNS-O according to ISO 10694 method (ISO,1995b).  
184 In order to estimate the organic and carbonate content in soil, the Loss of weight On Ignition  
185 (LOI) was determined by placing a soil sample in a muffle furnace at 950 °C for 6 h and  
186 measuring the percentage weight loss, as indicated by Heiri et al. (2001).

187 In this paper the soil PHEs concentrations were analysed using three different methodologies:  
188 i) extraction with a diethylenetriaminepentaacetic acid-based solution (hereafter DTPA)  
189 (element available for root uptake); ii) aqua regia extraction (hereafter AR) (pseudo-total  
190 element concentration); iii) X-Ray Fluorescence (hereafter XRF) (total element concentration)  
191 (Wang et al., 2021).

192 The bioavailable elements in soil (Cd, Cu, Fe, Ni, Pb and Zn) were extracted with a DTPA-  
193 based solution according to Lindsay and Norvell (1978) and ISO 14870 methods (ISO, 2001).  
194 Concentrations of Fe, Zn, Pb for all sites and Cd only in CS were measured by Flame Atomic  
195 Absorption Spectrometry (FAAS) using a Perkin-Elmer Analyst 100. Instead, concentration of  
196 Cu, Ni for all sites and Cd only in US were measured by Graphite Atomic Absorption  
197 Spectrometry (GFAAS) using a Perkin-Elmer HGS-800. The analyses were performed at  
198 Biological, Geological, and Environmental Sciences Department (BiGeA Dept.) of Bologna  
199 University using calibration standards from 0.2 to 5 mg/l and from 2 to 100 µg/l for FAAS and  
200 GFAAS, respectively. As reported by Kumpiene et al. (2017), this methodology is widely  
201 applied on non-acid soils and fitted perfectly to the US but it could be not appropriated for a  
202 few soil samples with low pH in CS. However, in order to obtain comparable results, DTPA  
203 extraction was adopted for all soil and tailing samples.

204 Aqua regia extraction was executed following the ISO 11047 method (ISO, 1998) and As, Cd,  
205 Co, Cr, Cu, Fe, Ni, Pb, V and Zn were quantified by Inductively Coupled Plasma Mass  
206 Spectrometry (ICP-MS) Perkin-Elmer ELAN DRC-eat at BiGeA Dept. of the Bologna University  
207 with calibration standards from 0.01 to 5 mg/l.

208 The total concentrations of major and trace elements in soils were determined by X-Ray  
209 Fluorescence (XRF) using a Panalytical Axios 4000 spectrometer, following the analytical  
210 methodology by Franzini et al. (1972) and Leoni and Saitta (1976) for matrix correction  
211 methods.

212 To ensure accuracy and precision in soil PHEs analysis, reagent blanks and certified reference  
213 materials were used. Quality control of DTPA-extractable PHEs was performed analysing the  
214 NCSDC85102a certified reference material. The obtained recoveries (mean value  $\pm$  standard



215 deviation in %) are the following: Cd  $102 \pm 2$ , Cu  $108 \pm 4$ , Fe  $90 \pm 1$ , Ni  $91 \pm 2$ , Pb  $80 \pm 3$ , Zn  
216  $70 \pm 1$ .

217

## 218 **2.3. Chemical analysis of plants**

219 After the separation into roots, stems and leaves, the plant samples were placed in an  
220 ultrasonic bath to remove soil particles, washed with deionized water, dried in a stove at 40 °C  
221 and minced.

222 For the determination of the total element concentrations, 250 mg of each plant part was  
223 digested with a mixture of 2 ml H<sub>2</sub>O<sub>2</sub> 30% and 6 ml HNO<sub>3</sub> 65% using a microwave Milestone  
224 mls 1200 Mega. After the filtration the digested solutions were stored in 50 ml volumetric flasks.  
225 In the plant's digested solutions Fe, Zn, Pb for all sites and Cd only in CS were analysed by  
226 FAAS (Perkin-Elmer Analyst 100) while Cu, Ni for all sites and Cd only in US were analysed  
227 by GFAAS (Perkin-Elmer HGS-800). The analyses were performed at BiGeA Dept. of the  
228 Bologna University using calibration standards from 0.2 to 5 mg/l and from 2 to 100 µg/l for  
229 FAAS and GFAAS, respectively.

230 As for soil, also for plant reagent blanks and certified reference materials were used.

231 Quality control on total PHEs concentrations in plants was performed analysing the IAEA – 359  
232 (Cabbage - Trace elements) certified reference material in three replicas. The obtained  
233 recoveries (mean value  $\pm$  standard deviation in %) are the following: Cd  $127 \pm 5$ , Cu  $126 \pm 8$ ,  
234 Ni  $96 \pm 14$ , Zn  $105 \pm 7$ .

235

## 236 **2.4 Data quality control and statistical analysis**

237 On the results, several statistical elaborations were performed using R Software version 3.3.2:  
238 the Shapiro Wilk test (Shapiro and Wilk, 1965) was adopted to verify the normal distribution of  
239 the data; a Mann-Whitney non-parametric test was performed both between the different  
240 sampling sites and the different plant tissues for the evaluation of the statistical significance of  
241 the difference between the medians. The Spearman Rank Correlation Coefficient (SRCCs)  
242 (Spearman, 1904) was applied in the determination of correlations between elements in soils  
243 (both total and bioavailable) and in plants from CS using the R software version 3.3.2. The  
244 obtained significance correlations were compared with  $< 0.05$  and  $< 0.01$  p-value.

245

246

## 247 **3 Results and discussion**

248

### 249 **3.1 Soils**

250 The main compositional and textural features for the analysed soils are summarized in Table  
251 1 (the entire dataset is available in Table S1 and S2 of the Supplementary Materials). The CS  
252 revealed a sandy loam texture with little silt and clay, whereas in the US, silt and sand were  
253 the dominant soil fractions, although a wide variation in texture was present ranging from sand  
254 to silty-loam. The CS showed a neutral-sub acid pH (5.4 - 6.7 as Min-Max range), while in the  
255 US subalkaline to alkaline pH were observed (7.6 – 9.1). EC was below 0.3 dS/m in all sites,  
256 while total lime was significant in the US and negligible in almost all the CS. In both the CS and  
257 US, TOC range was large (2 – 60 g/kg) with median values around 25 g/kg. Median values of  
258 TN content were identical (2 g/kg, for both CS and US), while the high TOC/TN ratio values (>  
259 12) could be affected by *H. italicum* plant residues that hardly decompose in soils (Brady and  
260 Weil, 2010).

261 The median concentrations of major elements such as Si (282938 mg/kg), Al (89741 mg/kg),  
262 Fe (40509 mg/kg) and K (30018 mg/kg), confirmed the abundance of sheet silicates in CS  
263 (Table 1). Compared with previous results by Buscaroli et al. (2017), the dominant silicates  
264 feature was confirmed with weak increase in Fe and LOI median values and weak decrease in  
265 Ti, Al, Mg, Na and K median values. Instead, median concentrations of Ca and LOI (129691  
266 and 211908 mg/kg, respectively) had the highest values in US reflecting the calcareous  
267 contribution of the substrate in the area as recorded in stream sediment by Lancianese and  
268 Dinelli (2015).

269 In terms of total concentration (XRF) the CS were characterized by high Pb and Zn contents  
270 (median 604 mg/kg and 11455 mg/kg, respectively). Compared with soils by Buscaroli et al.  
271 (2017) from the same area, soils from this study had slightly higher Pb and Cu median values,  
272 whereas Zn median concentration was double. The Ni total content (39 mg/kg) agrees with  
273 values found by Buscaroli et al. (2017). Compared with soil samples from the adjacent  
274 Ingurtosu mine area (Barbafieri et al., 2011), this study denoted around double Zn and half Pb  
275 content (7800 and 1800 mg/kg, respectively). Mean total PHEs concentrations found in this  
276 study area were comparable with concentrations of the nearby Barraxiutta mine district (De  
277 Agostini et al., 2020) and Campo Pisanu (Bacchetta et al., 2018), except for Cu and Pb that  
278 were 5 and 4 times lower, respectively. Overall, this high variability in values denoted the strong  
279 heterogeneity of mine tailings.

280 The US had low median total concentration of PHEs with only Cr, Ni, V and Zn ranging around  
281 100 mg/kg (Table 1). These results were aligned to the soils from the GEMAS project (Reimann  
282 et al., 2014), to the streambed sediments of the same area analysed by Lancianese and Dinelli  
283 (2015), and also to the natural background maps by Regional Soil Service, although related to  
284 the adjacent plain area (RER, 2016).

Results of the AR extraction showed Cd and Zn median values (67 and 9771 mg/kg, respectively) much higher than the Italian environmental legislation thresholds (Italian Government, 2006) for an industrial use in CS (15 and 1500 mg/kg, respectively). If the thresholds for public and residential destinations are considered (for As, Cd, Co, Cu, Pb and Zn are 10, 2, 20, 120, 100, and 150 mg/kg, respectively), all median values exceeded the limit and for Cd, Pb and Zn all samples were above the thresholds (bold values in Table 1). The US samples did not reveal concentrations exceeding the national environmental prescriptions, except for mean and maximum As concentration (16 mg/kg and 36 mg/kg, respectively). Being part of a dominant calcareous unit, the high As concentration in US is determined by a direct control of calcite abundance on As bio-accessibility (Raimondi et al., 2021).

Regarding bioavailable element concentrations (DTPA-extracted) (Table 1), the CS were characterized by high median concentrations of Zn followed by Cd, Pb, Fe and Cu (630, 9, 6, 4, 4 mg/kg, respectively). In particular, the median concentrations of Zn, Cd, Pb and Cu were respectively enriched by 3, 2, 1 and 1 orders of magnitude in CS compared to US; the Ni was double in CS compared to US, while Fe was comparable. Except for Pb, all the considered PHEs showed higher maximum values compared to Buscaroli et al. (2017). In comparison with this study Bacchetta et al. (2018) reported lower mean bioavailable concentrations of Cd, Pb and Zn (3.9, 13 and 117 mg/kg, respectively) in Campo Pisanu mine district. The lower bioavailable concentrations measured in US were aligned with unpolluted data by Buscaroli et al. (2017).

The SRCCs for elements in soils from CS revealed significant positive correlations only between bioavailable Cd and Ni (0.56), bioavailable Cd and Zn (0.66) and between total and bioavailable Zn (0.72). Negative significant correlation existed between bioavailable Fe and total Zn and bioavailable Zn (-0.69 and -0.68, respectively). The negative correlation between bioavailable Zn and Pb was weak but significant (-0.52) (Table S3 of the Supplementary Materials).

Table 1. Minimum, median, maximum and mean values for soils main characteristics in sampling areas. Contaminated sites (CS) (n = 22) and Uncontaminated sites (US) (n = 6). Bold values in the aqua regia results exceed the Italian environmental legislation (Italian Government, 2006) for soils in public and residential areas.

Areas		CS				US			
Statistics		Min	Median	Max	Mean	Min	Median	Max	Mean
Sand 2000-50 µm	g/kg	241	695	874	649	335	413	954	516
Silt 50-2 µm	g/kg	87	228	677	265	13	447	538	390
Clay < 2 µm	g/kg	39	83	138	86	33	52	176	93
Reaction in H <sub>2</sub> O	pH	5.4	6.2	6.7	6.1	7.6	8.0	9.1	8.2
EC	dS·m <sup>-1</sup>	0.2	0.3	0.3	0.3	0.1	0.2	0.3	0.2
Total lime	g/kg	0	0	67	8	3	157	291	146
TOC	g/kg	2	22	34	20	2	26	60	26
TN	g/kg	0	2	3	2	0	2	5	2
TOC/TN		5	13	20	13	9	12	31	15
Major elements and LOI									
Si	mg/kg	245288	282938	353616	293225	54917	164808	408674	275377
Ti	mg/kg	1352	4040	5085	3638	499	1615	3559	1844
Al	mg/kg	33902	89741	115742	80668	22022	41253	69146	43051
Fe	mg/kg	22962	40509	78330	43247	5875	16208	39645	20045
Mn	mg/kg	845	1396	3809	1641	152	552	697	462
Mg	mg/kg	3973	7593	11630	7692	2749	7112	26934	11770
Ca	mg/kg	1102	4483	48660	7089	18775	129691	407810	149083
Na	mg/kg	1077	5142	12834	5481	824	4554	8531	4576
K	mg/kg	12274	30018	43922	28494	5767	13933	23878	14454
P	mg/kg	271	558	1006	554	128	572	700	469
LOI	mg/kg	30032	55600	122774	67108	19900	211908	234504	152533
Total elements (XRF)									
As	mg/kg	19	94	191	103	2	6	50	13
Co	mg/kg	11	16	43	19	1	3	13	5
Cu	mg/kg	1	102	706	170	2	10	26	12
Cr	mg/kg	26	57	99	55	22	65	144	80
Ni	mg/kg	11	39	73	40	5	40	105	49
Pb	mg/kg	138	604	4619	1240	10	18	22	17
V	mg/kg	39	64	114	66	10	60	161	79
Zn	mg/kg	5020	11455	41200	14873	8	29	93	39
Extractable elements (AR)									
As	mg/kg	17	<b>80</b>	<b>333</b>	<b>98</b>	10	11	<b>36</b>	<b>16</b>
Cd	mg/kg	<b>5</b>	<b>67</b>	<b>228</b>	<b>67</b>	0	0	1	0
Co	mg/kg	16	<b>23</b>	<b>33</b>	<b>23</b>	1	3	19	6
Cr	mg/kg	6	28	49	31	3	30	127	49
Cu	mg/kg	12	<b>130</b>	<b>485</b>	<b>147</b>	1	8	40	13
Ni	mg/kg	7	34	62	36	2	20	91	36
Pb	mg/kg	<b>170</b>	<b>819</b>	<b>3442</b>	<b>1235</b>	5	8	16	10
V	mg/kg	4	53	149	63	14	42	88	42
Zn	mg/kg	<b>1227</b>	<b>9771</b>	<b>27286</b>	<b>9377</b>	10	23	63	31
Bioavailable elements (DTPA)									
Cd	mg/kg	1	9	29	9	0	0	0	0
Cu	mg/kg	0	4	26	5	0	1	2	1
Fe	mg/kg	0	4	39	7	1	4	8	4
Ni	mg/kg	0	1	3	1	0	0	1	0
Pb	mg/kg	0	6	143	29	0	1	1	1
Zn	mg/kg	101	630	894	555	0	1	2	1

The Table 2 reports the DTPA/total concentration ratios expressed as a percentage and used to evaluate the elements' behaviour. The percentages were generally low in CS, especially regarding Fe, although the maximum values reached 45% and 27% for Cu and Pb,

respectively. The differences between CS and US were limited and significant only for Fe in US and Ni in CS. Regardless of the soil conditions in CS and US, Cu and Pb showed highest bioavailability followed by Zn, Ni and Fe although no systematic order was observed. Previous work by Buscaroli et al. (2017) in Montevecchio district (in brackets in Table 2) reported lower median values of Cu, Pb and Zn, even if maximum values were aligned with the ones from this study. The US ratios from this study weakly differ from Buscaroli et al. (2017) confirming that North Appennine district was well characterized by the collected samples.

Table 2. Minimum, median and maximum bioavailable/total element ratio in soils, expressed as percentage (%). Contaminated sites (CS) (n = 22); Uncontaminated sites (US) (n = 6). The letters (C for Contaminated, U for Uncontaminated) indicate the presence of statistically significant differences between percentages of each element in each area according to the Mann–Whitney test. No letter means absence of statistically significant differences between areas. The considered significant levels are p-value < 0.1 (\*) and < 0.05 (\*\*). Values between brackets are by Buscaroli et al. (2017).

Areas	Statistics	Cu	Fe	Ni	Pb	Zn
CS	Min	0.67 (1.28)	0.0003 (0.003)	0.67 (0.48)	0.09 (1.2)	1.34 (3.87)
	Median	3.96 (6.66)	0.008 (0.01)	1.41 U** (1.03)	1.53 (5.51)	3.25 (7.07)
	Max	45 (9.2)	0.116 (0.05)	6.2 (1.79)	26.6 (23.3)	10.4 (10.82)
US	Min	0.41 (1.74)	0.02 (0.01)	0.13 (0.25)	1.44 (3.8)	1.53 (0.18)
	Median	5.38 (7.75)	0.03 C* (0.03)	0.64 (0.71)	5.47 (6.03)	3.14 (1.96)
	Max	19.5 (12.6)	0.04 (0.07)	1.32 (1.52)	9.45 (12.6)	8.19 (2.63)

## 3.2 Plants

The total PHEs concentrations in the different parts of *H. italicum* plant are shown in Table 3 grouped into CS and US, while the entire database is presented in Table S4 of the Supplementary Materials. The PHEs concentrations in plants were higher in CS than in US. In particular, the differences were statistically significant for Cd, Fe, Ni, Pb and Zn (“U” in Table 3), while there were no significant differences for Cu, although median concentrations in plants were slightly higher in CS than US.

The median concentrations in leaf were generally higher than the other plant parts, although statistical significance (p-value < 0.01) occurred only for Cu, Fe and Zn in CS (leaves marked with “r” and “s” in Table 3). In US no significant differences in element concentrations among plant parts were found. Only Cu in leaves was statistically different from Cu stem concentration. In both CS and US, the concentrations of PHEs in the stems were lower than in the other parts of the plant, although they were significantly different only for Cd (compared to root content), Cu, Fe, Pb and Zn in CS and for Cu in US, mainly compared to the leaves (Table 3). The ratio among median element concentrations in plant parts from CS and US revealed the following enrichment ranking: Pb > Zn > Cd > Fe > Ni > Cu (Table 3).

353 The median concentrations of Pb in plants were 40 mg/kg in roots, 27 mg/kg in stems and 64  
 354 mg/kg in leaves with the CS/US ratios ranging from 33 to 64 for stems and leaves, respectively  
 355 (Table 3). The median values of Zn in plants were 576 mg/kg in roots, 391 mg/kg in stems and  
 356 1206 mg/kg in leaves, this last proven statistically different from the others (Table 3).  
 357 *Helicrysum Italicum* plants showed from 13 to 26 times more Zn in CS than in US. The median  
 358 concentrations of Cd in plant were quite homogeneous: 3.1 mg/kg in stems, 6.0 mg/kg in leaves  
 359 and 5.7 mg/kg in roots, these last two significantly higher than in stems. These concentrations  
 360 of Cd were from 10 to 15 times more enriched in CS than in US. Median concentration of Fe  
 361 in leaves (1523 mg/kg) was significantly different from roots (701 mg/kg) and stems (62 mg/kg)  
 362 and *H. Italicum* plants were from 3 to 5 times more Fe-enriched in CS than in US. The median  
 363 concentrations of Ni were similar in plants: 4.4 mg/kg in roots, 3.5 mg/kg in stems, 4.6 mg/kg  
 364 in leaves. Plants from CS had only 2 – 3 times more Ni than the ones from US. The median  
 365 concentrations of Cu in CS plants were 13 mg/kg in roots, 10 mg/kg in stems and 20 mg/kg in  
 366 leaves, this last significantly different from roots and stems. No differences of Cu content  
 367 existed between CS and US (median ratio was 1 in Table 3).  
 368 Few authors studied element distribution in *H. italicum* plant parts including stems. In an  
 369 adjacent mine area, Barbafieri et al. (2011) showed similar Cd concentrations and distribution  
 370 in the same plant parts with stems as the lowest accumulation site. Instead, the same authors  
 371 reported increasing concentrations in Pb and Zn from roots to leaves, but compared to this  
 372 study mean concentrations were one order of magnitude higher for Pb and slightly lower for  
 373 Zn. Also Brunetti et al. (2018) evaluated Cd, Cu, Ni, Pb and Zn abundance in roots, stems and  
 374 leaves of *H. italicum* grown in a polluted soil in Apulia Region (Italy). Their abundances of Cu  
 375 and Ni in plants were aligned with the concentrations of this study, while Cd, Pb and Zn were  
 376 one order of magnitude lower. Brunetti et al. (2018) concluded that *H. italicum* stores PHEs in  
 377 the roots with stems as the least concentrated part. Although not clearly evident in this study,  
 378 the trend to accumulate Pb in roots rather than in other aerial parts is widely demonstrated in  
 379 *H. italicum* (Barbafieri et al., 2011; Brunetti et al., 2018) and also other plants such as *Oryza*  
 380 *sativa* (Ashraf et al., 2020) and *Crambe abyssinica* (Gonçalves et al., 2020), confirming the  
 381 poor Pb translocation.  
 382 In a pyrite-mine site in Tuscany (Italy), Bini et al. (2017) showed preferential accumulation of  
 383 PHEs in *H. italicum* roots. Compared to this study, Fe and Pb concentrations in plants were  
 384 aligned, Cd and Ni were 10 times lower, Zn was three orders of magnitude lower and only Cu  
 385 resulted 10 times higher. For *H. italicum* subsp. *tyrrhenicum* (*H. tyrrhenicum*), in the adjacent  
 386 mine area of Campo Pisanu, Bacchetta et al. (2018) detected higher concentrations in roots  
 387 and leaves for all considered elements (Cd, Pb and Zn) and, especially for Pb, the differences  
 388 with this study were quite important.

Pot trials at different soil contamination of Cu and Pb executed on *Helichrysum splendidum* Less revealed a reduction of chlorophyll content (phytotoxicity sign) only when the Cu leaves concentration was about 290 mg/kg, while for the Pb a constant chlorophyll reduction was evident starting from 90 mg/kg in leaves (Banda et al., 2021). *Helichrysum italicum* never reached high concentrations of Cu in this work, while maximum concentrations of Pb exceeded 90 mg/kg in CS plants. Since it was not within the aims of the work, no surveys regarding the health status of the plants were conducted, therefore the presence of *H. italicum* plants with phytotoxic symptoms in CS could not be excluded.

Table 3. Minimum, median, maximum and mean element concentration in plant parts for Contaminated Sites (CS, n = 22) and Uncontaminated Sites (US, n = 6). Concentrations are in mg/kg. The letters R, S and L represent roots, stems and leaves, respectively. According to the Mann–Whitney test, the Statistical Significant Difference (SSD) between the analysed plant tissues within each area was indicated by the letters r and s. According to the Mann–Whitney test, the letter U indicates statistically significant difference of the plant tissue from CS with US. No letter means absence of statistically significant differences. The considered significant levels are p-value < 0.1 (\*), < 0.05 (\*\*) and < 0.01 (\*\*\*).

Elements	Plant parts	CS					US					Median CS/US ratio
		Min	Median	Max	Mean	SSD	Min	Median	Max	Mean	SSD	
Cd	L	0.5	6.0	90.0	12.4	U**	0.1	0.4	1.3	0.5		15
	S	0.6	3.1	22.3	5.8	U**	0.1	0.3	1.1	0.4		10
	R	1.4	5.7	50.9	11.1	s**U**	0.1	0.5	1.7	0.6		11
Cu	L	7.8	20.2	98.4	26.0	r*s***	9.4	13.9	29.8	16.8	s*	1
	S	3.2	10.2	32.3	12.6		6.3	9.3	11.9	9.1		1
	R	6.5	12.9	103.2	22.4		6.3	9.8	19.0	10.9		1
Fe	L	359	1523	5991	1994	r**s***U**	174	290	677	343		5
	S	267	662	1905	748	U**	112	206	427	228		3
	R	142	701	7140	1245	U**	150	252	581	319		3
Ni	L	1.3	4.6	11.3	4.8	U**	0.6	1.7	4.1	1.9		3
	S	2.0	3.5	5.6	3.8	U**	0.4	1.3	2.1	1.2		3
	R	0.6	4.4	11.0	4.7	U*	0.8	1.8	2.4	1.7		2
Pb	L	5.9	63.7	288.4	74.5	s**U**	0.6	1.0	13.0	3.6		64
	S	7.8	26.5	181.4	37.4	U**	0.5	0.8	5.9	1.8		33
	R	5.2	39.5	384.4	76.5	U**	0.4	0.9	4.4	1.6		44
Zn	L	177	1206	9837	1764	r*s**U**	28	47	112	61		26
	S	132	391	1959	630	U**	19	30	51	32		13
	R	220	576	3337	936	U**	26	28	46	32		21

### 3.2.1 Interaction mechanisms of PHEs in *H. italicum* plants

The interaction mechanisms between PHEs could reveal synergistic or antagonistic effects able to improve or reduce element uptake and translocation in plant species. Only for plants collected in CS (n = 22), the SRCCs and their significance levels were calculated for each element (Cd, Cu, Fe, Ni, Pb and Zn) and for different plant parts and results are presented in Table 4.

Results of this study indicated that Zn in *H. italicum* was positively and significantly correlated with Cd in all considered plant parts (around 0.90 for R, S and L). Moreover, Zn and Cd were also themselves correlated with values of 0.92 and 0.9 between leaves and stems. The Zn and Cd, together with Pb were the most enriched elements in *H. italicum* compared to US (Table 3). Nevertheless, Zn and Cd were also positively and significantly correlated with all the other elements in leaves (from 0.55 to 0.7), except for Pb that showed correlations only with Cd in leaves (0.67). The relation between Zn and Cd is known in literature and depends by the element similarity (Fernández et al., 2017). Indeed, Cd plant uptake is hindered by high soil Zn concentrations (Choudhary et al., 1995; Oliver et al., 1994) because they share the same transportation protein in plants and Zn is selectively preferred (Hart et al., 2002). Kutrowska et al. (2017) documented the synergistic effect between Cd and Zn in *Brassica juncea* where Zn increased the accumulation of Cd in leaves. The same authors reported that Pb increased the Cd in stems, as identified also in the present study (Table 4).

In *H. italicum* grown at CS Cu was positively correlated with itself, especially between stems and leaves (0.74), and clearly positively correlated also with Zn for almost all plant parts with a peak of 0.86 in leaves. As Fe, also Cu and Zn are micronutrients for plants and serve in physiological processes; so, their synergism was expected for US, but it was found also at elevated Zn concentration in CS (Table 3). In other species like *Brassica juncea* antagonism between Zn and Cu was demonstrated and indicated as happening not in the roots, but later during xylem loading/unloading (Kutrowska et al., 2017). A previous study (An et al., 2004) revealed that Cu and Cd act antagonistically resulting in decreased accumulation of both metals in *Cucumis sativus*. In the present study Cu and Cd were weakly correlated with coefficients around 0.6 between Cu in leaves and Cd in roots, stems and leaves (Table 4).

The Fe was significantly correlated between stems and leaves (0.8), in leaves with Pb (0.71) and in leaves with Cd (0.73 and 0.72 for stems and leaves, respectively). Moreover, Fe and Ni resulted significantly correlated in all the *H. italicum* plant parts, with coefficients ranging around 0.70 and peaks for roots (0.74) and leaves (0.71) (Table 4). Khalid and Tinsley (1980), in *Lolium perenne*, reported a common increase of Ni and Fe concentrations in shoots with increasing rates of Ni. Same synergistic effect of Ni on Fe was detected in maize with highest evidence in roots and leaves by Torres et al. (2016).

Absence of correlation existed among Pb and Cu as well as Pb and Zn in the plant parts of *H. italicum* grown at CS. An antagonistic effect of Pb on the Cu accumulation was documented also in *Brassica juncea* and related to a competition between metals at the plant uptake site (Kutrowska et al., 2017). Also for Israr et al. (2011), Pb showed antagonistic effect on the accumulation of Cu, Ni and Zn in *S. drummondii* species, probably due to the competition between metals at the plant uptake sites. Yet, Wong et al. (1986) reported a reduced uptake



449 of Cu in the presence of Pb for *Brassica chinensis*. The inhibition of essential nutrient transfer  
450 (such as Cu and Zn) in plant biomass due to Pb elevated concentration has been also proposed  
451 by Yoon et al. (2006) for numerous plants grown in Florida contaminated site. In addition to all  
452 the side effects of Pb in plants, Pourrut et al. (2011) reported impaired uptake of essential  
453 elements, such as Mg and Fe. On the contrary, An et al. (2004) showed positive correlation of  
454 Zn and Pb in *Cucumis sativus* suggesting a synergistic effect.

455

456  
457  
458  
459  
460  
461

Table 4. Main Spearman Rank Correlation Coefficients (SRCCs) and their significance levels, calculated for each element (Cd, Cu, Fe, Ni Pb and Zn) between the different plant parts (R=roots, S=stems and L=leaves) and, for each of them, between the different elements. The SRCCs were calculated considering only the Contaminated Sites (CS) (n=22). The considered significant levels are p-value < 0.05 (\*\*) and < 0.01 (\*\*\*).

		R	S	L	R	S	L	R	S	L	R	S	L	R	S	L	R	S	L
		Cd			Cu			Fe			Ni			Pb			Zn		
R	Cd	1																	
S	Cd	0.76 ***	1																
L	Cd	0.81 ***	0.92 ***	1															
R	Cu	0.55 **	0.42	0.44 **	1														
S	Cu	0.22	0.30	0.31	0.60 ***	1													
L	Cu	0.66 ***	0.56 ***	0.66 ***	0.69 ***	0.74 ***	1												
R	Fe	0.37	0.19	0.20	0.57 ***	- 0.01	0.24	1											
S	Fe	0.36	0.64 ***	0.53 **	0.47 **	0.30	0.31	0.49 **	1										
L	Fe	0.55 **	0.73 ***	0.72 ***	0.47 **	0.30	0.49 **	0.33	0.80 ***	1									
R	Ni	0.27	0.08	0.17	0.55 **	- 0.01	0.24	0.74 ***	0.32	0.37	1								
S	Ni	0.39	0.46 **	0.51 **	0.65 ***	0.20	0.37	0.55 **	0.69 ***	0.68 ***	0.62 ***	1							
L	Ni	0.64 ***	0.66 ***	0.75 ***	0.48	0.01	0.41	0.31	0.52 **	0.71 ***	0.54 **	0.73 ***	1						
R	Pb	0.12	- 0.10	- 0.07	0.39	- 0.01	0.03	0.67 ***	0.17	0.08	0.56 ***	0.36	0.09	1					
S	Pb	0.40	0.40	0.48 **	0.15	0.09	0.23	0.42	0.47 **	0.46 **	0.42	0.35	0.41	0.53	1				
L	Pb	0.46 **	0.51 **	0.67 ***	0.20	0.08	0.29	0.33	0.53 **	0.71 ***	0.41	0.49 **	0.60 ***	0.36	0.86 ***	1			
R	Zn	0.90 ***	0.68 ***	0.70 ***	0.77 ***	0.33	0.70 ***	0.50 **	0.39	0.56 ***	0.46 **	0.52 **	0.64 ***	0.26	0.34	0.37	1		
S	Zn	0.78 ***	0.93 ***	0.87 ***	0.54 **	0.45 **	0.73 ***	0.24	0.61 ***	0.66 ***	0.12	0.43	0.60 ***	- 0.15	0.35	0.40	0.74 ***	1	
L	Zn	0.87 ***	0.82 ***	0.89 ***	0.60 ***	0.45 **	0.86 ***	0.27	0.38	0.59 ***	0.19	0.42	0.60 ***	- 0.09	0.28	0.42	0.82 ***	0.90 ***	1

462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473

3.3 Soil-plant relationship

The relationship between the soil bioavailable pool and the concentrations in plant parts of the investigated elements are represented by box plots in Figure 2. In terms of soil bioavailable pool, Fe and Ni showed comparable concentrations in CS and US even if Ni in US had a wider range of variation. At these low concentrations the *H. italicum* was able to absorb and concentrate the elements in the plant (2 and 1 order of magnitude for Fe and Ni, respectively). Although the soil bioavailable fractions were similar the plant concentrations between CS and US were statistically different (Table 3) with lower levels in the latter.

The Cu and Pb were enriched in CS compared to US, but due to the wider range of variations (Cu in US and Pb in CS) a clear separation was not evident. Regarding these elements the

474 plant was able to absorb and concentrate them compared to the soil bioavailable pool (1 order  
 475 of magnitude for Pb). The Cu concentrations in the plant were not statistically different between  
 476 CS and US (Table 3) indicating that CS could not be considered polluted by Cu. In fact,  
 477 Buscaroli et al. (2017) in Cu-contaminated Libiola mine and Brunetti et al. (2018) in Apulia soil  
 478 reported bioavailable Cu concentration greater than this study by 1 and 2 orders of magnitude,  
 479 respectively. The Pb in CS was absorbed and concentrated by *H. italicum*, while in US it was  
 480 absorbed but not concentrated compared to soil bioavailable fractions (Fig. 2).  
 481 Soil bioavailable concentration of Cd and Zn were more abundant in CS than US of 2 and 3  
 482 times, respectively. In CS bioavailable Cd and Zn were elevated (more than Bacchetta et al.  
 483 (2018) and Brunetti et al. (2018)) and *H. italicum* plant absorbed them, but weakly concentrated  
 484 Cd and Zn in the plant tissues. Instead, in US *H. italicum* was able to absorb and concentrate  
 485 Cd and Zn (Fig. 2), even if the plant concentrations remained significantly lower than CS (Table  
 486 3). Despite this behaviour within plants, *H. italicum* growing in CS accumulated two orders of  
 487 magnitude more Pb and one order of magnitude more Zn and Cd compared to US (Fig. 2).  
 488 Bacchetta et al. (2018) for *H. tyrrhenicum* in the adjacent mine area of Campo Pisanu, detected  
 489 higher concentrations in plant parts for all considered elements (Cd, Pb, and Zn). Yet,  
 490 bioavailable pools for Cd and Zn were a quarter the levels of the present study, while the  
 491 bioavailable pool of Pb was three times higher. Brunetti et al. (2018), starting from bioavailable  
 492 pools like Bacchetta et al. (2018), presented notably lower plant concentrations. This behaviour  
 493 could be related to the carbonate-soils studied by Brunetti et al. (2018) that contain high  
 494 exchangeable Ca. This latter could compete with heavy metals limiting their uptake. The  
 495 antagonistic effect of Ca on Cd, Cu, Fe, Ni, Pb and Zn uptake was found by Kabata-Pendias  
 496 (2010) and observed also for *D. viscosa* by Buscaroli et al. (2017).  
 497 Possible interaction mechanisms between elements in soil and plant from CS were  
 498 investigated through SRCCs and results are presented in Table S3 of the Supplementary  
 499 Materials. In general, the correlations between soil and plant concentrations were scarce.  
 500 The only positive significant correlation existed between total Pb in soil and Pb in roots (0.75).  
 501 There were weak significant negative correlations between total Ni and Cu in stems (-0.56)  
 502 and leaves (-0.59), between bioavailable Pb and Cd in stems (-0.53) and leaves (-0.52), and  
 503 Zn in stems (-0.54) (Table S3).  
 504




Figure 2. Bioavailable Cd, Cu, Fe, Ni, Pb and Zn in soils and their distribution in *H. italicum* plants of CS and US.

### 3.4 Root/bioavailable soil element concentration ratio

With the aim of studying the tolerance to PHEs of *H. italicum* and provide evidence of phytostabilization or phytoextraction capacity, the root/bioavailable soil concentration ratio (root/soil) was calculated for each element and the results are shown in Table 5. The mean values were presented only for literature comparison.

For all the elements in CS and US the medians for root/soil ratios were > 1 suggesting that *H. italicum* was able to absorb the bioavailable elements in roots. According to root/soil median values, in the CS the elements were absorbed following the ranking Fe >> Ni > Pb > Cu, whereas the ratios for Zn and Cd were close to 1. In the US the ranking resulted Fe > Zn > Cu > Cd > Ni and Pb only slightly above 1 (Table 5).

Out of the three most abundant bioavailable elements in CS (Zn, Pb and Cd, Table 1), Cd (1.21) and Zn (1.39) had the lowest ratios indicating a limited root uptake by *H. italicum* when soil available pool is abundant. Pb was more absorbed in CS (4.17) than in US (1.21). Same behaviour for Zn, Pb and Cd was detected in *D. viscosa* for the same mine area (Buscaroli et al., 2017).

The different ratio of Cu between CS and US depended from its wide variability in US, while Ni showed similar soil/root ratios in both conditions. The Fe is a fundamental micronutrient for plants strongly related to chlorophyll content and plant growth (Terry, 1980). Despite its low availability in soil (Tables 1 and 2), *H. italicum* showed an elevated root uptake capacity for Fe, especially in CS (Fig. 2).

Many other authors calculated the root/soil ratios of *H. italicum* (Barbafieri et al., 2011; Bini et al., 2017; Cao et al., 2004; Leita et al., 1989) but only Bacchetta et al. (2018) and Brunetti et al. (2018) applied the DTPA extraction, allowing a direct comparison with ratios from this study. In particular, Bacchetta et al. (2018) found ratios of 21, 29 and 9.6 for Zn, Pb and Cd, respectively, for the *H. tyrrhenicum* in a Sardinian mine site. These ratios were higher than this study, but with lower bioavailable concentrations and significantly higher root concentrations. Recalculated ratios by Brunetti et al. (2018) showed for Cu, Ni, Pb and Zn slightly lower values, under reduced bioavailable soil pool compared to CS of this study. Instead, Cd showed root/soil ratio one order of magnitude higher and bioavailable concentrations one order of magnitude lower compared to this study in CS, but comparable with values in US (Tables 1 and 5). This confirms that Cd is strongly incorporated in roots at low soil bioavailable concentrations (Brunetti et al., 2018), while its absorption is limited when soil concentrations increase (Fig. 2 and Bacchetta et al., 2018). This represents an excluding mechanism for Cd already documented in other plant species such as *Thlaspi arvense* (Martin et al., 2012) and *Arabidopsis thaliana* (Zhu et al., 2012), but not yet in *Helicrysum* spp.

Table 5. Minimum, median, maximum and mean root/soil values of selected elements in *H. italicum*. CS n=22 and US n=6.

Area	Statistics	Cd	Cu	Fe	Ni	Pb	Zn
CS	Min.	0.14	1.45	3.60	0.81	0.36	0.35
	Median	1.21	3.03	299	6.56	4.17	1.39
	Max.	7.95	28.9	6541	29.0	52.0	5.92
	Mean	1.76	5.47	988	8.84	11.57	1.92
US	Min.	4.43	3.22	23.0	1.33	0.40	15.7
	Median	12.1	19.1	91.2	4.67	1.21	26.3
	Max.	43.1	417	135	272	8.56	74.4
	Mean	18.47	107.73	89.06	70.56	2.54	38.99
	US/CS Median ratio	10	6.3	0.3	0.71	0.29	18.9

547

548

549

### 3.5 Leaf/root element concentration ratio

550 The capacity of *H. italicum* to transfer PHEs from roots to leaves has been quantified for each  
 551 element through the leaf/root concentration ratio (leaf/root ratio) and the results are shown in  
 552 Table 6. The leaf/root ratio was extensively applied for the evaluation of the phytoextraction  
 553 capacity of plants growing in mine soils (Buscaroli et al., 2017; Martínez-Sánchez et al., 2012;  
 554 Wang et al., 2019; Yoon et al., 2006) or in contaminated agricultural soils (Dinu et al., 2020;  
 555 Nadimi-Goki et al., 2014). As stated for root/soil ratio, also leaf/root ratio is widely used by  
 556 researchers, but under different names such as Translocation factor, Transfer factor or  
 557 Transportation index (Buscaroli et al., 2017).

558 The *H. italicum* in CS always showed leaf/root median ratios > 1, except for Cd and Ni that  
 559 were 0.75 and 1, respectively (Table 6). Iron reached a value of 2.36 followed by Zn (1.54)  
 560 then Pb (1.37) and Cu (1.34). The leaf/root median values in US were comparable with CS  
 561 except for Fe that was half. The ratios of Zn, Ni, and Cd were almost identical, while slightly  
 562 higher values were measured for Pb and Cu in US compared to CS (Table 6). However, the  
 563 maximum ratios were from 2 to 10 times higher in CS than US (Table 6) for all considered  
 564 elements, indicating an increased inclination of the plants to transfer elements in leaves when  
 565 growing on contaminated soils.

566 As all the *H. spp*, also *H. italicum* is a terpene-rich species, rarely appreciated as food by  
 567 wild or domestic herbivorous, reducing the possibilities of PHEs entering the food chain  
 568 (Rogosic et al., 2006). Moreover, metal accumulation in aerial parts is an evolutionary  
 569 adaptation that confers to plants also protection against herbivores or pathogens (Galeas et  
 570 al., 2008).

571 In presence of a tolerant plant (root/soil ratio > 1), the higher the leaf/root ratio (e.g., > 2), the  
 572 greater the capacity to transfer elements to aerial plant parts and the more adapt is the species  
 573 for phytoextraction strategy (Buscaroli et al., 2017; Yoon et al., 2006). The opposite indicates  
 574 the suitability of the species for phytostabilization (Rizzi et al., 2004; Yoon et al., 2006). Results  
 575 from this study indicated *H. italicum* as a tolerant species weakly able to concentrate Fe, Zn,

Pb and Cu in the aerial parts and unable to transfer Cd and Ni to leaves. The leaf/root ratio recalculated by Barbaferi et al. (2011) in *H. italicum* plants revealed almost identical values for Zn and Cd and double values for Pb compared to the present study. Identical values for the ratio were reported also by Bacchetta et al. (2018) in Campo Pisano mine site, but for *H. tyrrhenicum* and for the entire epigeal organs. Half values of the ratios for Cd, Cu, Pb and Zn and about one tenth for Fe were obtained by Bini et al. (2017) in a Tuscany mine district, while Brunetti et al. (2018) reported lower ratios for all elements (Cu, Ni, Pb and Zn). In the recent pot trials by Banda et al. (2021) leaves/root ratios of Cu and Pb were weakly above 1 in *Helichrysum splendidum* Less.

Only for comparison purposes, Boechat et al. (2016) reported leaf/root ratios for several Brazilian species well above 2. In particular, *Baccharis trimera* (Less) DC (5.48), *Cyperus eragrostis* Lam (3.54), *Eryngium horridum* Malme (2.91) and *Dicranopteris nervosa* (Kaulf.) (2.61) for Pb and *Senecio brasiliensis* (Spreng.) Less (2.93) for Cd.

589

Table 6. Minimum, median, maximum, and mean leaf/root total element concentration values of selected elements in *H. italicum*. CS n = 22 and US n = 6.

Area	Statistics	Cd	Cu	Fe	Ni	Pb	Zn
CS	Min.	0.22	0.41	0.17	0.39	0.12	0.80
	Median	0.75	1.34	2.36	1.02	1.37	1.54
	Max.	6.91	4.96	19.3	6.59	14.3	13.1
	Mean	1.20	1.61	3.39	1.46	2.46	2.27
US	Min.	0.40	0.73	0.38	0.30	0.71	1.04
	Median	0.74	1.78	1.08	1.04	1.87	1.51
	Max.	1.35	2.98	2.88	2.11	3.00	3.67
	Mean	0.82	1.70	1.33	1.12	1.83	1.90

592

593

### 3.6 Literature comparison and final remarks

The interest for the species *H. italicum* (subsp. *italicum* or *tyrrhenicum*) as a possible phytoremediation plant, especially in abandoned mine areas, goes long back in time. From Leita et al. (1989) up to Bacchetta et al. (2018) and Brunetti et al. (2018) many authors have studied the capacity of *H. italicum*, to uptake PHEs from contaminated soils. With the aim of summarizing the available literature on these species and outline their phytoremediation capacities, a literature research browsing the keyword “*Helichrysum italicum* and remediation” in the Web of Science (last time checked 08/02/2022) was performed. Only 3 publications appeared Bini et al. (2017), Brunetti et al. (2018) and Boi et al. (2020). The work of Boi et al. (2020) was dedicated to the seed germination and for this reason excluded. This literature research demonstrated that the publications already cited in this study represent the most updated articles dealing with the application of *H. italicum* for phytoremediation purpose. All

606 the available articles considering *H. italicum* for phytoremediation (N=6) (180 total citations and  
607 maximum number 75 for Barbafieri et al., 2011) have been reviewed and their main  
608 characteristics summarized in Table 7. Particular attention has been paid to the analytical  
609 methods applied on soil samples. The DTPA extraction method for the determination of the  
610 bioavailable soil fraction has been executed only by Bacchetta et al. (2018) and Brunetti et al.  
611 (2018). This last was a greenhouse study conducted on contaminated agricultural soil in Apulia  
612 Region and not on mine tailings as all the other considered studies (Table 7). The remaining  
613 studies adopted more aggressive soil extraction techniques and, although presented, they  
614 cannot be considered in the discussion.

615

616



Table 7. Locations of the studies, quantity of analysed samples, adopted methodologies for soil and plant analysis, considered elements and *Helicrysum Spp.* from the literature review for the *H. italicum Spp.*. “n.a.” means not available information.

Literature	N° of CS samples	Methodology for PHEs determination in contaminated soils	N° of plant	Methodology for PHEs determination in plants	Considered elements	Studied <i>Helicrysum</i> subspp.	Location in Italy
Current study	22	DTPA extraction, Lindsay and Norvell (1978)	22	0.25 g plant +2 ml H <sub>2</sub> O <sub>2</sub> + 6 ml HNO <sub>3</sub>	Cd, Cu, Fe, Ni, Pb, Zn	<i>H. italicum</i>	Montevecchio mine district (Sardinia)
Barbafieri et al., 2011	3	SEP <sup>(1)</sup> with H <sub>2</sub> O, KNO <sub>3</sub> , EDTA	9	HNO <sub>3</sub> /HClO <sub>4</sub> in 2.5/1 ratio	Cd, Pb, Zn	<i>H. italicum</i>	Ingurtosu mine district (Sardinia)
Bacchetta et al., 2018	5	DTPA extraction, Lindsay and Norvell (1978)	5	0.5 g plant + 9 ml HNO <sub>3</sub> + 0.5 ml HF	Cd, Pb, Zn	<i>H. tyrrhenicum</i>	Campo Pisano mine district (Sardinia)
Brunetti et al., 2018	10	DTPA extraction, Lindsay and Norvell (1978)	10	HNO <sub>3</sub> :H <sub>2</sub> O <sub>2</sub> :HCl mixture (5:1:1 v/v)	Cd, Co, Cr, Cu, Ni, Pb, Zn	<i>H. italicum</i>	Agricultural area Alta Murgia (Apulia)
Bini et al., 2017	n.a.	0.2 g of soil + 5 ml of aqua regia (37% HCl+65% HNO <sub>3</sub> , 1:3) +1 ml of 48% HF + 1 ml of cold supersaturated H <sub>3</sub> BO <sub>3</sub>	n.a.	0.5 g of plants + 5 ml 65% HNO <sub>3</sub> + 3 ml 30% H <sub>2</sub> O <sub>2</sub> in open vessels on the hot plat	Cd, Co, Cr, Cu, Fe, Mn, Ni, Pb, Zn	<i>H. italicum</i>	Nocciola mine district (Tuscany)
Cao et al., 2004	n.a.	SEP <sup>(1)</sup> with H <sub>2</sub> O, KNO <sub>3</sub> , EDTA	n.a.	Aqua Regia	Pb, Zn	<i>H. italicum</i>	Montevecchio mine district (Sardinia)
Leita et al., 1989	3	10 g soil + 50 ml of 0.05 M EDTA.	3	1 g plants digested in concentrated HNO <sub>3</sub> - HCl 3:1 at 150 °C	Cd, Cu, Pb, Zn	<i>H. italicum</i>	Wide Iglesias mine district (Sardinia)

(1): Sequential Extraction Procedure

620


Concerning concentrations in plants, the literature review showed that only Zn and Pb were considered in all 7 studies (including this), Cd was measured in 5, while Cu, Ni and Fe in 4, 3 and 2 studies, respectively (Fig. 3). Only this study, Barbafieri et al. (2011) and Brunetti et al. (2018) measured PHEs in stems and leaves separately. All the other studies were limited to roots and leaves and Leita et al. (1989) considered only leaves.

Mean element concentrations identified Zn, Pb and Fe as most abundant in *H. italicum* plant with concentrations around 10<sup>3</sup> mg/kg, 5\*10<sup>2</sup> mg/kg and 10<sup>3</sup> mg/kg, respectively (Fig. 3). Instead, Cu, Cd and Ni, were poorly absorbed by the plant and their concentrations ranged around 10 mg/kg for Cu and below 10 mg/kg for Cd and Ni. The stems were the most impoverished parts for all considered elements. Based on the thresholds provided by Van der Ent et al. (2013), *H. italicum* cannot be considered an hyperaccumulator species.

Considering only studies from Sardinia region that shared the same soil element abundance ranking Zn > Pb > Cd, both *H. italicum* or *H. tyrrhenicum* (our study, Bacchetta et al., 2018; Barbafieri et al., 2011; Cao et al., 2004; Leita et al., 1989) showed similar abundances for Cd, Pb and Zn, with *H. tyrrhenicum* reporting higher concentrations. The experiment of Brunetti et al. (2018) was conducted on compost-contaminated clay-loam soils at basic pH with the

637 following contaminant soil ranking: Zn > Cu > Pb > Ni > Cd, while Bini et al. (2017), in Tuscany,  
638 had different geological settings and the contaminant rank in soil was Fe > Mn > Pb > Zn > Cu  
639 > Ni > Cd. These different experimental conditions justified the evident differences in terms of  
640 element abundances in plant parts of Cd, Zn and Pb, as well as the alignment of Fe and Cu  
641 (Fig. 3).

642



643

644 Figure 3. Mean concentration of the analysed PHEs in *H. Spp.* plant parts (r= roots; s = stems; l = leaves) from this study  
645 and from the works presented in Table 7.

646

647 As stated above, the root/soil ratio can be compared only against works that adopted the same  
648 DTPA extraction on soil samples, even if ratios from Cao et al. (2004) and Barbafieri et al.  
649 (2011) are also reported (empty bars) in Figure 4A. The uptake capacity was evident for *H.*  
650 *italicum* because all values were > 1 as reported here and also by Brunetti et al. (2018) and  
651 Bacchetta et al. (2018). The ratio values rarely exceed 10 (except for Fe, but it has a biological  
652 function) and when it happens, it is for *H. tyrrhenicum* subsp. for Pb and Zn (Bacchetta et al.,  
653 2018) or for Cd at low bioavailability in soils by Brunetti et al. (2018).

654 *Helicrysum italicum* can be defined as a tolerant species for Cd, Cu, Fe, Ni, Pb and Zn, but it  
655 is not suitable for phytoextraction since it shows little capacity to accumulate elements in  
656 leaves. Indeed, Cd was equally distributed in the plant and leaf/root ratio was 1 both in the  
657 present study and Barbafieri et al. (2011) and Bacchetta et al. (2018), while it was 0.6 in  
658 Tuscany mine of Bini et al. (2017) (Fig. 4B). Although the studies conducted in the Sardinia  
659 region showed an accumulation of Zn and Pb in leaves with mean leaf/root ratios around 2,  
660 and even higher in Barbafieri et al. (2011) (Fig. 4B), the high variability of the leaf/root ratios in  
661 this study does not allow a clear and univocal indication on translocation capacity of *H. italicum*  
662 (Table 6).

663 In this study, Fe resulted the most absorbed element (Fig. 4A) and contemporarily also the  
664 most translocated from roots to leaves (2.36 leaf/root median ratio in Table 6) because of its  
665 biological function in photosynthetic process. Despite this, the Fe leaf/root ratio by Bini et al.  
666 (2017) was widely < 1 indicating a root accumulation (Fig. 4B). Bioavailable Fe in CS and US  
667 were aligned (Table 1) and the higher absorption and translocation in CS was probably related  
668 to the lower pH and lime content in respect to US, as reported by Buscaroli et al. (2017). In the  
669 mine site investigated by Bini et al. (2017), total Fe was the most abundant element and plants  
670 preferred to store it in roots. Similar behaviour has been reported in rice species that are able  
671 to oxidise Fe at the root surface, leading to the formation of iron plaques (Green and  
672 Etherington, 1977) or accumulating Fe as ferric hydroxides (goethite and lepidocrocite) in roots  
673 (Bacha and Hossner, 1977). In both cases Fe precipitation in roots could later influence the  
674 uptake of other elements (Armstrong and Armstrong, 1988).

675  
676




Figure 4. Mean root/bioavailable soil element concentration ratio (A) and mean leaf/root element concentration ratio (B) in *H. Spp.*, calculated from this study and from the other papers presented in Table 7.

680

681 Plants suitable for phytoextraction should possess multiple abilities: first of all absorb (root/soil)  
682 and translocate (leaf/root) to aerial parts heavy metals then rusticity, fast growth, high biomass  
683 yield and easy harvesting (Jabeen et al., 2009).

684 Based on the literature review, the content of all investigated elements, in plant tissues makes  
685 *H. italicum* a tolerant species (especially in respect of Pb and Zn), but it does not reach the  
686 concentrations to be defined an hyperaccumulator plant (Baker et al., 2000): 100 mg/kg of Cd  
687 (of the leaf dry weight), 1000 mg/kg for Ni, Cu and Pb and 10000 mg/kg for Zn. The root/soil  
688 bioavailable ratio > 1 for all elements suggests the use of *H. italicum* for phytostabilization in  
689 mine areas as a pioneering strategy. Although the median leaf/root ratio was > 1 for Zn and Pb  
690 in plants grown in Sardinia mine districts, contrasting mean ratios were achieved for *H. italicum*  
691 in different contaminated sites not allowing a clear evidence of its phytoextraction ability.

692 Moreover, Boi et al. (2020) argued that few kilograms per hectare (6 - 11 kg/ha) can be  
693 recovered by *H. tyrrhenicum* (the most performant subspp. as shown in Fig. 3 and 4) and given  
694 the actual price of Zn, it does not allow economic sustainability.

695 Since *H. italicum* i) is a spontaneous and perennial species, tolerant to PHEs; ii) guarantees  
696 the canopy cover all throughout the year, preventing wind dispersion and water erosion; iii)

697 influences the soil retention capacity and can itself rehabilitate the vegetation cover,  
698 reactivating pedological processes; iv) can be propagated sowing directly seeds on  
699 contaminated soils allowing cheaper propagation; v) permits the stabilization of mine tailing  
700 also from land management point of view, it can be indicated for phytostabilization in  
701 abandoned mine districts, reducing the impact of PHEs on the mine sites and surrounding  
702 environments (Barbafieri et al., 2011; Bacchetta et al., 2018; Boi et al., 2019; 2020).

703

704

## 705 **4. Conclusions**

706 This study aimed at evaluating phytoremediation properties of *H. italicum* for PHEs (Cd, Cu,  
707 Fe, Ni, Pb and Zn) by the determination of elements concentration on the roots, stems and  
708 leaves, and by the related root/soil and leaf/root ratios on plants collected from mine tailing  
709 deposits in contaminated sites (CS, Montevecchio mine, Sardinia) and in uncontaminated sites  
710 (US). Moreover, a literature review on the phytoremediation properties of *Helicrysum* Spp. has  
711 been executed in order to elucidate its phytoremediation potentiality.

712 The Cd, Pb and Zn resulted to be the most bioavailable PHEs in soils collected from  
713 Montevecchio mine district compared to US, also exceeding the thresholds of the Italian  
714 environmental legislation.

715 Element concentrations in plants from CS were higher than those from US. Leaf element  
716 concentrations were higher compared to stems and roots, although statistically significant only  
717 for Cu, Fe and Zn in CS. Interaction mechanisms (synergistic effects) between Cd and Zn, Fe  
718 and Ni, and Cu and Zn in *H. italicum* plants grown on CS were detected.

719 The medians for root/soil ratio were > 1 for Cu, Fe, Ni and Pb in both CS and US, meaning that  
720 *H. italicum* was able to accumulate bioavailable elements in roots. Cadmium and Zn in CS had  
721 root/soil ratio close to one (1.39 and 1.21, respectively), suggesting their limited uptake when  
722 soil bioavailable concentrations are elevated. Based on the root/soil ratios *H. italicum* is  
723 considered a metal tolerant species.

724 The medians leaf/root ratio in CS were 2.36, 1.54, 1.37 and 1.34 for Fe, Zn, Pb and Cu,  
725 respectively, while ratios were < 1 for Cd (0.75) and Ni (1). Similar ratios were also calculated  
726 for US proving that *H. italicum* has a weak phytoextraction capacity.

727 The literature review on the phytoremediation potentiality of *H. italicum* confirmed the analytical  
728 findings of this study. Indeed, Zn, Pb and Fe were the most abundant elements in *H. italicum*  
729 plants grown on contaminated soils with concentrations around  $10^3$  mg/kg,  $5 \cdot 10^2$  mg/kg and  
730  $10^3$  mg/kg, respectively. The Cu, Cd and Ni, were poorly absorbed by plants and their  
731 concentrations ranged around 10 mg/kg of Cu and below 10 mg/kg of Cd and Ni. In light of

732 this, *H. Italicum* cannot be considered a hyperaccumulator species. Overall, the root/soil  
733 bioavailable ratio > 1 for all elements suggested the use of *H. italicum* for phytostabilization in  
734 mine areas as a pioneering strategy of remediation.

735 Given that, *H. italicum* is a spontaneous and perennial species, which guarantees the canopy  
736 cover all throughout the year, rehabilitates the vegetation cover and it can be propagated by  
737 directly sowing seeds on contaminated soil, it can be recommended for phytostabilization of  
738 abandoned mine districts and for stabilization of mine tailing from land management point of  
739 view.

740

741

## 742 **Acknowledgements**

743 Thanks to the Interdisciplinary Lab-based Course within the Master Program in Environmental  
744 Assessment and Management at University of Bologna – Ravenna Campus for the support  
745 during the sampling campaign. A particular thanks also to Sardinian local authorities allowing  
746 sampling in mining areas and to anonymous reviewers that helped in improving the general  
747 quality of the manuscript.

748

749

## 750 **Funding sources**

751 This research did not receive any specific grant from funding agencies in public, commercial,  
752 or not-for-profit sectors.

753

754

## 755 References

- 756 Abreu, M.M., Tavares, M.T., Batista, M.J., 2008. Potential use of *Erica andevalensis* and *Erica australis* in  
757 phytoremediation of sulphide mine environments: São Domingos, Portugal. Journal of Geochemical  
758 Exploration 96, 210-222. Doi: 10.1016/j.gexplo.2007.04.007.
- 759 An, Y. J., Kim, Y. M., Kwon, T. I., & Jeong, S. W., 2004. Combined effect of copper, cadmium, and lead upon  
760 *Cucumis sativus* growth and bioaccumulation. Science of the Total Environment, 326(1-3), 85-93. Doi:  
761 10.1016/j.scitotenv.2004.01.002.
- 762 Arbogast, B.F., Knepper, D.H., Langer, W.H., 2000. The human factor in mining reclamation (Vol. 1191). US  
763 Department of the Interior, US Geological Survey.
- 764 Armstrong, J., Armstrong, W., 1988. *Phragmites australis*—A preliminary study of soil-oxidizing sites and internal  
765 gas transport pathways. New phytologist 108(4), 373-382. Doi: 10.1111/j.1469-8137.1988.tb04177.x.
- 766 Ashraf, U., Mahmood, M.H.U.R., Hussain, S., Abbas, F., Anjum, S.A., Tang, X., 2020. Lead (Pb) distribution and  
767 accumulation in different plant parts and its associations with grain Pb contents in fragrant rice. Chemosphere  
768 248, 126003. Doi: 10.1016/j.chemosphere.2020.126003.
- 769 Bacchetta, G., Boi, M.E., Cappai, G., De Giudici, G., Piredda, M., Porceddu, M., 2018. Element tolerance  
770 capability of *Helichrysum microphyllum* Cambess. subsp. *tyrrhenicum* Bacch., Brullo & Giusso: A candidate  
771 for phytostabilization in abandoned mine sites. Bulletin of environmental contamination and toxicology  
772 101(6), 758-765. Doi: 10.1007/s00128-018-2463-9.
- 773 Bacha, R.E., Hossner, L.R., 1977. Characteristics of coatings formed on rice roots as affected by iron and  
774 manganese additions. Soil Science Society of America Journal 41(5), 931-935. Doi:  
775 10.2136/sssaj1977.03615995004100050025x.
- 776 Baker, A.J.M., McGrath, S.P., Reeves, R.D., & Smith, J. A. C. 2000. Metal hyperaccumulator plants: A review of  
777 the ecology and physiology of a biological resource for phytoremediation of metal-polluted soils. p. 85– 107.  
778 In N. Terry and G. Bañuelos (ed.) Phytoremediation of contaminated soil and water. Lewis Publishers,  
779 Boca Raton, FL.
- 780 Banda, M. F., Mokgalaka, N. S., Combrinck, S., & Regnier, T. 2021. Five-weeks pot trial evaluation of  
781 phytoremediation potential of *Helichrysum splendidum* Less. for copper- and lead-contaminated soils. Int. J.  
782 Environ. Sci. Technol. Doi: 10.1007/s13762-021-03243-z
- 783 Barbafieri, M., Dadea, C., Tassi, E., Bretzel, F., Fanfani, L., 2011. Uptake of heavy elements by native species  
784 growing in a mining area in Sardinia, Italy: discovering native flora for Phytoremediation. International Journal  
785 of Phytoremediation 13, 985-997. Doi: 10.1080/15226514.2010.549858.
- 786 Bianchini, A., Santoni, F., Paolini, J., Bernardini, A.F., Mouillot, D., Costa, J., 2009. Partitioning the relative  
787 contributions of inorganic plant composition and soil characteristics to the quality of *Helichrysum italicum*  
788 subsp. *italicum* (Roth) G. Don fil. essential oil. Chemistry & biodiversity 6(7), 1014-1033. Doi:  
789 10.1002/cbdv.200800328.
- 790 Bini, C., Wahsha, M., 2014. Potentially harmful elements and human health. In PHEs, Environment and Human  
791 Health (pp. 401-463). Springer, Dordrecht.

792 Bini, C., Maleci, L., Tani, C., Wahsha, M., 2017. Preliminary Observations On The Element Tolerance And  
793 Resilience Capacity Of *Helichrysum italicum* (Roth) G. Don Growing On Mine Soils. *EQA - International*  
794 *Journal of Environmental Quality* 21(1), 41-50. Doi: 10.6092/issn.2281-4485/6599.

795 Boechat, C. L., Pistóia, V. C., Gianelo, C., & de Oliveira Camargo, F. A. 2016. Accumulation and translocation of  
796 heavy metal by spontaneous plants growing on multi-metal-contaminated site in the Southeast of Rio Grande  
797 do Sul state, Brazil. *Environmental Science and Pollution Research*, 23(3), 2371-2380. Doi: 10.1007/s11356-  
798 015-5342-5.

799 Boi, M.E., Porceddu, M., Cappai, G., De Giudici, G., Bacchetta, G., 2019. Effects of zinc and lead on seed  
800 germination of *Helichrysum microphyllum subsp. tyrrhenicum*, an element-tolerant plant. *International*  
801 *Journal of Environmental Science and Technology* 1-12. Doi: 10.1007/s13762-019-02589-9.

802 Boi, M.E., Medas, D., Aquilanti, G., Bacchetta, G., Birarda, G., Cappai, G., Carlomagno, I., Casu, M.A.,  
803 Gianoncelli, A., Meneghini, C., Piredda, M., Podda, F., Porceddu, M., Rimondi, V., Vaccari, L., De Giudici,  
804 G., 2020. Mineralogy and Zn Chemical Speciation in a Soil-Plant System from an Element-Extreme  
805 Environment: A Study on *Helichrysum microphyllum subsp. tyrrhenicum* (Campo Pisano Mine, SW Sardinia,  
806 Italy). *Minerals*, 10(3), 259. Doi: 10.3390/min10030259.

807 Bolan, N.S., Park, J.H., Robinson, B., Naidu, R., Huh, K.Y., 2011. Phytostabilization: a green approach to  
808 contaminant containment. *Advances in agronomy* 112, 145-204. Doi: 10.1016/B978-0-12-385538-1.00004-  
809 4.

810 Boni, M., Costabile, S., De Vivo, B., Gasparrini, M., 1999. Potential environmental hazard in the mining district of  
811 southern Iglesiente (SW Sardinia, Italy). *Journal of geochemical exploration* 67(1-3), 417-430. Doi.  
812 10.1016/S0375-6742(99)00078-3.

813 Brady, N.C., Weil, R.R., 2010. *Element of the nature and properties of soil* 3<sup>rd</sup> Edition. Prentice Hall, New Jersey,  
814 US. <https://fratstock.eu/sample/Solutions-Manual-Nature-Properties-of-Soils-14th-Edition-Brady.pdf>.

815 Brunetti, G., Ruta, C., Traversa, A., D'Ambruso, G., Tarraf, W., De Mastro, F., De Mastro, G., Coccozza, C., 2018.  
816 Remediation of a heavy elements contaminated soil using mycorrhized and non-mycorrhized *Helichrysum*  
817 *italicum* (Roth) Don. *Land degradation & development* 29(1), 91-104. Doi: 10.1002/ldr.2842.

818 Bullitta S., Piluzza G., Usai M., 2010 Determinazione di metalli pesanti accumulati in *Helichrysum italicum* (Roth)  
819 *G. Don subsp. microphyllum* (Willd.) Nyman vegetante su aree minerarie dismesse. *Fitomed* 2010, 116,  
820 Cagliari.

821 Buscaroli, A., 2017. An overview of indexes to evaluate terrestrial plants for phytoremediation purposes.  
822 *Ecological Indicators* 82, 367-380. Doi: 10.1016/j.ecolind.2017.07.003.

823 Buscaroli, A., Zannoni, D., Menichetti, M., Dinelli, E., 2017. Assessment of element accumulation capacity of  
824 *Dittrichia viscosa* (L.) Greuter in two different Italian mine areas for contaminated soils remediation. *Journal*  
825 *of Geochemical Exploration* 182, 123-131. Doi: 10.1016/j.gexplo.2016.10.001.

826 Cao, A., Cappai, G., Carucci, A., Muntoni, A., 2004. Selection of Plants for Zinc and Lead Phytoremediation.  
827 *Journal of Environmental Science and Health* 39(4), 1011-1024. Doi: 10.1081/ESE-120028410.

828 Caboi, R., Cidu, R., Cristini, A., Fanfani, L., Massoli-Novelli, R., Zuddas, P., 1993. The abandoned Pb-Zn mine of  
829 Ingurtosu, Sardinia (Italy). *Engineering Geology* 34, 211-218. Doi: 10.1016/0013-7952(93)90090-Y.



830 Camizuli, E., Monna, F., Scheifler, R., Amiotte-Suchet, P., Losno, R., Beis, P., Bohard, B., Chateau, C., Alibert,  
831 P., 2018. Trace elements from historical mining sites and past metallurgical activity remain bioavailable to  
832 wildlife today. *Scientific reports* 8(1), 1-11. Doi: 10.1038/s41598-018-20983-0

833 Carbone, C., Dinelli, E., Marescotti, P., Gasparotto, G., Lucchetti, G., 2013. The role of AMD secondary minerals  
834 in controlling environmental pollution: Indications from bulk leaching test. *Journal of Geochemical Exploration*  
835 132 (2), 188-200. Doi: 10.1016/j.gexplo.2013.07.001.

836 Choudhary, M., Bailey, L.D., Grant, C.A., Leisle, D., 1995. Effect of Zn on the concentration of Cd and Zn in plant  
837 tissue of two durum wheat lines. *Canadian Journal of Plant Science* 75(2), 445-448. Doi: 10.4141/cjps95-  
838 074.

839 Concas, A., Ardaù, C., Cristini, A., Zuddas, P., Cao, G., 2006. Mobility of heavy elements from tailings to stream  
840 waters in a mining activity contaminated site. *Chemosphere* 63 (2), 244-253.  
841 Doi:10.1016/j.chemosphere.2005.08.024.

842 Conesa, H.M., Faz, Á., Arnaldos, R., 2006. Heavy element accumulation and tolerance in plants from mine tailings  
843 of the semiarid Cartagena–La Unión mining district (SE Spain). *Science of the total environment* 366(1), 1-  
844 11. Doi: 10.1016/j.scitotenv.2005.12.008.

845 Conesa, H.M., Maria-Cervantes, A., Alvarez-Rogel, J., Gonzalez-Alcaraz, M.N., 2011. Influence of soil properties  
846 on trace element availability and plant accumulation in a Mediterranean salt marsh polluted by mining wastes:  
847 Implications for phytomanagement. *Science of the Total Environment* 409, 4470-4479. Doi:  
848 10.1016/j.scitotenv.2011.07.049.

849 Cuccuru, S., Naitza, S., Secchi, F., Puccini, A., Casini, L., Pavanetto, P., Linnemann, U., Hofmann, M., Oggiano,  
850 G., 2016. Structural and metallogenic map of late Variscan Arbus Pluton (SW Sardinia, Italy). *Journal of*  
851 *Maps* 12 (5), 860-865. Doi: 10.1080/17445647.2015.1091750.

852 Day, P.R., 1965. Particle fractionation and particle-size analysis, in Black, C.A. (Eds.), *Methods of soil analysis*,  
853 Part I. American Society of Agronomy Inc., Madison, pp. 545-567.

854 De Agostini, A., Caltagirone, C., Caredda, A., Cicatelli, A., Cogoni, A., Farci, D., Guarino, F., Garau, A., Labra,  
855 M., Lussu, M., Piano, D., Sanna, C., Tommasi, N., Vacca, A., Cortis, P., 2020. Heavy metal tolerance of  
856 orchid populations growing on abandoned mine tailings: A case study in Sardinia Island (Italy). *Ecotoxicology*  
857 *and Environmental Safety* 189, 110018. Doi: 10.1016/j.ecoenv.2019.110018.

858 Dinelli, E., Lucchini, F., Fabbri, M., Cortecchi, G., 2001. Element distribution and environmental problems related  
859 to sulfide oxidation in the Libiola copper mine area (Ligurian Appennines, Italy). *Journal of Geochemical*  
860 *Exploration* 74, 141-152. Doi: 10.1016/S0375-6742(01)00180-7.

861 Dinu, C., Vasile, G.G., Buleandra, M., Popa, D. E., Gheorghe, S., Ungureanu, E.M., 2020. Translocation and  
862 accumulation of heavy metals in *Ocimum basilicum* L. plants grown in a mining-contaminated soil. *Journal*  
863 *of Soils and Sediments* 20(4), 2141-2154. Doi: 10.1007/s11368-019-02550-w.

864 Dore, E., Fancello, D., Rigonat, N., Medas, D., Cidu, R., Da Pelo, S., Frau, F., Lattanzi, P., Marras, P.A.,  
865 Meneghini, C., Podda, F., Rimondi, V., Runkel, R.L., Kimball, B., Wanty, R.B., De Giudici, G., 2020. Natural  
866 attenuation can lead to environmental resilience in mine environment. *Applied Geochemistry* 104597. Doi:  
867 10.1016/j.apgeochem.2020.104597.

Esringü, A., Turan, M., Güneş, A., Karaman, M.R., 2014. Roles of *Bacillus megaterium* in remediation of boron, lead, and cadmium from contaminated soil. *Communications in soil science and plant analysis* 45(13), 1741-1759. Doi: 10.1080/00103624.2013.875194.

Eurostat, 2017. Land cover statistics. Accessed September 2020.

Fernández, S., Poschenrieder, C., Marcenò, C., Gallego, J. R., Jiménez-Gámez, D., Bueno, A., Afif, E., 2017. Phytoremediation capability of native plant species living on Pb-Zn and Hg-As mining wastes in the Cantabrian range, north of Spain. *Journal of Geochemical Exploration* 174, 10-20. Doi: 10.1016/j.gexplo.2016.05.015.

Fitamo, D., Leta, S., 2010. Assessment of plants growing on gold mine wastes for their potential to remove heavy metals from contaminated soils. *International journal of environmental studies* 67(5), 705-724. Doi: 10.1080/00207233.2010.513587.

Franzini, M., Leoni, L., Saitta, M., 1972. A simple method to evaluate the matrix effects in X-Ray fluorescence analysis. *X-Ray Spectrometry* 1, 151-154.

Galbany-Casals, M., Blanco-Moreno, J.M., Garcia-Jacas, N., Breitwieser, I., Smissen, R.D., 2011. Genetic variation in Mediterranean *Helichrysum italicum* (Asteraceae; Gnaphalieae): do disjunct populations of *subsp. microphyllum* have a common origin?. *Plant Biology* 13(4), 678-687. Doi: 10.1111/j.1438-8677.2010.00411.x.

Galeas, M. L., Klamper, E. M., Bennett, L. E., Freeman, J. L., Kondratieff, B. C., Quinn, C. F., & Pilon-Smits, E. A., 2008. Selenium hyperaccumulation reduces plant arthropod loads in the field. *New Phytologist*, 177(3), 715-724. Doi: 10.1111/j.1469-8137.2007.02285.x.

García, G., Conesa, H.M., Faz, A., 2002. Phytoremediation of zinc polluted soils by Mediterranean plant species: Usefulness of bioaccumulation and tolerance capabilities. In M. Mench and B. Mocquot (Eds.), *COST Action 837. 4th WG2 Workshop, Bordeaux'2002 – Risk assessment and sustainable land management using plants in trace metal-contaminated soils*, Villenave d'Ornon, France (p. 140).

García, G., Zanuzzi, A.L., Faz, Á., 2005. Evaluation of heavy metal availability prior to an in situ soil phytoremediation program. *Biodegradation* 16(2), 187-194.

Gonçalves, A.C., Schwantes, D., Braga de Sousa, R.F., Benetoli da Silva, T.R., Guimarães, V.F., Campagnolo, M.A., Soares de Vasconcelos, E., Zimmermann, J., 2020. Phytoremediation capacity, growth and physiological responses of *Crambe abyssinica* Hochst on soil contaminated with Cd and Pb. *Journal of Environmental Management* 262, 110342. Doi: 10.1016/j.jenvman.2020.110342.

Green, M.S., Etherington, J.R., 1977. Oxidation of ferrous iron by rice (*Oryza sativa* L.) roots: a mechanism for waterlogging tolerance?. *Journal of Experimental Botany* 28(3), 678-690. Doi: 10.1093/jxb/28.3.678.

Hart, J.J., Welch, R.M., Norvell, W.A., Kochian, L.V., 2002. Transport interactions between cadmium and zinc in roots of bread and durum wheat seedlings. *Physiologia Plantarum* 116(1), 73-78. Doi: 10.1034/j.1399-3054.2002.1160109.x.

Heiri, O., Lotter, A.F., Lemcke, G., 2001. Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results. *Journal of Paleolimnology* 25, 101-110. Doi: 10.1023/A:1008119611481.

906 Hesami, R., Salimi, A., Ghaderian, S.M., 2018. Lead, zinc, and cadmium uptake, accumulation, and  
 907 phytoremediation by plants growing around Tang-e Douzan lead–zinc mine, Iran. *Environmental Science*  
 908 *and Pollution Research* 25(9), 8701-8714. Doi: 10.1007/s11356-017-1156-y.

909 ISO, 1995a. ISO 10693 - Soil quality - Determination of carbonate content - Volumetric method. First edition.

910 ISO, 1995b. ISO 10694 – Soil quality – Determination of organic and total carbon after dry combustion (elementary  
 911 analysis). First edition.

912 ISO, 1998. ISO 11047 - Soil quality — Determination of cadmium, chromium, cobalt, copper, lead, manganese,  
 913 nickel and zinc in aqua regia extracts of soil — Flame and electrothermal atomic absorption spectrometric  
 914 methods. First edition.

915 ISO, 2001. ISO 14870 - Soil quality - Extraction of trace metals by buffered DTPA solution. First edition.

916 ISPRA, <http://sgi2.isprambiente.it/arcgis/rest/services/servizi/cartageologica500k/MapServer>. Accessed  
 917 15/04/2021.

918 Italian Government, 2001. DM 468/2001. Supplemento ordinario alla Gazzetta Ufficiale n. 13 del 16 gennaio 2002  
 919 - Serie generale. Ministero Dell'ambiente E Della Tutela Del Territorio. Decreto 18 settembre 2001, n. 468.  
 920 Regolamento recante: Programma nazionale di bonifica e ripristino ambientale.  
 921 <https://www.gazzettaufficiale.it/eli/id/2002/01/16/001G0506/sg>. Accessed 26/03/2021.

922 Italian Government, 2006. Decreto Legislativo 152/2006. Norme in Materia Ambientale - Testo Unico Ambientale.  
 923 Gazzetta Ufficiale Serie Generale n.88 del 14-04-2006 – Suppl. Ordinario n. 96. Parlamento della Repubblica  
 924 Italiana. <https://www.gazzettaufficiale.it/dettaglio/codici/materiaAmbientale>. Accessed 02/05/2021.

925 Jabeen, R., Ahmad, A., & Iqbal, M., 2009. Phytoremediation of heavy metals: physiological and molecular  
 926 mechanisms. *The Botanical Review*, 75(4), 339-364. Doi: 10.1007/s12229-009-9036-x.

927 Kabata-Pendias, A., 2010. Trace elements in soils and plants. 4<sup>th</sup> edition, CRC Press, Taylor and Francis Group.

928 Khalid, B. Y., Tinsley, J., 1980. Some effects of nickel toxicity on rye grass. *Plant and Soil*, 55(1), 139-144. Doi:  
 929 10.1007/BF02149717.

930 Khaokaew, S., Landrot, G., 2014. A field-scale study of cadmium phytoremediation in a contaminated agricultural  
 931 soil at Mae Sot District, Tak Province, Thailand: (1) Determination of Cd-hyperaccumulating plants.  
 932 *Chemosphere*. Doi: 10.1016/j.chemosphere.2014.09.108.

933 Koosaletse-Mswela, P., Przybyłowicz, W.J., Cloete, K.J., Barnabas, A.D., Nelson Torto, N., Jolanta Mesjasz-  
 934 Przybyłowicz, J., 2015. Quantitative mapping of elemental distribution in leaves of the metallophytes  
 935 *Helichrysum candolleanum*, *Blepharis aspera*, and *Blepharis diversispina* from Selkirk Cu–Ni mine,  
 936 Botswana. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with*  
 937 *Materials and Atoms* 363, 188-193. Doi.org/10.1016/j.nimb.2015.09.005.

938 Kumpiene, J., Giagnoni, L., Marschner, B., Denys, S., Mench, M., Adriaensen, K., Vangronsveld, J.,  
 939 Puschenreiter, M., Renella, G., 2017. Assessment of methods for determining bioavailability of trace  
 940 elements in soils: a review. *Pedosphere* 27(3), 389-406. Doi: 10.1016/S1002-0160(17)60337-0.

941 Kutrowska, A., Małecka, A., Piechalak, A., Masiakowski, W., Hanć, A., Barańkiewicz, D., Andrzejewskac, B.,  
 942 Zbierska, J., Tomaszewska, B., 2017. Effects of binary metal combinations on zinc, copper, cadmium and

943 lead uptake and distribution in *Brassica juncea*. Journal of Trace Elements in Medicine and Biology, 44, 32-  
944 39. Doi: 10.1016/j.jtemb.2017.05.007.

945 Lancianese, V., Dinelli, E., 2015. Different spatial methods in regional geochemical mapping at high density  
946 sampling: An application on stream sediment of Romagna Apennines, Northern Italy. Journal of Geochemical  
947 Exploration 154, 143-155. Doi: 10.1016/j.gexplo.2014.12.014.

948 Leita, L., De Nobili, M., Pardini, G., Ferrari, F., Sequi, P., 1989. Anomalous contents of heavy metals in soils and  
949 vegetation of a mine area in SW Sardinia, Italy. Water, Air, and Soil Pollution 48(3-4), 423-433. Doi:  
950 10.1007/BF00283340.

951 Leoni, L., Saitta, M., 1976. X-ray fluorescence analysis of 29 trace elements in rock and mineral standard.  
952 Rendiconti Società Italiana di Mineralogia e Petrologia 32, 497-510.

953 Li, M.S., Luo, Y.P., Su, Z.Y., 2007. Heavy element concentrations in soils and plant accumulation in a restored  
954 manganese mineland in Guangxi, South China. Environmental Pollution 147, 168-175. Doi:  
955 10.1016/j.envpol.2006.08.006.

956 Lin, G C., Ho, S.P., 2003. China's land resources and land-use change: insights from the 1996 land survey. Land  
957 use policy 20(2), 87-107. Doi: 10.1016/S0264-8377(03)00007-3

958 Lindsay, W.L., Norvell, W.A., 1978. Development of a DTPA Soil Test for Zinc, Iron, Manganese and Copper. Soil  
959 Science Society American Journal 42, 421-428. Doi: 10.2136/sssaj1978.03615995004200030009x.

960 Lombini, A., Dinelli, E., Ferrari, C., Simoni, A., 1998. Plant-soil relationships in the serpentinite screes of Mt.  
961 Prinzera (Northern Apennines, Italy). Journal of Geochemical Exploration 64, 19-33. Doi: 10.1016/S0375-  
962 6742(98)00017-X.

963 Mahar, A., Wang, P., Ali, A., Awasthi, M.K., Lahori, A.H., Wang, Q., Li, R., Zhang, Z., 2016. Challenges and  
964 opportunities in the phytoremediation of heavy metals contaminated soils: a review. Ecotoxicology and  
965 environmental safety 126, 111-121. Doi: 10.1016/j.ecoenv.2015.12.023.

966 Marescotti, P., Carbone, C., De Capitani, L., Grieco, G., Lucchetti, G., Servida, D., 2008. Mineralogical and  
967 geochemical characterization of open pit tailing and waste rock dumps from the Libiola Fe-Cu sulphide mine  
968 (Eastern Liguria, Italy). Environmental Geology 53, 1613-1626. Doi: 10.1007/s00254-007-0769-8.

969 Martin, S.R., Llugany, M., Barceló, J., Poschenrieder, C., 2012. Cadmium exclusion a key factor in differential Cd-  
970 resistance in *Thlaspi arvense* ecotypes. Biologia Plantarum 56(4), 729-734. Doi: 10.1007/s10535-012-0056-  
971 8.

972 Martínez-Sánchez, M.J., García-Lorenzo, M.L., Pérez-Sirvent, C., Bech, J., 2012. Trace element accumulation in  
973 plants from an aridic area affected by mining activities. Journal of Geochemical Exploration 123, 8-12. Doi:  
974 10.1016/j.gexplo.2012.01.007.

975 Melito, S., Petretto, G. L., Podani, J., Foddai, M., Maldini, M., Chessa, M., Pintore, G., 2016. Altitude and climate  
976 influence *Helichrysum italicum subsp. microphyllum* essential oils composition. Industrial Crops and Products  
977 80, 242-250. Doi: 10.1016/j.indcrop.2015.11.014.

978 Mendez, M.O., Maier, R.M., 2008. Phytoremediation of mine tailings in temperate and arid environments. Review  
979 Environment Science Biotechnology 7, 47-59. Doi: 10.1007/s11157-007-9125-4.

- 980 Meuser, H., 2013. Soil Remediation and Rehabilitation: Treatment of Contaminated and Disturbed Land.  
981 Environmental Pollution, vol. 23, Springer Science + Business Media, Dordrecht, The Netherlands. Doi:  
982 10.1007/978-94-007-5751-6\_1
- 983 Mills, M.P., 2020. Mines, minerals, and “green” energy: a reality check. Manhattan Institute Report. Energy &  
984 Environment Regulations. [https://www.manhattan-institute.org/mines-minerals-and-green-energy-reality-](https://www.manhattan-institute.org/mines-minerals-and-green-energy-reality-check)  
985 check Accessed 10 September 2020.
- 986 Moroni, M., Naitza, S., Ruggieri, G., Aquino, A., Costagliola, P., De Giudici, G., Caruso, S., Ferrari, E., Fiorentini,  
987 M.L., Lattanzi, P., 2019. The Pb-Zn-Ag vein system at Montevecchio-Ingurtosu, southwestern Sardinia, Italy:  
988 A summary of previous knowledge and new mineralogical, fluid inclusion, and isotopic data. Ore Geology  
989 Reviews 115, 103194. Doi: 10.1016/j.oregeorev.2019.103194.
- 990 Nadimi-Goki, M., Wahsha, M., Bini, C., Yorichio, K., Vianello, G., Vittori Antisari, L., 2014. Assessment of total soil  
991 and plant elements in rice-based production systems in NE Italy. Journal of Geochemical Exploration 147,  
992 200-214. Doi: 10.1016/j.gexplo.2014.07.008.
- 993 Nkoane, B.B.M, Sawula, G.M., Wibetoe, G., Lund, W., 2005. Identification of Cu and Ni indicator plants from  
994 mineralised locations in Botswana. Journal of Geochemical Exploration 86, 130-142. Doi:  
995 10.1016/j.gexplo.2005.03.003.
- 996 Nkoane, B.B.M., Wibetoe, G., Lund, W., Abegaz, B.M., Torto, N., 2007. Examination of *Blepharis aspera* as a  
997 possible Cu–Ni indicator plant. South African Journal of Sciences 103, 363-364
- 998 Ninčević, T., Grdiša, M., Šatović, Z., Jug-Dujaković, M., 2019. *Helichrysum italicum* (Roth) G. Don: Taxonomy,  
999 biological activity, biochemical and genetic diversity. Industrial crops and products 138, 111487. Doi:  
1000 10.1016/j.indcrop.2019.111487.
- 1001 Oliver, D.P., Hannam, R., Tiller, K.G., Wilhelm, N.S., Merry, R.H., Cozens, G.D., 1994. The effects of zinc  
1002 fertilization on cadmium concentration in wheat grain (Vol. 23, No. 4, pp. 705-711). American Society of  
1003 Agronomy, Crop Science Society of America and Soil Science Society of America. Doi:  
1004 10.2134/jeq1994.00472425002300040013x.
- 1005 Pandey, V.C., Pandey, D.N., Singh, N., 2015. Sustainable phytoremediation based on naturally colonizing and  
1006 economically valuable plants. Journal of cleaner Production 86, 37-39. Doi: 10.1016/j.jclepro.2014.08.030.
- 1007 Pandey, V.C., Bajpai, O., Singh, N., 2016. Energy crops in sustainable phytoremediation. Renewable and  
1008 Sustainable Energy Reviews 54, 58-73. Doi: 10.1016/j.rser.2015.09.078.
- 1009 Pandey, V.C., Bajpai, O., 2019. Phytoremediation: from theory toward practice. In Phytomanagement of Polluted  
1010 Sites (pp. 1-49). Elsevier. Doi: 10.1016/B978-0-12-813912-7.00001-6.
- 1011 Perez, J., 2012. The soil remediation industry in Europe: the recent past and future perspectives. Ernst & Young  
1012 Research Report. [https://ec.europa.eu/environment/archives/soil/pdf/may2012/08%20-](https://ec.europa.eu/environment/archives/soil/pdf/may2012/08%20-%20Julien%20Perez%20-%20final.pdf)  
1013 %20Julien%20Perez%20-%20final.pdf. Accessed 26/03/2021.
- 1014 Poschenrieder, C., Llugany, M., Lombini, A., Dinelli, E., Bech, J., Barceló, J., 2012. *Smilax aspera* L. an evergreen  
1015 Mediterranean climber for phytoremediation. Journal of Geochemical Exploration 123, 41–44. Doi:  
1016 10.1016/j.gexplo.2012.07.012.

- 1017 Pourrut, B., Shahid, M., Dumat, C., Winterton, P., & Pinelli, E., 2011. Lead uptake, toxicity, and detoxification in  
1018 plants. *Reviews of environmental contamination and toxicology* volume 213, 113-136. Doi: 10.1007/978-1-  
1019 4419-9860-6\_4.
- 1020 Rimondi, V., Costagliola, P., Lattanzi, P., Catelani, T., Fornasaro, S., Medas, D., Morelli, G. & Paolieri, M., 2021.  
1021 Bioaccessible arsenic in soil of thermal areas of Viterbo, Central Italy: implications for human health risk.  
1022 *Environmental Geochemistry and Health*, 1-21. Doi: 10.1007/s10653-021-00914-1.
- 1023 Reimann, C., Birke, M., Demetriades, A., Filzmoser, P., O'Connor, P. (Editors), 2014. Chemistry of Europe's  
1024 agricultural soils-Part B: General background information and further analysis of the GEMAS data set.  
1025 *Geologisches Jahrbuch (Reihe B 103)*, Schweizerbarth, Hannover, 352 pp.
- 1026 Remon, E., Bouchardon, J.L., Faure, O., 2007. Multi-tolerance to heavy metals in *Plantago Arenaria* Waldst. &  
1027 Kit.: Adaptative versus constitutive characters. *Chemosphere* 69, 41-47. Doi:  
1028 10.1016/j.chemosphere.2007.04.067.
- 1029 RER, 2016. Carta del fondo naturale dei metalli pesanti della pianura emiliano-romagnola.  
1030 [http://mappegis.regione.emilia-romagna.it/gstatico/documenti/dati\\_pedol/CARTA\\_PEDOGEOCHIMICA.pdf](http://mappegis.regione.emilia-romagna.it/gstatico/documenti/dati_pedol/CARTA_PEDOGEOCHIMICA.pdf).  
1031 Accessed 27 October 2020.
- 1032 Rizzi, L., Petruzzelli, G., Poggio, G., Guidi, G.V., 2004. Soil physical changes and plant availability of Zn and Pb  
1033 in a treatability test of phytostabilization. *Chemosphere* 57(9), 1039-1046. Doi:  
1034 10.1016/j.chemosphere.2004.08.048.
- 1035 Robinson, B.H., Leblanc, M., Petit, D., Brooks, R.R., Kirkman, J.H., Gregg, P.E.H., 1998. The potential of *Thlaspi*  
1036 *caerulescens* for phytoremediation of contaminated soils. *Plant and Soil* 203, 47-56. Doi:  
1037 10.1023/A:1004328816645.
- 1038 Rogosic, J., Pfister, J. A., Provenza, F. D., Grbesa, D., 2006. Sheep and goat preference for and nutritional value  
1039 of Mediterranean maquis shrubs. *Small Ruminant Research*, 64(1-2), 169-179. Doi:  
1040 10.1016/j.smallrumres.2005.04.017.
- 1041 Shapiro, S.S., Wilk, M.B., 1965. An analysis of variance test for normality (complete samples). *Biometrika* 52, pp.  
1042 591.
- 1043 Sidhu, G.P.S., Singh, H.P., Batish, D.R., Kohli, R.K., 2017. Tolerance and hyperaccumulation of cadmium by a  
1044 wild, unpalatable herb *Coronopus didymus* (L.) Sm. (Brassicaceae). *Ecotoxicology and Environmental Safety*  
1045 135, 209–215. Doi: 10.1016/j.ecoenv.2016.10.001.
- 1046 Singh, B.R., Gupta, S.K., Azaizeh, H., Shilev, S., Sudre, D., Song, W.Y., Martinoia, E., Mench, M., 2011. Safety  
1047 of food crops on land contaminated with trace elements. *Journal of the Science of Food and Agriculture*  
1048 91(8), 1349-1366. Doi: 10.1002/jsfa.4355.
- 1049 Spearman, C., 1904. The proof and measurement of association between two things. *The American Journal of*  
1050 *Psychology* 15, 72-101.
- 1051 Terry, N., 1980. Limiting factors in photosynthesis: I. Use of iron stress to control photochemical capacity in vivo.  
1052 *Plant Physiology* 65(1), 114-120. Doi: 10.1104/pp.65.1.114.

1053 Torres, G. N., Camargos, S. L., Weber, O. L. D. S., Maas, K. D. B., Scaramuzza, W. L., Pereira, M., 2016. Growth  
1054 and micronutrient concentration in maize plants under nickel and lime applications. *Revista Caatinga*, 29(4),  
1055 796-804. Doi: 10.1590/1983-21252016v29n403rc.

1056 USEPA, 2000. Introduction to phytoremediation. EPA 600/R-99/107. U.S. Environmental Protection Agency,  
1057 Office of Research and Development, Cincinnati, OH.

1058 Vamerali, T., Bandiera, M., Mosca, G., 2010. Field crops for phytoremediation of metal-contaminated land. A  
1059 review. *Environmental Chemistry Letters* 8(1), 1-17. Doi: 10.1007/s10311-009-0268-0.

1060 Van der Ent, A., Baker, A.J.M., Reeves, R.D., Pollard, A.J., Schat, H., 2013. Hyperaccumulators of metal and  
1061 metalloids trace elements: Facts and fiction. *Plant Soil* 362, 319-334. Doi: 10.1007/s11104-012-1287-3.

1062 Wang, B., Chen, S., Chen, Y., Belzile, N., Zheng, R., Yang, Y., Fu, K., Chen, Y., Lin, B., Liu Z., Sun, J., 2021. The  
1063 geochemical behavior of trace metals and nutrients in submerged sediments of the Three Gorges Reservoir  
1064 and a critical review on risk assessment methods. *Environmental Science and Pollution Research*, 1-16. Doi:  
1065 10.1007/s11356-021-12827-8.

1066 Wang, Z., Liu, X., Qin, H., 2019. Bioconcentration and translocation of heavy elements in the soil-plants system  
1067 in Machangqing copper mine, Yunnan Province, China. *Journal of Geochemical Exploration* 200, 159-166.  
1068 Doi: 10.1016/j.gexplo.2019.02.005.

1069 Wong, M.K., Chuah, G.K., Ang, K.P., & Koh, L.L., 1986. Interactive effects of lead, cadmium and copper  
1070 combinations in the uptake of metals and growth of *Brassica chinensis*. *Environmental and experimental*  
1071 *botany*, 26(4), 331-339. Doi: 10.1016/0098-8472(86)90020-1.

1072 Wu, Q., Leung, J.Y.S., Huang, X., Yao, B., Yuan, X., Ma, J., Guo, S., 2015. Evaluation of the ability of black  
1073 nightshade *Solanum nigrum* L. for phytoremediation of thallium-contaminated soil. *Environmental Science*  
1074 *and Pollution Research*, 22(15), 11478-11487 Doi: 10.1007/s11356-015-4384-z.

1075 Yao, Z., Li, J., Xie, H., Yu, C., 2012. Review on remediation technologies of soil contaminated by heavy metals.  
1076 *Procedia Environmental Sciences* 16, 722-729. Doi: 10.1016/j.proenv.2012.10.099.

1077 Yoon, J., Cao, X., Zhou, Q., Ma, Q.L., 2006. Accumulation of Pb, Cu and Zn in native plants growing on a  
1078 contaminated Florida site. *Science of the Total Environment* 368, 456-464. Doi:  
1079 10.1016/j.scitotenv.2006.01.016.

1080 Zhu, X.F., Lei, G.J., Jiang, T., Liu, Y., Li, G.X., Zheng, S.J., 2012. Cell wall polysaccharides are involved in P-  
1081 deficiency-induced Cd exclusion in *Arabidopsis thaliana*. *Planta* 236(4), 989-997. Doi: 10.1007/s00425-012-  
1082 1652-8.

1083

1084