

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Is mimicry a diversification-driver in ants? Biogeography, ecology, ethology, genetics and morphology define a second West-Palaeartic Colobopsis species (Hymenoptera: Formicidae)

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Schifani, E., Giannetti, D., Csoz, S., Castellucci, F., Luchetti, A., Castracani, C., et al. (2022). Is mimicry a diversification-driver in ants? Biogeography, ecology, ethology, genetics and morphology define a second West-Palaeartic Colobopsis species (Hymenoptera: Formicidae). ZOOLOGICAL JOURNAL OF THE LINNEAN SOCIETY, 194(4), 1424-1450 [10.1093/zoolinnean/zlab035].

Availability:

This version is available at: <https://hdl.handle.net/11585/896741> since: 2024-05-15

Published:

DOI: <http://doi.org/10.1093/zoolinnean/zlab035>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

Mimicry as a diversification driver in ants? Biogeography, ecology, ethology, genetics, and morphology define a second West-Palearctic *Colobopsis* species (Hymenoptera: Formicidae)

ENRICO SCHIFANI^{1,†,*}, DANIELE GIANNETTI^{1,†}, SÁNDOR CSÖSZ^{2,†}, FILIPPO CASTELLUCCI³, ANDREA LUCHETTI³, CRISTINA CASTRACANI¹, FIORENZA A. SPOTTI¹, ALESSANDRA MORI^{1,‡}, DONATO A. GRASSO^{1,‡}

¹ Department of Chemistry, Life Sciences & Environmental Sustainability, University of Parma, Parco Area delle Scienze, 11/a, 43124 Parma, Italy

² MTA-ELTE-MTM, Ecology Research Group, Pázmány Péter sétány 1C, H-1117 Budapest, Hungary

Evolutionary Ecology Research Group, Institute of Ecology and Botany, Centre for Ecological Research, Vácrátót, Hungary

³ Department of Biological, Geological and Environmental Sciences – University of Bologna, via Selmi 3, 40126 Bologna, Italy

† co-first authors

‡ co-last authors

* corresponding author: enrico.schifani@unipr.it

ORCID NUMBERS: ES - 0000-0003-0684-6229; DG - 0000-0003-1622-1329; SC - 0000-0002-5422-5120; FC - 0000-0002-9944-2196; AL - 0000-0002-2986-721X; CC - 0000-0001-8802-9614; FAS - 0000-0002-4270-7229; AM - 0000-0001-6544-3038; DAG: 0000-0001-9334-4280.

Running title: Mimicry-driven diversification in *Colobopsis* ants?

ACKNOWLEDGEMENTS

We are deeply grateful to the people who helped us during our investigation, particularly those who donated the *Colobopsis* specimens they collected: Rumsaïs Blatrix (CEFE/University of Montpellier, France), Marco Forcieri (Palermo, Italy), Vincenzo Gentile (Torre Annunziata, Italy), Kiko Gómez (Barcelona, Spain), Mattia Menchetti (Institut de Biologia Evolutiva, Spain), Guido Maniscalco (Palermo, Italy), Massimiliano Napoli (Palermo, Italy), Joaquin Reyes-López (University of Córdoba, Spain), Daniel Sánchez García (Polish Academy of Science, Poland), and Ahmed Taheri (Chouaib Doukkali University, Morocco). We wish to thank the museum curators who kindly allowed us to access the relevant type material we examined: Fulvio Giachino (Museo Regionale di Scienze Naturali - Turin, Italy) and Dominique Zimmermann (Natural History Museum - Vienna, Austria). We are thankful to Piergiorgio di Pompeo (Bologna, Italy) and Virgilio Caleca (University of Palermo, Italy) for identifying some of the arthropods mentioned in this paper. Our thanks also go to Lech Borowiec (University of Wroclaw, Poland) and Herbert C. Wagner (University of Innsbruck, Austria) for some interesting personal communications, and Ghania Barech (University Mohamed Boudiaf - M'sila, Algeria) for pictures of Algerian *Colobopsis* specimens. Finally, we wish to thank Brian L. Fisher and Michele Esposito (California Academy of Sciences) for uploading AntWeb pictures and Anila R. Scott-Monkhouse (University of Parma) for proofreading our manuscript.

FUNDING

This work was supported through grants from the University of Parma (FIL-2018, 2019) assigned to D.A.G. and A.M., by grant No. K 135795 of the National Research, Development and Innovation fund assigned to S.C. and Canziani funding assigned to A.L. The work also benefited from the equipment and framework of the COMP-HUB Initiative, funded by the 'Departments of Excellence' program of the Italian Ministry for Education, University and Research (MIUR, 2018-2022).

ABSTRACT

The West-Palearctic *Colobopsis* ant populations have long been considered a single species (*Colobopsis truncata*). We studied the diversity of this species by employing a multidisciplinary approach and combining data from our surveys, museum and private collections, and citizen science platforms. As a result, we have revealed the existence of a second species, which we describe as *Colobopsis imitans* sp. nov., distributed allopatrically from *C. truncata* and living in the Maghreb, Sicily and Southern Iberia. While the pigmentation of *C. truncata* is reminiscent of *Dolichoderus quadripunctatus*, that of *C. imitans* sp. nov. is very similar to *Crematogaster scutellaris*, with which *C. imitans* sp. nov. lives in close spatial association, and whose foraging trails it habitually follows, similarly to *Camponotus lateralis* and other ant-mimicking ants. The isolation between *C. imitans* sp. nov. and *C. truncata* seems to have occurred relatively recently because of the significant, yet not extreme morphometric differentiation, and to mtDNA polyphyly. Both *C. imitans* sp. nov. and *C. truncata* appear to employ mimicry of an unpalatable or aggressive ant species as an important defensive strategy; this ‘choice’ of a different model species is motivated by biogeographic reasons and appears to act as a critical evolutionary driver of their diversification.

ADDITIONAL KEYWORDS: ADAPTATION – BATESIAN MIMICRY – COI MTDNA – DISCRIMINANT-FUNCTION ANALYSIS – MEDITERRANEAN – MULTIVARIATE STATISTICS – NORTH AFRICA – SIBLING SPECIES – SPECIATION.

INTRODUCTION

Ants (Insecta: Formicidae) are a hyperdiverse group of organisms which counts about 13,860 species (Bolton, 2021) and is extraordinarily successful in most terrestrial ecosystems (Hölldobler & Wilson, 1990; Gibb *et al.*, 2017; Seifert, 2017). Such high diversification stems from several evolutionary strategies and lifestyles, and enables even hundreds of different ant species to coexist in the same habitat (Hölldobler & Wilson, 1990; 2008). However, only one or very few dominant species generally characterize even the most species-rich ant communities: these species form very populous colonies, with large, often permanent foraging trails, and they defend territories that may extend over hectares (Hölldobler & Wilson, 1990; Andersen, 1995; 1997; Grasso *et al.*, 1998; 1999; 2005; Arnan *et al.*, 2018). These ants are exposed to higher predation risk compared the ones whose workers forage solitarily or in small groups, and are accordingly equipped with effective defensive mechanisms (Buschinger & Maschwitz, 1984; Hölldobler & Wilson, 1990; Dornhaus & Powell, 2010; Seifert, 2018). Most ant species live in small colonies and forage solitarily or in small groups, only occasionally form trails, and develop evasive anti-predatory strategies (e.g. Hölldobler & Wilson, 1990; Tautz *et al.*, 1994; Andersen, 1995; Dornhaus & Powell, 2010; Helms *et al.*, 2014; Larabee & Suarez, 2015; Seifert, 2018; Grasso *et al.*, 2020). Of the species that live in small colonies, the only ones armed with dangerous defences are some predatory ants which retain the primitive feature of a powerful functional stinger to hunt (Buschinger & Maschwitz, 1984; Hölldobler & Wilson, 1990; Dornhaus & Powell, 2010).

Well-armed ant species, in particular the ones that build large colonies, are a very good model for several mimicking organisms, mostly arthropods. Some of these mimics are myrmecophilous organisms: commonly ant predators or parasites, they have adapted to live within or close to ant colonies by relying on chemical or acoustic mimicry (e.g. Geiselhardt *et al.*, 2007; Barbero *et al.*, 2009; Cushing, 2012; Parker & Grimaldi, 2014; Parker, 2016; Scarparo *et al.*, 2019). On the other hand, myrmecomorph species resemble their ant model thanks to morphological and/or behavioural adaptations (e.g. Komatsu 1961; Jackson & Drummond, 1974; Oliveira & Sazima, 1984; Oliveira 1988; Cobben, 1986; McIver, 1987; McIver & Stonedahl, 1993; Trjapitzin & Trjapitzin, 1995; Cassis & Wall, 2010; Chandler, 2010; Durkee *et al.*, 2011; Huang *et al.*, 2011; Cushing, 2012; Pekár, 2014; Corcobado *et al.*, 2016; Pekár *et al.*, 2017; Harvey *et al.*, 2018; de L. Nascimento & Perger, 2018; Gnezdilov, 2019). The main aim of myrmecomorphism is predation avoidance: compared to the mimics, models usually possess superior defensive mechanisms and are also more numerous. As such, they are usually interpreted as Batesian mimics (e.g. Jackson & Drummond, 1974; McIver, 1987; Durkee *et al.*, 2011; Huang *et al.*, 2011; Cushing, 2012; Harvey *et al.*, 2018), whose evolution can be favoured by model abundance (Kikuchi & Pfenning, 2010).

64 Many ant mimics are ants themselves. Some are parasites (e.g. inquilines) that act similarly to
65 myrmecophilous organisms (Buschinger, 2009), relying on chemical adaptations to interact with the
66 host species. However, there are some free-living ant species that act as mimics of more aggressive
67 or dominant ant species and are therefore interpreted as Batesian mimics. Nonetheless, solid empirical
68 evidence to reject the alternative hypothesis of Müllerian mimicry (see Müller, 1879; Pasteur, 1982;
69 Ritland, 1991) is rarely available (Ito *et al.*, 2004; Wagner, 2014). In these species, chromatic mimicry
70 is the prevalent mechanism, while behavioural or morphological adaptations are more rarely
71 documented (Emery, 1886; Forel, 1886; Santschi, 1919; Gobin *et al.*, 1998; Merrill & Elgar, 2000; Ito
72 *et al.*, 2004; Ward, 2009; Gallego-Roperro & Feitosa, 2014; Powell *et al.*, 2014; Pekár *et al.*, 2017;
73 Rasoamanana *et al.*, 2017; Seifert, 2019a). In addition, recurrent behaviour among mimicking ant
74 species is interspecific trail-following, which consists in the mimics regularly infiltrating into the
75 foraging trails of the model and may lead to parasitic behaviour with regard to food resources (Emery,
76 1886; Santschi, 1919; Gobin *et al.*, 1998; Ito *et al.*, 2004; Menzel *et al.*, 2010; Powell *et al.*, 2014).
77 Unlike specialized parasites (e.g. Visicchio *et al.*, 2001; Buschinger, 2009; de la Mora *et al.*, 2020),
78 no advanced mechanisms of chemical deception exist in most of these cases, so in the eventuality of
79 direct encounters the model species recognizes and attacks the mimic, which however is well-
80 equipped to escape (Goetsch, 1942; 1951; Kaudewitz, 1955; Gobin *et al.*, 1998; Ito *et al.*, 2004;
81 Menzel *et al.*, 2010). The vast majority of the hitherto well-documented cases come from the tropics:
82 mimics mainly belong to diverse lineages from the Formicinae tribe Camponotini (mostly
83 *Camponotus* Mayr, 1861), while their models are phylogenetically scattered, including
84 Ectatomminae, Myrmeciinae, Myrmicinae (Crematogastrini and Stenammini) and
85 Pseudomyrmecinae. Only one case of ant-mimicking ant species is well-documented in the West-
86 Palearctic zone, i.e. *Camponotus lateralis* (Olivier, 1792). It chromatically mimics the similar
87 *Crematogaster* species *Cr. ionia* Forel, 1911, *Cr. scutellaris* (Olivier, 1792) and *Cr. schmidtii* (Mayr,
88 1853), and follows their trails (Emery, 1886; Baroni Urbani, 1969; Menzel *et al.*, 2010; Wagner,
89 2014; Seifert, 2018; 2019a). The three species are closely related with neighbouring geographic
90 ranges and differ slightly chromatically (Blaimer, 2012). Interestingly, there seems to be a geographic
91 trend in the chromatic pattern of *Ca. lateralis*, allowing it to better resemble these three
92 *Crematogaster* species in the regions of sympatric occurrence (Wagner, 2014; Seifert, 2019a).

93 The ant genus *Colobopsis* Mayr, 1861 (Formicinae: Camponotini), recently separated from
94 *Camponotus* (Ward *et al.*, 2016), currently counts 95 valid species and 21 subspecies (Bolton, 2021).
95 It is distributed across the Holarctic, Indomalayan and Australasian regions, and is most diversified
96 in the latter two regions (Janicki *et al.*, 2016; Guénard *et al.*, 2017). *Colobopsis* species are usually
97 arboreal ants that nest in dead wood, form small-sized colonies and behave timidly towards other ants

98 (Wheeler, 1904; Ward *et al.*, 2016). In the Western-Palearctic region, *Colobopsis truncata* (Spinola,
 99 1808) is the only recognized species of its genus. The queen caste was described by Spinola (1808)
 100 from north-western Italy (Liguria region), while the other castes were described later (Dufour &
 101 Perris, 1840; Forel, 1874; Emery, 1916). Another species, *Co. fuscipes* (Mayr, 1853) was described
 102 from Austria by Mayr (1853), but was later reclassified as a junior synonym of *Co. truncata* (Emery
 103 & Forel, 1879). *Colobopsis truncata* is therefore considered to have a wide geographic distribution,
 104 from the Caucasus to Iberia and from Central Europe to the Maghreb (Seifert, 2018; Janicki *et al.*,
 105 2016; Guénard *et al.*, 2017). It is an arboreal-nesting species, as is typical of the genus, and it
 106 preferably nests on broadleaved trees, where it lives in monogynous and often polydomous colonies,
 107 rarely exceeding 500 workers (Seifert, 2018). Queens and soldiers are specialized for phragmosis,
 108 and soldiers may also function as repletes (living containers of liquid food), seldom leaving the safety
 109 of the nests they guard (Brun 1924; Goetsch, 1950; 1953; Seifert, 2018). Minor workers are usually
 110 active outside the nest during both day and night, forage solitarily, do not recruit nest mates to food
 111 sources, and perform very quick evasive movements when encountering other ants (Seifert, 2018).

112 During field observations across Italy we encountered marked divergences between *Colobopsis*
 113 colonies: workers of some colonies resembled *Cr. scutellaris* and followed its trails, while others
 114 resembled *Dolichoderus quadripunctatus* (Linnaeus, 1771), two species having remarkably different
 115 appearance. This is reflected by several contradictory reports which however contain no comment on
 116 these incongruences. Forel (1874) first referred to Swiss ants by considering *Co. truncata* a Batesian
 117 mimic of *D. quadripunctatus*. He later suggested that the two species show pre-adaptations to
 118 parabiogenic nest-sharing (Forel, 1903). Zimmermann (1934) instead studied ants in Croatia and
 119 considered the occasional relationship between *Co. truncata* and *Cr. scutellaris* similar to the one
 120 between *Ca. lateralis* and *Cr. scutellaris*. Goetsch (1942) stated that in Spain *Co. truncata* behaves
 121 similarly to *Ca. lateralis*, following the trails of *Cr. scutellaris*, , yet is also chromatically very
 122 different and shows no adaptation to mimicry. Baroni Urbani (1971) reported on a case of trail-
 123 following between a *Co. truncata* queen and a *Cr. scutellaris* trail from central Italy. More recently,
 124 Tinaut (1991) claimed that in southern Iberia *Co. truncata* can be easily confused with *D.*
 125 *quadripunctatus* during field surveys due to their similarity. However, working in the same area,
 126 Carpintero *et al.* (2001; 2005) instead affirmed that *Co. truncata* is a visual mimic of *Cr. scutellaris*
 127 and follows its trails, and even speculated that after nuptial flights *Co. truncata* queens specifically
 128 choose trees occupied by *Cr. scutellaris* to found their colony. In reviewing the distribution of
 129 *Colobopsis* in Iberia, García (2020) mentioned possible chromatic similarity of *C. truncata* with *Cr.*
 130 *scutellaris* and its mimic *Ca. lateralis*, but also *D. quadripunctatus*. More recently, there have been
 131 reports on trail-following by *Co. truncata* on *Cr. scutellaris* ants in Italy (mentioned by Seifert 2018,

no locality specified; Lake Garda according to Wagner HC, personal communication). Wagner (2019) described a close association between *Co. truncata* and *D. quadripunctatus* in Austria (Vienna), highlighted morphological and chromatic similarity between the two, and reported trail-following of *D. quadripunctatus* trails by *Co. truncata*.

We aimed to test whether the abovementioned diversity of traits represented intraspecific variation or indicated the existence of multiple *Colobopsis* species in the West-Palearctic. In order to address the different biological dimensions of this problem, we used a multidisciplinary approach which involves the description of the patterns of diversity within the traditional notion of *Co. truncata*. We relied on the principles of integrative taxonomy (Schlick-Steiner *et al.*, 2010), involving morphological, genetic, ecological, ethological and biogeographical data.

MATERIALS AND METHODS

We combined qualitative morphology through chromatic pattern evaluation, quantitative morphology through the multivariate analysis of morphometric data and genetics in the form of mtDNA (COI) sequencing, ecological surveys to study the association between *Colobopsis* and its putative model species and recorded ethological data to document cross-species trail following. Finally, we compared our results with the known biogeographic patterns of other ant species. We chose this quantitative morphological approach as it is widely regarded as the most practical and reliable single source of evidence for cryptic ant species delimitation, and as a cornerstone in integrative approaches on cryptic species complexes of ants (Seifert, 2009; 2018; Seifert *et al.*, 2014; Wagner *et al.*, 2017; Steiner *et al.*, 2011; 2018; Csősz *et al.*, 2020). Moreover, mtDNA sequencing represents a widespread and cost-effective method to gain preliminary information on species identification, biogeography and cryptic speciation (Hebert *et al.*, 2003; 2016; Ratnasingham & Hebert, 2007), which has developed into an aid to myrmecological faunistic, biogeographic and taxonomic studies (Steiner *et al.*, 2005; 2018; Csősz *et al.*, 2015; Seifert *et al.*, 2017; Schär *et al.*, 2018; 2020; Blatrix *et al.*, 2020). Ecological data on species associations and ethological data are rarely used in ant taxonomy but appear to be highly relevant to the specific case we are investigating, while biogeography is important to understand species diversity.

For our morphological and molecular analyses, we gathered type material of *Co. truncata* and *Co. fuscipes* as well as additional non-type material of *Colobopsis* from the Mediterranean region, and relied on our own efforts and the generous contribution of colleagues to achieve a satisfactory geographic coverage. In particular, the type series of *Co. truncata* consists in a single queen with the label “*Polyergus* (?) | *F.ca truncata* | Spin. in Ligur. | Genova || 6571”. This queen could be safely identified as the type since it is the sole *Colobopsis* queen in the Spinola collection at the Museo di

165 Scienze Naturali in Turin (Italy), and matched the description given by the author (Spinola, 1808).
166 Concerning *Co. fuscipes*, at least two syntypes are stored in the Museum für Naturkunde, University
167 of Berlin (Germany), and their pictures are available on AntWeb (AntWeb.org, codes FOCOL2496
168 and FOCOL2497): these are labelled “Oesterreich | Coll. Rhd || *Colobopsis fuscipes* Mayr || Type ||
169 29812 || GBIF-D/FoCol | 2496 | specimen + label | data documented” and “Oesterreich | Coll. Rhd ||
170 *Colobopsis fuscipes* Mayr || Type || GBIF-D/FoCol | 2497 | specimen + label | data documented”.
171 Although the label is unlikely to be an original by Gustav Mayr (B. Seifert, pers. comm.), we deem
172 their status as types credible. We also retrieved a worker labelled *Co. fuscipes* in Mayr’s collection at
173 the Natural History Museum of Vienna, but with no explicit indication ensuring its type status. In
174 order to gather information on chromatic variation of Euro-Mediterranean *Colobopsis*, we relied on
175 AntWeb pictures, images from scientific papers or monographs (Glaser, 2009; Wagner, 2014; 2019;
176 Lebas *et al.*, 2016; Seifert, 2018; Scupola, 2018; García, 2020; Salata *et al.*, 2020; Tăușan *et al.* 2020),
177 and on georeferenced photographs uploaded on citizen science platforms (iNaturalist.org,
178 biodiversidadvirtual.org) and on biodiversity-related Facebook groups. A complete list of the material
179 examined, their depositories and collecting data is available as a Supplementary Material file to this
180 paper. Ecological and behavioural data were obtained through field surveys across Italy.

181 PIGMENTATION: CHROMATIC MIMICRY

182 Preliminary observations highlighted that chromatic pattern provides the most evident difference
183 between allopatric *Colobopsis* populations which resemble two different model ant species in Italy.
184 We preemptively described the two chromatic forms and investigated whether these were consistently
185 able to represent *Colobopsis* diversity across the Mediterranean basin, and whether they occurred
186 intracolony or sympatrically, and also checked for the possible existence of third forms. The two
187 *Colobopsis* model patterns were pre-emptively established by observing ten workers per each form
188 (10 from Sicily and 10 from mainland Italy), and all subsequent investigations were carried out by
189 inspecting specimens of well-preserved pigmentation primarily belonging to the worker caste.
190 Queens and soldiers were also examined and identified only if closely resembling one worker model
191 pattern, while males were not considered due to their very different pigmentation. Furthermore, to
192 better describe the differences between the two models, a ratio was calculated between head color
193 and mesosoma color by taking dorsal pictures of specimens and calculating the average value of red
194 (RGB colors) between 5 randomly selected pixels of the head and 5 of the mesosoma via software
195 ImageJ (Schneider *et al.*, 2012). Calculating a ratio rather than considering the absolute values greatly
196 reduces the variation produced by different light conditions and camera settings among different
197 pictures, thereby allowing comparison of pictures from various sources. The same ratio was also
198 calculated for the two putative model species *Cr. scutellaris* and *D. quadripunctatus*. Chromatic ratios

199 were calculated on 20 workers per chromatic pattern or species from across their respective
200 geographic range using both directly inspected specimens and images from citizen science platforms.
201 Any differences were statistically tested by using the software R 4.0.3 and R Studio 1.3.1056 (R Core
202 Team, 2021), and employing an ANOVA test and subsequent Tukey Post-hoc test for pairwise
203 comparisons.

204 The visual examination to verify correspondence to either of the two chromatic models was conducted
205 on 79 directly observed *Colobopsis* colonies (76 of which containing workers) plus images of 136
206 further specimens (including 76 isolated queens), for a total of 310 workers and 79 queens covering
207 a total of 16 countries from across the W-Palearctic *Colobopsis* distribution (see supplementary
208 material).

209 The two model patterns are defined as follows:

210 *Cr. scutellaris*-like pattern (CSL pattern): head, or head and anterior part of the mesosoma (rarely
211 most of it) uniformly red, rest of the body evidently darker and mostly black. White stripes or dots
212 on the second gastral tergite often absent or weak (present in 10% of the examined workers). See Fig.
213 1.

214 *D. quadripunctatus*-like pattern (DQL pattern): head, mesosoma and appendages from reddish to
215 blackish (therefore chromatically more variable than the *Cr. scutellaris*-like model), head at least
216 slightly darker than the mesosoma or less frequently concolour, gaster black. White stripes or dots on
217 the second gastral tergite often present (80% of examined workers) and more obvious. Phragmotic
218 heads of soldiers or queens are always reddish in their anterior, heavily sculptured part (approximately
219 one half of the head), while the rest follows the same scheme of workers. See Fig. 1.



Figure 1. The model species and the two detected chromatic model patterns of *Colobopsis*: a) *Cr. scutellaris*; b) CSL *Colobopsis* from Sicily; c) *D. quadripunctatus* from Tuscany; d) DQL *Colobopsis* from Tuscany.

NUMERIC MORPHOLOGY: MULTIVARIATE ANALYSES OF MORPHOMETRIC DATA

A total of 12 continuous morphometric traits were defined following Seifert (2018) (Tab. 1) and measured on 115 *Colobopsis* workers from 44 nest samples (considering minor workers only, and not the soldiers). All measurements were made in μm by using a pin-holding stage, which allowed rotations around the X, Y, and Z axes. An Olympus SZX9 stereomicroscope was used at x150 magnification for each character; however, with characters larger than the field of view x75 magnification was applied. Due to the low number of the much rarer queens, males and soldiers in our possession, we recorded only a reduced set of 7 morphometric traits aimed at providing a brief description of these castes without using them in the following statistical analyses. Morphometric data are provided in μm throughout the whole paper.

Repeatability of the recorded size parameters were evaluated via Intraclass Correlation Coefficients (ICC) by using Package ICC (Wolak *et al.*, 2012), see Tab. 1. Variables were tested via matrix scatterplots and Pearson product-moment correlation coefficients for error variance and outliers. Each character resulted highly repeatable, except for NOL, which was considered moderately repeatable.

Exploratory analyses through NC-PART clustering

238 The prior species hypothesis was generated based on workers through combined application of NC
 239 clustering (Seifert *et al.*, 2014) and Partitioning Based on Recursive Thresholding (PART) (Nilsen &
 240 Lingjaerde, 2013). The script for NC-clustering combined with PART was written in R and can be
 241 found in Appendix S1 in Csősz & Fisher (2016). Our exploratory data analysis approach follows the
 242 protocol described by Csősz & Fisher (2016) with the following specific settings: bootstrap iterations
 243 in PART were set to ‘b=1000’, and the minimum size of clusters was set to ‘minSize=5’ for both
 244 ‘hclust’ and ‘kmeans’. The optimal number of clusters and the partitioning of samples are accepted
 245 as the preliminary species hypothesis in every case in which the two clustering methods, ‘hclust’ and
 246 ‘kmeans’ through PART, have yielded the same conclusion.

247 *Exploratory analyses via PCA using allometrically corrected data*

248 An alternative prior species hypothesis has been generated via the ordinating Principal Component
 249 Analysis (PCA) that searches for discontinuities in continuous morphometric data and display plots
 250 in a graphic.

251 Using raw data (without removal of allometric variance) in PCA may lead to weaker performance in
 252 ordination because the first vector of the PCA often describes the size component, which is a useless
 253 information when cryptic species of similar size have to be separated, hence, in PCA residuals were
 254 used. Residuals, in which the head length (CL) was used as covariate, were calculated via a linear
 255 regression model according to the following steps: a) scaling properties, intercept and steepness were
 256 calculated for each nest sample separately (note: nest samples constituted by a singleton were not
 257 involved in this phase); b) a grand average for steepness and intercept was calculated from scaling
 258 properties of each nest sample; c) residuals are calculated for every nest sample (including singletons)
 259 based on the grand average. Residuals of every trait calculated against head length (CL) are given
 260 (Tab. 1). In contrast to NC-PART clustering, the PCA has no estimation on the number of clusters
 261 and “classification” of objects has been made based on subjective decision. The coefficients (x any
 262 intercept) for removal allometric variance for each trait are given in supplementary material.

263 *Hypothesis testing by confirmatory analysis*

264 The validity of the prior species hypothesis was tested via Linear Discriminant Analysis (LDA).
 265 Classification hypotheses were imposed for all samples that were congruently classified by
 266 partitioning methods, while wild-card settings (i.e. no prior hypothesis imposed on its classification)
 267 were given to samples that were incongruently classified by the two partitioning methods. Statistical
 268 analyses were conducted through the software R 3.6.3 (R Core Team 2021).

Abbr.	Description of the trait	ICC (R)
-------	--------------------------	---------

CL	Maximum median length of head capsule. The head must be carefully tilted so the maximum length is positioned in the measuring plane.	0.982
CW	Maximum head with including compound eyes. The largest distance between profiles of the two compound eyes in full-face view.	0.951
EL	Eye length. Maximum diameter of the compound eye.	0.967
dAN	Minimum distance of the inner margins of antennal socket rings.	0.985
ML	Diagonal length of the alitrunk in profile. Measured in lateral view from the anteriormost point of anterior pronotal slope to the caudalmost point of the lateral metapleural lobe.	0.969
MW	Maximum width of pronotum.	0.989
NOL	Petiole node length; measured in lateral view, from the center of the petiolar spiracle to the posterior profile.	0.890
PeW	Petiole width. The maximum width of petiole in dorsal view.	0.994
PreOC	Preocular distance. Use a cross-scaled ocular micrometer and adjust the head to the measuring position of CL. Frontal measuring point: median clypeal margin; caudal measuring point: reference line between the frontalmost border of the two compound eyes.	0.951
SL	Scape length. The maximum straight-line scape length excluding the articular condyle.	0.971
HTL	Hind tibia length. Measured from the distalmost point of the tibia to the proximal end where the tibia is narrowest in profile.	0.968
PeSH	Petiole scale height measured from the center of petiolar spiracle to top of the crest.	0.959

Table. 1 Abbreviation (Abbr.) of morphometric characters, definition of measurements, and ICC (R) a metric for repeatability parameter are given. Definitions of morphometri characters follow Seifert (2018).

GENETICS: MITOCHONDRIAL COI SEQUENCES

Total genomic DNA was extracted from leg tissues using the NucleoSpin DNA Insect kit (Macherey-Nagel, Düren, Germany), following the manufacturer's protocol. A 700 bp region of mitochondrial gene cytochrome *c* oxidase subunit I (COI) was amplified using the primer couple LCO1490/HCO2198 (Folmer *et al.*, 1994). PCR was carried out in 25 µL reactions using the following profile: initial denaturation step at 95 °C for 5 minutes, 35 amplification cycles (denaturation at 95 °C for 30 seconds, annealing at 52 °C for 30 seconds, elongation at 72 °C for 45 seconds), final elongation at 72 °C for 7 minutes. PCR products were sent to MacroGen Europe (Amsterdam, Netherlands) for Sanger sequencing. Chromatograms were checked and edited using SeqTrace (Stucky, 2012). Sequences were aligned using the MUSCLE (Edgar, 2004) algorithm as implemented in AliView (Larsson, 2014). Model selection and Maximum Likelihood phylogenetic analysis were performed on the IQ-TREE web server (Trifinopoulos *et al.*, 2016) using the Eastern-Palearctic *Co. nipponica* (Wheeler, W.M., 1928) and *Co. shohki* (Terayama, 1999) and the Indomalayan *Colobopsis* nr. *saundersi* (Emery, 1889) (GenBank accession numbers AB019417, AB019418 and KU975365, respectively) as outgroups. Ten separate runs were launched, each with 1000 replicates of ultrafast bootstrap, and the tree with the best likelihood value out of the ten was

287 chosen. Twenty-three colony samples were sequenced, consisting in 41 workers from 6 countries and
288 18 localities. Obtained sequences have been submitted to Genbank, under accession numbers
289 MW462045–MW462085 (see supplementary material).

290 ECOLOGY: COEXISTENCE WITH MODEL SPECIES

291 Field surveys were conducted in the Italian Peninsula (Emilia-Romagna, Tuscany; n sites = 5, DQL
292 pattern) and Sicily (n sites = 8, CSL pattern) to test whether the local *Colobopsis* populations, showing
293 a DQL and a CSL phenotype respectively, actually lived in close proximity with either of the two
294 species indicated as probable mimicry models (see Supplementary material). In each site, we searched
295 for the presence of *Colobopsis* workers on trees until a tree occupied by a *Colobopsis* colony was
296 detected. Then, we performed a 10 minutes-long continuous sampling within a 1.5 m radius of the
297 point of the tree trunk where *Colobopsis* was firstly observed, recording the eventual presence of *Cr.*
298 *scutellaris* or *D. quadripunctatus* workers.

299 Occurrences of *Cr. scutellaris* or *D. quadripunctatus* on trees occupied also by *Colobopsis* according
300 to the different DQL and CSL models were statistically tested by using the software IBM SPSS
301 statistics, Italian version 24 and the chi-squared test. All data are presented in the Supplementary
302 Material.

303 ETHOLOGY: INTERSPECIFIC TRAIL-FOLLOWING BEHAVIOUR

304 Field surveys were conducted in the Italian Peninsula (Emilia Romagna, Tuscany) and Sicily (sites
305 as in the section before, also see supplementary material) with the aim of quantifying the occurrence
306 of trail-following behaviour performed by *Colobopsis* ants in relation to *Cr. scutellaris* or *D.*
307 *quadripunctatus* trails. We selected trees where *Colobopsis* colonies coexisted with either *Cr.*
308 *scutellaris*, *D. quadripunctatus* or both. In accordance with the relevant literature (Gobin *et al.*, 1988;
309 Ito *et al.*, 2004; Menzel *et al.*, 2010; Powell *et al.*, 2014), trail-following was defined as the event of
310 *Colobopsis* workers moving along an established pheromone trail of *Cr. scutellaris* or *D.*
311 *quadripunctatus* within 1 cm from the trail itself. A 10 minutes continuous sampling was used to
312 record the presence or absence of this behaviour on each of the examined tree.

313 To study trail-following on *Cr. scutellaris* trails, we selected a total of 59 trees inhabited by this
314 species: 29 trees hosted *Colobopsis* colonies exhibiting the CSL pattern (Sicily, 4 sites) and 30 hosted
315 *Colobopsis* with the DQL pattern (Emilia-Romagna and Tuscany, 5 sites). Observations on *D.*
316 *quadripunctatus* trails could be performed only in 23 *Colobopsis* colonies exhibiting the DQL pattern
317 (Emilia-Romagna and Tuscany): since no *D. quadripunctatus* colonies could be found in the studied
318 sites in Sicily (where the species is known to be very rare, see Schifani & Alicata, 2018), no

319 *Colobopsis* colonies exhibiting the CSL pattern could be tested in this regard. All data are summarized
320 in the supplementary material.

321 SPECIES CONCEPT

322 Integration of the evidence provided by different complementary disciplines into an evolutionarily
323 credible species hypothesis is performed according to the principles emphasized by Schlick-Steiner
324 *et al.* (2010), i.e. resolving eventual disagreements by invoking solid evolutionary explanations.
325 Biogeography is here treated as an additional source of information, holding an important advisory
326 role to the formation of the final species-hypothesis. We abide by the universal Gene and Gene
327 Expression (GAGE) species concept proposed by Seifert (2020), which, although recently
328 formulated, convincingly summarizes the main theoretical and practical formulae most commonly
329 adopted during the last few decades as a rigorous approach on alpha taxonomy of cryptic ants,
330 especially in Europe.

331 RESULTS


332 PIGMENTATION: CHROMATIC MIMICRY

333 All the examined colonies are safely assignable to one of the two models and no transitional or third
334 forms are detected (see Supplementary Material). No intracolonyal coexistence of the two models is
335 detected either. The type series of both *Co. truncata* and *Co. fuscipes* show the DQL pattern (Fig. 2).
336 The two models occur strictly allopatrically according to the examined material. Samples from the
337 south-western Mediterranean basin, namely Algeria, Sicily (Italy), Morocco, southern Portugal and
338 Andalusia (southern Spain) are assigned to the CSL pattern. All the rest is assigned to the DQL
339 pattern, that is samples from Austria, Bulgaria, Croatia, Czech Republic, France, Germany, Greece,
340 Hungary, Israel, Italian peninsula (Apulia, Campania, Emilia-Romagna, Liguria, Tuscany), Romania,
341 Serbia, Slovenia, Switzerland, the rest of Spain (Aragon, Catalonia, Balearic Islands, Castilla-La
342 Mancha), and Turkey. In addition, photographs of *Co. truncata* specimens present in the European
343 ant fauna guides by Lebas *et al.* (2016) and Seifert (2018), regional faunistic guides by Glaser (2009)
344 (Liechtenstein), Wagner (2014) (Austria's Carinthia), Scupola (2018) (Italy's Veneto) as well as in
345 the Crete's ant fauna monograph by Salata *et al.* (2020) and in the papers by Wagner (2019) (Austria),
346 García (2020) (Spain) and Tăușan *et al.* (2020) (Romania) all show the DQL pattern.



Figure 2. Type material of the so far described West-Palearctic *Colobopsis*, all adhering to the “*D. quadripunctatus*-like” pattern. **a)** holotype queen of *Colobopsis truncata* from Liguria, Italy, preserved at the Turin Natural History Museum (Italy). **b)** syntype worker of *Colobopsis fuscipes* from Austria (picture from AntWeb.org, FOCOL2496; photographer: Christiana Klingenberg), preserved at the Museum für Naturkunde der Humboldt-Universität Berlin (Berlin, Germany). Note that the queen’s red color in the anterior heavily sculptured part of the phragmotic head is not relevant to evaluate its chromatic pattern. Scale bars: 0.5 mm.

The head red/mesosoma red ratio is statistically different among the two *Colobopsis* patterns and their models ($F_{3,76} = 152.4$, $p < 0.001$) (also see supplementary material). Pairwise comparisons show no statistically significant difference between CSL *Colobopsis* and *Cr. scutellaris* ($p = 0.817$; mean \pm sd = 1.96 ± 0.36 for CSL *Colobopsis*; mean \pm sd = 2.05 ± 0.38 for *Cr. scutellaris*) and between DQL *Colobopsis* and *D. quadripunctatus* ($p = 0.299$; mean \pm sd = 0.60 ± 0.27 for DQL *Colobopsis*; mean \pm sd = 0.43 ± 0.20 for *D. quadripunctatus*), while all other comparisons are significantly different ($p < 0.001$) (Fig. 3).



361

362 Figure. 3. Chromatic ratios calculated from pictures of the *Colobopsis* CSL and DQL patterns and from pictures of the
 363 two putative model species *Cr. scutellaris* and *D. quadripunctatus* (N=20 for each species or chromatic form). Boxplots
 364 show mean and standard deviation, while whiskers represent minimum and maximum values. Dots are measured
 365 individuals. Their dispersal on the x-axis is a randomized graphical effect to avoid overlaps.

366

NUMERIC MORPHOLOGY: MULTIVARIATE ANALYSES OF MORPHOMETRIC DATA

367 Two morphological clusters are identified via NC-clustering combined with “kmeans”, and “hclust”
 368 (Figs. 4). These two clusters correspond to the CSL pattern and DQL pattern specimens, respectively.
 369 All but two samples are congruently classified via both partitioning methods. The two incongruently
 370 placed samples (ITA:Mondello-VillaMercadante_col-12, ITA:Mondello-VillaMercadante_col-16;
 371 both CSL pattern from Sicily) are classified as belonging to the CSL cluster (posterior p = 0.85 and
 372 0.81, geometric means of 3 workers each). Without running samples as wild-cards, the overall
 373 classification success is 96.3% using all variables in the analysis.





Figure 4. Figure 4. Dendrogram comparing the results of “kmeans”, and “hclust” in NC Clustering of Colobopsis morphometric raw data. Two samples (4.5% of the total) are misplaced by both the dendrogram and one of the partitioning analyses, NC-part.kmeans; partially different samples being affected in each of the three analyses. The

378 other partitioning analysis, NC-part.hclust returned the same sample assignment as the LDA did.



379
380 Figure 5. Principal Component Analyses of morphometric data of analyzed *Colobopsis* nest samples according to the
381 two clusters evidenced by NC-PART Clustering. Each small dot represents a colony sample. Large dots represent
382 centroids.

383 T-tests were calculated to assess significant differences (p) of body size ratios between specimens of
384 the two different clusters, resulting in significant differences for 7 ratios (Tab. 2). Unfortunately, there
385 is not a single numeric body size ratio available for reliable separation of these clusters on individual
386 level (Tab. 2): the most reduced multivariate function that can reach the goal of attaining an
387 acceptably high rate of classification success (>95%) requires a minimum 6 morphometric characters
388 achieved via backward stepwise method.

389 The most simple D(6) function that yields 4.3% of error rate at the individual level is as follows:

390
$$D(6) = 0.03501 * CW - 0.03384 * SL - 0.03144 * HTL - 0.01762 * ML + 0.03653 * PeSH +$$

391
$$0.07458 * EL + 16.61469$$

392 D(6) scores for CSL pattern cluster (n = 55) = mean - 1.59 [- 4.17, + 0.66] ± 1.05

393 D(6) scores for DQL pattern cluster (n = 60) = mean - 1.54 [- 0.37, + 3.76] ± 0.96

394

395


character	CSL (n = 55)	p	DQL (n = 60)
CS	897 ± 51 [725, 1025]	0.924	898 ± 55 [803, 1042]
PreOc/CL	0.546 ± 0.01 [0.517, 0.571]	0.000	0.537 ± 0.01 [0.517, 0.570]
CL/CW	1.152 ± 0.02 [1.113, 1.198]	0.052	1.145 ± 0.02 [1.106, 1.189]
dAN/CS	0.387 ± 0.01 [0.345, 0.431]	0.128	0.391 ± 0.02 [0.358, 0.426]
SL/CS	0.872 ± 0.03 [0.798, 0.931]	0.000	0.846 ± 0.02 [0.798, 0.906]
MW/CS	0.682 ± 0.02 [0.648, 0.717]	0.476	0.680 ± 0.02 [0.648, 0.723]
PeW/CS	0.332 ± 0.02 [0.268, 0.372]	0.257	0.336 ± 0.02 [0.306, 0.395]
HTL/CS	0.931 ± 0.02 [0.888, 0.978]	0.000	0.902 ± 0.02 [0.845, 0.957]
ML/CS	1.485 ± 0.03 [1.395, 1.543]	0.000	1.460 ± 0.02 [1.410, 1.526]
NOL/CS	0.139 ± 0.01 [0.117, 0.165]	0.008	0.134 ± 0.01 [0.112, 0.157]
PeSH/CS	0.239 ± 0.02 [0.193, 0.284]	0.000	0.253 ± 0.02 [0.203, 0.293]
EL/CS	0.312 ± 0.01 [0.290, 0.335]	0.000	0.321 ± 0.01 [0.304, 0.343]

397 Table 2. Mean of morphometric ratios calculated for CSL pattern and DQL pattern clusters based on individuals (raw
398 data). Morphometric traits are divided by cephalic size (CS), namely the arithmetic mean of CL and CW. The upper row
399 in each data field gives arithmetic mean ± standard deviation, the lower one, in square brackets, lower and upper extremes.
400 Significant differences are highlighted in bold.

401 GENETICS: MITOCHONDRIAL COI SEQUENCES

402 The Maximum Likelihood phylogenetic analysis on mtDNA COI sequences identifies four main
403 clusters with good nodal support (Fig. 6). The clade A is formed by specimens exhibiting the CSL
404 pattern and collected from Andalusia (Spain) and Morocco. The specimens of the clade B exhibited
405 the DQL pattern and were sampled from Bulgaria, Castilla La Mancha (Spain), Hungary, the Italian

406 Peninsula, and the Occitanic region of France. The clade C groups specimens with the DQL pattern,
 407 sampled in the Spanish regions of Aragona and Catalonia, and from the Balearic Islands. Finally, the
 408 clade D is formed by all specimens from Sicily, showing the CSL pattern, and one of the Spanish
 409 specimens from Catalonia actually exhibiting the DQL pattern.



410
 411 Figure 6. Maximum likelihood phylogenetic tree based on the barcode fragment of the mtCOI gene from the sequenced
 412 *Colobopsis* specimens.

413 ECOLOGY: COEXISTENCE WITH MODEL SPECIES

414 *Crematogaster scutellaris* is present in 97% of the investigated trees occupied by CSL *Colobopsis*
 415 and in 20% of those occupied by DQL *Colobopsis*, the difference is statistically significant ($\chi^2_1 =$
 416 26.23, $p < 0.001$). *Dolichoderus quadripunctatus* was never detected in trees occupied by CSL
 417 *Colobopsis*. On the other hand, *D. quadripunctatus* occurs on 40% of the investigated trees occupied
 418 by DQL *Colobopsis* (6% of which also hosted *Cr. scutellaris*). Results are illustrated in Fig. 7, and
 419 detailed data is provided in the supplementary material.




Figure 7. Coexistence between *Colobopsis* and their putative model species on the same tree.

ETHOLOGY: INTERSPECIFIC TRAIL-FOLLOWING BEHAVIOUR

During field observations, trail-following behaviour was never performed by *Colobopsis* colonies exhibiting the DQL pattern (neither to *Cr. scutellaris* nor *D. quadripunctatus* trails). Conversely, 77% of the observed *Colobopsis* colonies exhibiting a CSL pattern had workers following the *Cr. scutellaris* trails (Fig. 8). Detailed data are shown in the supplementary material.

CSL *Colobopsis* followed trails of *Cr. scutellaris* by either walking directly on them (more scarcely populated worker trails with considerable gaps) or slightly sideways (crowded trails without or with very small gaps only). If coming into contact with a *Cr. scutellaris* worker, they immediately performed sudden accelerations and evasive movements. Trail-following often began a few moments after the *Colobopsis* worker left its nest and encounter a *Cr. scutellaris* trail on the tree trunk and ended with the *Colobopsis* worker leaving the trail and directing towards some specific twigs, no longer following *Cr. scutellaris* workers.




Figure 8. Trail-following behaviour on *Crematogaster scutellaris* trails by other ants (indicated with arrows). On the left (a,c,e) CSL *Colobopsis*, on the right (b,d,f) *Camponotus lateralis* observed in the same locality performing the same behaviour (photos taken in Palermo (Sicily) during field surveys).

BIOGEOGRAPHY, EVIDENCE DISCUSSION AND FINAL SPECIES HYPOTHESIS

CSL and DQL chromatic patterns are found to effectively split into two the Mediterranean *Colobopsis* into two populations, occurring allopatrically and each covering a vast geographic region (Fig. 9). The DQL pattern characterizes almost the entire European distribution of *Colobopsis* in addition to

Western Asia, while the CSL pattern occurs mainly in the Maghreb region (North-Western Africa), a well-recognized ant biodiversity hotspot (Borowiec, 2014), and in the European regions of greater biogeographic proximity to it (Sicily and Southern Iberia) (e.g. Alicata & Schifani, 2019; Tinaut & Ruano, 2021). In particular, this distribution mirrors strikingly well those of some camponotine ant species such as *Camponotus barbaricus* Emery, 1905, *Ca. micans* (Nylander, 1856) and *Ca. ruber* Emery, 1925 (Fig. 9; for their distribution see de Haro *et al.*, 1996; Forel, 1890; 1905; Santschi, 1925; Finzi, 1940; Menozzi, 1940; Cagniant, 1968; 1996; Collingwood & Yarrow, 1969; Baroni Urbani, 1971; Cagniant & Espadaler, 1993; Schembri & Collingwood, 1995; Janicki *et al.*, 2016; Guénard *et al.*, 2017; Schär *et al.*, 2020) and to a slightly lesser extent the distribution of myrmicine ants as the *Aphaenogaster crocea* species group, *A. sardoa* Mayr, 1855 or the *Temnothorax algiricus-mediterraneus* complex (see Mayr, 1853; Emery, 1880; Forel, 1909; Santschi, 1929; Galkowski & Cagniant, 2017; Alicata & Schifani, 2019). Ecological and behavioural field surveys across different Italian regions suggest that the two chromatic patterns are related to quite different lifestyles: the CSL pattern often coexists with *Cr. scutellaris* and very often follows its trails, while the DQL pattern is associated with *D. quadripunctatus* without the involvement of frequent trail-following. The CSL pattern characterized specimens from the Spanish locality where mimicry, close nesting association and extensive trail-following of *Cr. scutellaris* were described by Carpintero *et al.* (2001; 2005). On the other hand, the DQL pattern characterized samples from the Austrian region where Wagner (2019) based his suggestions of close association and mimicry between *Colobopsis* and *D. quadripunctatus*. A survey conducted in the region of Vienna (Austria) employing similar methodologies to ours estimated that 36% of the investigated *Colobopsis* colonies (n = 110) nested on trees occupied by *D. quadripunctatus*, confirming the trend observed in our study (Wagner, pers. comm.). However, interspecific trail-following between DQL pattern *Colobopsis* and either *D. quadripunctatus* or *Cr. scutellaris* as reported in Wagner (2019; personal communication) was never observed during our surveys and seems to represent a considerably less frequent phenomenon. Such differences between the two groups in chromatic pattern, biogeography and life history traits could arguably be sufficient to suggest a separation of the West-Palearctic *Colobopsis* into two species even according to a conservative classical taxonomical approach. Moreover, examined specimens from the two chromatic patterns are also classified into two morphometric clusters, whose separation reaches a significant threshold indicating heterospecificity according to the current procedures of cryptic ant species separation (Seifert, 2020). At the same time, the morphometric separation between the two clusters is relatively narrow, possibly indicating that the two species may have separated quite recently. Concerning the mtDNA phylogenetic analysis, each clade is unambiguously monophyletic with respect to morphometric and chromatic evidences (A and D = CSL pattern; B and C = DQL pattern),

476 with only one misplaced DQL specimen (a 2.4% error rate). On the other hand, with respect to
477 mtDNA, CSL and DQL patterns resulted in polyphy. This can be explained with possible retention
478 of ancestral polymorphisms and/or introgression of mtDNA (see Chan & Levin, 2005; Willis *et al.*,
479 2013). These phenomena appear, in fact, largely responsible for the actual estimate of paraphyly
480 emerging from mtDNA phylogenies analyses in about 20% of animal species (Funk & Omland, 2003;
481 Ross, 2014). Mitochondrial DNA introgression is, like in other eukaryotic groups, quite frequent in
482 ants (e.g. Darras & Aron 2015; Beresford et al. 2017; Seifert, 2018), and coalescence during
483 speciation commonly results in species undergoing through phases of polyphyly and paraphyly –
484 averagely longer in arthropods than in other groups – before normally reaching monophyly due to the
485 stochastic process of complete lineage sorting (Avise, 2004; Ross 2014). In the presently analyzed
486 taxa, this would support the hypothesis of the recent divergence. Due to their geographic origins, the
487 ambiguous placement of a few specimens during morphometric or genetic analyses also seems better
488 supported by this hypothesis than by hybridization (despite the latter being relatively frequent in
489 European ants, e.g. Steiner *et al.*, 2011; Seifert, 2018; 2019b).

490 In conclusion, the CSL and DQL *Colobopsis* clusters are considerable separate species in accordance
491 with the good practices of ants' alpha-taxonomy: all available sources of evidence suggest monophyly
492 with the exception of mtDNA, whose advisory role to infer species boundaries may be relatively
493 weak in comparison with nuclear genes or nuclear genes' expression products for the arguments given
494 by Seifert (2020). As a result, the formal naming of CSL and DQL *Colobopsis* species holds a key
495 informative value over their biology and life history traits. The type material of *Co. truncata*,
496 consisting of a single queen, could not be part of the morphometric or genetic analyses, but shows
497 very clearly the DQL pattern and its geographic origin is unambiguous (with the type locality at
498 mountains of Orero, near Genoa, in Italy's Liguria, placed in the middle of a highly investigated area
499 within the DQL *Colobopsis* geographic range and about 780 km away from the closest area inhabited
500 by CSL *Colobopsis*). The same arguments of safe chromatic identification apply for *Co. fuscipes*, and
501 in this case they are supplemented by an even stronger biogeographic argument. As a result, the
502 *Colobopsis* characterized by the CSL pattern is an undescribed species. Accordingly, a formal
503 description is provided below.




Figure 9. Above: distribution map of examined *Colobopsis* samples – countries where *Colobopsis* presence is known from literature are highlighted in grey. Below: approximate distributions of other Camponotini (*Camponotus barbaricus*, of *C. micans* and of *C. ruber*) which resemble that of CSL *Colobopsis*.

DESCRIPTION OF *COLOBOPSIS IMITANS* SP. NOV.


Etymology: imitans is the present participle of the latin verb imitor, meaning “imitating”, and is here used in apposition. It refers to the interpretation that this species resembles *Cr. scutellaris*.

Type series: 1 holotype worker (Figure 10) and 14 paratype workers from Mondello, Sicily (Italy), 38.1953, 13.3354, 5 m, 14.X.2018, E. Schifani leg. The holotype is stored in the Hungarian Natural History Museum collection.

Worker description: Morphometric indexes are shown in Tab. 2. Head subrectangular, on all sides rounded. A straight, central furrow runs from the frontal triangle to the level at which the frontal carinae end. Eyes large, ocelli extremely reduced. Antennae of 12 segments, without a distinct antennal club. Pronotum significantly wider than the rest of the mesosoma. In lateral profile, pronotum and mesonotum gently convex, propodeum profile often showing a central concavity thus

519 having a saddle-like appearance. Petiolar scale profile anteriorly roundly concave and posteriorly
520 straight, its dorsal crest excavated in frontal view. Promesonotal and mesoepinotal sutures as well as
521 metathoracic and propodeal spiracles well-visible. All legs with well-developed tibial spurs, but more
522 so in the anterior legs which are characterized by strikingly large femurs (identical to *Co. truncata*,
523 function unknown). Pigmentation as described in the CSL model. Very fine alveolate to areolate
524 sculpture covering the whole body and appendages. Few erect hairs near the posterior margin of the
525 vertex, between the frons and on the clypeus, and few others on the gaster tergites. See Figs 1, 8, 10,
526 13.

527 **Soldier (= phragmotic major worker) description:** Measurements (2 specimens from Sicily): CL
528 = 1327, 1419; CW = 1262, 1470; SL = 865, 942; ML = 1752, 1774; MW = 921, 1103; EW = 264,
529 331; EL = 409, 459; CS = 1294, 1444; CL/CW = 0.96, 1.05; SL/CS = 0.65, 0.67; ML/CS = 1.23,
530 1.35. Large phragmotic head with a cylindrical shape, and a flattened anterior part formed by the
531 mandibles, part of the clypeus and of the genae. In the distalmost half, it is characterized by a strong
532 areolate-rugose sculpture and a dense coverage of thick and short erect hairs. Rest of the shape,
533 sculpture and pigmentation generally similar to the worker but white dots or stripe on the first gastral
534 tergite sometimes very evident. See Fig. 10.




535

536 Figure 10. *Colobopsis imitans* sp. nov.: a,b,e) worker (holotypus), c,d,f,g) soldier (specimen from the type
 537 locality). Scale bars: 0.5 mm. Pictures also available on AntWeb.org database, specimen codes:
 538 ANTWEB1041481 and ANTWEB1041482.

539 **Queen description:** Measurements (3 specimens from Sicily): CL = 1437-1531; CW = 1281-1406;
 540 SL = 1156-1218; ML = 2687-3031; MW = 1281-1312; EW = 325-362; EL = 525-537; CS = 1359-
 541 1468; CL/CW = 1.08-1.11; SL/CS = 0.83-0.87; ML/CS = 1.93-2.22. Large phragmotic head very
 542 similar to the soldier not only in shape but also in size (despite larger body size), but well-developed
 543 ocelli, eyes much larger and much longer scapi. Immediately distinguishable by the larger, dorsally
 544 flatter mesosoma, which is largely unsculptured and shiny. Propodeum profile similar to the end of
 545 soldiers' propodeum. Head red as in the worker, but the mesosoma is brownish and the white dots or
 546 stripe on the first gaster tergite are/is evident. See Fig. 11.

547 **Male description:** Measurements (3 specimens from Sicily): CL = 875-1093; CW = 781-1000; SL
 548 = 937-1001; ML = 2281-2437; MW = 1062-1218; EW = 300-387; EL = 462-525; CS = 828-1046;
 549 CL/CW = 1.09-1.17; SL/CS = 0.92-1.20; ML/CS = 2.32-2.75. Small, subrectangular head with large
 550 ocelli and very large eyes protruding laterally. Toothless mandibles. Relatively large mesosoma,

551 propodeum more gently rounded than in queens or workers. Petiolar node very low and round.
552 Sculpture very weak, mesosoma shiny. Mandibles very hairy, other hairs on clypeus and gaster. Entire
553 body ferruginous or brownish, gaster blackish. See Fig. 11. Genitalia as in Fig. 12.



554
555 Figure 11. *Colobopsis imitans* sp. nov.: a,b,e,f) queen, c,d,g) male. Specimens from the type locality. Scale
556 bars: 0.5 mm. Pictures also available on AntWeb.org database, specimen codes: ANTWEB1041483 and
557 ANTWEB1041484.



558

559 Figure 12. Male genitalia of *Colobopsis imitans* sp. nov. in ventral and dorsal view, specimen from the type
560 locality. Scale bars = 0.25 mm.

561 **Diagnosis:** Generally easy to determine on the basis of worker material due to strong chromatic
562 differentiation from *Co. truncata* and allopatric distribution (although further investigation is required
563 for possible contact regions in southern Iberia and southern Italy). Some very small worker specimens
564 may appear almost completely black, therefore lacking the typical chromatic pattern, and workers
565 with entirely red mesosoma can seldom be observed. Tentative identifications of isolated soldier or
566 queen specimens should be much more cautious, although their chromatic appearance may sometimes
567 appear to be very explicit. The low number of males and the lack of particularly evident distinctive
568 characters from *Co. truncata* do not allow a safe species-level identification of this caste based on
569 morphology. In respect to workers, the morphometric linear discriminant function provided in the
570 results section should be helpful to determine dubious cases including decolored specimens. Finally,
571 DNA barcoding, which can be used for the same purpose and also be employed on the other castes,
572 shows a relatively low error rate but may present some risks due to the polyphyletic pattern that we
573 observed.

574 **Biological, ecological and phenological notes:** Relatively thermophilous, in Sicily occurring from
575 few meters above the sea level to at least 1015 m a.s.l., in Morocco ascending up to 1290 m a.s.l. and
576 in Spain so far known from coastal lowland areas. Probably very common but also heavily under-
577 recorded due to cryptic arboreal lifestyle, low colony population (most-likely monogynous), effective
578 mimicry and long periods of inactivity during the most arid and coldest times of the year. Workers
579 and soldiers are unlikely to descend to the ground but were observed to do so at least once, following
580 a sparsely populated *Cr. scutellaris* trail. Soldiers in general are very difficult to be found outside the
581 nest and usually seen acting as gatekeepers of the nest entrance. Founding queens were observed to
582 do the same. Nests are hidden in minute holes on the dead parts of arboreal trunks, where *Co. imitans*

583 sp. nov. often seems to act as a secondary user of cavities excavated by xylophagous insects. It
584 exploits *Andricus quercustozae* (Bosc, 1792) oak galls as nests (occupying about 15% of galls
585 collected in Sicily's Bosco della Ficuzza in a recent survey, authors' unpublished data), in a similar
586 way to *Co. truncata* (see Giannetti *et al.*, 2019; 2021; Fürjes-Mikó *et al.*, 2020). Polydomy appears
587 probable due to repeated findings of groups of workers without queens within oak-galls. Observed
588 nesting on several and diverse plant species, including at least: *Citrus reticulata* Blanco, 1837, *Ci.*
589 *sinensis* (L.) Osbeck, 1765, *Laurus nobilis* L., *Olea europaea* L., *Quercus ilex* L., *Q. pubescens*-
590 group, *Q. suber* L., *Pyrus communis* L., *Pittosporum tobira* (Thunb.) W.T.Aiton. Apparently dense
591 populations were found in old *Citrus* orchards and relatively sparse *Q. suber* woods, but also in
592 deciduous oak forests. However, it occurs in a broad range of habitats from cities to agricultural lands
593 to natural forest habitats, but information available so far is insufficient to depict a satisfactory picture
594 of habitat preferences. Despite the earlier claim by Carpintero *et al.* (2005), there is currently no
595 evidence backing the fascinating hypothesis that *C. imitans* sp. nov. foundress queens prefer trees
596 hosting *Cr. scutellaris* to found their colonies. A focused investigation on this topic would be
597 interesting. Nuptial flights for *Co. imitans* sp. nov. occur approximately in the same period of *Co.*
598 *truncata* (alates in Sicily observed from June 30 to July 13, n = 5, see supplementary material).
599 Winged queens and males were repeatedly seen attracted by artificial lights at night.

600 FINAL REMARKS

601 Body pigmentation pattern is the only qualitative character that makes *Co. imitans* sp. nov.
602 identifiable without recurring to quantitative data, as it is otherwise morphologically extremely
603 similar to *Co. truncata* up to a significant level of crypsis (see Wagner *et al.*, 2018). These
604 pigmentation differences among West-Palearctic *Colobopsis* so far went completely unnoticed, the
605 sole exception being a brief statement by Santschi (1929) noting that the chromatic aspect of the
606 Moroccan *Co. truncata* is different than the typus one by its lighter head color. The case we
607 documented can be considered one of the few where such element is important for species
608 discrimination in European ants. While body pigmentation has been used without scientific rigor by
609 some past ant taxonomists (see the example described by Boer, 2008), it can be important for the
610 morphological identification of species such as *Formica clara* Forel, 1886 and *F. cunicularia*
611 Latreille, 1798 or even fundamental for many *Temnothorax* spp. (Seifert & Schulz, 2009; Seifert,
612 2018) and should not be overlooked in multi-character approaches for taxonomic purposes. Under
613 these conditions, checking pictures uploaded on citizen science platforms and social media proved to
614 be significantly helpful to obtain data on these species distribution, evidencing once more the
615 uncovered potentials of citizen science in the study of ant distribution (e.g. Lucky *et al.*, 2014; Zhang

616 *et al.*, 2019; Castracani *et al.*, 2020; Sheard *et al.*, 2020) and more in general of platforms hosting
617 these kind of data in the study of insect distribution (e.g. Schifani & Paolinelli 2018; Hochmair *et al.*,
618 2020; Ruzzier *et al.*, 2020; Winterton, 2020). Moreover, behavioural data are seldom considered in
619 integrative taxonomic approaches dealing with ants, but they may prove valuable in some cases (see
620 also Ronque *et al.*, 2016). Finally, while mtDNA has a decent identification performance, our data
621 clearly support the idea that it should not be used as the primary source of information to take
622 taxonomic decision on species delimitation (see Seifert 2020).

623 The taxonomic status of the West-Palearctic *Colobopsis* populations appears now well-resolved. Still,
624 our analyses missed data from what the existing literature describes as the easternmost distribution
625 of *Co. truncata* east to the Mediterranean region, which reaches to the Kopet Dag in Turkmenistan
626 (Dlussky *et al.*, 1990; Gratiashvili & Barjadze, 2008; Dubovikoff & Yusupov, 2018; Bračko, 2019;
627 Samin *et al.*, 2020). In biogeographic terms, they are extremely unlikely to represent a disjunct *Co.*
628 *imitans* sp. nov. population, while conspecificity with *Co. truncata* appears likely due to the existence
629 of several ant species with similar distributions (e.g. Wagner *et al.*, 2017; Seifert, 2018). Within the
630 Mediterranean, the range limits of *Co. imitans* sp. nov. and *Co. truncata* or their eventual sympatry
631 in contact zones should be appropriately investigated in areas of biogeographic transition (southern
632 Iberia, Sicily, Calabria and perhaps Sardinia, see Alicata & Schifani, 2019; García, 2020; Schifani *et*
633 *al.*, 2020; 2021; Tinaut & Ruano, 2021).


634 The fact that *Co. imitans* sp. nov. and *Co. truncata* greatly differ chromatically is interesting if one
635 considers that phylogenetics and morphometry suggest a recent differentiation. In evolutionary terms,
636 the most likely interpretation is to link such differentiation to a shared strategy based on ant-mimicry
637 modulated according to the presence or absence of certain good models across different
638 Mediterranean regions. Both *D. quadripunctatus* and *Cr. scutellaris* have much more populous
639 colonies than *Co. imitans* sp. nov. and *Co. truncata*, while both are likely less palatable for predators
640 and armed with effective toxic substances (Cavill & Hinterberger, 1960; Wagner, 2019). Therefore,
641 even though only *Cr. scutellaris* is truly recognized as an aggressive and dominant species (Santini
642 *et al.*, 2007; Frizzi *et al.*, 2015; Castracani *et al.*, 2017; Seifert, 2018), both appear to possess the
643 required traits to be considered good Batesian models to the non-aggressive and relatively unarmed
644 *Colobopsis* (which still possess some formic acid). However, across the distribution range of *Co.*
645 *imitans* sp. nov., *D. quadripunctatus* is almost completely absent: it does not inhabit the Maghreb, its
646 Iberian distribution is concentrated to the North and in Sicily it is considered to be very rare (Schifani
647 & Alicata, 2018; Cabanillas *et al.*, 2019). Yet it is interesting to note that the opposite is not true for
648 *Co. truncata*: the latter is not only sympatric with *D. quadripunctatus* along its entire range (including
649 in the hypothesis that easternmost *Colobopsis* are *Co. truncata*: see Reznikova, 2003; Ghahari *et al.*,

2015), but also sympatric with *Cr. scutellaris* in south-western Europe and with *Cr. schmidtii* in the east. Finally, it is worth noting that the white dots or stripe that have been linked to mimicry of *D. quadripunctatus* in *Co. truncata* (Forel, 1886; Wagner, 2019) are/is absent or hardly visible in *Co. imitans* sp. nov. workers but at the same time well-visible in at least a few soldiers and especially queens that we inspected. Following the mimicry interpretation of the chromatic patterns, it is imaginable that this character is an ancestral remnant but that selective (predatory) pressures leading to perfect mimicry are stronger on workers than on queens or soldiers which rarely leave the safety of their nest.

Mimicry may be considered as a third defensive strategy of *Colobopsis* unique or very rare among ants after suicidal autothysis and phragmosis (Emery, 1925; Maschwitz & Maschwitz, 1974; Davidson *et al.*, 2012; Shorter & Rueppel, 2012; Ward *et al.*, 2016; Laciny *et al.*, 2018). Apart from the two species we treated, the only existing claims of mimicry in the genus come from morphologically very different and likely unrelated species from Fiji Islands (Santschi 1928; Wheeler 1934). However, since several other Palearctic *Colobopsis* species share a general morphological similarity with *Co. imitans* sp. nov. and *Co. truncata*, likely belonging to the same evolutionary lineage, it is possible that some of them represent yet undiscovered mimics. It also appears that mimicry may have played a powerful role driving phenotypic diversification of West-Palearctic *Colobopsis*: *Co. imitans* sp. nov. and *Co. truncata* can be considered as the only well-documented example among ants that suggests mimicry-driven phenotypic divergence of sister species. In comparison, the intraspecific case of mimicry pattern divergence in *Ca. lateralis* is one of much more modest differentiation (Wagner, 2014; Seifert, 2019a). Similar accounts are not particularly common in other organisms either, but recently an interesting scenario of strong diverging aposematic patterns coupled with minimal genetic differentiation was described by for a group of frogs (Tarvin *et al.*, 2017).

It is still unknown which visual predators may have been responsible for determining the selective pressures that lead to the emergence of ant-mimicry across different ant lineages. Birds and lizards appear to be good candidates to start with (Ito *et al.*, 2004; Wagner, 2014). Our surveys around the colonies and trails of *Cr. scutellaris* and *D. quadripunctatus* led us to find several possible mimics of either species that belong to different insect and spider groups already known for ant mimicry (Fig. 13). In particular, Santschi (1919) suggested in the the Canarian relative of *Ca. ruber* (*Ca. guanchus* Santschi, 1908) the existence of an association similar to that between *Ca. lateralis* and *Cr. scutellaris*, while Harvey *et al.* (2018) described the anti-predatory function of ant-mimicry in *Gelis* spp., Komatsu (1961) reported on *Phrurolithus*-ant associations, Corcobado *et al.* (2016) reported on *Leptorchestes*-ant associations, and finally Chandler (2010) mentions myrmecomorphism among

684 Anthicidae. Although some of these findings may have been coincidental and deserve further
685 investigation, it is imaginable that *Co. imitans* sp. nov. and *Co. truncata* are each part of a larger
686 cohort of different arthropods that evolved mimicry to resemble *Cr. scutellaris* or *D. quadripunctatus*
687 in response to visually hunting generalist insectivores, similarly to the “golden mimicry complex”
688 described by Pekár *et al.* (2017). Further investigation is also required to understand whether the
689 advantages of mimicry for *Co. imitans* sp. nov. and *Co. truncata* may lay in a dilution effect, if
690 Batesian mimicry is truly implied and if Müllerian mimicry also plays a role (see Speed, 1999; Pekár
691 *et al.*, 2017) –, keeping in mind that different evolutionary relations may exist between the same prey
692 and different predators.



693

694 Figure 13. Above *Cr. scutellaris* and species showing a very similar chromatic pattern that were collected near *Cr.*
 695 *scutellaris* trails in Sicily (Italy): a) *Co. imitans* sp. nov. (worker from Mondello), b) *Ca. lateralis* (worker from Monte
 696 Pellegrino), c) *Gelis* sp. (Hymenoptera: Braconidae) from Monte Petroso, d) *Cr. scutellaris* from Levanzo island, e)
 697 *Phrurolithus* sp. (Araneae: Phrurolitidae) from Mondello, f) *Ca. ruber* (worker from Monte Pellegrino) and g)
 698 *Leptorchestes* sp. (Araneae: Salticidae) from Monte Petroso. Below, *D. quadripunctatus* and species with a very similar
 699 chromatic pattern that were collected near its trails or in the same trees in the Italian Peninsula: h) *Co. truncata* (specimen

700 from Bulgaria, AntWeb code CASENT0280000, photographer Michele Esposito), i) *D. quadripunctatus* (specimen from
701 Czech Republic, AntWeb code CASENT0179916, photographer Michele Esposito), j) *Formicomus pedestris* (Rossi,
702 1790) (Coleoptera: Anthicidae) from Parma (Italy).

703 Interspecific but intrageneric trail-following described for some ants is likely relatable to
704 phylogenetic proximity and morphofunctional and behavioural similarities or similar foraging
705 strategies among species (e.g. Grasso *et al.*, 2002 and references therein). On the other hand, the
706 significance of the recorded *Colobopsis-Crematogaster* trail-following is not yet fully clear to us.
707 Similar cases (such as that of *Ca. lateralis*) have often been referred to as parabiosis, but we avoided
708 this term since quite different interpretations of its meaning coexist creating ambiguity: it is
709 sometimes used to simply indicate trail-following but in other occasions it implies also nest-sharing
710 (see Forel, 1898; Swain, 1980; Vantaux *et al.*, 2007; Menzel *et al.*, 2008; 2010; 2014a; 2014b; Seifert,
711 2018). Outside of *Co. imitans* sp. nov. and *Co. truncata*, many other camponotine ants follow
712 *Crematogaster* trails without always acting as mimics (Ito *et al.*, 2004; Vantaux *et al.*, 2007; Menzel
713 *et al.*, 2008; 2014). Baroni Urbani (1969) and Menzel *et al.* (2014) speculated that the compounds
714 used as trails pheromones by *Crematogaster* are generally easily perceived by *Camponotus*, this
715 capacity representing an important pre-adaptation to trail-following. During our field surveys, we
716 unexpectedly observed several workers of *Camponotus piceus* (Leach, 1825) (a relative of *Ca.*
717 *lateralis* with no resemblance of *Cr. scutellaris*, see Seifert, 2019a) easily following part of a *Cr.*
718 *scutellaris* trail to the canopy of a tangerine tree while avoiding *Cr. scutellaris* attacks. It may be
719 possible that many other similar camponotine ants rarely perform the same without possessing a
720 specific mimicry adaptation, which can partly explain the occasional observations of trail-following
721 between *Co. truncata* and *Cr. scutellaris* (Zimmermann 1934; Goetsch 1942; Baroni Urbani 1969;
722 Wagner 2014). Still, in the overwhelming majority of the documented cases inter-specific trail-
723 following is associated either with mimicry (as for *Colobopsis imitans* sp. nov., see Gobin *et al.*,
724 1998; Ito *et al.*, 2004; Menzel *et al.*, 2010; Powell *et al.*, 2014) or with nest-sharing (Vantaux *et al.*,
725 2007; Menzel *et al.*, 2008; 2014). For mimics, it appears to be primarily a way to better hide within
726 the ranks of the model species, an example of dilution effect (Lehtonen & Jaatinen, 2016), which may
727 also apply to a certain degree non-mimic ant species as well. However, the trail-followers may be
728 able to obtain additional benefits in their success of locating trophic resources, sometimes even
729 establishing somewhat parasitic relationships (see Vantaux *et al.* 2007; Menzel *et al.* 2010; 2014a;
730 2014b). It is unclear whether *Colobopsis* ants may also benefit from a similar mechanism although a
731 parasitic aspect of its trail-following behaviour has been suggested by Baroni Urbani (1969).

732 REFERENCES

733 Alicata A, Schifani E. 2019. Three endemic *Aphaenogaster* from the Siculo-Maltese archipelago and
734 the Italian Peninsula: part of a hitherto unrecognized species group from the Maghreb?
735 (Hymenoptera: Formicidae: Myrmicinae). *Acta Entomologica Musei Nationalis Pragae* 59: 1–16.

736 Andersen AN. 1995. A classification of Australian ant communities, based on functional groups
737 which parallel plant life-forms in relation to stress and disturbance. *Journal of Biogeography* 22: 1–
738 29.

739 Andersen A. 1997. Functional groups and patterns of organization in North American ant
740 communities: a comparison with Australia. *Journal of Biogeography* 24: 433–460.

741 Arnan X, Andersen AN, Gibb H, Parr CL, Sanders NJ, Dunn RR, Angulo E, Baccaro FB, Bishop TR,
742 Boulay R, Castracani C, Cerdá X, Del Toro I, Delsinne T, Donoso DA, Elten EK, Fayle TM,
743 Fitzpatrick MC, Gómez C, Grasso DA, Grossman BF, Guénard B, Gunawardene N, Heterick B,
744 Hoffmann BD, Janda M, Jenkins CN, Klimes P, Lach L, Laeger T, Leponce M, Lucky A, Majer J,
745 Menke S, Mezger D, Mori A, Moses J, Munyai TC, Paknia O, Pfeiffer M, Philpott SM, Souza JLP,
746 Tista M, Vasconcelos HL, Retana J. 2018. Dominance–diversity relationships in ant communities
747 differ with invasion. *Global Change Biology* 24: 4614–4625.

748 Avise JC. 2004. Molecular markers, natural history and evolution. Sinauer Associates Inc.,
749 Sunderland, MA, USA..

750 Barbero F, Bonelli S, Thomas JA, Balletto E, Schönrogge K. 2009. Acoustical mimicry in a predatory
751 social parasite of ants. *Journal of Experimental Biology* 212: 4084–4090.

752 Baroni Urbani C. 1969. Trail sharing between *Camponotus* and *Crematogaster*: some comments and
753 ideas. Proceedings of the VI Congress of the International Union for the Study of Social Insects
754 (IUSI), Bern, 15–20 September 1969.

755 Baroni Urbani C. 1971. Catalogo delle specie di Formicidae d'Italia (Studi sulla mirmecofauna d'Italia
756 X). *Memorie della Società Entomologica Italiana* 50: 5–287.

757 Beresford J, Elias M, Pluckrose L, Sundström L, Butlin RK, Pamilo P, Kulmuni J. 2017. Widespread
758 hybridization within mound-building wood ants in Southern Finland results in cytonuclear
759 mismatches and potential for sex-specific hybrid breakdown. *Molecular Ecology* 26: 4013–4026.

760 Blaimer BB. 2012. A subgeneric revision of *Crematogaster* and discussion of regional species-groups
761 (Hymenoptera: Formicidae). *Zootaxa* 3482: 47–67.

Blatrix R, Aubert C, Decaëns T, Berquier C, Andrei-Ruiz M, Galkowski C. 2020. Contribution of a DNA barcode to an assessment of the specificity of ant taxa (Hymenoptera: Formicidae) on Corsica. *European Journal of Entomology* 117: 420–429.

Boer P. 2008. *Plagiolepis obscuriscapa* Santschi, 1923, a junior synonym of *Plagiolepis pygmaea* (Latreille, 1798) (Hymenoptera: Formicidae: Formicinae) and the use of pigmentation as discriminating character in ant taxonomy. *Zoologische Mededelingen* 82: 485–488.

Bolton B. 2021. *An online catalog of the ants of the world*. Available from <https://antcat.org> (accessed on 03.01.2021).

Borowiec L. 2014. Catalogue of ants of Europe, the Mediterranean Basin and adjacent regions (Hymenoptera: Formicidae). *Genus* 25: 1–340.

Bračko G. 2019. New data on the ant fauna (Hymenoptera: Formicidae) of Azerbaijan. *Caucasian Entomological Bulletin* 15: 165–175.

Brun R. 1924. *Das Leben der Ameisen*. Leipzig, Berlin: Verlag und Druck von B. G. Teubner.

Buschinger A, Maschwitz U. 1984. Defensive behavior and defensive mechanisms in ants. In: Hermann HR, ed. *Defensive mechanisms in social insects*, Santa Barbara: Praeger Publishers.

Buschinger A. 2009. Social parasitism among ants: a review (Hymenoptera: Formicidae). *Myrmecological News* 12: 219–235.

Cabanillas D, Narro-Martín AJ, Fernández-Martínez JA. 2019 Ampliación de la distribución de *Dolichoderus quadripunctatus* (Linnaeus, 1771) (Formicidae, Dolichoderinae) en la Península Ibérica. *Iberomyrmex* 11: 9–12.

Cagniant H. 1968. Liste préliminaire de fourmis forestières d'Algérie. Résultats obtenus de 1963 à 1966. *Bulletin de la Societe d'Histoire Naturelle de Toulouse* 104: 138–147.

Cagniant H, Espadaler X. 1993. *Camponotus guanchus* Satschi, 1908, stat. nov. et étude des populations de *Camponotus sicheli* Mayr, 1866 (Hymenoptera: Formicidae). *Journal of African Zoology* 107: 419–438.

Cagniant H. 1996. Les *Camponotus* du Maroc (Hymenoptera: Formicidae): clé et catalogue des espèces. *Annales de la Société entomologique de France* 32: 87–100.

790 Carpintero S, Tinaut A, Reyes J, de Reyna LA. 2001. Estudio faunístico de los formícidos
791 (Hymenoptera, Formicidae) del Parque Nacional de Doñana. *Boletín de la Asociación Española de*
792 *Entomología* 25: 133–152.

793 Carpintero S, Reyes-López J, De Reyna LA. 2005. Impact of Argentine ants (*Linepithema humile*)
794 on an arboreal ant community in Donana National Park, Spain. *Biodiversity & Conservation* 14: 151–
795 163.

796 Cassis G, Wall MA. 2010. Systematics and phylogeny of the hatchet head plant bug genus
797 *Myrmecoroides* Gross (Insecta: Heteroptera: Miridae: Orthotylinae). *Entomologica Americana* 116:
798 29–49.

799 Castracani C, Bulgarini G, Giannetti D, Spotti FA, Maistrello L, Mori A, Grasso DA. 2017. Predatory
800 ability of the ant *Crematogaster scutellaris* on the brown marmorated stink bug *Halyomorpha halys*.
801 *Journal of Pest Science* 90: 1181–1190.

802 Castracani C, Spotti FA, Schifani E, Giannetti D, Ghizzoni M, Grasso DA, Mori A. 2020. Public
803 Engagement Provides First Insights on Po Plain Ant Communities and Reveals the Ubiquity of the
804 Cryptic Species *Tetramorium immigrans* (Hymenoptera, Formicidae). *Insects* 11: 678.

805 Cavill GWK, Hinterberger H. 1960. The chemistry of ants. IV. Terpenoid constituents of some
806 *Dolichoderus* and *Iridomyrmex* species. *Australian Journal of Chemistry* 13: 514–519.

807 Chan KM, Levin SA. 2005. Leaky prezygotic isolation and porous genomes: rapid introgression of
808 maternally inherited DNA. *Evolution* 59: 720–729.

809 Chandler DS. 2010: Anthicidae Latreille, 1819. In: Leschen RAB, Beutel RG, Lawrence JF, eds.
810 *Handbuch der Zoologie / Handbook of Zoology, Vol. IV (Arthropoda: Insecta), Part 38 Coleoptera,*
811 *Beetles. Volume 2: Morphology and systematics (Elateroidea, Bostrichiformia, Cucujiformia*
812 *partim.)*. Berlin: Walter de Gruyter.

813 Cobben RH. 1986. A Most strikingly myrmecomorphic Mirid from Africa, with some notes on ant-
814 mimicry and chromosomes in *Hallodapines* (Miridae, Heteroptera). *Journal of the New York*
815 *Entomological Society* 94: 194–204.

816 Collingwood CA, Yarrow, IHH. 1969. A survey of Iberian Formicidae. *Revista Española de*
817 *Entomología* 44: 53–101.

818 Corcobado G, Herberstein ME, Pekár, S. 2016. The role of ultraviolet colour in the assessment of
819 mimetic accuracy between Batesian mimics and their models: a case study using ant-mimicking
820 spiders. *The Science of Nature* 103: 90.

821 Csősz S, Heinze J, Mikó I. 2015. Taxonomic synopsis of the Ponto-Mediterranean ants of
822 *Temnothorax nylanderi* species-group. *PloS One* 10: e0140000.

823 Csősz S, Fisher BL. 2016: Taxonomic revision of the Malagasy members of the *Nesomyrmex*
824 *angulatus* species group using the automated morphological species delineation protocol NC-PART-
825 clustering. *PeerJ* 4: e1796.

826 Csősz S, Seifert B, Mikó I, Boudinot BE, Borowiec ML, Fisher BL, Prebus M, Puniamoorthy J,
827 Rakotonirina JC, Rasoamanana N, Schultz R, Trietsch C, Ulmer JM, Elek Z. 2020. Insect
828 Morphometry is Reproducible Under Average Investigation Standards. *Ecology and Evolution* 1–13.

829 Cushing PE. 2012. Spider-ant associations: an updated review of myrmecomorphy, myrmecophily,
830 and myrmecophagy in spiders. *Psyche* 1–23.

831 Darras H, Aron S. 2015. Introgression of mitochondrial DNA among lineages in a hybridogenetic
832 ant. *Biology letters* 11: 20140971.

833 Davidson DW, Salim KA, Billen J. 2012. Histology of structures used in territorial combat by
834 Borneo's 'exploding ants'. *Acta Zoologica* 93: 487–491.

835 de Haro A, Collingwood CA. 1992. Prospección mirmecológica por Extremadura (España) y Sao
836 Brás-Almodovar, Alcácer do Sal, Serra da Estrela (Portugal). *Boletim da Sociedade Portuguesa de*
837 *Entomologia* 3: 95–104.

838 de la Mora A, Sankovitz M, Purcell J. 2020. Ants (Hymenoptera: Formicidae) as host and intruder:
839 recent advances and future directions in the study of exploitative strategies. *Myrmecological News*
840 30: 53–71.

841 de L Nascimento FE, Perger R. 2018. Genus *Pseudolepturges* Gilmour (1957)(Coleoptera:
842 Cerambycidae: Lamiinae): a new species from Bolivia, key to the species of the genus and first reports
843 of a possible *Pseudomyrmex* ant mimic in longhorn beetles. *Journal of Natural History* 52: 1463–
844 1471.

845 Dlussky GM, Soyunov OS, Zabelin SI. 1990. *Ants of Turkmenistan*. Ashkabad: Ylym Press.

846 Dornhaus A, Powell S. 2010. Foraging and defence strategies. In: Lach L, Parr CL, Abbott K, eds.
847 *Ant ecology*. New York: Oxford University Press.

848 Dufour L, Perris E. 1840. Sur les insectes Hyménoptères qui nichent dans l'intérieur des tiges sèches
849 de la Ronce. *Annales de la Société Entomologique de France* 9: 5–53.

850 Dubovikoff DA, Yusupov ZM. 2018. Family Formicidae - Ants. In: Belokobylskij SA, Lelej AS, eds.
851 *Annotated catalogue of the Hymenoptera of Russia. Proceedings of the Zoological Institute of the*
852 *Russian Academy of Sciences* 6: 197–210.

853 Durkee CA, Weiss MR, Uma DB. 2011. Ant mimicry lessens predation on a North American jumping
854 spider by larger salticid spiders. *Environmental Entomology* 40: 1223–1231.

855 Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput.
856 *Nucleic Acids Research* 32: 1792–1797.

857 Emery C. 1925. Hymenoptera. Fam. Formicidae. Subfam. Formicinae. *Genera Insectorum* 183: 1–
858 302.

859 Emery C, Forel A. 1879. Catalogue des Formicides d'Europe. *Mitteilungen der Schweizerischen*
860 *Entomologischen Gesellschaft* 5: 441–481.

861 Emery C. 1880. Crociera del Violante, comando dal capitano armatore Enrico d'Albertis, durante
862 l'anno 1877. Formiche. *Annali del Museo Civico di Storia Naturale Giacomo Doria* 15: 389–398.

863 Emery C. 1886. Mimetismo e costumi parassitari del *Camponotus lateralis* Ol. *Bollettino della*
864 *Società Entomologica Italiana* 18: 412–13.

865 Emery C. 1916. Fauna Entomologica Italiana. I. Hymenoptera.-Formicidae. *Bollettino della Società*
866 *Entomologica Italiana* 47: 79–275.

867 Finzi B. Formiche della Libia. *Memorie della Società Entomologica Italiana* 18: 155–166.

868 Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. 1994. DNA primers for amplification of
869 mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular*
870 *Marine Biology and Biotechnology* 3: 294–299.

871 Forel A. 1874. Les fourmis de la Suisse. Systématique, notices anatomiques et physiologiques,
872 architecture, distribution géographique, nouvelles expériences et observations de mœurs. *Neue*
873 *Denkschriften der Allgemeinen Schweizerischen Gesellschaft für die Gesamten*
874 *Naturwissenschaften* 26: 1–452.

875 Forel A. 1886. Diagnoses provisoires de quelques espèces nouvelles de fourmis de Madagascar,
876 récoltées par M. Grandidier. *Annales de la Société Entomologique de Belgique* 30: 101–152.

- 877 Forel A. 1890. Fourmis de Tunisie et de l'Algérie orientale. *Annales de la Société Entomologique de*
878 *Belgique* 34: 61–76.
- 879 Forel A. 1898. La parabiose chez les fourmis. *Bulletin de la Société Vaudoise des Sciences Naturelles*
880 34: 380–384.
- 881 Forel A. 1903. Faune myrmécologique des noyers dans le canton de Vaud. *Bulletin de la Société*
882 *Vaudoise des Sciences Naturelles* 39: 83–94.
- 883 Forel A. 1905. Miscellanea myrmécologiques II. *Annales de la Société Entomologique de Belgique*
884 49: 155–185.
- 885 Forel A. 1909. Études myrmécologiques en 1909. Fourmis de Barbarie et de Ceylan. Nidification des
886 *Polyrhachis*. *Bulletin de la Société Vaudoise des Sciences Naturelles* 45: 369–407.
- 887 Frizzi F, Ciofi C, Dapporto L, Natali C, Chelazzi G, Turillazzi S, Santini G. 2015. The rules of
888 aggression: how genetic, chemical and spatial factors affect intercolony fights in a dominant species,
889 the mediterranean acrobat ant *Crematogaster scutellaris*. *PLoS One* 10: e0137919.
- 890 Funk DJ, Omland KE. 2003. Species-level paraphyly and polyphyly: frequency, causes, and
891 consequences, with insights from animal mitochondrial DNA. *Annual Review of Ecology, Evolution,*
892 *and Systematics* 34: 397–423.
- 893 Fürjes-Mikó F, Csősz S, Csóka G. 2020. Ants inhabiting oak Cynipid galls in Hungary. *North-*
894 *Western Journal of Zoology* 16: 95–98.
- 895 Galkowski C, Cagniant H. 2017. Contribution à la connaissance des fourmis du groupe *angustulus*
896 dans le genre *Temnothorax* (Hymenoptera, Formicidae). *Revue de l'Association Roussillonnaise*
897 *d'Entomologie* 26: 180–191.
- 898 Gallego-Roper MC, Feitosa RM. 2014. Evidences of batesian mimicry and parabiosis in ants of the
899 Brazilian Savanna. *Sociobiology* 61: 281–285.
- 900 García F. 2020. *Colobopsis truncata* (Spinola, 1808) en Galicia, NO Iberia. Estudio morfológico,
901 descripción da larva e distribución ibérica (Hymenoptera, Formicidae). *Archivos Entomológicos* 22:
902 401–416.
- 903 Geiselhardt SF, Peschke K, Nagel P. 2007. A review of myrmecophily in ant nest beetles (Coleoptera:
904 Carabidae: Paussinae): linking early observations with recent findings. *Naturwissenschaften* 94: 871–
905 894.

- 906 Ghahari H, Sharaf MR, Aldawood AS, Collingwood C. 2015. A contribution to the study of the ant
907 fauna (Hymenoptera: Formicidae) of Eastern Iran. *Contributions to Entomology* 65: 341–359.
- 908 Giannetti D, Castracani C, Spotti FA, Mori A, Grasso DA. 2019. Gall-Colonizing Ants and Their
909 Role as Plant Defenders: From 'Bad Job' to 'Useful Service'. *Insects* 10: 392.
- 910 Giannetti D, Mandrioli M, Schifani E, Castracani C, Spotti F, Mori A, Grasso DA. 2021. First report
911 of the acrobat ant *Crematogaster scutellaris* storing live aphids inside its oak-gall nests. *Insects* (in
912 press).
- 913 Gibb H, Dunn RR, Sanders NJ, Grossman BF, Photakis M, Abril S, Agosti D, Andersen AN, Angulo
914 E, Armecht I, Arnan X, Baccaro FB, Bishop TR, Boulay R, Brühl C, Castracani C, Cerda X, Del Toro
915 I, Delsinne T, Diaz M, Donoso DA, Ellison AM, Enriquez ML, Fayle TM, Feener DH Jr, Fisher BL,
916 Fisher RN, Fitzpatrick MC, Gómez C, Gotelli NJ, Gove A, Grasso DA, Groc S, Guénard B,
917 Gunawardene N, Heterick B, Hoffmann B, Janda M, Jenkins C, Kaspari M, Klimes P, Lach L, Laeger
918 T, Lattke J, Leponce M, Lessard JP, Longino J, Lucky A, Luke SH, Majer J, McGlynn TP, Menke S,
919 Mezger D, Mori A, Moses J, Munyai TC, Pacheco R, Paknia O, Pearce-Duvet J, Pfeiffer M, Philpott
920 SM, Resasco J, Retana J, Silva RR, Sorger MD, Souza J, Suarez A, Tista M, Vasconcelos HL,
921 Vonshak M, Weiser MD, Yates M, Parr CL. 2017. A global database of ant species abundances.
922 *Ecology* 98: 883–884.
- 923 Glaser F. 2009. Ameisen. Naturkundliche Forschung im Fürstentum Liechtenstein, Band 6.
924 Liechtenstein: Amt für Umwelt.
- 925 Gnezdilov VM. 2019. A new species of the myrmecomorphic planthopper genus *Formiscurra*
926 (Fulgoroidea: Caliscelidae) from Ethiopia. *Acta Entomologica Musei Nationalis Pragae* 59: 17–22.
- 927 Gobin B, Peeters C, Billen J, Morgan ED. 1998. Interspecific trail-following and commensalism
928 between the ponerine ant *Gnamptogenys menadensis* and the formicine ant *Polyrhachis rufipes*.
929 *Journal of Insect Behavior* 11: 361–369.
- 930 Goetsch W. 1942. Beiträge zur Biologie spanischer Ameisen. *Revista Española de Entomología* 18:
931 175–241.
- 932 Goetsch W. 1950. Beiträge zur Biologie und Verbreitung der Ameisen in Kärnten und in den
933 Nachbargebieten. *Österreichische Zoologische Zeitschrift* 2: 39–69.
- 934 Goetsch W. 1951. Ameisen- und Termiten-Studien in Ischia, Capri und Neapel. *Zoologische*
935 *Jahrbücher Abteilung für Systematik Ökologie und Geographie der Tiere* 80: 64–98.

936 Goetsch W. 1953. Die Staaten der Ameisen. Berlin, Göttingen, Heidelberg: Springer-Verlag.

937 Grasso DA, Mori A, Le Moli F. 1998. Chemical communication during foraging in the harvesting
938 ant *Messor capitatus* (Hymenoptera, Formicidae). *Insectes Sociaux* 45: 85–96.

939 Grasso DA, Mori A, Le Moli F. 1999. Recruitment and trail communication in two species of *Messor*
940 ants (Hymenoptera, Formicidae). *Italian Journal of Zoology* 66: 373–378.

941 Grasso DA, Mori A, Le Moli F. 2002. Behavioural investigation of trail signals specificity in three
942 sympatric species of *Messor* ants (Hymenoptera, Formicidae). *Italian Journal of Zoology* 69: 147–
943 151.

944 Grasso DA, Sledge MF, Le Moli F, Mori A, Turillazzi S. 2005. Nest-area marking with faeces: A
945 chemical signature that allows colony-level recognition in seed harvesting ants (Hymenoptera,
946 Formicidae). *Insectes Sociaux* 52: 36–44.

947 Grasso DA, Giannetti D, Castracani C, Spotti FA, Mori A. 2020. Rolling away: a novel context-
948 dependent escape behavior discovered in ants. *Scientific Report* 10: 3784.

949 Gratiashvili N, Barjadze S. 2008. Checklist of the ants (Formicidae Latreille, 1809) of Georgia.
950 *Proceedings of the Institute of Zoology* 23: 130–146.

951 Guénard B, Weiser M, Gomez K, Narula N, Economo EP. 2017. The Global Ant Biodiversity
952 Informatics (GABI) database: a synthesis of ant species geographic distributions. *Myrmecological*
953 *News* 24: 83–89.

954 Harvey JA, Visser B, Lammers M, Marien J, Gershenzon J, Ode PJ, Heinen R, Gols R, Ellers J. 2018.
955 Ant-like traits in wingless parasitoids repel attack from wolf spiders. *Journal of Chemical Ecology*
956 44: 894–904.

957 Hebert PD, Cywinska A, Ball SL, deWaard JR. 2003. Biological identifications through DNA
958 barcodes. *Proceedings of the Royal Society of London, Series B: Biological Sciences* 270: 313–321.

959 Hebert PD, Hollingsworth PM, Hajibabaei M. 2016. From writing to reading the encyclopedia of life.
960 *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 371: 20150321.

961 Helms JA, Peeters C, Fisher BL. 2014. Funnels, gas exchange and cliff jumping: natural history of
962 the cliff dwelling ant *Malagidris sofina*. *Insectes Sociaux* 61, 357–365.

963 Hochmair HH, Scheffrahn RH, Basille M, Boone M. 2020. Evaluating the data quality of iNaturalist
964 termite records. *PLoS One* 15: e0226534.

965 Hölldobler B, Wilson EO. 1990. *The ants*. Cambridge: Harvard University Press.

966 Hölldobler B, Wilson EO. 2008. *The Superorganism: The Beauty, Elegance and Strangeness of Insect*
967 *Societies*. Cambridge: Harvard University Press.

968 Huang JN, Cheng RC, Li D, Tso IM. 2011. Salticid predation as one potential driving force of ant
969 mimicry in jumping spiders. *Proceedings of the Royal Society B: Biological Sciences* 278: 1356–
970 1364.

971 Ito F, Hashim R, Huei YS, Kaufmann E, Akino T, Billen J. 2004. Spectacular Batesian mimicry in
972 ants. *Naturwissenschaften* 91: 481–484.

973 Janicki J, Narula N, Ziegler M, Guénard B, Economo EP. 2016. Visualizing and interacting with
974 large-volume biodiversity data using client–server web-mapping applications: The design and
975 implementation of antmaps. org. *Ecological Informatics* 32: 185–193.

976 Jackson JF, Drummond BA. 1974. A Batesian Ant-Mimicry Complex from the Mountain Pine Ridge
977 of British Honduras, with an Example of Transformational Mimicry. *American Midland Naturalist*
978 91: 248–251.

979 Kaudewitz F. 1955. Zum Gastverhältnis zwischen *Crematogaster scutellaris* Ol. mit *Camponotus*
980 *lateralis bicolor* Ol. *Biologisches Zentralblatt* 74: 69–87.

981 Kikuchi DW, Pfennig DW. 2010. High-model abundance may permit the gradual evolution of
982 Batesian mimicry: an experimental test. *Proceedings of the Royal Society B: Biological Sciences* 277:
983 1041–1048.

984 Komatsu T. 1961. Notes on spiders and ants. *Acta Arachnologica* 17: 25–27.

985 Laciny A, Zettel H, Kopchinskiy A, Pretzer C, Pal A, Salim KA, Rahimi MJ, Hoenigsberger M, Lim
986 L, Jaitrong W, Druzhinina IS. 2018. *Colobopsis explodens* sp. nov., model species for studies on
987 “exploding ants” (Hymenoptera, Formicidae), with biological notes and first illustrations of males of
988 the *Colobopsis cylindrica* group. *ZooKeys* 751: 1–40.

989 Larabee FJ, Suarez AV. 2015. Mandible-powered escape jumps in trap-jaw ants increase survival
990 rates during predator-prey encounters. *PLoS One* 10: e0124871.

991 Larsson A. 2014. AliView: a fast and lightweight alignment viewer and editor for large data sets.
992 *Bioinformatics* 30: 3276–3278.

993 Lehtonen J, Jaatinen K. 2016. Safety in numbers: the dilution effect and other drivers of group life in
994 the face of danger. *Behavioral Ecology and Sociobiology* 70: 449–458.

995 Lebas C, Galkowski C, Blatrix R, Wegnez P. 2016. *Fourmis d'Europe Occidentale. Le premier guide*
996 *complet d'Europe*. Paris: Delachaux et Niestlé.

997 Lleonart J, Salat J, Torres GJ. 2000. Removing allometric effects of body size in morphological
998 analysis. *Journal of Theoretical Biology* 205: 85–93.

999 Lucky A, Savage AM, Nichols LM, Castracani C, Shell L, Grasso DA, Mori A, Dunn RR. 2014.
1000 Ecologists, educators, and writers collaborate with the public to assess backyard diversity in The
1001 School of Ants Project. *Ecosphere* 5: 1–23.

1002 Mayr G. 1853. Beschreibungen einiger neuer Ameisen. *Verhandlungen der Zoologisch-Botanischen*
1003 *Vereins in Wien* 3: 277–286.

1004 Maschwitz U, Maschwitz E. 1974. Platzen der Arbeiterinnen: eine neue Art der Feindabwehr bei
1005 sozialen Hautflüglern. *Oecologia* 14: 289–294.

1006 McIver JD. 1987. On the myrmecomorph *Coquillettia insignis* Uhler (Hemiptera: Miridae): arthropod
1007 predators as operators in an ant-mimetic system. *Zoological Journal of the Linnean Society* 90: 133–
1008 144.

1009 McIver JD, Stonedahl G. 1993. Myrmecomorphy: Morphological and Behavioral Mimicry of Ants.
1010 *Annual Review of Entomology* 38: 351–377.

1011 Menozzi C. 1940. Contributo alla fauna della Tripolitania. *Bollettino del Laboratorio di Zoologia*
1012 *Generale e del Regio Istituto Superiore Agrario in Portici* 31: 244–273.

1013 Menzel F, Linsenmair KE, Blüthgen N. 2008. Selective interspecific tolerance in tropical
1014 *Crematogaster*–*Camponotus* associations. *Animal Behavior* 75: 837–846.

1015 Menzel F, Woywod M, Blüthgen N, Schmitt T. 2010. Behavioral and chemical mechanisms behind
1016 a Mediterranean ant–ant association. *Ecological Entomology* 35: 711–720.

1017 Menzel F, Orivel J, Kaltenpoth M, Schmitt T. 2014a. What makes you a potential partner? Insights
1018 from convergently evolved ant–ant symbioses. *Chemoecology* 24: 105–119.

1019 Menzel F, Kriesell H, Witte V. 2014b. Parabiocotic ants: the costs and benefits of symbiosis. *Ecological*
1020 *Entomology* 39: 436–444.

1021 Merrill DN, Elgar MA. 2000. Red legs and golden gasters: Batesian mimicry in Australian ants.
1022 *Naturwissenschaften* 87: 212–215.

1023 Müller F. 1879. *Ituna* und *Thyridia*: Ein merkwürdiges Beispiel von Mimicry bei Schmetterlingen.
 1024 *Kosmos* 5: 100–108.

1025 Nilsen G, Lingjaerde OC. 2013. clusterGenomics: Identifying clusters in genomics data by recursive
 1026 partitioning. R package version 1.0. <http://CRAN.R-project.org/package=clusterGenomics>.

1027 Oliveira PS, Sazima I. 1984. The adaptive bases of ant-mimicry in a neotropical aphantochilid spider
 1028 (Araneae: Aphantochilidae). *Biological Journal of the Linnean Society* 22: 145–155.

1029 Oliveira PS. 1988. Ant-mimicry in some Brazilian salticid and clubionid spiders (Araneae: Salticidae,
 1030 Glubionidae). *Biological Journal of the Linnean Society* 33: 1–15.

1031 Parker J, Grimaldi DA. 2014. Specialized Myrmecophily at the Ecological Dawn of Modern Ants.
 1032 *Current Biology* 24: 2428–2434.

1033 Parker J. 2016. Myrmecophily in beetles (Coleoptera): evolutionary patterns and biological
 1034 mechanisms. *Myrmecological news* 22: 65–108.

1035 Pekár S. 2014. Is inaccurate mimicry ancestral to accurate in myrmecomorphic spiders (Araneae)?.
 1036 *Biological Journal of the Linnean Society* 113: 97–111.

1037 Pekár S, Petráková L, Bulbert MW, Whiting MJ, Herberstein ME. 2017. The golden mimicry
 1038 complex uses a wide spectrum of defence to deter a community of predators. *Elife* 6: e22089.

1039 Pasteur G. 1982. A classificatory review of mimicry systems. *Annual Review of Ecology and*
 1040 *Systematics* 13: 169–199.

1041 Powell S, Del-Claro K, Feitosa RM, Brandao CRF. 2014. Mimicry and eavesdropping enable a new
 1042 form of social parasitism in ants. *American Naturalist* 184: 500–509.

1043 Prestwich GD. 1984. Defense mechanisms of termites. *Annual review of entomology* 29: 201–232.

1044 R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for
 1045 Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.

1046 Rasoamanana N, Csősz S, Fisher BL. 2017. Taxonomic revision of imitating carpenter ants,
 1047 *Camponotus* subgenus *Myrmopytia* (Hymenoptera, Formicidae) of Madagascar, using morphometry
 1048 and qualitative traits. *ZooKeys* 681: 119–152.

1049 Ratnasingham S, Hebert PD. 2007. Bold: the barcode of life data system
 1050 (<http://www.barcodinglife.org>). *Molecular Ecology Notes* 7: 355–364.

1051 Reznikova ZI. 2003. Distribution patterns of ants in different natural zones and landscapes in
 1052 Kazakhstan and West Siberia along a meridian trend. *Euroasian Entomological Journal* 2: 235–342.

1053 Ritland DB. 1991. Revising a classic butterfly mimicry scenario: demonstration of Müllerian mimicry
 1054 between Florida viceroys (*Limenitis archippus floridensis*) and queens (*Danaus gilippus berenice*).
 1055 *Evolution* 45: 918–934.

1056 Ronque MU, Azevedo-Silva M, Mori GM, Souza AP, Oliveira PS. 2016. Three ways to distinguish
 1057 species: using behavioural, ecological, and molecular data to tell apart two closely related ants,
 1058 *Camponotus renggeri* and *Camponotus rufipes* (Hymenoptera: Formicidae). *Zoological Journal of*
 1059 *the Linnean Society* 176: 170–181.

1060 Ross HA. 2014. The incidence of species-level paraphyly in animals: a re-assessment. *Molecular*
 1061 *Phylogenetics and Evolution* 76: 10–17.

1062 Ruzzier E, Menchetti M, Bortolotti L, Selis M, Monterastelli E, Forbicioni L. 2020. Updated
 1063 distribution of the invasive *Megachile sculpturalis* (Hymenoptera: Megachilidae) in Italy and its first
 1064 record on a Mediterranean island. *Biodiversity Data Journal* 8: e57783.

1065 Salata S, Borowiec L, Trichas A. 2020. Review of ants (Hymenoptera: Formicidae) of Crete, with
 1066 keys to species determination and zoogeographical remarks. *Monographs of the Upper Silesian*
 1067 *Museum* 12: 5–296.

1068 Samin N, Yusupov Z, Navaeian M, Sakenin H. 2020. A contribution to ants (Hymenoptera:
 1069 Formicidae) from North and Northwestern regions of Iran. *Natura Somogyiensis* 35: 29–36.

1070 Santini G, Tucci L, Ottonetti L, Frizzi F. 2007. Competition trade-offs in the organisation of a
 1071 Mediterranean ant assemblage. *Ecological Entomology* 32: 319–326.

1072 Santschi F. 1919. Fourmis d'Espagne et des Canaries. *Boletín de la Real Sociedad Española de*
 1073 *Historia Natural* 19: 241–248.

1074 Santschi F. 1925. Fourmis d'Espagne et autres espèces paléartiques. *Revista Española de*
 1075 *Entomología* 1: 339–360.

1076 Santschi F. 1928. Fourmis de îles Fidji. *Revue Suisse de Zoologie* 35: 67–74.

1077 Santschi F. 1929. Fourmis du Maroc, d'Algerie et de Tunisie. *Annales de la Société Entomologique*
 1078 *de Belgique* 69: 138–165.

1079 Scarparo G, d’Ettorre P, Di Giulio A. 2019. Chemical Deception and Structural Adaptation in
1080 *Microdon* (Diptera, Syrphidae, Microdontinae), a Genus of Hoverflies Parasitic on Social Insects.
1081 *Journal of Chemical Ecology* 45: 959–971.

1082 Schär S, Talavera G, Espadaler X, Rana JD, Andersen Andersen A, Cover SP, Vila R. 2018. Do
1083 Holarctic ant species exist? Trans-Beringian dispersal and homoplasy in the Formicidae. *Journal of*
1084 *Biogeography* 45: 1917–1928.

1085 Schär S, Menchetti M, Schifani E, Hinojosa JC, Platania L, Dapporto L, Vila R. 2020. Integrative
1086 biodiversity inventory of ants from a Sicilian archipelago reveals high diversity on young volcanic
1087 islands (Hymenoptera: Formicidae). *Organisms Diversity & Evolution* 20: 405–416.

1088 Schembri S, Collingwood CA. 1995. The myrmecofauna of the Maltese Islands. Remarks and
1089 additions (Hymenoptera Formicidae). *Bollettino della Società Entomologica Italiana* 127: 153–158.

1090 Schifani E, Alicata A. 2018. Exploring the myrmecofauna of Sicily: thirty-two new ant species
1091 recorded, including six new to Italy and many new aliens (Hymenoptera, Formicidae). *Polish Journal*
1092 *of Entomology* 87: 323–348.

1093 Schifani E, Paolinelli R. 2018. Forums and social media help to discover exotic species in Europe
1094 and monitor their spread the case of *Exaireta spinigera* (Wiedemann, 1830) (Diptera, Stratiomyidae)
1095 in the Italian peninsula and Sicily. *Graellsia* 74: e079.

1096 Schifani E, Scupola A, Alicata A. 2020. Morphology, ecology and biogeography of *Myrmecina sicula*
1097 André, 1882, rediscovered after 140 years (Hymenoptera, Formicidae). *Biogeographia* 35: 105–116.

1098 Schifani E, Nalini E, Gentile V, Alamanni F, Ancona C, Caria M, Cillo D, Bazzato E. 2021. Ants of
1099 Sardinia: an updated checklist based on new faunistic, morphological and biogeographical notes.
1100 *Redia* (in press).

1101 Schlick-Steiner BC, Steiner FM, Seifert B., Stauffer C, Christian E, Crozier RH. 2010. Integrative
1102 Taxonomy: A Multisource Approach to Exploring Biodiversity. *Annual Review of Entomology* 55:
1103 421–438.

1104 Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis.
1105 *Nature Methods* 9: 671–675.

1106 Scupola A. 2018. Le formiche del Veneto, The Ants of Veneto. Verona: World Biodiversity
1107 Association.

1108 Seifert B. 2008. Removal of allometric variance improves species separation in multi-character
1109 discriminant functions when species are strongly allometric and exposes diagnostic characters.
1110 *Myrmecological News* 11: 91–105.

1111 Seifert B. 2009. Cryptic species in ants (Hymenoptera: Formicidae) revisited: we need a change in
1112 the alpha-taxonomic approach. *Myrmecological News* 12: 149–166.

1113 Seifert B, Schultz R. 2009. A taxonomic revision of the *Formica rufibarbis* Fabricius, 1793 group
1114 (Hymenoptera: Formicidae). *Myrmecological News* 12: 255–272.

1115 Seifert B, Ritz M, Csősz S. 2014. Application of exploratory data analyses opens a new perspective
1116 in morphology-based alpha-taxonomy of eusocial organisms. *Myrmecological News* 19: 1–15.

1117 Seifert B. 2017. The ecology of Central European non-arboreal ants – 37 years of a broad-spectrum
1118 analysis under permanent taxonomic control. *Soil Organisms* 89: 1–67.

1119 Seifert B, d'Eustacchio D, Kaufmann B, Centorame M, Lorite P, Modica M. 2017. Four species within
1120 the supercolonial ants of the *Tapinoma nigerrimum* complex revealed by integrative taxonomy
1121 (Hymenoptera: Formicidae). *Myrmecological News* 24: 123–144.

1122 Seifert B. 2018. *The ants of Central and North Europe*. Tauer: Lutra Verlags-und
1123 Vertriebsgesellschaft.

1124 Seifert B. 2019a. A taxonomic revision of the members of the *Camponotus lateralis* species group
1125 (Hymenoptera: Formicidae) from Europe, Asia Minor and Caucasia. *Soil Organisms* 91: 7–32.

1126 Seifert B. 2019b. Hybridization in the European carpenter ants *Camponotus herculeanus* and *C.*
1127 *ligniperda* (Hymenoptera: Formicidae). *Insectes Sociaux*, 66: 365–374.

1128 Seifert B. 2020. The Gene and Gene Expression (GAGE) species concept—an universal approach for
1129 all eukaryotic organisms. *Systematic Biology* 69: 1033–1038.

1130 Sheard JK, Sanders NJ, Gundlach C, Schär S, Larsen RS. 2020. Monitoring the influx of new species
1131 through citizen science: the first introduced ant in Denmark. *PeerJ* 8: e8850.

1132 Shorter JR, Rueppell O. 2012. A review on self-destructive defense behaviors in social insects.
1133 *Insectes Sociaux* 59: 1–10.

1134 Speed MP. 1999. Batesian, quasi-Batesian or Müllerian mimicry? Theory and data in mimicry
1135 research. *Evolutionary Ecology* 13: 755–776.

- 1136 Spinola M. 1808. *Insectorum Liguria species novae aut rariores, quae in agro ligustico nuper*
 1137 *detexit, descripsit et iconibus illustravit Maximilianus Spinola, adjecto catalogo specierum*
 1138 *auctoribus jam enumeratarum, quae in eadam regione passim occurrent. Tom. II. Fasc. 4.* Genova:
 1139 Y. Gravier.
- 1140 Steiner FM, Schlick-Steiner BC, Sanetra M, Ljubomirov T, Antonova V, Christian E, Stauffer C.
 1141 2005. Towards DNA-aided biogeography: an example from *Tetramorium* ants (Hymenoptera,
 1142 Formicidae). *Annales Zoologici Fennici* 42: 23–25.
- 1143 Steiner FM, Seifert B, Grasso DA, Le Moli F, Arthofer W, Stauffer C, Crozier RH, Schlick-Steiner,
 1144 BC. 2011. Mixed colonies and hybridisation of *Messor* harvester ant species (Hymenoptera:
 1145 Formicidae). *Organisms Diversity & Evolution* 11: 107–134.
- 1146 Steiner FM, Csősz S, Markó B, Gamisch A, Rinnhofer L, Folterbauer C, Hammerle S, Stauffer C,
 1147 Arthofer W, Schlick-Steiner BC. 2018. Turning one into five: integrative taxonomy uncovers
 1148 complex evolution of cryptic species in the harvester ant *Messor* “*structor*”. *Molecular Phylogenetics*
 1149 *and Evolution* 127: 387–404.
- 1150 Stucky BJ. 2012. SeqTrace: A Graphical Tool for Rapidly Processing DNA Sequencing
 1151 Chromatograms. *Journal of Biomolecular Techniques* 23: 90–93.
- 1152 Swain RB. 1980. Trophic competition among parabiotic ants. *Insectes Sociaux* 27: 377–390.
- 1153 Tarvin RD, Powell EA, Santos JC, Ron SR, Cannatella DC. 2017. The birth of aposematism: High
 1154 phenotypic divergence and low genetic diversity in a young clade of poison frogs. *Molecular*
 1155 *Phylogenetics and Evolution* 109, 283–295.
- 1156 Tăușan I, Pintilioaie A, Milea D, Zachy M, Țicu S. 2020. First record of *Colobopsis truncata*
 1157 (Hymenoptera: Formicidae) from Moldova region of Romania. *Travaux du Muséum National*
 1158 *d’Histoire Naturelle “Grigore Antipa”* 63: 169–173.
- 1159 Tautz J, Hölldobler B, Danker T. 1994. Te ants that jump: diferent techniques to take off. *Zoology-*
 1160 *Analysis of Complex Systems* 98: 1–6.
- 1161 Tinaut A. 1991. Contribución al conocimiento de los formícidos del Parque Nacional de Doñana
 1162 (Hymenoptera, Formicidae). *Boletín de la Asociación Española de Entomología* 15: 57–63.
- 1163 Tinaut A., Ruano F. Biogeography of Iberian Ants (Hymenoptera: Formicidae). *Diversity* 13: 88.
- 1164 Trjapitzin VA, Trjapitzin SV. 1995. A new species of the genus *Coelaspidia* Timberlake 1923
 1165 (Insecta Hymenoptera Encyrtidae) from Cuba. *Tropical Zoology* 8: 341–346.

1166 Trifinopoulos J, Nguyen LT, von Haeseler A, Minh BQ. 2016. W-IQ-TREE: a fast online
1167 phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Research* 44: 232–235.

1168 Vantaux A, Dejean A, Dor A, Orivel J. 2007. Parasitism versus mutualism in the ant-garden
1169 parabiosis between *Camponotus femoratus* and *Crematogaster levior*. *Insectes Sociaux* 54: 95–99.

1170 Visicchio R, Mori A, Grasso DA, Castracani C, Le Moli F. 2001. Glandular sources of recruitment,
1171 trail, and propaganda semiochemicals in the slave-making ant *Polyergus rufescens*. *Ethology Ecology*
1172 *and Evolution* 13: 361–372.

1173 Wagner HC. 2014. Die Ameisen Kärntens. Verbreitung, Biologie, Ökologie und Gefährdung.
1174 Klagenfurt am Wörthersee: Naturwissenschaftlicher Verein für Kärnten.

1175 Wagner HC, Arthofer W, Seifert B, Muster C, Steiner FM, Schlick-Steiner BC. 2017. Light at the
1176 end of the tunnel: Integrative taxonomy delimits cryptic species in the *Tetramorium caespitum*
1177 complex (Hymenoptera: Formicidae). *Myrmecological News* 25: 95–129.

1178 Wagner HC, Gamisch A, Arthofer W, Moder K, Steiner FM, Schlick-Steiner BC. 2018. Evolution of
1179 morphological crypsis in the *Tetramorium caespitum* ant species complex (Hymenoptera:
1180 Formicidae). *Scientific Reports* 8: 1–10.

1181 Wagner HC. 2019. Wiener Ameisenbeobachtungen (Hymenoptera: Formicidae). *Beiträge zur*
1182 *Entomofaunistik* 20: 143–159.

1183 Ward PS. 2009. The ant genus *Tetraaponera* in the Afrotropical Region: the *T. grandidieri* group
1184 (Hymenoptera: Formicidae). *Journal of Hymenopteran Research* 18: 285–304.

1185 Ward PS, Blaimer BB, Fisher BL. 2016. A revised phylogenetic classification of the ant subfamily
1186 Formicinae (Hymenoptera: Formicidae), with resurrection of the genera *Colobopsis* and
1187 *Dinomyrmex*. *Zootaxa* 4072: 343–357.

1188 Wheeler WM. 1904. The American ants of the subgenus *Colobopsis*. *Bulletin of the American*
1189 *Museum of Natural History* 20: 139–158.

1190 Wheeler WM. 1934. Some aberrant species of *Camponotus* (*Colobopsis*) from the Fiji Islands. *Annals*
1191 *of the Entomological Society of America* 27: 415–424.

1192 Willis SC, Farias IP, Ortí G. 2014. Testing mitochondrial capture and deep coalescence in Amazonian
1193 cichlid fishes (Cichlidae: *Cichla*). *Evolution* 68: 256–268.

1194 Winterton SL. 2020. A new bee-mimicking stiletto fly (Therevidae) from China discovered on
1195 iNaturalist. *Zootaxa* 4816: 361–369.

- 1196 Wolak ME, Fairbairn DJ, Paulsen YR. 2012. Guidelines for Estimating Repeatability. *Methods in*
1197 *Ecology and Evolution* 3: 129–137.
- 1198 Zhang YM, Vitone T, Storer CG, Payton AC, Dunn RR, Hulcr J, McDaniel SF, Lucky A. 2019. From
1199 Pavement to Population Genomics: Characterizing a Long-Established Non-native Ant in North
1200 America Through Citizen Science and ddRADseq. *Frontiers in Ecology and Evolution* 7: 453.
- 1201 Zimmermann S. 1934. Beitrag zur Kenntnis der Ameisenfauna Süddalmatiens. *Verhandlungen der*
1202 *Zoologisch-Botanischen Gesellschaft in Wien* 84: 1–65.