

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

The impact of psoriasis on quality of life of children and their caregivers: an Italian experience

This is the submitted version (pre peer-review, preprint) of the following publication:

Published Version:

Bardazzi F., A Chessa M., Patrizi A., Iommi M., Rucci P., M Ali F., et al. (2021). The impact of psoriasis on quality of life of children and their caregivers: an Italian experience. EUROPEAN JOURNAL OF DERMATOLOGY, 31(5), 649-650 [10.1684/ejd.2021.4127].

Availability:

This version is available at: <https://hdl.handle.net/11585/898892> since: 2022-11-02

Published:

DOI: <http://doi.org/10.1684/ejd.2021.4127>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the pre-print version of:

Bardazzi F, A Chessa M, Patrizi A, Iommi M, Rucci P, M Ali F, Filippi F, Tengattini V, Neri I.

The impact of psoriasis on quality of life of children and their caregivers: an Italian experience.

Eur J Dermatol. 2021 Oct 1;31(5):649-650.

The final published version is available online at:

<https://doi.org/10.1684/ejd.2021.4127>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Title: Impact of psoriasis on quality of life of children and their caregivers: an Italian experience

Authors: F. Bardazzi MD^{1*}, MA Chessa MD^{1*}, A. Patrizi Prof.¹, M. Iommi PhD Student², P. Rucci PhD², F.M. Ali MD³, F. Filippi MD, V Tengattini MD^{1*}, I. Neri MD^{1*}.

Affiliation:

***These authors have equally contributed.**

¹ Department of Specialized, Diagnostic and Experimental Medicine, Division of Dermatology, University of Bologna, Italy.

² Department of Biomedical and Neuromotor Sciences, University of Bologna, Italy

³ Institute of Infection and Immunity, School of Medicine, Cardiff University

Correspondence: Annalisa Patrizi, Department of Specialized, Diagnostic and Experimental Medicine, Division of Dermatology, University of Bologna, Via Massarenti 1, 40138 Bologna, Italy. Email: annalisa.patrizi@unibo.it

Keywords: psoriasis, caregiver, quality of life, pediatric

Running short title: Quality of life in Italian children with psoriasis and their caregivers.

Word count: 700

Figure count: 1

References: 10

Consent for publication is from all authors.

Conflict of interest from all authors: none

Funding sources: none

The authors have no conflict of interest to disclose

Statement approval from CE: Prot/n/2411/04.07.19)

ACKNOWLEDGMENTS

We are grateful to Prof. Andrew Finlay for kindly granting us the permission to use CDLQI and PFI-14. We are grateful also to APIAFCO (ASSOCIAZIONE PSORIASICI ITALIANI AMICI della FONDAZIONE CORAZZA) for helping with the project.

In recent years there is an increasing interest in childhood psoriasis^{1,2}. In pediatric patients the impact of psoriasis on quality of life(QoL) is not limited to the patient himself, but inevitably also extends to the caregiver, by the chronic course of the disease^{3,4}.

The Children's Dermatology Life Quality Index (CDLQI) was developed to measure the impact of any skin disease on QoL in children⁵ while The Family Dermatology Life Quality Index and Psoriasis Family Index(PFI-14) were developed to measure the "secondary impact" of psoriasis on the caregivers^{6,7}.

The aim of this study was to investigate the impact of psoriasis on QoL in Italian children and their caregivers and to correlate demographic, clinical and therapeutic features with QoL.

Patients and caregivers were consecutively recruited in dermatology pediatric unit of Bologna from December 2018 to September 2020. Inclusion criteria were: (i)diagnosis of psoriasis in a child aged from 4 to 16 years; (ii)written consent form from at least one parent. Exclusion criteria were:(i)poor understanding of the Italian language, (ii)children with other systemic or inflammatory dermatological condition.

During the visit, were collected information about age, gender, age at the onset of the disease, family history, type of psoriasis, involved areas, severity of the disease (Psoriasis Area and Severity Index-PASI, Body Surface Area-BSA) and the impact on QoL by CDLQI. PASI was always rated by the same investigator (MAC-coauthor,dermatologist). Caregivers completed the PFI-14 questionnaire.

Patients were also classified into two groups according to treatment status (topical versus systemic agents). The study protocol was approved by the Local Research Ethic Committee of Bologna, Italy. Statistical analyses were conducted using SPSS Statistics Version 25.0. All tests were two-tailed, and the significance level was set at $p<0.05$.

Fifty patients and caregivers met the inclusion criteria and completed CDLQI and PFI-14 (Table 1). Thirty percent of patients were males, the mean age was 10.7 ± 3.6 years, the mean age at onset of psoriasis was 6.6 ± 2.8 years and the mean of psoriasis duration in our sample was 4 ± 2.9 years. There

were 44 patients(88%) with topic therapy, and 6(12%) with systemic therapy. Mean PASI score was 3.6 ± 2.6 (median=3) and mean BSA score was $2.6\pm 2.4\%$ (median=1); 96% had BSA and/or PASI below 10, that denotes mild psoriasis according to the rule of tens⁸. Mean CDLQI score was 7.5 ± 4.0 points(median=5).

According to CDLQI scores, in 2.0% of patients, psoriasis had no effect on their quality of life, in 60.0% a small effect, in 26.0% a moderate, effect, in 12.0% a very large effect.

Concerning caregivers, mean age was 37.7 ± 7.2 , 48 were the mother of the patient and 2 were the father, and 14 caregivers were single parents (28%). The mean PFI-14 score was 11.2 ± 6.9 points (median=9).

The total CDLQI score was significantly and moderately correlated with PASI (**figure 1a**) but not with patient age, sex, BSA, family history, alone child, type of psoriasis, disease duration, or age at disease onset.

Furthermore, there was a significant but weak correlation between CDLQI and PFI-14 (**figure 1b**).

The multiple linear regression model showed a positive and significant relation of CDLQI score with PFI-14 score ($\beta=0.722$; $p=0.012$), while PASI and BSA score had no significant effect on PFI-14 (**figure 1 c**).

Moreover, patients treated with topical agents had significantly lower CDLQI scores compared to those treated with systemic agents (**figure 1d**). The PFI-14 scores distribution was similar between the two groups.

The impact of many skin diseases is not limited to the patient but may extend to the rest of the family.

In our series the mother plays a central role in the care of children in 96% of cases. These findings are not surprising in Italian population where mothers spent more time with children than fathers.

Our study shows that impact of psoriasis on caregiver is significantly correlated with the children's QoL but not with PASI and BSA^{4,6,7}.

Noteworthy, 98% of pediatric patients reported that psoriasis has an effect on their life. The significant correlation between PASI and CLDQI are in accordance with results found in a recent metaanalysis and Turkish study^{9,10}.

PASI and BSA scores were unrelated with PFI-14 score, suggesting that patient QoL influences PFI-14 independently of disease severity. This finding confirming that QoL is a complex concept, with disease severity being merely one contributing factor^{6,9}. In this point of view dermatologists should address caregiver QoL in an integrated approach offering psychological support.

REFERENCES

1. Toscano P, Chiodini P, Ametrano O, Moscarella E. Childhood psoriasis: a survey among pediatricians in Italy. *G Ital Dermatol Venereol* 2018;153:473-476.
2. Burden-Teh E, Thomas KS, Ratib S, Grindlay D, Adaji E, Murphy R. The epidemiology of childhood psoriasis: a scoping review. *Br J Dermatol* 2016; 174:1242-57.
3. Beattie PE, Lewis-Jones MS. A comparative study of impairment of quality of life in children with skin disease and children with other chronic childhood diseases. *Br J Dermatol* 2006;155:145-51.
4. Gånemo A, Wahlgren CF, Svensson Å. Quality of life and clinical features in Swedish children with psoriasis. *Pediatr Dermatol* 2011;28:375-9.
5. Finlay AY, Lewis-Jones MS, Sharp JL, Dykes PJ. Childrens' Dermatology Life Quality Index: cartoon version validation. *Br J Dermatol* 1998; S67.
6. Eghlileb AM, Basra MK, Finlay AY. The Psoriasis Family Index: preliminary results of validation of a quality of life instrument for family members of patients with psoriasis. *Dermatology* 2009; 219:63-70.
7. Basra MK, Sue-Ho R, Finlay AY. The Family Dermatology Life Quality Index: measuring the secondary impact of skin disease. *Br J Dermatol* 2007;156 :528-38.
8. Finlay AY. Current severe psoriasis and the rule of tens. *Br J Dermatol* 2005;152:861-7.
9. Tekin B, Gurel MS, Topkarci Z, Topaloglu Demir F, Aytakin S, et al. Assessment of quality of life in Turkish children with psoriasis and their caregivers. *Pediatr Dermatol* 2018;35:651-659.
10. Randa H, Todberg T, Skov L, et al. Health-related quality of life in children and adolescents with psoriasis: a systematic review and meta-analysis. *Acta Derm Venereol* 2017;97:555-563.

Figure 1. (a) Scatterplot showing the relationship between CDLQI and PASI scores (Spearman coefficient $\rho=0.471$; $p<0.001$; $n = 50$); on the contrary CDLQI was not correlated with patient age, sex, BSA, family history, alone child, type of psoriasis, disease duration, or age at disease onset. **(b)** Scatterplot showing the relationship between Child Dermatology Life Quality Index (CDLQI) and PFI-14 scores (Spearman coefficient $\rho=0.363$; $p=0.01$; $n = 50$). **(c)** Multiple linear regression model on PFI-14 total score. Adjusted $R^2=0.171$, $F=4.370$, $p=0.009$. **(d)** Patients treated with topical agents had significantly lower CDLQI scores (median 5, IQR [4-7]) compared to those treated with systemic agents (median 12, IQR [8-15]) (Mann-Whitney U test=229.5, $p=0.002$), in particular in the subscale Treatment (Mann-Whitney U test=214; $p=0.012$).




TABLE 1. Multiple linear regression model on PFI-14 total score. Adjusted $R^2=0.171$, $F=4.370$, $p=0.009$.

Model	Coefficients	p-value	95% CI
Intercept	6.416	0.001	[2.815; 10.016]
CDLQI score	0.722	0.012	[0.168; 1.276]
PASI score	-0.637	0.238	[-1.710; 0.436]

BSA score	0.883	0.103	[-0.185; 1.952]
-----------	-------	-------	-----------------
