

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

How to predict aortic events after acute type A dissection repair: a matter of 'core' architecture

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Murana G., Di Marco L., Pacini D. (2022). How to predict aortic events after acute type A dissection repair: a matter of 'core' architecture. EUROPEAN JOURNAL OF CARDIO-THORACIC SURGERY, 62(3), 1-2 [10.1093/ejcts/ezac252].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/902844> since: 2022-11-15

*Published:*

DOI: <http://doi.org/10.1093/ejcts/ezac252>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

## Editorial comment

### How to predict aortic events after acute type A dissection repair: a matter of “core” architecture

Giacomo Murana<sup>1</sup> MD, PhD; Luca Di Marco<sup>1</sup> MD, PhD; Davide Pacini<sup>1,2</sup> MD, PhD

<sup>1</sup> Division of Cardiac Surgery, Cardiac Surgery Department, IRCCS, Azienda Ospedaliero-Universitaria di Bologna

<sup>2</sup> Department of Experimental, Diagnostic and Specialty Medicine, DIMES, University of Bologna, Bologna

**Key words:** aortic, dissection, computed tomographic, remodelling, false lumen.

**Word count:** 1114

**Conflict of Interest:** none declared

#### Corresponding author

Dr. Giacomo Murana

Cardiac Surgery Department

S. Orsola Hospital,

University of Bologna,

Via Massarenti 9, 40138, Bologna, Italy

Tel: 0039 051 2143361

Fax: 0039 051 345990

e-mail: [g.murana@hotmail.com](mailto:g.murana@hotmail.com); [giacomo.murana@aosp.bo.it](mailto:giacomo.murana@aosp.bo.it)

29 Ancient pagodas are architectural landmarks in many oriental countries. They usually consist in polygonal  
30 structures made of stone or wood built with sophisticated symmetries (both reflectional and rotational) due to  
31 the oriental cultural background [1]. As the height of the slice increases, the estimated radius will change  
32 rapidly so that the slice of the roof, eaves, and main bodies (multiple floors) can be identified.

33 The aorta can be imagined like a pagoda, both characterized by being long structures constructed over three  
34 multifunctioning layers that have the function to protect its integrity over decades.

35 Although these two fascinating creations present some architectural similarities, there is one big difference  
36 among them. The pagoda's secret is a core pillar that runs from the ground to the ceiling which is able to  
37 increase the pagoda's strength and protect it from collapse. Hence, an acute aortic dissection is able to alter  
38 the aortic architecture more than an earthquake is capable of damaging a Pagoda because the aortic "core" is  
39 missing.

40 For certain, distal aortic remodelling after dissection depends on multiple anatomic elements (including  
41 number and location of entries, aortic diameters, patency of FL, etc.) where the possibility to resect the  
42 primary entry tear represents the first fundamental step towards a positive remodeling of the downstream  
43 aorta [2,3].

44 In a very interesting study of Sang yoon Kim et al. from Seoul [4], they try to answer an ambitious question:  
45 can preoperative computed tomographic (CT) features predict residual false lumen remodelling after tear-  
46 oriented limited resection for acute type I aortic dissection. With this aim, the authors evaluate 101 patients  
47 who underwent ascending aorta with and without proximal partial arch replacement, analyzing preoperative  
48 and postoperative late CT scans. Preoperative CT parameters at the level of distal anastomosis were  
49 compared between patients who presented late positive remodeling of proximal portion of descending  
50 thoracic aorta and those who did not.

51 The main results of the study demonstrated that a positive remodelling of proximal descending thoracic aorta  
52 was observed in almost half of the patients and it was strongly correlated with small FL area ratio (<50%),  
53 with narrow FL width (<20%) at the distal anastomotic zone, and with not leaving residual arch branches  
54 having patent FL [4]. Based on these results, the authors concluded with the statement that a favorable  
55 remodelling can be expected in patients with limited cross-sectional extent of dissection after tear-oriented

56 limited resection and some elderly and non-Marfan patients can be spared from the aggressive total arch  
57 replacement with FET.

58 These very interesting findings raised two major reflections:

59 - The presence of residual entry in the downstream aorta usually influences blood fluid dynamic and  
60 patency of the false lumen affecting on distal aortic event-free survival [5]. The study did not  
61 investigate the impact of distal re-entry tears, neither the anastomotic technique for handling the  
62 distal anastomotic stump which is an important determinant for new anastomotic entry tear.

63 On the other hand, the authors respected the important principle of resecting every tear found in the  
64 operative field and focused the attention on cross-sectional extent in the preoperative CT image in  
65 the proximal descending thoracic aorta, which may affect the chance of creation of a new tear at the  
66 anastomosis site.

67 - Initial limited aortic resection is associated with an unsatisfactory long-term prognosis as  
68 demonstrated by a 70% occurrence of late distal aortic complications, including aneurysmal  
69 degeneration, rupture, malperfusion, and the need for extensive secondary interventions [6]. The  
70 possibility to predict distal aortic remodelling after total arch replacement (Zone II or III  
71 anastomosis) with a classical or a frozen elephant trunk (FET) was not considered because the  
72 authors wanted to focus just on the remodelling features following less extensive procedures.  
73 However, the FET become a widely adopted procedure supported by strong evidences reporting up  
74 to 85% positive remodelling in the proximal descending thoracic aorta [7]. Certainly, total arch  
75 replacement could have obscured the significance of other relevant preoperative factors but it still  
76 remains a crucial issue that would deserve further analyses.

77 In summary, we might argue that the only certain way to reduce later reinterventions is to resect or cover the  
78 primary entry tear in the arch at the time of first intervention. However, the study from the group of Seoul  
79 enhances the utility of computed tomographic features in predicting postoperative changes of the remaining  
80 descending thoracic aorta. In the same direction a future mission, similar to the “core pillar” of pagodas,  
81 should be to gain engineeristic capabilities regarding the use of geometric models and artificial intelligence  
82 systems to anticipate catastrophic events.

83

84     **References**

- 85           1.   Chan TO, Xia L, Chen Y, Lang W, Chen T, Sun Y, et al. Symmetry Analysis of Oriental Polygonal  
86           Pagodas Using 3D Point Clouds for Cultural Heritage. *Sensors (Basel)*. 2021 Feb 9;21(4):1228.
- 87           2.   Pacini D, Murana G, Di Marco L, Di Bartolomeo R. Aortic remodeling in acute type A aortic  
88           dissection after frozen elephant trunk implantation. *Indian J Thorac Cardiovasc Surg*. 2019  
89           Jun;35(Suppl 2):112-117.
- 90           3.   Berezowski M, Scheumann J, Beyersdorf F, Jasinski M, Plonek T, Siepe M, et al. Early aortic  
91           growth in acute descending aortic dissection. *Interact Cardiovasc Thorac Surg*. 2022 Jan 19:ivab351.  
92           Online ahead of print.
- 93           4.   Kim SY, Chang HW, Lee JH, Kim JS, Lim C, Park KH. Can preoperative features predict residual  
94           false lumen remodelling after tear-oriented limited resection for acute type I dissection? *Eur J*  
95           *Cardiothorac Surg*. 2022 Mar 9:ezac097. Online ahead of print.
- 96           5.   Kawahito K, Aizawa K, Kimura N, Yamaguchi A, Adachi H. Influence of residual primary entry  
97           following the tear-oriented strategy for acute type A aortic dissection. *Eur J Cardiothorac Surg*. 2021  
98           Nov 25:ezab456. Online ahead of print.
- 99           6.   Inoue Y, Matsuda H, Omura A, Seike Y, Uehara K, Sasaki H, et al. Comparative study of the frozen  
100           elephant trunk and classical elephant trunk techniques to supplement total arch replacement for acute  
101           type A aortic dissection *Eur J Cardiothorac Surg* 2019; 56: 579-586.
- 102           7.   Di Bartolomeo R, Murana G, Di Marco L, Pantaleo A, Alfonsi J, Leone A, et al. Frozen versus  
103           conventional elephant trunk technique: application in clinical practice. *Eur J Cardiothorac Surg*.  
104           2017 Jan;51(suppl 1):i20-i28.