

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

Dermoscopic Findings of Discoid Lupus Erythematosus Involving the Eyelids.

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Cortez de Almeida RF, F.B. (2022). Dermoscopic Findings of Discoid Lupus Erythematosus Involving the Eyelids. SKIN APPENDAGE DISORDERS, 8(5), 424-426 [10.1159/000524239].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/902974> since: 2022-11-15

*Published:*

DOI: <http://doi.org/10.1159/000524239>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Cortez de Almeida RF, Ferreira BFM, Melo DF, Benez M, Vincenzi C, Starace M, Tosti A.

*Dermoscopic Findings of Discoid Lupus Erythematosus Involving the Eyelids.*

Skin Appendage Disord. 2022 Sep; 8(5): 424-426.

The final published version is available online at: [10.1159/000524239](https://doi.org/10.1159/000524239)

#### Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

**When citing, please refer to the published version.**

## ***Novel Insights from Clinical Practice***

### ***Dermoscopic findings of Discoid Lupus Erythematosus involving the eyelids***

Rita Fernanda Cortez de Almeida<sup>a</sup>, Bianca de Franco Marques Ferreira<sup>a</sup>, Daniel Fernandes Melo<sup>a</sup>,  
Marcela Benez<sup>a</sup>, Colombina Vincenzi<sup>b</sup>, Michela Starace<sup>c</sup>, Antonella Tosti<sup>d</sup>

<sup>a</sup> Department of Dermatology, State University of Rio de Janeiro (UERJ), Rio de Janeiro, RJ, Brazil

<sup>b</sup> Department of Dermatology, Private Hospital Nigrisoli, Bologna, Italy

<sup>c</sup> Dermatology-IRCCS, Policlinico Sant'Orsola, Department of Specialized, Experimental and Diagnostic Medicine, Alma Mater Studiorum, University of Bologna, Bologna, Italy

<sup>d</sup> Dr. Phillip Frost Department of Dermatology and Cutaneous Surgery, University of Miami School of Medicine, Miami, FL, USA

**Short Title:** Dermoscopic findings of Lupus involving the eyelids

#### **Corresponding Author:**

Rita Fernanda Cortez de Almeida  
Department of Dermatology  
State University of Rio de Janeiro (UERJ)  
Boulevard 28 de Setembro, 77  
Rio de Janeiro, Rio de Janeiro, 20551-030, Brazil  
Tel: +55 21 28688478  
E-mail: ritaacortez@hotmail.com

**Tables:** 0

**Figures:** 1

**Word count:** 474

**Keywords:** Alopecia; Lupus; Lupus erythematosus; Eyelids; Dermoscopy

1    **Established Facts and Novel Insights**

Established Facts

- The eyelid involvement is uncommon in discoid lupus erythematosus (DLE).
- Dermoscopy is a valuable tool for diagnosing DLE and the literature is poorly described when affecting eyelids.

2

Novel Insights

- We report the main dermoscopic findings seen in two patients diagnosed with DLE, who presented an uncommon involvement of the eyelids.
- We reinforce the importance of considering DLE when facing inflammatory lesions and hair loss in the eyelids, especially using dermoscopy.

3

4

5    **Abstract**

6            Although facial involvement in discoid lupus erythematosus (DLE) is common, eyelid  
7 involvement is atypical. Identifying this condition is challenging due to misdiagnosis and it is important  
8 to avoid potential deformities of the eyelid margin. The aim of this paper is to report the main  
9 dermoscopic findings in two female patients with a confirmed diagnosis of DLE, who presented eyelids  
10 involvement. This study highlights the importance of performing a dermoscopy examination to help  
11 physicians to obtain an early diagnosis of DLE.

## Introduction

The eyelid involvement is uncommon in DLE, affecting only 5–6% of the patients [1]. As it is an atypical presentation, it may easily lead to misdiagnosis [2]. Dermoscopy is a valuable tool for diagnosing DLE in several locations. Its findings correspond to histopathology and disease duration [1,3]. However, the literature remains scarce about the dermoscopy of DLE affecting eyelids.

This article aims to report the dermoscopic features in two DLE patients with eyelid involvement and confirmed by histopathology, reiterating dermoscopy as an adjunctive tool for this diagnosis.

## Case Presentation

The first patient was a 38-year-old Afro-descendant female diagnosed with DLE, presenting an erythematous scaly patch in the lower eyelids and ciliary madarosis for 4 years. Dermoscopy showed hair loss, dystrophic hair, telangiectasias, polymorphic vessels, red globules (red lacunes), milky red area, yellowish scales, whitish structureless areas, reduced follicular openings, erythematous background, and scattered brown discoloration. Additionally, we referred the patient to an ophthalmologist evaluation to avoid a possible eye dysfunction.

The second patient was an 87-year-old Caucasian female, diagnosed with DLE, who presented a 2-year history of an erythematous and scaling patch on the upper left eyelid without the involvement of eyelashes. Dermoscopy revealed telangiectasias, erythematous background, and yellow dots. The main dermoscopic findings of both cases are presented in **Figure 1**.

## Discussion/Conclusion

The features found in the two patients follow the most common in scalp DLE, a site with well-established dermoscopic criteria [4]. As described in the literature for scalp and non-scalp DLE, red dots, milky red areas, scattered brown discoloration, and blue-grey dots are referred to as active DLE, whereas white structureless areas, telangiectasias, thin arborizing vessels on yellow dots, and loss of follicular openings are found in late inactive DLE [4].

DLE may lead to disfiguring scarring and permanent hair loss in the affected areas. Particularly, the unusual involvement of the eyelids may be even more challenging due to misdiagnosis and it is important to avoid potential deformities of the eyelid margin.

This report summarized the dermoscopic findings of eyelid DLE, which were very similar to those previously described in other areas. Furthermore, we highlight the importance of considering

DLE when facing inflammatory lesions and hair loss in the eyelids, especially using dermoscopy. In conclusion, despite the lack of criteria for this anatomical region, dermoscopy remains an auxiliary tool to help physicians obtain an early diagnosis of DLE.

## **Statements**

These data have not been previously presented. This work has not been submitted to another journal.

## **Statement of Ethics**

The study complies with the internationally accepted standards for research practice and reporting. Subjects have given their written informed consent to publish photos and details of the case. Ethical approval was not required for this study following national guidelines.

## **Conflict of Interest Statement**

Dr. Antonella Tosti acts as consultant for: Monat Global, Almirall, Tirthy Madison, Eli Lilly, Bristol Myers Squibb, P&G, Pfizer and Myovant. The other authors have no conflict of interest to declare.

## **Funding Sources**

This article has no funding source.

## **Author Contributions**

D.F.M. and A.T. conceived the study. R.F.C.A., B.F.M.F., D.F.M., M.B. and C.V. wrote the manuscript and collected the data. M.S. and A.T. reviewed the text. All authors discussed and reviewed the final manuscript.

## **Data Availability Statement**

All data generated or analyzed during this study are included in this article. Further enquiries can be directed to the corresponding author.

## References

1. Wang X, Wang P, Wang M, Huang X, Wang C, Qu X, et al. Discoid lupus erythematosus solely involving the eyelids: case report and literature review. *Int J Dermatol*. 2021; Oct;60(10):1190-1198.
2. Tosti A, Tosti G, Giovannini A. Discoid lupus erythematosus solely involving the eyelids: report of three cases. *J Am Acad Dermatol*. 1987; Jun;16(6):1259-60.
3. Costamilan LZ, Cerci FB, Werner B. Erythematous Plaque on the Inferior Eyelid. *JAMA Dermatol*. 2018; Aug;154(8):957-958.
4. Żychowska M, Żychowska M. Dermoscopy of discoid lupus erythematosus - a systematic review of the literature. *Int J Dermatol*. 2021; Jul;60(7):818-828

## Figure Legends

89 Fig. 1. Dermoscopic findings in Discoid Lupus Erythematosus involving the eyelids **(a)** presence of hair  
90 loss, reduced follicular openings, red globules (like red lacunes), milky red area, yellowish scales,  
91 whitish structureless areas, and scattered brown discoloration on dermoscopy of the right lower eyelid  
92 from patient 1. 3Gen DermLite® DL4 at 20-fold magnification **(b)** presence of dystrophic hair, hair loss,  
93 reduced follicular openings, red globules, telangiectasias, yellowish scales and whitish structureless  
94 areas on dermoscopy of the left lower eyelid from patient 1. 3Gen DermLite® DL4 at 20-fold  
95 magnification **(c)** close-up view showing the polymorphic vessels on dermoscopy of the left lower  
96 eyelid from patient 1. Fotofinder Dermoscope Germany at 20-fold magnification. **(d)** presence of  
97 telangiectasias, erythematous background and yellow dots on dermoscopy of the left upper eyelid  
98 from patient 2. Fotofinder Dermoscope Germany at 20-fold magnification.