

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Flow cytometry characterization of porcine seminal extracellular vesicles (EVs) based on the content of EV-specific protein markers

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Isabel Barranco, A.Á. (2022). Flow cytometry characterization of porcine seminal extracellular vesicles (EVs) based on the content of EV-specific protein markers. REPRODUCTION IN DOMESTIC ANIMALS, 57(S5), 143-143 [10.1111/rda.14227].

Availability:

This version is available at: <https://hdl.handle.net/11585/909789> since: 2022-12-16

Published:

DOI: <http://doi.org/10.1111/rda.14227>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the accepted version of an abstract published:

Isabel Barranco, Alberto Álvarez-Barrientos, Lorena Padilla, Ana Parra, Xiomara Lucas, Jordi Roca

Flow cytometry characterization of porcine seminal extracellular vesicles (EVs) based on the content of EV-specific protein markers

Reproduction in Domestic Animals. 2022; 57 (S5), 143.

The final published version is available online at:

<https://doi.org/10.1111/rda.14227>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Flow cytometry characterization of porcine seminal extracellular vesicles (EVs) based on the content of EV-specific protein markers

I. Barranco^{1,*}; A. Álvarez-Barrientos²; L Padilla^{3,4}; A. Parra⁴; X. Lucas⁴; J. Roca⁴

¹Department of Veterinary Medical Sciences, University of Bologna, Ozzano dell'Emilia, Bologna, Italy; ²Applied Bioscience Facility, University of Extremadura, Badajoz, Spain; ³Biotechnology of Animal and Human Reproduction (TechnoSperm), Department of Biology, Faculty of Sciences, Institute of Food and Agricultural Technology, University of Girona, Girona, Spain; ⁴ Department of Medicine and Animal Surgery, Faculty of Veterinary Science, University of Murcia, Murcia, Spain

Porcine seminal plasma (SP) contains a heterogeneous population of EVs that when isolated by a protocol based on size exclusion chromatography (SEC) can be grouped into two subsets showing size differences, namely large (L) and small (S) EVs. This study evaluated the expression of EV-specific proteins in S- and L- pig SP-EVs. 105 SP-samples (one per boar) were randomly arranged in 21 pools (5 SP-samples/pool) and EV-subsets were isolated by sequential centrifugations, SEC and ultrafiltration. The S- and L-EVs were characterized by total protein (BCATMassay), morphology (TEM), size distribution (DLS) and purity (flow cytometry determination of albumin). The profile of the EV-specific proteins CD9, CD44, CD63, CD81, and HSP90 β was analyzed using a flow cytometer (CytoFLEX S). S- and L- SP-EVs showed similar total protein concentration, morphology, and purity (albumin <7%). DLS revealed differences ($P < 0.0001$) in size (median, 25–75th percentiles; 120.8, 115.3–126.2 nm and 260.8, 235.3–300.7 nm to S- and L-EVs, respectively). There was no difference between S- and L-EVs in the expression of EV-specific proteins. Both EV-subsets showed high positivity for CD44 (mean \pm SEM, 98.82 \pm 0.11 vs 98.43 \pm 0.10 to S- and L-EVs, respectively); HSP90 β (83.67 \pm 3.11 vs 84.03 \pm 2.60) and CD63 (83.11 \pm 1.31 vs 77.35 \pm 2.21); and low for CD9 (26.32 \pm 5.54 vs 28.56 \pm 5.12) and CD81 (29.66 \pm 3.45 vs 22.66 \pm 2.42). These results would indicate that both pig SP-EV-subsets show similar EV-specific protein profiles and CD44, CD63 and HSP90 β labelling allows for accurate characterization. Fundings: MCIN/AEI/10.13039/501100011033 (PID2020-113493RB-I00); EC (H2020-MSCA-IF-2019-891382).