

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Background independence and field redefinitions in quantum gravity

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Casadio, R., Kamenshchik, A., Kuntz, I. (2023). Background independence and field redefinitions in quantum gravity. ANNALS OF PHYSICS, 449, 1-8 [10.1016/j.aop.2022.169203].

Availability:

This version is available at: <https://hdl.handle.net/11585/910870> since: 2022-12-31

Published:

DOI: <http://doi.org/10.1016/j.aop.2022.169203>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

Casadio, R., Kamenshchik, A., & Kuntz, I. (2023). Background independence and field redefinitions in quantum gravity. *Annals of Physics*, 449

The final published version is available online at
<https://dx.doi.org/10.1016/j.aop.2022.169203>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Background independence and field redefinitions in quantum gravity

Roberto Casadio^{ab*}, Alexander Kamenshchik^{ab†}, and Iberê Kuntz^{c‡}

^a*Dipartimento di Fisica e Astronomia, Università di Bologna
via Irnerio 46, 40126 Bologna, Italy*

^b*I.N.F.N., Sezione di Bologna, I.S. FLAG
viale B. Pichat 6/2, 40127 Bologna, Italy*

^c*Departamento de Física, Universidade Federal do Paraná
PO Box 19044, Curitiba – PR, 81531-980, Brazil*

Abstract

It is generally believed that a full-fledged theory of quantum gravity should exhibit background independence and diffeomorphism invariance. In its most general form, the latter comprises field redefinitions, which are diffeomorphisms in configuration space. We show that any path-integral approach to quantum gravity leads to a tension between these properties, such that they cannot hold simultaneously.

1 Introduction

Background independence and covariance are the basic building blocks of general relativity. Both concepts are intertwined with the diffeomorphism invariance of gravity and has led to a long history of confusion [1–3]. Anderson [3] was the first to put such concepts into a more precise language by demarcating the differences of absolute and dynamical structures. Although not free of inconsistencies [4], Anderson’s approach delineates the role played by the diffeomorphism group in gravitational and non-gravitational theories, which is a stepstone for defining covariance and background-independence rigorously.

*E-mail: casadio@bo.infn.it

†E-mail: kamenshchik@bo.infn.it

‡E-mail: kuntz@fisica.ufpr.br

These concepts are also believed to play a fundamental role in the quantization of gravity [5,6]. The conventional perturbative approach of quantum field theory singles out a background metric required for the vacuum structure of the theory. Because this background is not dynamical, it violates the background independence of general relativity. Allowing for this possibility, but insisting on the invariance under diffeomorphisms, is the cornerstone of string theory [7–10]. On the other hand, Canonical Quantum Gravity and some of its variants, including Loop Quantum Gravity, has been developed as a background-independent theory *ab initio* [11–16]. Nonetheless, the absence of spacetime structures makes it difficult to recover well-tested results of quantum field theory at low energies.

Quantum field theories of gravity can also be formulated, to some extent, in a background-independent manner. This is indeed implemented with the background field method [17], where a separation of quantum perturbations from (dynamical) background fields is performed. The background field method was introduced as a convenient way to deal with gauge theories, in such a way that the gauge symmetry could be preserved at the background level while fixing the gauge for the quantum perturbations. The resulting action is therefore gauge invariant, which eases tremendously some calculations, such as the proof of renormalizability.

The background field, despite its name, need not be chosen *a priori*. It need not even satisfy the classical equations of motion. Indeed, the background field should solve the effective equations of motion. It is in this sense that a quantum field theory of gravity can be background independent [18]. Nonetheless, this construction requires the definition of the asymptotic vacuum state, thus restricting the set of background metric solutions to the ones with sufficient asymptotic symmetry. We stress, however, that it is the asymptotic boundary conditions rather than the background metric that are fixed beforehand.

When the background field is not chosen as a solution to the field equations, the effective action turns out to depend on the gauge. One thus reaches an impasse, where either a background is picked up *a priori*, violating the background independence of general relativity but whose predictions are gauge-independent, or the background is solved for *a posteriori*, which respects background independence but suffers from gauge dependence. This issue is solved within the Vilkovisky-DeWitt formalism [19, 20], where a Levi-Civita connection in configuration space is introduced to cancel out the gauge-dependence while maintaining the background fields (including the metric) arbitrary. The cancellation of the gauge dependence also extends the invariance of on-shell scattering amplitudes under field redefinitions to off-shell correlation functions. We note that field redefinitions are nothing else than diffeomorphisms in configuration space.¹ In the remainder of this article, covariance shall thus

¹An example of field redefinitions often treated in gravity and cosmology is the transition between the Jordan frame and the Einstein frame. It is important to emphasize that this transition includes not only the conformal transformation of the metric turning the nonminimal coupling between the scalar field and the curvature into the minimal one, but also the redefinition of the scalar field to make its kinetic term canonical [21]. The question of the classical and quantum equivalence between two frames was extensively discussed in the literature (see e.g. [22] and [23] and the references therein). We think that these frames are equivalent mathematically, in the sense that one can describe everything using any frame (just like one can use inertial and non-inertial frames in Newtonian mechanics). At the the same time, physically, observers connected with different frames can see different phenomena. For example, in one frame one can have de

mean both spacetime and configuration-space covariance.

The purpose of this paper is to show that quantum field theories of gravity, even within the formalism of Vilkovisky-DeWitt, cannot consistently satisfy both background-independence (absence of *any* non-dynamical structure) and covariance (under spacetime diffeomorphisms and field redefinitions) at the same time. As we shall show, the latter necessarily requires absolute structures to be introduced, which in turn violates the former, whereas a theory that satisfies the former does not possess the correct transformation properties, thus violating the latter.

2 The geometrical effective action

A valuable property of quantum field theory regards the invariance of the S-matrix under field redefinitions [25]. Scattering amplitudes are indeed the observables of interest in Particle Physics, thus their invariance under field reparameterizations seems to suggest some profound principle. Observables in Astrophysics and Cosmology are, however, correlation functions and these do not transform covariantly in the standard formulation of quantum field theory. One could argue that this proves the absence of such a general principle. Nonetheless, in a gauge theory, field-parameterization dependence induces gauge dependence in the observables. The former is, therefore, a genuine issue.

The origin of this gauge dependence is most easily seen by the naive effective action

$$\exp\left(\frac{i}{\hbar}\Gamma_N[\phi]\right) = \int \mathcal{D}\varphi \exp\left\{\frac{i}{\hbar}\left[S[\varphi] + (\varphi^i - \phi^i)\frac{\delta\Gamma_N}{\delta\phi^i}\right]\right\}, \quad (2.1)$$

where $\varphi^i = \varphi^I(x)$ collectively denotes an arbitrary set of fields with classical action $S[\varphi]$ and ϕ^i is the mean field. The set of all φ^i is the configuration space, formally defined by the infinite-dimensional space

$$\mathcal{C}(M) = \prod_x \mathcal{N}(x), \quad (2.2)$$

where $\mathcal{N}(x)$ denotes the finite-dimensional space formed by all $\varphi^I(x)$ evaluated at the same spacetime point $x \in M$. Under gauge transformations or field redefinitions, neither the path-integral measure nor the second term within the curly brackets in Eq. (2.1) remains invariant. One can impose invariance in the former by introducing a configuration-space metric G_{ij} , whereas the latter calls for a configuration-space connection Γ_{ij}^k compatible with G_{ij} . This procedure results in the Vilkovisky-DeWitt effective action [19, 20].

The configuration-space metric G_{ij} is usually identified as the bilinear operator from the classical kinetic term [19].² However, this procedure to fix G_{ij} is not required nor physically well-motivated. It is a pure mathematical procedure. For this reason, we shall see G_{ij} as part of the definition of the theory, together with a choice for the classical action (but not

Sitter expansion, while in another frame one sees other law of the expansion or even some kind of bounce [24].

²We recall that the canonical quantisation of metric theories of gravity is based on the Bergmann-Komar group [26] of the super-momenta and super-Hamiltonian, whose explicit form depends on the spatial part of the DeWitt metric, hence the choice of field dynamics.

derived from it). One can then determine G_{ij} using symmetry and an expansion on the energy scale, in the same spirit of effective field theories [27].

For the sake of simplicity, in this paper we shall omit the connection-dependent term and focus on the functional measure.³ This is justified because of the compatibility property of the aforementioned connection, which then provides no additional absolute structures (to be defined in Section 3) other than the already present metric G_{ij} , thus not altering our argument of Section 4. The improved effective action shall then be written as

$$\exp\left(\frac{i}{\hbar}\Gamma_{\text{VD}}[\phi]\right) = \int \mathcal{D}\varphi \sqrt{\text{Det } G_{ij}} \exp\left(\frac{i}{\hbar}S[\varphi]\right), \quad (2.3)$$

where Det denotes the functional determinant.

3 Background independence and covariance

In this section, we shall review Anderson's efforts [3] to define covariance rigorously. In the notations of [4], a theory's equations of motion are generally of the form

$$\mathcal{F}(\gamma, \phi, \Sigma) = 0, \quad (3.1)$$

where γ denotes all structures defined by maps into spacetime (e.g. a particle's trajectory) and ϕ denotes collectively all structures represented by maps from spacetime (e.g. fields). Here Σ is a given non-dynamical structure defined on the spacetime M for which the equations of motion (3.1) can be solved for (γ, ϕ) . A theory is called covariant iff, for every $f \in \text{Diff}(M)$,

$$\mathcal{F}(f \cdot \gamma, f \cdot \phi, f \cdot \Sigma) = 0 \iff \mathcal{F}(\gamma, \phi, \Sigma) = 0, \quad (3.2)$$

where the dot product $f \cdot X$ denotes the action of f , by means of pullback/pushforward, on the object X . On the other hand, a theory is said to be invariant iff, for all $f \in \text{Diff}(M)$,

$$\mathcal{F}(f \cdot \gamma, f \cdot \phi, \Sigma) = 0 \iff \mathcal{F}(\gamma, \phi, \Sigma) = 0. \quad (3.3)$$

The subtle difference lies on the transformation of the non-dynamical structure Σ . Covariance is the requirement for the theory to be well-defined geometrically, i.e. for the theory to live on M . Thus, when some structure is transformed under a diffeomorphism, all other structures of the theory are transformed along to counteract the differences and keep their relation the same. Invariance, on the other hand, requires solutions of the theory to be transformed into other solutions for the *same* Σ . The latter is much more restrictive and constitutes an important guiding principle in physics.

The structure Σ is called an absolute structure if it is a field that either is not dynamical or whose solutions are all locally diffeomorphic to each other. One can then define a background-independent theory as a theory which contains no absolute structure. Notice that, in this case, covariance equals invariance.

³For this reason, our argument also applies to any construction of the path integral where a functional measure is required, like in non-linear sigma models.

In special relativity, for instance, the absolute structure is the Minkowski metric $\Sigma = \{\eta_{\mu\nu}\}$. The passage to general relativity is characterized by making the metric dynamical $\eta_{\mu\nu} \rightarrow g_{\mu\nu}$. Now $g_{\mu\nu}$ is not given, thus $\Sigma = \emptyset$, but is found *a posteriori* as a solution of Einstein's field equations, hence $\phi = \{g_{\mu\nu}\}$.

As we anticipated in the Introduction, Anderson's definitions are not flawless. To see this, one can write the dynamics of a scalar field in special relativity in a diffeomorphism-invariant form as

$$g^{\mu\nu} \nabla_\mu \nabla_\nu \Phi = 0 , \quad (3.4)$$

$$R_{\mu\nu\rho\sigma} = 0 . \quad (3.5)$$

According to Anderson's definitions, this would also make special relativity background independent, which is an obvious flaw. First of all, Anderson's definition of an absolute structure does not entail the action-reaction principle. While $\eta_{\mu\nu}$ is obtained dynamically, thereby acting on Φ , it is not acted upon by Φ . A fully dynamical field should have a symmetrical role where it acts on and is also acted upon by other fields ⁴. Interactions are indeed always symmetric.

The source of the problem lies in the fact that Eq. (3.5) is not a dynamical equation, but rather a constraint. In fact, the most general solution $g_{\mu\nu} = \alpha \eta_{\mu\nu}$, for any non-zero constant α , does not contain enough integration constants to fully account for the initial conditions, hence the Cauchy evolution is ill-defined. Furthermore, Eq. (3.5) does not follow from any diffeomorphism-invariant action unless a Lagrange multiplier is introduced, which stresses the fact that this equation is a constraint.

In view of the above issues, we shall refine the definitions of covariance, invariance and background independence. Rather than basing our definitions on the equations of motion, our proposal shall be performed at the level of the effective action. This not only avoids the issues outlined above, but generalises the discussion to the quantum realm, which is our main concern. Moreover, the equations of motion do not capture important physical effects contained in the boundary terms (vide black hole entropy). The effective action, on the other hand, contains information about all physics at classical or quantum scales.

We shall define an absolute structure Σ as any local object that does not contain a corresponding kinetic term in the effective action. A theory is then called covariant iff, for every $f \in \text{Diff}(\mathcal{C}(M))$, the effective action transforms as

$$\Gamma[f \cdot \phi, f \cdot \Sigma] = \Gamma[\phi, \Sigma] , \quad (3.6)$$

whereas we say that it is invariant iff, for every $f \in \text{Diff}(\mathcal{C}(M))$, we have ⁵

$$\Gamma[f \cdot \phi, \Sigma] = \Gamma[\phi, \Sigma] . \quad (3.7)$$

⁴This property is somewhat related to Mach's principle [28, 29].

⁵More generally, a symmetry group $G \subset \text{Diff}(\mathcal{C}(M))$ of some $\text{Diff}(\mathcal{C}(M))$ -covariant theory is the group that leaves the absolute structure Σ invariant. An invariant theory is thus invariant under the largest symmetry group $G = \text{Diff}(\mathcal{C}(M))$.

Because we are interested in a fundamental theory, where all interactions are entirely described with fields, we have not included γ in our definitions. We should also stress that, like in Anderson’s formulation, Σ is really any non-dynamical object, not necessarily a spacetime object. A theory is then said to be background independent whenever $\Sigma = \emptyset$.

We stress that some absolute global framework shall always be needed in order to define a workable mathematical theory of physical phenomena. For instance, the spacetime $(M, g_{\mu\nu})$ should be globally hyperbolic in order to give rise to well-posed Cauchy problems in classical general relativity [30], and similarly in (classical or quantum) field theories, although such restrictions are sometimes lifted or overlooked. This means that properties like the dimension, topology, and differential structures of the spacetime (or field space) must be fixed to some extent from the onset when describing specific systems. In practical terms, our discussion of background independence will therefore concern the existence of a fixed (local) metric at some level of mathematical abstraction, as we will further clarify in the following. Points, dimension, topological and differential structures are therefore not included in our definition of background independence.

Before we proceed, it is important to recall that observables in general relativity are mathematically given by scalars built out of (pairs of) tensors pertaining to the quantity to be measured and the measuring apparatus (observer).⁶ The prototype is given by the energy of a particle, which is obtained by the contraction of the particle’s 4-velocity u^μ with the observer’s 4-velocity U^μ , that is $E = -g_{\mu\nu} u^\mu U^\nu$, where $g_{\mu\nu}$ is the background metric tensor. Similarly, one might notice that fields in configuration space have no direct physical meaning. This makes fields like spacetime coordinates but also as tensor quantities in spacetime: proper observables, like scattering cross sections, are built out of fields by (more or less) involved “scalarizations” involving the measuring apparatus.

4 Background independence and field redefinitions

At the level of the naive effective action $\Gamma^{(0)}[\phi] \equiv \Gamma_N[\phi]$, no absolute structure exists, thus $\Sigma = \emptyset$ and the covariance (3.6) collapses into invariance (3.7). However, like we saw in Section 2, $\Gamma^{(0)}[\phi]$ depends on the field parameterization and, therefore, does not satisfy invariance (3.7).

This can be amended by the introduction of a configuration-space metric G_{ij} , which is used to define an invariant path-integral measure and ultimately results in the Vilkovisky-DeWitt effective action $\Gamma^{(1)}[\phi, \Sigma] \equiv \Gamma_{\text{VD}}[\phi, G]$. Although $\Gamma^{(1)}[\phi, \Sigma]$ satisfies invariance (3.7), we now have an absolute structure $\Sigma = \{G_{ij}\}$ that renders the theory background dependent.

Let us make the theory background independent by promoting G_{ij} to a dynamical structure. On assuming ultralocality,

$$G_{ij} = G_{IJ} \delta(x, x') , \tag{4.1}$$

⁶This is the main idea behind the “relational” approach to (quantum) physics (see [31] and references therein).

one could envisage doing so by introducing the action

$$S = \int d^n x \sqrt{-g} \mathcal{L}(G_{IJ}, R_{IJMN}) , \quad (4.2)$$

where R_{IJMN} is the Riemann tensor of the finite-dimensional space \mathcal{N} and $\mathcal{L}(G_{IJ}, R_{IJMN})$ is some Lagrangian invariant under spacetime and configuration-space diffeomorphisms. The precise form of the action for G_{IJ} is not essential to our argument as the only crucial point to the following is the existence of dynamics (regardless of the particular kind of evolution). One could, for example, assume that such a Lagrangian takes the Einstein-Hilbert form

$$\mathcal{L}(G_{IJ}, R_{IJMN}) = G^{IJ} R_{IJ} , \quad (4.3)$$

where $R_{IJ} = G^{MN} R_{MINJ}$ is the configuration-space Ricci tensor.

With a dynamical configuration space, there is no other absolute structure present, thus the theory recovers its background independence $\Sigma = \emptyset$. However, this came at the cost of introducing the additional degree of freedom G_{ij} in the theory, which must be quantised along with φ^i . In the absence of external sources, the quantum theory then reads

$$\exp \left(\frac{i}{\hbar} \Gamma^{(2)}[\phi] \right) = \int \mathcal{D}\phi \mathcal{D}G_{ij} \sqrt{\text{Det } G_{ij}} \exp \left(\frac{i}{\hbar} S[\phi, G] \right) , \quad (4.4)$$

The functional measure

$$\mathcal{D}\phi \mathcal{D}G_{ij} \sqrt{\text{Det } G_{ij}} = \prod_{x, I < J} d\phi dG_{IJ} \sqrt{\det G_{IJ}} \quad (4.5)$$

does not transform covariantly. To see this, it is sufficient to perform an infinitesimal field redefinition

$$G_{ij} \rightarrow G'_{ij} = G_{ij} + \xi_{ij}(G) , \quad (4.6)$$

under which the functional measure transforms as

$$\mathcal{D}G'_{ij} \sqrt{\text{Det } G'_{ij}} = \mathcal{D}G_{ij} \sqrt{\text{Det } G_{ij}} \left(1 + G^{mn} \xi_{mn} + \frac{\delta \xi_{mn}}{\delta G_{mn}} \right) . \quad (4.7)$$

Promoting G_{ij} to a dynamical field thus brings back the violation of invariance under field redefinitions. To correct this, we must find a different measure that transforms covariantly. This unavoidably requires the introduction of some absolute structure $\mathcal{S}^{(2)}$ that transforms in such a way that ⁷

$$\mathcal{D}\phi \mathcal{D}G'_{ij} \sqrt{\text{Det } G'_{ij}} \mathcal{S}^{(2)'} = \mathcal{D}\phi \mathcal{D}G_{ij} \sqrt{\text{Det } G_{ij}} \mathcal{S}^{(2)} , \quad (4.8)$$

⁷Should the geometrical construction be maintained, $\mathcal{S}^{(2)} = \sqrt{\text{Det } G_{i_2 j_2}^{(2)}}$ would correspond to the determinant of a second-order functional metric $G_{i_2 j_2}^{(2)}$ defined on a second-order configuration space $\mathcal{C}^{(2)}(M)$, where now the indices i_2, j_2 include the new field G_{ij} .

which recovers background dependence since $\Sigma = \{\mathcal{S}^{(2)}\}$. We thus see that promoting $\mathcal{S}^{(n)}$ to a dynamical field requires, upon quantization, that an absolute structure $\Sigma = \{\mathcal{S}^{(n+1)}\}$ be introduced, hence the effective action $\Gamma^{(n)}[\phi, \Sigma]$ can never satisfy invariance (3.7) and background independence $\Sigma = \emptyset$ at the same time.

In summary, although we have been able to recover background independence by making G_{ij} dynamical, we lost invariance in the full quantum theory where G_{ij} belongs to the spectrum. To regain invariance we would need to include another layer of abstraction (a sort of *über*-field space) with yet another higher-order absolute structure, which would then face the problems with background dependence and so on and so forth. This process continues *ad infinitum*, proving the existence of a tension between background independence and covariance in quantum gravity.

5 Conclusions

In this paper, we have considered extended notions of diffeomorphism invariance and background independence related to the (non) existence of some fixed metric structure at the quantum level. In this context, diffeomorphism invariance also comprises invariance of observables under field redefinitions, which are required to enforce gauge independence in the correlation functions. Likewise, a metric structure in field space is naturally introduced by any given dynamical theory.

We have then shown that diffeomorphism invariance and background independence, in the extended sense of the absence of any fixed metric structure at any level, cannot coexist. While the former is unavoidable for any sensible physical theory, the latter does not seem to be rooted in any physical property other than an aesthetic appeal (to some). Therefore, one possible and natural implication of our finding is that quantum gravity is actually background dependent (at some level in the hierarchy of mathematical structures described at the end of the last section). We stress that our definition of background independence concerns only (local) geometrical objects, such as metric structures in the *über*-field spaces. As we already stated, points, dimension, topological and differential structures are not included in this definition. On the other hand, should background independence be revealed as an intrinsic aspect of Nature, models of quantum gravity based on quantum field theory, such as asymptotic safety and causal dynamical triangulation, would become effective rather than fundamental theories.

Acknowledgments

R.C. and A.K. are partially supported by the INFN grant FLAG. The work of R.C. has also been carried out in the framework of activities of the National Group of Mathematical Physics (GNFM, INdAM).

References

- [1] J. D. Norton, Rep. Prog. Phys. **56** 791 (1993) 791.
- [2] O. Pooley, Einstein Stud. **13** (2017) 105 [arXiv:1506.03512 [physics.hist-ph]].
- [3] J. L. Anderson, “Principles of Relativity Physics”, Academic Press, 1967.
- [4] D. Giulini, Lect. Notes Phys. **721** (2007) 105 [arXiv:gr-qc/0603087 [gr-qc]].
- [5] D. Colosi, L. Doplicher, W. Fairbairn, L. Modesto, K. Noui and C. Rovelli, Class. Quant. Grav. **22** (2005) 2971 [arXiv:gr-qc/0408079 [gr-qc]].
- [6] L. Smolin, “The Case for background independence,” [arXiv:hep-th/0507235 [hep-th]].
- [7] J. Polchinski, “String theory. Vol. 1: An introduction to the bosonic string,” Cambridge University Press, 2007.
- [8] J. Polchinski, “String theory. Vol. 2: Superstring theory and beyond,” Cambridge University Press, 2007.
- [9] E. Witten, “Quantum background independence in string theory,” [arXiv:hep-th/9306122 [hep-th]].
- [10] M. Rozali, Adv. Sci. Lett. **2** (2009) 244 [arXiv:0809.3962 [gr-qc]].
- [11] B. S. DeWitt, Phys. Rev. **160** (1967) 1113.
- [12] P. G. Bergmann, Phys. Rev. **144** (1966) 1078.
- [13] P. A. M. Dirac, Proc. Roy. Soc. Lond. A **246** (1958) 326; P. A. M. Dirac, “Lectures on quantum mechanics,” Dover, 1964.
- [14] C. Kiefer, “Quantum Gravity,” Oxford University Press, 2007.
- [15] C. Rovelli, “Quantum gravity,” Cambridge University Press, 2004.
- [16] A. Ashtekar and J. Lewandowski, Class. Quant. Grav. **21** (2004) R53 [arXiv:gr-qc/0404018 [gr-qc]].
- [17] L. F. Abbott, Acta Phys. Polon. B **13** (1982) 33.
- [18] M. Reuter, Phys. Rev. D **57** (1998) 971 [arXiv:hep-th/9605030 [hep-th]]; M. Becker and M. Reuter, Phys. Rev. D **102** (2020) 125001 [arXiv:2008.09430 [gr-qc]]; M. Becker and M. Reuter, Phys. Rev. D **104** (2021) 125008 [arXiv:2109.09496 [hep-th]].
- [19] G. A. Vilkovisky, Nucl. Phys. B **234** (1984) 125.
- [20] B. S. DeWitt, “The effective action”, in Batalin, I.A. (Ed.) et al.: Quantum field theory and quantum statistics, Vol. 1, 191-222, CRC Press (1987).

- [21] R.W. Wagoner, Phys. Rev. D **1** (1970) 3209-3216
- [22] A.Yu. Kamenshchik and C.F. Steinwachs, Phys. Rev. D **91** (2015) 8, 084033 [arXiv:1408.5769 [gr-qc]]
- [23] K. Falls and M. Herrero-Valea, Eur. Phys. J. C **79** (2019) 595 [arXiv:1812.08187 [hep-th]].
- [24] A.Yu. Kamenshchik, E.O. Pozdeeva, A. Tronconi, G. Venturi and S.Yu. Vernov, Class. Quant. Grav. **31** (2014) 105003 [arXiv:1312.3540 [hep-th]]
- [25] A. Rebhan, Nucl. Phys. B **298** (1988) 726.
- [26] P. G. Bergmann and A. Komar, Int. J. Theor. Phys. **5** (1972) 15.
- [27] R. Casadio, A. Kamenshchik and I. Kuntz, Nucl. Phys. B **971** (2021) 115496 [arXiv:2102.10688 [hep-th]].
- [28] J. Ehlers, “Machian Ideas and General Relativity”, in Barbour, J.B. (Ed.) et al.: Mach’s principle - from Newton’s bucket to quantum gravity, 458-473, Birkhäuser (1995).
- [29] A. Einstein, “The Meaning of Relativity”, Princeton University Press (1922).
- [30] K. Landsman, “Reopening the Hole Argument,” [arXiv:2206.04943 [gr-qc]].
- [31] C. Rovelli, “The Relational Interpretation of Quantum Physics,” [arXiv:2109.09170 [quant-ph]].