

Alma Mater Studiorum Università di Bologna  
Archivio istituzionale della ricerca

'Rotor free-wheeling' in impaired F1FO-ATPase induces congenital hypermetabolism

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

*Published Version:*

Nesci S., Romeo G. (2023). 'Rotor free-wheeling' in impaired F1FO-ATPase induces congenital hypermetabolism. *TRENDS IN ENDOCRINOLOGY AND METABOLISM*, 34(2), 63-65 [10.1016/j.tem.2022.12.002].

*Availability:*

This version is available at: <https://hdl.handle.net/11585/911708> since: 2023-04-20

*Published:*

DOI: <http://doi.org/10.1016/j.tem.2022.12.002>

*Terms of use:*

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).  
When citing, please refer to the published version.

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

**'Rotor free-wheeling' in impaired  $F_1F_0$ -ATPase induces congenital hypermetabolism.**

**Nesci S, Romeo G. Trends in Endocrinology and Metabolism. 2023, 34: 63-65.**

The final published version is available online at:  
**<https://doi.org/10.1016/j.tem.2022.12.002>**

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

[Click here to view linked References](#)

# “Rotor free-wheeling” in impaired F<sub>1</sub>F<sub>0</sub>-ATPase causes congenital hypermetabolism

Salvatore Nesci<sup>1\*</sup> and Giovanni Romeo<sup>2</sup>

<sup>1</sup>Department of Veterinary Medical Sciences, University of Bologna, 40064 Ozzano Emilia, Italy.

<sup>2</sup>Medical Genetics Unit, Sant’Orsola-Malpighi University Hospital, 40126 Bologna, Italy.

\*Correspondence: [salvatore.nesci@unibo.it](mailto:salvatore.nesci@unibo.it)

**Keywords:** F<sub>1</sub>F<sub>0</sub>-ATPase; congenital hypermetabolism; mitochondrial bioenergetics; ATP production; mutation.

## Abstract

A *de novo* heterozygous variant in the catalytic subunit of mitochondrial F<sub>1</sub>F<sub>0</sub>-ATPase arises as the main cause of an autosomal dominant syndrome of hypermetabolism associated with defective ATP production as recently discovered by Ganetzky *et al.* We describe how the “rotor free-wheeling” causes this F<sub>1</sub>F<sub>0</sub>-ATPase dysfunction in primary congenital hypothyroidism.

## Main text

Two monozygotic twin boys with a new type of congenital euthyroid hypermetabolism characterized by caloric intake, inability to gain weight, and sporadic hyperthermia were recently described by Ganetzky *et al.* [1]. The Authors discovered a *de novo* heterozygous variant with c.1004 T→C transition in the nuclear *ATP5F1B* gene of mitochondrial F<sub>1</sub>F<sub>0</sub>-ATPase. Accordingly, the mutation p.Leu<sub>335</sub>Pro influences the catalytic β subunit of the enzyme (Fig. 1A). The clinical features might be associated with the molecular evidence of pathological conditions that characterize the mitochondrial uncoupling syndromes. Indeed, the dominant-negative effect on bioenergetic parameters at the mitochondrial level highlighted a defective chemiosmotic coupling of oxidative phosphorylation (OXPHOS), which is typical of depolarized mitochondria in the presence of increased oxygen consumption rate without ATP production [1].

F<sub>1</sub>F<sub>0</sub>-ATPase is embedded in the inner mitochondrial membrane (IMM) working as a unique bifunctional enzyme in nature. Indeed, the electrochemical gradient of proton (H<sup>+</sup>) or protonmotive force (Δ*p*) through the IMM, built during the substrate oxidation by the mitochondrial respiratory complexes, is exploited by F<sub>1</sub>F<sub>0</sub>-ATPase to synthesize ATP. However, under patho(physio)logical conditions the enzyme in the absence of oxygen working in reverse as H<sup>+</sup> pump and hydrolyzing ATP can polarize the IMM [2]. A hydrophilic F<sub>1</sub> domain directed into the mitochondria matrix and a hydrophobic membrane-localized F<sub>0</sub> domain form the F<sub>1</sub>F<sub>0</sub>-ATPase. The former is responsible for ATP catalysis, *i.e.* ATP synthesis/hydrolysis, whereas the latter is involved in the H<sup>+</sup> translocation through the IMM [3]. The two domains are functionally coupled by the central stalk, which penetrates inside the F<sub>1</sub> inducing a mechanical conformational change in the catalytic sites by its

rotation, and a peripheral stalk connecting laterally  $F_1$  with membrane subunits of  $F_0$  that avoids the rotation driven by the rotor [4].

In the  $F_0$  domain of the enzyme, an asymmetric half-channels system of  $\alpha$  subunit dissipates the  $\Delta p$  by  $H^+$  translocation. The  $H^+$  flow across the IMM consists of a mechanism of protonation and deprotonation of the  $H^+$  binding site of  $c$  subunits that drives the  $c$ -ring rotation. The torsion generated by  $H^+$  translocating down the  $\Delta p$  is transmitted via the central stalk to power the mechanism of binding change conformation of catalytic sites into  $F_1$  domain. Therefore, in energized mitochondria, the  $F_1F_0$ -ATPase can perform ATP synthesis as “a splendid molecular machine” [5]. The chemomechanical mechanism of  $F_1$  domain produces three ATP molecules with one complete revolution of the rotor accordingly to the three-fold symmetric structure of the  $F_1$  domain. Indeed, three catalytic  $\beta$  subunits and three non-catalytic  $\alpha$  subunits are arranged in alternation to form a hexagonal structure  $(\alpha\beta)_3$  with the cavity filled with the asymmetric shape of the central stalk  $\gamma$  subunit [6]. The  $(\alpha\beta)_3$  hexamer has three adenine nucleotide (AN) binding sites placed on the three catalytic subunits at the interfaces with the homologous non-catalytic subunit that in response to the curvature of the  $\gamma$  subunit during the rotor rotation, adopt a different conformation with different nucleotide-binding and catalytic properties. These are identified in the order of transformation as empty  $\beta_E$  structure,  $\beta_{DP}$ , which contains bound ADP plus  $P_i$  and, in the last conformation of the cycle, as  $\beta_{TP}$  that binds ATP. Each binding change, which drives the simultaneous interchange of the  $\beta_E \rightarrow \beta_{DP} \rightarrow \beta_{TP}$  forms, requires a  $120^\circ$  rotation of the  $\gamma$  subunit with respect to  $\beta$  subunits and triggers the synthesis of an ATP molecule [7] (Fig. 1B).

The new Leu<sub>335</sub>Pro variant affects the mitochondrial bioenergetics causing an increase in basal oxygen consumption rate almost reaching maximal cell respiration level. This means that the low level of ATP production and depolarized IMM are inconsistent with the stimulated activity of substrate oxidation in mitochondria. The impaired chemiosmotic mechanism of OXPHOS dwelling in the failing energy transduction phenomenon is due to an intrinsic uncoupling of the  $F_1F_0$ -ATPase. In other terms, the enzyme depletes the  $\Delta p$  without a mutual increase in ATP synthesis. The inhibition of  $H^+$  translocation by an inhibitor of  $F_0$   $H^+$  transport, *i.e.* oligomycin, can therefore block the uncontrolled  $H^+$  flow increasing the membrane potential and the oxygen consumption rate stopping the passive  $H^+$  slip of  $F_1F_0$ -ATPase, whereas protonophore molecules maximize mitochondrial respiration. The results of Ganetzky *et al.* [1] found in mutated mitochondria highlight a synergistic action in dropping the mitochondrial membrane potential in the absence of ATP synthesis and in polarizing the IMM inhibiting the  $F_0$  domain. We may conclude that the mitochondrial dysfunction lies in the chemo-mechanical activity of  $F_1$  domain. The damage in the nuclear *ATP5F1B* gene can modify the affinity of catalytic sites for adenine nucleotides and/or may give rise to a  $\gamma$  subunit “rotor free-wheeling” in the catalytic hexamer without causing the simultaneous interchange of the  $\beta$  subunits (Fig 1B).

The asymmetric bulge of the  $\gamma$  subunit has three communication points with the catalytic sites on the  $\beta$  subunits. Starting from the apex of the  $F_1$  domain going towards the  $F_0$  domain, a hydrophobic loop sleeve is near the tip of the coiled-coil  $\alpha$ -helices of the  $\gamma$  subunit. In the middle of the cavity of  $(\alpha\beta)_3$  structure lies the Leu<sub>335</sub>Pro mutation on a loop connecting a hydrophobic portion supporting the interaction with rotor rotation. Finally, in the C-terminal domain of the  $\beta$  subunit a critical helix-turn-helix motif known as “DELSSEED” is presumably involved in the coupling between catalysis and rotation.

*Cis-trans*-isomerization of the peptide bond is a feature of a secondary amine group, called imine, of Pro residue. Relevance on Leu<sub>335</sub>Pro mutation might be the *cis-trans* isomerization during protein folding by torsional peptide bond that could displace the hydrophobic communication points gearing inside the  $(\alpha\beta)_3$  from  $\gamma$  subunit. Accordingly, the torque generation of the central stalk required for the mechanochemical power of the  $F_1$ -ATPase domain [8] is obtained through a “free-wheeling of the rotor” (Fig 1B).

In general, the non-ohmic  $H^+$  conductance of IMM is stimulated by  $\Delta p$  and uncoupling protein 1 and ADP/ATP carrier have the physiological role of dissipation of the excessive membrane potential by  $H^+$  leak [9]. The

mechanochemical destruction of  $F_1$  energy transduction is the result of intrinsic  $F_1F_0$ -ATPase uncoupling [10]. During the  $H^+$  translocation, the enzyme misses the ATP synthesis dropping the membrane potential and depicting the biochemical phenomenon of congenital hypermetabolism symptoms. On balance, the  $H^+$  slip of  $F_1F_0$ -ATPase characterizes several primary mitochondrial disorders stemming from pathological “rotor free-wheeling” (Fig. 1B).

## Declaration of Interests

All authors declare no competing interests.

## References

1. Ganetzky, R.D. *et al.* (2022) Congenital Hypermetabolism and Uncoupled Oxidative Phosphorylation. *N Engl J Med* 387, 1395–1403
2. Nesci, S. *et al.* (2020) Mitochondrial F-type ATP synthase: multiple enzyme functions revealed by the membrane-embedded FO structure. *Crit. Rev. Biochem. Mol. Biol.* 55, 309–321
3. Pinke, G. *et al.* (2020) Cryo-EM structure of the entire mammalian F-type ATP synthase. *Nat Struct Mol Biol* 27, 1077–1085
4. Murphy, B.J. *et al.* (2019) Rotary substates of mitochondrial ATP synthase reveal the basis of flexible  $F_1$ - $F_0$  coupling. *Science* 364
5. Boyer, P.D. (1997) The ATP synthase--a splendid molecular machine. *Annu. Rev. Biochem.* 66, 717–749
6. Furuike, S. *et al.* (2008) Axle-less  $F_1$ -ATPase rotates in the correct direction. *Science* 319, 955–958
7. Hahn, A. *et al.* (2016) Structure of a Complete ATP Synthase Dimer Reveals the Molecular Basis of Inner Mitochondrial Membrane Morphology. *Mol. Cell* 63, 445–456
8. Suzuki, T. *et al.* (2014) Chemomechanical coupling of human mitochondrial  $F_1$ -ATPase motor. *Nat. Chem. Biol.* 10, 930–936
9. Bertholet, A.M. and Kirichok, Y. (2022) Mitochondrial  $H^+$  Leak and Thermogenesis. *Annu Rev Physiol* 84, 381–407
10. Nesci, S. *et al.* (2021) Molecular and Supramolecular Structure of the Mitochondrial Oxidative Phosphorylation System: Implications for Pathology. *Life* 11, 242

## Figure

Figure 1. Mechanochemistry dysfunction of  $F_1F_0$ -ATPase caused by Leu<sub>335</sub>Pro mutation. A) Transition T→C (red square), as well as the Leu<sub>335</sub>Pro mutation highlighted during the transcription and translation of *ATP5F1B* gene. B) Representation of  $F_1F_0$ -ATPase drawn as ribbon representations obtained from modified Protein Data Bank identifier 6TT7. The differently coloured letters identify the subunits, drawn in the same colour as the letter. On the left is the *wild-type* enzyme during ATP synthesis and in the insert the mechanochemical coupling of the  $F_1$  domain. On the right is  $F_1F_0$ -ATPase with mutated  $\beta$  subunit missing the coupling of  $H^+$  translocation generation of rotor rotation to the binding chance mechanism of ATP production. Leu<sub>335</sub>Pro mutation induces a rotor free-wheeling in the energy transduction mechanism of the  $F_1F_0$ -ATPase. The three dotted squares on the  $\beta$  subunits show in the space-filling model the three mutated amino acid residues, which are coloured in red. This figure was created using BioRender (<https://biorender.com/>).

