

Alma Mater Studiorum Università di Bologna
Archivio istituzionale della ricerca

Real-world effectiveness of pembrolizumab as first-line therapy for cisplatin-ineligible patients with advanced urothelial carcinoma: the ARON-2 study

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Santoni, M., Myint, Z.W., Büttner, T., Takeshita, H., Okada, Y., Lam, E.T., et al. (2023). Real-world effectiveness of pembrolizumab as first-line therapy for cisplatin-ineligible patients with advanced urothelial carcinoma: the ARON-2 study. *CANCER IMMUNOLOGY, IMMUNOTHERAPY*, 72(9), 2961-2970 [10.1007/s00262-023-03469-5].

Availability:

This version is available at: <https://hdl.handle.net/11585/927473> since: 2023-05-30

Published:

DOI: <http://doi.org/10.1007/s00262-023-03469-5>

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>).
When citing, please refer to the published version.

(Article begins on next page)

This is the accepted manuscript of:

Santoni M, Myint ZW, Büttner T, Takeshita H, Okada Y, Lam ET, Gilbert D, Küronya Z, Tural D, Pichler R, Grande E, Crabb SJ, Kemp R, Massari F, Scagliarini S, Iacovelli R, Vau N, Basso U, Maruzzo M, Molina-Cerrillo J, Galli L, Bamias A, De Giorgi U, Zucali PA, Rizzo M, Seront E, Popovic L, Caffo O, Buti S, Kanavaras R, Kopecky J, Kucharz J, Zeppellini A, Fiala O, Landmesser J, Ansari J, Giannatempo P, Rizzo A, Zabalza IO, Monteiro FSM, Battelli N, Calabrò F, Porta C.

Real-world effectiveness of pembrolizumab as first-line therapy for cisplatin-ineligible patients with advanced urothelial carcinoma: the ARON-2 study.

Cancer Immunol Immunother. 2023 Sep;72(9):2961-2970.

The final published version is available online at: <https://doi.org/10.1007/s00262-023-03469-5>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

Real-world effectiveness of pembrolizumab as first-line therapy for cisplatin-ineligible patients with advanced Urothelial Carcinoma: the ARON-2 study

Matteo Santoni¹, Zin W. Myint², Thomas Büttner³, Hideki Takeshita⁴, Yohei Okada⁴, Elaine T Lam⁵, Zsófia Küronya⁶, Deniz Tural⁷, Renate Pichler⁸, Enrique Grande⁹, Simon J Crabb¹⁰, Francesco Massari^{11*}, Sarah Scagliarini¹², Roberto Iacovelli¹³, Nuno Vau¹⁴, Umberto Basso¹⁵, Javier Molina-Cerrillo¹⁶, Luca Galli¹⁷, Aristotelis Bamias¹⁸, Ugo De Giorgi¹⁹, Paolo Andrea Zucali^{20,21}, Mimma Rizzo²², Emmanuel Seront²³, Lazar Popovic²⁴, Orazio Caffo²⁵, Sebastiano Buti²⁶, Ravindran Kanesvaran²⁷, Jindrich Kopecky²⁸, Jakub Kucharz²⁹, Annalisa Zeppellini³⁰, Ondřej Fiala³¹, Johannes Landmesser³², Jawaher Ansari³³, Patrizia Giannatempo³⁴, Ignacio Ortego Zabalza⁹, Fernando Sabino M. Monteiro^{35,36}, Nicola Battelli¹, Fabio Calabrò^{37°}, Camillo Porta^{22,38°}

°co-Senior Authors

¹Oncology Unit, Macerata Hospital, via Santa Lucia 2, 62100, Macerata, Italy;

²Markey Cancer Center, University of Kentucky, Lexington, KY, 40536-0293, USA;

³Department of Urology, University Hospital Bonn (UKB), 53127 Bonn, Germany;

⁴Department of Urology, Saitama Medical Center, Saitama Medical University, Saitama, Japan; ⁵University of Colorado Anschutz Medical Campus, Aurora, CO, USA;

⁶Department of Genitourinary Medical Oncology and Clinical Pharmacology, National Institute of Oncology, Budapest, Hungary; ⁷Department of Medical

Oncology, Bakirköy Dr. Sadi Konuk Training and Research Hospital, Zuhuratbaba District, Tevfik Saglam St. No: 11, Bakirkoy, Istanbul, Turkey; ⁸Department of Urology, Medical University of Innsbruck, Anichstrasse 35, 6020 Innsbruck, Austria; ⁹Department of Medical Oncology, MD Anderson Cancer Center Madrid, Madrid, Spain; ¹⁰Southampton Experimental Cancer Medicine Centre, University of Southampton, Southampton, United Kingdom; ¹¹Medical Oncology, IRCCS Azienda Ospedaliero-Universitaria di Bologna, Via Albertoni - 15, Bologna – Italia; ¹²UOC di Oncologia, Azienda Ospedaliera di Rilievo Nazionale Cardarelli di Napoli, Naples, Italy; ¹³Oncologia Medica, Fondazione Policlinico Universitario Agostino Gemelli IRCCS, Roma, Italy; ¹⁴Urologic Oncology, Champalimaud Clinical Center, 1400-038 Lisbon, Portugal; ¹⁵Medical Oncology 1 Unit, Department of Oncology, Istituto Oncologico Veneto IOV IRCCS, 35128 Padova, Italy; ¹⁶Department of Medical Oncology, Hospital Ramón y Cajal, Madrid, Spain; ¹⁷Oncology Unit 2, University Hospital of Pisa, Pisa 56126, Italy; ¹⁸2nd Propaedeutic Dept of Internal Medicine, ATTIKON University Hospital, National and Kapodistrian University of Athens, School of Medicine, Athens, Greece; ¹⁹Department of Medical Oncology, IRCCS Istituto Romagnolo per lo Studio dei Tumori (IRST) "Dino Amadori", Meldola, Italy; ²⁰Department of Biomedical Sciences, Humanitas University, Pieve Emanuele, Milan, Italy; ²¹Department of Oncology, IRCCS Humanitas Research Hospital, Rozzano - Milan, Italy; ²²Division of Medical Oncology, A.O.U. Consorziale Policlinico di Bari, Piazza G. Cesare 11, 70124 Bari, Italy; ²³Department of Medical Oncology, Centre Hospitalier de Jolimont, Haine Saint Paul, Belgium; ²⁴Oncology Institute of Vojvodina, Faculty of Medicine, University Novi Sad, Novi Sad, Serbia; ²⁵Medical Oncology Unit, Santa Chiara Hospital, Trento, Italy; ²⁶Medical Oncology Unit, University Hospital of

Parma – Department of Medicine and Surgery, University of Parma, Parma, Italy;

²⁷Division of Medical Oncology, National Cancer Centre Singapore, Singapore;

²⁸Department of Clinical Oncology and Radiotherapy, University Hospital Hradec Kralove, Hradec Kralove, Czech Republic; ²⁹Department of Uro-oncology, Maria Sklodowska-Curie National Research Institute of Oncology Warsaw, Poland;

³⁰Niguarda Cancer Center, Grande Ospedale Metropolitano Niguarda, Milan, Italy;

³¹Department of Oncology and Radiotherapeutics, Faculty of Medicine and University Hospital in Pilsen, Charles University, Pilsen, Czech Republic; ³²Klinik für Urologie, Ratzeburger Allee 160, 23538 Lübeck, Germany; ³³Medical Oncology, Tawam Hospital, Al Ain, United Arab Emirates; ³⁴Dipartimento di Oncologia Medica, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy; ³⁵Latin American Cooperative Oncology Group – LACOG; ³⁶Oncology and Hematology Department, Hospital Santa Lucia, SHLS 716 Cj. C, Brasília, DF 70390-700, Brazil; ³⁷Department of Oncology, San Camillo Forlanini Hospital, Rome, Italy; ³⁸Chair of Oncology, Interdisciplinary Department of Medicine, University of Bari “Aldo Moro”, Bari, Italy.

***Correspondence to:**

Matteo Santoni, MD, PhD

Oncology Unit, Macerata Hospital, via Santa Lucia 2, 62100, Macerata, Italy. Phone number: +3907332572960; FAX number: +3907332573783; e-mail: mattymo@alice.it

Abstract

To be completed according to Journal Instructions.

Keywords: ARON-2 study; Immunotherapy; NCT05290038; Pembrolizumab; Real-world data; Survival; Tumor Response; Urothelial Cancer.

Introduction

The American Cancer Society has estimated 164,190 new cancer cases of the urinary system only in the United States in 2022 [1]. Approximately the 50% of these cases consist in tumors of the urinary bladder, which represents the 6% of all cancer diagnoses and the 4% of cancer-related deaths in men [1].

Urothelial cancer (UC) is the most prevalent histologic subtype of tumors of the upper and lower urinary tracts, accounting for approximately 90% of all cases [2]. About 25% of patients with UC present with metastatic disease, reporting a 5-year survival rate of only 7.7% [3].

In the past three decades, the management of first-line advanced UC has consisted in the administration of platinum-based chemotherapy [4]. The advent of immune-checkpoint inhibitors (ICIs) have challenged previous treatment paradigms of advanced UC in the post-platinum setting as well as in the first-line setting for cisplatin-ineligible patients [5,6], defined by the presence of at least one of the criteria published by Galsky *et al.* in 2011 [7]: (1) Eastern Cooperative Oncology Group (ECOG)-Performance Status (PS) of 2, (2) Creatinine clearance of less than 60 mL/minute, (3) Common Terminology Criteria for Adverse Events (CTCAE) of at least grade 2 hearing loss, (4) CTCAE of at least grade 2 neuropathy and (5) New York Heart Association Class III heart failure.

The approval of pembrolizumab by the Food and Drugs Administration (FDA) in cisplatin-unfit patients followed the results of the Keynote-052 phase 2 trial [8]. In this study, patients previously treated with prior systemic chemotherapy for unresectable or metastatic UC were excluded, while patients receiving platinum-based chemotherapy for tumor recurrence >12 months since completion of

adjuvant or neoadjuvant therapy were eligible. The primary endpoint was objective response; 374 patients were enrolled; at a median follow-up of 5 months, Overall Response Rate (ORR) was 24%. In 2020, Vuky *et al.* [9] published the results of the long-term outcomes of patients included in the Keynote-052 study, with a minimum follow-up of 2 years. The ORR was 28.6%, with 8.9% of complete remissions, with median overall survival (OS) and duration of response of 11.3 months (95% CI, 9.7 to 13.1 months) and 30.1 months (95% CI, 18.1 months to not reached [NR]), respectively.

The ARON project was designed to globally share and analyze real-world data from patients with genitourinary tumors treated by immunotherapy. In particular, the ARON-2 study included UC patients receiving pembrolizumab as first or subsequent line therapy. In this study, we investigated the effectiveness of pembrolizumab as first-line treatment for cisplatin-ineligible patients affected by advanced UC.

Patients and Methods

Study population

The ARON-2 study retrospectively collected data from patients aged ≥ 18 years having a cytological and/or histological confirmed diagnosis of cisplatin-ineligible UC receiving first-line pembrolizumab (200 mg i.v. flat dose, every three weeks until clear radiological or clinical progression of disease) from January 1st 2017 to September 1st 2022. Cisplatin ineligibility was defined according to the Galsky criteria [7]. Thirty-three Institutions from 18 countries were involved in the ARON-2 study.

The ARON-2 dataset included clinical and laboratory data extracted from patients'

paper and electronic charts on age, gender, ECOG-PS, tumor histology, surgery, sites of metastases, and response to therapy were retrospectively collected. Patients with insufficient data on tumor response to therapy were not included in the ARON-2 study.

Computed tomography (CT) or magnetic resonance imaging (MRI) scans were performed following standard local procedures every 8–12 weeks. Physical and laboratory tests were usually carried out every 4–6 weeks during patients' follow-up.

Study endpoints

The primary endpoint of the study was the effectiveness, assessed as ORR, duration of response (DoR), progression-free survival (PFS) and OS. Objective Response Rate (ORR) was calculated as the proportion of patients who achieved a complete (CR) or partial response (PR) defined by RECIST 1.1 criteria [10]. DoR was defined as the time from CR or PR to progression. Overall Survival (OS) was the time from treatment initiation to death from any cause. Progression-Free Survival (PFS) was calculated from the start of first-line pembrolizumab to progression or death from any cause, whichever occurred first. Patients without disease progression or death or lost at follow-up at the time of the analysis were censored at the last follow-up visit.

Statistical Analysis

The Kaplan-Meier statistical methods with Rothman's 95% confidence intervals (CI) were used to estimate median OS, PFS and DoR. Cox proportional hazards models were used to compare the multivariable effects on patients' survival and to calculate hazard ratios (HRs) and 95% CIs. Receiver operating characteristic (ROC)

analysis was also used to understand the sensitivity and specificity profile and to identify potential cut-offs that better stratify patients in risk groups. The chi-square test was used to compare each group for categorical variables. Statistical differences were considered significant when the p -value was <0.05 , and all p values were two-sided.

The upper tract was defined as the renal pelvis or ureter, and the lower tract as the bladder or urethra. Bajorin prognostic classification (including both the presence of visceral/bone metastases and Eastern Cooperative Oncology Group-Performance Status, ECOG-PS), was used to stratify patients according the number of risk factors [11].

MedCalc version 19.6.4 (MedCalc Software, Broekstraat 52, 9030 Mariakerke, Belgium) was used for the statistical analyses.) for all analyses. This study is registered with ClinicalTrials.gov, number NCT05290038.

Results

Baseline characteristics

Our analysis included 162 patients. The median follow-up time was 18.9 months (95%CI 15.3–76.9); 117 patients (74%) were males. Median age was 71y (range 43–94). ECOG-PS was ≥ 2 in 27 patients (17%). Tumors of the upper urinary tract accounted for the 16% of all cases. Tumor histology was pure UC in 142 patients (88%), squamous in 8 (5%), micropapillary in 3 (2%), glandular in 2 (1%), nested in 2 (1%), sarcomatoid in 2 (1%), undifferentiated in 2 (1%), and clear cell in 1 (1%). The 27% of patients presented metastatic disease at UC diagnosis. Lymph node and visceral metastases were identified in 112 (69%) and 90 (56%) patients, respectively.

Stratifying by Bajorin prognostic classification [11], 63 patients (39%) presented 0 factors, 81 patients (50%) 1 factor and 18 patients (11%) both factors.

Seventy-three patients (45%) had died at time of the analysis. Treatment with pembrolizumab was ongoing in 74 patients (46%). Twenty-nine (33%) of the 88 patients progressed during first-line pembrolizumab were treated with second-line therapies. Patients' baseline characteristics are summarized in Table 1.

Overall Survival and Progression-Free Survival analyses

In the overall study population, the median OS was 15.8 months (95%CI 11.3–32.4, Figure 1). The median OS was significantly longer in males vs females (21.2 months, 95%CI 12.3–45.9, vs 11.7 months, 95%CI 5.9–16.8, $p=0.031$, Figure 2). Otherwise, no statistically significant differences were observed between patients aged <65y vs ≥ 65 y (16.8 months, 95%CI 10.5–45.9 vs 15.8 months, 95%CI 10.5–32.4, $p=0.614$) and between smokers and non-smokers (16.8 months, 95%CI 11.1–45.9, vs 13.4 months, 95%CI 8.8–29.4, $p=0.594$).

Patients with ECOG-PS ≥ 2 showed worst median OS compared to ECOG-PS 0-1 (7.4 months, 95%CI 6.1–14.3, vs 19.4 months, 95%CI 11.7–45.9, $p=0.044$, Figure 2).

Patients with pure UC histology showed a median OS of 16.8 months (95%CI 11.2–45.9), while in patients with mixed histology was 12.5 months (95%CI 3.3–32.4, $p=0.438$). Analogously, no statistically significant differences were found between patients with tumors of the upper tract (29.4 months, 95%CI 10.5–29.4) vs lower tract (14.3 months, 95%CI 11.1–33.4, $p=0.160$).

Metachronous metastatic disease was associated with longer median OS (19.4 months, 95%CI 11.3–45.9, vs 11.7 months, 95%CI 4.3–32.4, $p=0.038$, Figure 3). Patients

with lymph node metastases only showed longer median OS compared to those with visceral metastases (29.4 months, 95%CI 13.4–45.9, vs 11.1 months, 95%CI 7.7–17.0, $p=0.026$, Figure 3). By stratifying patients according to sites of metastasis, a statistically significant difference was observed between patients with or without bone metastases (7.6 months, 95%CI 3.6–16.8, vs 21.2 months, 95%CI 12.5–45.9, $p=0.016$, Figure 3).

According to Bajorin risk factors [11], the median OS resulted 32.4 months (95%CI 15.8–45.9) in patients with 0 risk factors, 12.3 months (95%CI 7.7–17.0) in patients with 1 risk factor and 11.1 months (95%CI 1.7–14.3) in patients with 2 risk factors ($p=0.019$, Figure 3).

Furthermore, no significant differences were found between patients treated with first-line pembrolizumab who had received or not previous adjuvant or neoadjuvant chemotherapy (14.2 months, 95%CI 10.5–32.4, vs 17.0 months, 10.5–45.9, $p=0.791$).

In the overall study population, the median PFS was 10.2 months (95%CI 6.2–17.6, Figure 1). The median PFS was longer in males vs females, but the difference was not statistically significant (11.4 months, 95%CI 8.2–28.6, vs 5.1 months, 95%CI 2.7–26.3, $p=0.155$). Analogously, no statistically significant differences were observed between patients aged <65y vs ≥65y (9.0 months, 95%CI 6.2–36.7 vs 11.3 months, 95%CI 4.8–26.3, $p=0.688$), smokers vs non-smokers (9.1 months, 95%CI 6.2–17.0, vs 12.2 months, 95%CI 4.3–28.6, $p=0.734$), ECOG-PS≥2 vs <2 (9.3 months, 95%CI 1.9–15.2, vs 10.2 months, 95%CI 6.2–24.2, $p=0.585$), pure vs mixed UC histology (11.4 months, 95%CI 6.2–24.2, vs 4.8 months, 95%CI 2.0–13.4, $p=0.111$), upper vs lower urinary tract (18.6 months, 95%CI 2.2–28.6, vs 10.2 months, 95%CI 6.2–17.6, $p=0.790$), and previous adjuvant or neoadjuvant chemotherapy vs no previous treatments (7.5 months,

95%CI 4.8–11.3, vs 15.2 months, 6.6–28.6, $p=0.120$).

Synchronous metastatic disease was associated with shorter median PFS (5.3 months, 95%CI 2.3–11.7, vs 11.7 months, 95%CI 7.5–28.6, $p=0.017$, Figure 4). Patients with lymph node metastases only showed longer median PFS compared to those with visceral metastases (26.3 months, 95%CI 11.3–33.7, vs 5.5 months, 95%CI 3.8–9.1, $p=0.002$, Figure 4). By stratifying patients according to sites of metastasis, patients with bone metastases showed a significantly shorter median PFS (3.4 months, 95%CI 1.9–5.5, vs 17.0 months, 95%CI 9.4–28.6, $p<0.001$, Figure 4).

Finally, according to Bajorin risk classification, the median PFS resulted 26.3 months (95%CI 9.4–33.7) in patients with 0 risk factors, 8.6 months (95%CI 5.1–17.0) in patients with 1 risk factor and 4.8 months (95%CI 1.2–13.4) in patients with 2 risk factors ($p=0.022$, Figure 4).

Response to therapy

According to Recist 1.1 criteria, 26 patients (16%) experienced CR, 32 (20%) PR, 39 (24%) SD and 55 (34%) PD. Patients with 0, 1 or 2 Bajorin risk factors showed an ORR of 43%, 33%, and 22%, respectively ($p=0.007$).

The median OS resulted significantly different according to the type of response, being NR (95%CI NR–NR), 45.9 months (95%CI 17.0–45.9), 13.4 months (95%CI 7.4–19.4) and 6.4 months (95%CI 2.9–21.2) in patients with CR, PR, SD, and PD, respectively ($p<0.001$, Figure 5). In the 58 patients who presented CR or PR, 44 were still ongoing at time of the analysis. The median DoR was NR (95%CI NR–NR).

Role of prognostic factors

At univariate analysis, gender, synchronous metastatic disease and Bajorin risk classification were significant predictors of OS, and their prognostic role was confirmed at multivariate analysis (Table 2).

Furthermore, IMDC group, synchronous metastatic disease and Bajorin risk classification significantly correlated with PFS at univariate analysis, although only Bajorin risk factors proved to be associated with PFS at multivariate analysis (Table 2).

Discussion

Cisplatin-unfit patients represent at least 30–40% of patients with metastatic UC [12]. Owing to a median age at diagnosis of 70 years, and being smoking an associated risk factor, many patients have pulmonary and/or cardiovascular diseases that lead to an accelerated deterioration of renal function. Age-related decrease in glomerular filtration rate impairs the patients' possibility to receive cisplatin, as well as impaired ECOG-PS.

A randomized EORTC study [13] comparing M-CAVI and gemcitabine plus carboplatin in patients cisplatin-unfit not only proved acceptable Response Rate of the gemcitabine plus carboplatin arm (38%), but also supported the importance of stratification parameters and Bajorin risk groups on RR. Indeed, patients with 0, 1 or 2 Bajorin risk factors showed a RR of 47%, 39% and 20%, respectively.

More recently, Martini *et al.* [14] led a meta-analysis to compare chemotherapy and immunotherapy in the first-line setting of advanced UC. They observed no OS benefit for patients treated with first-line immune checkpoint inhibition compared to

chemotherapy among the overall population, cisplatin-ineligible patients, and PD-L1-high patients.

The ARON-2 study was designed to assess the effectiveness of pembrolizumab as first or successive line therapy in patients with advanced UC and the presence of factors influencing the prognosis of these patients. In this sub-analysis, we focused on the efficacy of pembrolizumab in the first-line setting of patients unable to receive cisplatin. We showed that median OS and PFS were 15.8 and 10.2 months, respectively. The presence of synchronous metastatic disease, visceral or bone metastases and a higher number of Bajorin risk factors were significantly associated with worst OS and PFS. The ORR was 36% with 16% of CR and type of tumor response according to RECIST 1.1 criteria was a significant predictor of OS.

Our results in terms of both OS and PFS are more favorable than those previously published in the Keynote-052 study investigating pembrolizumab in the same context. This difference can be explained by the presence of a lower percentage of patients with ECOG-PS $\geq$ 2 (17% vs 42%) and visceral metastases (56% vs 85%) included in the ARON-2 study population. This means that in our population we have 83% of patients considered cisplatin unfit due to reasons other than PS (i.e. renal failure, hearing loss, neuropathy and heart failure), probably reflecting better the population in real life who need first-line treatment in this setting. Of note, the recent presentation of novel cisplatin-unfit criteria at ASCO 2022 [15] may influence the selection of patients candidate to receive first-line pembrolizumab and should be prospectively investigated.

Our study presents several limitations, mainly due to its retrospective nature. We had no available data neither on the criteria of eligibility of cisplatin-unfit patients nor on

concomitant medication or other comorbidities that could affect the efficacy of first-line therapy. As a consequence, our results should be interpreted with caution and are in need of a larger prospective validation.

Despite a centralized review of radiological imaging was not performed, the strong correlation between response and OS suggest a reliable assessment of RECIST 1.1 in the present study.

Nevertheless, our data show that pembrolizumab was effective as first-line therapy for cisplatin-unfit patients. Further studies investigating the biological and immunological characteristics of UC patients are warranted in order to optimize the outcome of patients receiving immunotherapy in this setting.

Conflicts of Interest

Matteo Santoni has received research support and honoraria from Janssen, Bristol Myers Squibb, Ipsen, MSD, Astellas and Bayer, all unrelated to the present paper.

R. Kanesvaran has received fees for speaker bureau and advisory board activities from the following companies; Pfizer, MSD, BMS, Eisai, Ipsen, Johnson and Johnson, Merck, Amgen, Astellas and Bayer.

Enrique Grande has received honoraria for speaker engagements, advisory roles or funding of continuous medical education from Adacap, AMGEN, Angelini, Astellas, Astra Zeneca, Bayer, Blueprint, Bristol Myers Squibb, Caris Life Sciences, Celgene, Clovis-Oncology, Eisai, Eusa Pharma, Genetracer, Guardant Health, HRA-Pharma, IPSEN, ITM-Radiopharma, Janssen, Lexicon, Lilly, Merck KGaA, MSD, Nanostring Technologies, Natera, Novartis, ONCODNA (Biosequence), Palex, Pharmamar, Pierre Fabre, Pfizer, Roche, Sanofi-Genzyme, Servier, Taiho, and Thermo Fisher

Scientific. EG has received research grants from Pfizer, Astra Zeneca, Astellas, and Lexicon Pharmaceuticals

Fernando Sabino Marques Monteiro has received research support from Janssen, Merck Sharp Dome and honoraria from Janssen, Ipsen, Bristol Myers Squibb and Merck Sharp Dome.

Camillo Porta has received honoraria from Angelini Pharma, AstraZeneca, BMS, Eisai, General Electric, Ipsen and MSD and acted as a Protocol Steering Committee Member for BMS, Eisai and MSD.

Sebastiano Buti received honoraria as speaker at scientific events and advisory role by BMS, Pfizer, MSD, Ipsen, AstraZeneca, Merck.

The other authors declare to have no conflicts of interest.

Fundings

None to declare.

References

- [1] Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2022. *CA Cancer J Clin*. 2022;72(1):7-33.
- [2] Mohanty SK, Lobo A, Cheng L. The 2022 revision of World Health Organization classification of tumors of the urinary system and male genital organs: advances and challenges. *Hum Pathol*. 2022:S0046-8177(22)00224-6.
- [3] Cronin KA, Lake AJ, Scott S, Sherman RL, Noone AM, Howlader N, et al. Annual report to the Nation on the status of cancer, part I: National cancer statistics. *Cancer* 124(13): 2785-2800

- [4] Galsky MD, Chen GJ, Oh WK, Bellmunt J, Roth BJ, Petrioli R, Dogliotti L, Dreicer R, Sonpavde G. Comparative effectiveness of cisplatin-based and carboplatin-based chemotherapy for treatment of advanced urothelial carcinoma. *Ann Oncol* 2012;23(2):406-10..
- [5] Mollica V, Rizzo A, Montironi R, Cheng L, Giunchi F, Schiavina R, Santoni M, Fiorentino M, Lopez-Beltran A, Brunocilla E, Brandi G, Massari F. Current Strategies and Novel Therapeutic Approaches for Metastatic Urothelial Carcinoma. *Cancers (Basel)* 2020;12(6):1449.
- [6] Powles T, Smith K, Stenzl A, Bedke J. Immune checkpoint inhibition in metastatic urothelial cancer. *Eur Urol* 2017;72:477–481.
- [7] Galsky MD, Hahn NM, Rosenberg J, Sonpavde G, Hutson T, Oh WK, et al. A consensus definition of patients with metastatic urothelial carcinoma who are unfit for cisplatin-based chemotherapy. *Lancet Oncol* 2011;12(3):211-4.
- [8] Balar AV, Castellano D, O'Donnell PH, Grivas P, Vuky J, Powles T, Plimack ER, Hahn NM, de Wit R, Pang L, Savage MJ, Perini RF, Keefe SM, Bajorin D, Bellmunt J. First-line pembrolizumab in cisplatin-ineligible patients with locally advanced and unresectable or metastatic urothelial cancer (KEYNOTE-052): a multicentre, single-arm, phase 2 study. *Lancet Oncol* 2017;18(11):1483-1492.
- [9] Vuky J, Balar AV, Castellano D, O'Donnell PH, Grivas P, Bellmunt J, et al. Long-Term Outcomes in KEYNOTE-052: Phase II Study Investigating First-Line Pembrolizumab in Cisplatin-Ineligible Patients with Locally Advanced or Metastatic Urothelial Cancer. *J Clin Oncol* 2020;38(23):2658-2666.

- [10] Schwartz LH, Litière S, de Vries E, Ford R, Gwyther S, Mandrekar S, et al. RECIST 1.1-Update and clarification: From the RECIST committee. *Eur J Cancer* 2016; 62:132-7.
- [11] Bajorin DF, Dodd PM, Mazumdar M, Fazzari M, McCaffrey JA, Scher HI, et al. Long-term survival in metastatic transitional-cellcarcinoma and prognostic factors predicting outcomeof therapy. *J Clin Oncol* 1999;17(10):3173-81.
- [12] de Wit R. Overview of bladder cancer trials in the European Organization for Research and Treatment. *Cancer* 2003;97:2120–26.
- [13] de Santis M, Bellmunt J, Mead G, Kerst M, Leahy M, Maroto P, et al. Randomized phase II/III trial assessing gemcitabine/carboplatin and methotrexate/carboplatin/vinblastine in patients with advanced urothelial cancer ‘unfit’ for cisplatin-based chemotherapy: phase II – results of EORTC study 30986. *J Clin Oncol* 2009;27:5634–9.
- [14] Martini A, Raggi D, Fallara G, Nocera L, Schultz JG, Belladelli F, et al. Immunotherapy versus chemotherapy as first-line treatment for advanced urothelial cancer: A systematic review and meta-analysis. *Cancer Treat Rev.* 2022;104:102360.
- [15] Gupta S, Bellmunt J, Plimack ER, Sonpavde GP, Grivas P, Apolo AB. Defining “platinum-ineligible” patients with metastatic urothelial cancer (mUC). *J Clin Oncol* 2022; 40:4577-4577.

Table Legends

Table 1. Patients' characteristics.

	Patients (n=162)
Sex	
Male	117 (72)
Female	45 (28)
Age, years (y)	
Median	71
Range	43–94
ECOG Performance Status	
0	53 (33)
1	81 (50)
2	25 (16)
3	2 (1)
Current or former smokers	114 (70)
Primary tumour location	
Upper urinary tract	26 (16)
Lower urinary tract	136 (84)
Tumor histology	
Pure urothelial carcinoma	142 (88)
Variants	20 (12)
Metastatic disease	
Synchronous	44 (27)
Metachronous	119 (73)
Common sites of metastasis	
Lymph nodes	112 (69)
Lung	40 (25)
Liver	21 (13)
Bone	40 (25)
Brain	3 (2)
Visceral metastases	
Yes	90 (56)
No	72 (44)
Previous adjuvant or neoadjuvant platinum based chemotherapy	62 (38)
Second-line therapy	
Carboplatin + Gemcitabine	10 (6)
Vinflunine	5 (3)
Paclitaxel	5 (3)
Gemcitabine	2 (1)
Enfortumab vedotin	2 (1)
Clinical trials	5 (3)

Table 2. Univariate and Multivariate analyses of predictors of Progression-Free Survival and Overall Survival in cisplatin-unfit UC patients treated with first-line pembrolizumab. Statistically significant values were reported in bold.

Overall Survival	Univariate Cox Regression		Multivariable Cox regression	
	HR (95%CI)	p-value	HR (95%CI)	p-value
Gender (females vs males)	1.68 (1.04–2.72)	0.033	1.71 (1.05–2.77)	0.031
Age (≥65y vs <65y)	1.14 (0.69–1.86)	0.614		
Smokers vs no-smokers	0.88 (0.54–1.43)	0.876		
Histology (mixed vs pure UC)	1.30 (0.67–2.54)	0.440		
Upper vs Lower urinary tract	0.59 (0.28–1.24)	0.165		
Synchronous metastatic disease (yes vs no)	1.94 (1.20–3.15)	0.007	1.61 (0.97–2.66)	0.048
Bajorin risk factors	1.66 (1.16–2.37)	0.005	1.57 (1.09–2.24)	0.014
Previous neoadjuvant or adjuvant chemotherapy	1.07 (0.66–1.71)	0.791		
Progression-Free Survival	Univariate Cox Regression		Multivariable Cox regression	
	HR (95%CI)	p-value	HR (95%CI)	p-value
Gender (females vs males)	1.38 (0.88–2.18)	0.161		
Age (≥65y vs <65y)	1.10 (0.70–1.73)	0.691		
Smokers vs no-smokers	1.09 (0.68–1.74)	0.725		
Histology (mixed vs pure UC)	1.56 (0.90–2.73)	0.115		
Upper vs Lower urinary tract	0.92 (0.51–1.66)	0.790		
Synchronous metastatic disease (yes vs no)	1.71 (1.09–2.67)	0.019	1.49 (0.93–2.37)	0.094
Bajorin risk factors	1.55 (1.12–2.13)	0.007	1.43 (1.03–1.98)	0.033
Previous neoadjuvant or adjuvant chemotherapy	1.40 (0.91–2.14)	0.122		
ECOG-PS = Eastern Cooperative Oncology Group-Performance Status; UC = Urothelial Carcinoma				

Figure Legends

Figure 1. Median Overall Survival and Progression-Free Survival in cisplatin-unfit UC patients treated with first-line pembrolizumab.




Figure 2. Median Overall Survival in cisplatin-unfit UC patients treated with first-line pembrolizumab stratified by sex and ECOG-PS.




Figure 3. Median Overall Survival in cisplatin-unfit UC patients treated with first-line pembrolizumab stratified by synchronous or metachronous metastatic disease, visceral metastases, bone metastases and Bajorin risk classification.




Figure 4. Median Progression-Free Survival in cisplatin-unfit UC patients treated with first-line pembrolizumab stratified by synchronous or metachronous metastatic disease, visceral metastases, bone metastases and Bajorin risk classification.





Figure 5. Overall Survival by tumor response to therapy (RECIST 1.1).



Number at risk

Group: CR	26	26	21	15	10	5	5
Group: PR	32	27	21	14	9	6	4
Group: SD	49	29	15	10	4	3	2
Group: PD	55	20	5	2	0	0	0