

# Alma Mater Studiorum Università di Bologna Archivio istituzionale della ricerca

Pseudo-affinity capture of K. phaffii host cell proteins in flow-through mode: Purification of protein therapeutics and proteomic study

This is the final peer-reviewed author's accepted manuscript (postprint) of the following publication:

Published Version:

Sripada S.A., Elhanafi D., Collins L.B., Williams T.I., Linova M.Y., Woodley J.M., et al. (2023). Pseudoaffinity capture of K. phaffii host cell proteins in flow-through mode: Purification of protein therapeutics and proteomic study. SEPARATION AND PURIFICATION TECHNOLOGY, 326, 1-14 [10.1016/j.seppur.2023.124777].

Availability:

This version is available at: https://hdl.handle.net/11585/941760 since: 2023-09-14

Published:

DOI: http://doi.org/10.1016/j.seppur.2023.124777

Terms of use:

Some rights reserved. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (https://cris.unibo.it/). When citing, please refer to the published version.

(Article begins on next page)

- 1 Pseudo-affinity capture of *K. phaffii* host cell proteins in flow-through mode: purification of protein
- 2 therapeutics and proteomic study
- 3 Sobhana A. Sripada<sup>1</sup>, Driss Elhanafi<sup>2</sup>, Leonard B. Collins<sup>3</sup>, Taufika I. Williams<sup>3</sup>, Marina Y. Linova<sup>4</sup>, John M.
- 4 Woodley<sup>4</sup>, Cristiana Boi<sup>1,2,5,†</sup>, Stefano Menegatti<sup>1,2,†</sup>
- 5 1. Department of Chemical and Biomolecular Engineering, North Carolina State University, 911 Partners
- 6 Way, Raleigh, NC 27695, USA
- 7 2. Biomanufacturing Training and Education Center (BTEC), 850 Oval Drive, Raleigh, NC 27606, USA
- 8 3. Molecular Education, Technology, and Research Innovation Center (METRIC), North Carolina State Uni-
- 9 versity, 2620 Yarbrough Dr., Raleigh, NC 27607, USA
- 4. Department of Chemical and Biochemical Engineering, Technical University of Denmark, 240, 2800Kongens Lyngby, Denmark
- 12 5. DICAM, Alma Mater Studiorum-Università di Bologna, via Terracini 34, 40131 Bologna, Italy
- 13 <sup>†</sup> Corresponding authors: <u>cristiana.boi@unibo.it</u>, <u>smenega@ncsu.edu</u>.
- 14

15 Abstract. K. phaffii is a versatile expression system that is increasingly utilized to produce biological ther-16 apeutics - including enzymes, engineered antibodies, and gene-editing tools - that feature multiple sub-17 units and complex post-translational modifications. Two major roadblocks limit the adoption of K. phaffii 18 in industrial biomanufacturing: its proteome, while known, has not been linked to downstream process 19 operations and detailed knowledge is missing on problematic host cell proteins (HCPs) that endanger pa-20 tient safety or product stability. Furthermore, the purification toolbox has not evolved beyond the capture 21 of monospecific antibodies, and few solutions are available for engineered antibody fragments and other 22 protein therapeutics. To unlock the potential of yeast-based biopharmaceutical manufacturing, this study 23 presents the development and performance validation of a novel adsorbent – PichiaGuard – functional-24 ized with peptide ligands that target the whole spectrum of K. phaffii HCPs and designed for protein puri-25 fication in flow-through mode. The PichiaGuard adsorbent features high HCP binding capacity (~25 g per 26 liter of resin) and successfully purified a monoclonal antibody and an ScFv fragment from clarified K. 27 phaffii harvests, affording >300-fold removal of HCPs and high product yields (70 – 80%). Notably, Pich-28 iaGuard outperformed commercial ion exchange and mixed-mode resins without salt gradients or opti-29 mization in removing high-risk HCPs – including aspartic proteases, ribosomal subunits, and other pepti-30 dases – thus demonstrating its value in modern biopharmaceutical processing. 31

- Keywords. Komagataella phaffii, chromatography, protein purification, monoclonal antibodies, antibody
   fragment, proteomics.
- 34

35 1. Introduction. Strain engineering of the Pichia genus and similar yeasts has been pursued for more than 36 two decades - recently, as alternative hosts to mammalian cell lines for biomanufacturing due to benefits 37 in enabling sustainability [1,2] – by introducing plasmid designs, promoters, and signal peptides that im-38 prove productivity and fidelity. In particular, the species Komagataella phaffii (K. phaffii, synonymous to 39 Pichia pastoris) is a methylotrophic yeast that is currently utilized to express a variety of therapeutic pro-40 teins including monoclonal antibodies - either whole and engineered fragments - as well as fusion pro-41 teins, virus-like-particles, etc. [3–12]. These efforts have been in concert with systems biology to better 42 understand K. phaffii's genetic repertoire as well as the temporal evolution of mRNA and host cell proteins 43 (HCPs) across the various phases of cell growth under different conditions [7,13-20]. A recent SWOT anal-44 ysis on the adoption of K. phaffii as a platform for protein production highlighted the need to intensify 45 efforts towards improving secretion rates and reducing protease-mediated product degradation [21]. Bi-46 opharmaceutical benchmarks tracked the growth in the use of K. phaffii from its early days [22] to its

current adoption in producing five clinically approved products [23]. While capable of both intra- and
 extra-cellular protein production, *K. phaffii* has been noted for its innately slow secretory trafficking
 [24,25]. Secretion rates can be increased by tuning gene regulation [26,27] or developing strains with
 lower proteases content [28], although these modifications can impact cell growth rates and viability [29].
 Therefore, strain engineering and cultivation must be adjuvated by purification technologies capable of
 removing *K. phaffii* HCPs with enzymatic and/or immunogenic activity as early as possible in the manufacturing pipeline [29].

8 Current biotherapeutic formulations contain a final average of ~20 ng HCP per mg drug product 9 [30,31]. Although K. phaffii naturally produces fewer HCPs than its mammalian counterparts, failure to 10 reduce their titer below the prescribed limit puts at risk both process yield and product stability, and thus patient safety. Decades of production of therapeutic monoclonal antibodies (mAbs) in Chinese hamster 11 12 ovary (CHO) cells has pointed at the presence of HCPs that combine an inherent toxic or immunogenic 13 activity with the ability to escape purification by coeluting with the mAb product. Recent research has 14 assessed the harmful potential of HCPs secreted by non-mammalian cell lines and microbial strains em-15 ployed – of forthcoming – in biopharmaceutical manufacturing [32,33]. However, the identity and thresh-16 old values of potentially detrimental K. phaffii HCPs, and their ability to evade clearance by current chro-17 matographic technology have been minimally investigated [34]. For example, variants of K. phaffii have 18 been engineered with higher Kex2 expression [35] or the concomitant expression of isomerases and pro-19 teases to improve secretion; the CHO counterparts of these HCPs, however, are known as immunogenic 20 and difficult-to-remove, while other evidence points to the interference of similar HCPs with downstream 21 processing [26,27,36,37].

22 Current processes for the purification of K. phaffii-secreted proteins rely on a train of operations in-23 cluding cell separation, multiple chromatographic steps, final concentration or filtration [37-40]. An initial 24 ultrafiltration/diafiltration step is often included to adjust buffer conditions due to the methanolic growth 25 conditions as well as pH adjustment or product concentration [31,41-43]. As in the CHO pipeline, the 26 chromatographic segment typically comprises an affinity-based capture step followed by polishing via ion exchange and hydrophobic interaction [36,44-48] (the purification processes of clinically approved prod-27 28 ucts expressed by K. phaffii are outlined in **Table S1**). As the cells are harvested at high density, a liter of 29 culture fluid may contain up to few grams of HCPs [49], whose distribution of physicochemical and bio-30 chemical properties can vary with product identity and culture conditions. Such process- and product-31 dependent diversity of the K. phaffii secretome, combined with the limited knowledge of its impact on 32 purification [29,37] and the lack of affinity technology dedicated to the removal of K. phaffii HCPs, both 33 may increase the number of purification/polishing steps in the downstream train, and increases the risk 34 of contaminants present in the final product.

35 Developing a true platform process, capable of sustaining the diversity in titer and biomolecular make-36 up of industrial feedstocks, can benefit significantly by adopting resins with affinity-driven capture of HCPs 37 and host cell DNA (hcDNA). In prior work, we introduced the paradigm of "flow-through affinity chroma-38 tography" and developed an adsorbent – named LigaGuard™ [50] – functionalized with peptide ligands 39 designed to target the CHO proteome spectrum. Our team demonstrated LigaGuard<sup>™</sup> performance by 40 purifying therapeutic mAbs from an ensemble of seven CHO harvests: when utilized alone, LigaGuard™ afforded a logarithmic HCPs removal value (HCP LRV) ranging between 1.5-2.5; when coupled with an 41 42 affinity resin operated in bind-and-elute mode, the two-step process afforded cumulative HCP and hcDNA 43 LRVs ~ 4, corresponding to a final HCP level of 8 ppm. Additionally, LigaGuard™ managed to clear high-44 risk and hard-to-remove HCPs – such as Cathepsins and Phospholipases – thus demonstrating its potential 45 to de-risk and simplify the chromatographic pipeline of protein purification.

1 In this study, we sought to expand our efforts towards yeast expression systems, with *K. phaffii* as the 2 model organism. To expand our bioprocess-relevant understanding of *K. phaffii* HCPs, we conducted a 3 proteomics survey of their cultivation harvests to identify species that can cause product degradation via 4 proteolysis or instability via excipient degradation, or present the potential to trigger an immunogenic 5 response in patients. While such data has previously been curated for other commonly used cell lines used 6 for therapeutic production [51,52], we utilized this list pertaining to *K. phaffii* as a guidance to validate 7 the development of a novel purification strategy (described below).

8 We present the development of a pseudo-affinity adsorbent – PichiaGuard – for removing K. phaffii 9 HCPs and hcDNA. We leveraged the knowledge of HCP physicochemical properties to design and screen 10 a peptide library against the K. phaffii secretome to identify HCP-targeting peptides under competing con-11 ditions with a model product (herein, polyclonal IgG). Following the identification of the peptide ligands 12 and the formulation of PichiaGuard, we evaluated its global HCP binding capacity and conducted prote-13 omics analyses of the flow-through fractions to evaluate the removal of individual HR-HCPs. We finally 14 demonstrated the application of PichiaGuard by purifying ScFv and a full mAb in flow-through mode from 15 clarified K. phaffii harvests. Globally, our results indicate that PichiaGuard outperforms state-of-the-art 16 chromatographic resins in terms of both capture of K. phaffii HCPs – notably, among them, aspartic pro-17 teases, peptidases, ribosomal subunit proteins, heat shock proteins etc. – demonstrating potential for the 18 purification of high-value biotherapeutics produced by yeast hosts.

19 20

22

# 21 2. Materials and Methods

23 2.1. Materials. Fmoc-protected amino acids Fmoc-Gly-OH, Fmoc-Ser(tBu)-OH, Fmoc-Ile-OH, Fmoc-Ala-24 OH, Fmoc-Phe-OH, Fmoc-Tyr(tBu)-OH, Fmoc-Asp(OtBu)-OH, Fmoc-His(Trt)-OH, Fmoc-Arg(Pbf)-OH, Fmoc-25 Lys(Boc)-OH, Fmoc-Asn(Trt)-OH, Fmoc-Glu(OtBu)-OH, Fmoc-Pro-OH, Fmoc-Trp(Boc)-OH, Fmoc-Cys(Trt)-26 OH, and Fmoc-Leu-OH, the coupling agent azabenzotriazole tetramethyl uronium hexafluorophosphate 27 (HATU), diisopropylethylamine (DIPEA), piperidine, and trifluoroacetic acid (TFA) were sourced from 28 ChemImpex International (Wood Dale, IL, USA). The Toyopearl AF-Amino-650M (TP-650M) and HW-50F 29 (TP-50F) resins were obtained from Tosoh Bioscience (Tokyo, Japan). Ammonium bicarbonate, phosphate 30 buffered saline at pH 7.4, sodium deoxycholate, iodoacetamide (IAA), calcium chloride (CaCl<sub>2</sub>), triiso-31 propylsilane (TIPS), 1,2-ethanedithiol (EDT), anisole, Zwittergent 3-16, Protease Inhibitor Cocktail, Kaiser 32 test kits, and 3 kDa and 10 kDa MWCO centrifugal filters were from Millipore Sigma (St. Louis, MO, USA). 33 N,N'-dimethylformamide (DMF), dichloromethane (DCM), methanol, N-methyl-2-pyrrolidone (NMP), so-34 dium phosphate (monobasic), sodium phosphate (dibasic), hydrochloric acid, sodium acetate, glacial ace-35 tic acid, formic acid, Tris-HCl at pH 7.5 and 8, trypsin (100 µg), Dithiothreitol (DTT), 660 nm Histidine-rich 36 protein quantification and bicinchoninic acid (BCA) assay reagents were obtained from Fisher Chemicals 37 (Hampton, NH, USA). Human IgG was obtained from Athens Bio (Atlanta, GA). Vici Jour PEEK chromatog-38 raphy columns (2.1 mm ID, 30 mm length, 0.1 mL volume) and polyethylene frits were obtained from VWR 39 international (Radnor, PA, USA). The 10-20% Tris-Glycine HCl SDS-PAGE gels, urea, and Coomassie blue 40 stain were purchased from Bio-Rad Life Sciences (Carlsbad, CA, USA). Alexa Fluor 488 and 594 dyes, and 41 Pierce dye removal columns were obtained from Thermo Fisher Scientific. HiPrep Desalting 26/10 column 42 and Sepharose CM ion exchange resin were obtained from Cytiva.

43

44 2.2. Proteomic analysis. A label-free relative quantification workflow was developed to analyze the *K*.
 45 phaffii secretome using a bottom-up approach. Filter-aided sample preparation (FASP) [53] was per 46 formed using 30 kDa MWCO filtration units passivated with 20 μL of 0.1 M Tris HCl at pH 8. A BCA assay

was initially conducted to determine the total protein titer in each 200  $\mu$ L sample. Addition of 15  $\mu$ L of 50 1 2 mM dithiothreitol or DTT (in 0.1 M Tris HCl at pH 8) to each sample - with incubation for 30 minutes at 3 56°C - served to cleave disulfide bonds in the proteins. This was followed by further protein denaturation 4 by adding 200  $\mu$ L of 8 M urea (in 0.1 M Tris HCl at pH 8) to each sample, with vortex and spinning down. 5 Each sample was then transferred onto a separate, passivated, 30 kDa MWCO filtration unit for FASP, and 6 concentrated by centrifugation at 12,000g for 15 minutes at 21°C; the flow-through was discarded. An 7 additional 200 µL of 8 M urea was added to each filtration unit, and immediately followed by the addition 8 of 64  $\mu$ L of 200 mM iodoacetamide or IAA (in 8 M urea), resulting in a total solution volume of 264  $\mu$ L and 9 IAA concentration of 50 mM. Incubation for 60 minutes at room temperature in the dark ensured the 10 alkylation of free sulfhydryl groups of cysteine residues displayed by the HCPs. This was followed by centrifugation at 12,000g for 15 minutes at 21°C. Subsequent washing steps were performed involving mul-11 tiple rounds of centrifugation (12,000g for 15 minutes at 21°C) and resuspension into Tris-based buffers 12 13 (3 rounds of 2 M Urea and 10 mM CaCl<sub>2</sub> in 0.1 M Tris HCl at pH 8 and 3 rounds in digestion buffer 0.1 M 14 Tris HCl pH 7.5) before the addition of trypsin to the MWCO filters (1:50 enzyme/protein ratio – based on BCA assay results – in 0.1 M Tris HCl at pH 7.5) for overnight proteolytic digestion. All flow-through solu-15 16 tions up to this point were discarded.

17 The overnight digestion was disrupted using a solution of 0.001% v/v Zwittergent in aqueous 1% v/v 18 formic acid (quench buffer). This quench buffer was added in steps (with centrifugation at 12,000g for 15 19 minutes at 21°C in between steps, combining and collecting the flow-through solutions from each sample) 20 up to a total volume of 450 µL of tryptic peptide solution per sample. All samples were then lyophilized 21 and stored at -20°C until nanoLC-MS/MS. Upon reconstitution in 200 μL of Mobile Phase A (MPA: 98% v/v 22 water, 1.9% v/v acetonitrile, 0.1% v/v formic acid), reverse-phase separation of the tryptic peptide frag-23 ments was carried out using a PepMap 100 C18 trap column (3 µm particle size, 75 µm ID, 20 mm length) 24 in series with an EASY-Spray C18 analytical column (2 μm particle size, 75 μm ID, 250 mm length) - in a 25 'trap-and-elute' configuration - installed on an EASY nanoLC-1200 instrument (Thermo Scientific, San Jose, 26 CA) interfaced with an Orbitrap-Exploris-480 (Thermo Scientific, Bremen, Germany). The flowrate was 27 maintained at 300 nL/min. Peptides were eluted using a 105 minutes solvent gradient, ramping from 5% 28 to 25% Mobile Phase B (MPB: 80% v/v acetonitrile, 19.9% v/v water, 0.1% v/v formic acid) over 75 min, 29 followed by another ramp to 40% MPB over 10 min, and then a steep ramp to 95% MPB in 1 min, at which 30 point MPB was maintained at 95% for 17 minutes for column washing. Eluted tryptic peptides were ion-31 ized by subjecting them to 1.9 kV in the ion source for electrospray ionization. The ion transfer tube tem-32 perature was maintained at 275°C. The peptides were interrogated by full MS scan and data-dependent 33 acquisition (DDA) MS/MS. DDA was performed as a 105-minute nanoLC-MS/MS method. Scan parameters 34 for full MS were: 120,000 resolution, RF Lens of 40%, maximum injection time of 120 msec, scan range of 35 375-1,600 m/z, with normalized AGC setting at 300%; dynamic exclusion (with 20 sec exclusion duration) 36 was used to minimize re-interrogation of pre-sampled precursors; cycle time was set at 1.5 sec; charge 37 states 2-6 were included in the analysis; and the intensity threshold of 1.5 10<sup>4</sup> was set for ddMS<sup>2</sup>. Scan 38 parameters for ddMS<sup>2</sup> were: 15,000 resolution, inclusion window of 1.5 m/z, maximum injection time of 39 21 msec, HCD collision energy of 30% (fixed), with normalized AGC setting at 100%. Quality control in 40 nanoLC-MS/MS analyses involved blank runs, as well as standard BSA digest and standard HeLa digest 41 runs at regular intervals within the length of the sample queue. 42 Post-acquisition data analysis and automated LFQ was performed by Proteome Discoverer v. 2.4 (PD, 43 Thermo Scientific, San Jose, CA) with appropriate grouping variables. The raw nanoLC-MS/MS files were

44 interrogated against the *Pichia pastoris / Komagataella phaffii* protein database (4,983 sequences) down-

45 loaded from UniProt. The data was also searched against a contaminants database (69 sequences) to

identify potential contaminants during the experiments. These databases were searched with the follow-1 2 ing parameters: trypsin (full) as the digesting enzyme, a maximum of 3 missed trypsin cleavage sites al-3 lowed, 5 ppm precursor mass tolerance, 0.02 Da fragment mass tolerance, dynamic modifications on N-4 terminus (acetyl), methionine (oxidation, met-loss+acetyl, and met-loss), and static carbamidomethyl 5 modifications on cysteine residues. The SEQUEST HT algorithm was employed in data interrogation: the 6 algorithm compares the observed peptide MS/MS spectra and theoretically derives spectra from the tar-7 get database(s) to assign appropriate quality scores. These scores and other important predictors are 8 combined in the algorithm, which assigns a composite score to each peptide. The overall confidence in 9 protein identification is increased with increasing number of distinct amino acid sequences identified. 10 Therefore, proteins are normally categorized into a different priority group depending on whether they 11 have only one or multiple unique tryptic peptide sequences of the required peptide identification confi-12 dence. Percolator peptide validation was based on the q-value (adjusted p-value) and minimal false dis-13 covery rate (FDR) < 0.01 was selected as a condition for successful tryptic peptide assignments.

14

15 2.3. Design of K. phaffii strain X-33 producing ScFv13R4. The DNA sequence encoding the ScFv13R4 pro-16 tein [54] was first codon-optimized for efficient expression in Pichia pastoris wild type X-33 (Life Technol-17 ogies Ltd, Grand Island, NY). The gene was amplified via PCR using primers scFv13R4  $\alpha$ F1 18 (GCATGAATTCATGGCAGAAGTCCAATTAG) and scFv13R4 αR1 (GAATGCGGCCGCTCATAA-19 GACTGCCAACTTAGTACC) carrying the extra restriction sites EcoRI and NotI (New England Biolabs, Beverly, 20 MA), respectively. The PCR product was cloned in the vector pPICZ $\alpha$ -A (Life Technologies Ltd, Grand Island, 21 NY) at EcoRI and NotI sites generating a fusion protein ScFv13R4 to both His and C-myc tags present on 22 the expression vector. After electroporation in E. coli Stbl4 and selection on LB agar plates supplemented 23 with zeocin at 25 µg/mL, positive recombinant plasmids were identified using colony PCR. One recombi-24 nant plasmid was linearized with the restriction enzyme SacI and introduced into Pichia pastoris X-33 via 25 electroporation. Selection was performed on YPD plates supplemented with zeocin at a concentration of 26 1000  $\mu$ g/mL and positive clones were screened with colony PCR.

27

28 2.4. Cultivation of K. phaffii WT (null-X-33) and ScFv13R4 cultures. Overnight culture was initiated from 29 a single colony in 3 mL of YPD medium and incubated at 30°C under agitation at 350 rpm. A volume of 1 30 mL of this culture was used as the inoculum to initiate a new 50 mL culture in YPD medium, which was 31 grown at the same conditions for 24 hours. In order to adapt the cells to the production medium, cells 32 were centrifuged at 3000g for 10 minutes and resuspended in 50 mL of BFM21 media containing glycerol 33 at 40 g/L and supplemented with trace elements. This media was developed and optimized [55] as a part 34 of BTEC's training and education program; its full composition is provided in Table S4. Cell growth was 35 allowed to proceed for 24 hours at 30°C under agitation at 350 rpm. This culture was in turn used to 36 inoculate a 1-L working volume bioreactor containing BFM21 supplemented with PTM4 trace metal solu-37 tion (12 ml/L) medium and glycerol (40 g/L) starting at low optical density (OD<sub>600nm</sub> ~ 2). Fermentation was 38 conducted in multiple bioreactors using the following parameters: pH controlled at 5.0 using 28% v/v 39 ammonium hydroxide and 80% v/v phosphoric acid; temperature set point at 30°C; and dissolved oxygen 40 maintained at 30% in cascade with agitation. The initial glycerol was exhausted within 24 hours of fer-41 mentation, after which a glycerol feed phase was initiated, and the fermentation was continued until the 42 fluid reached an  $OD_{600nm} \sim 100$ . At this point, the reactor was harvested and centrifuged at 4000g for 30 43 minutes to remove the wet cell mass, while ensuring minimal cellular rupture. For ScFv13R4 production, 44 a methanol feed phase was started in a bioreactor grown similarly, by adding it at a concentration of 0.1% 45 v/v at 0.3 mL/min while maintaining the dissolved oxygen at 30%, to induce ScFv13R4 production. After 46 48 hours of fermentation in methanol and upon achieving a cell density of OD<sub>600nm</sub> ~ 300, the cells were

harvested via centrifugation at 4000g for 30 minutes. The supernatant was then filtered through a 0.45
 μm filter before further use. For experiments with mAbs, a known concentration of human polyclonal IgG

- 3 was spiked into the aforementioned null culture fluid to simulate real in-process CCF.
- 4

5 2.5. Peptide synthesis. A tetrameric one-bed one-peptide (OBOP) library in the format X<sub>1</sub>-X<sub>2</sub>-X<sub>3</sub>-X<sub>4</sub>-GSG 6 was prepared on HMBA-ChemMatrix resin via direct solid-phase Fmoc/tBu peptide synthesis using a Syro 7 I automated peptide synthesizer (Biotage, Uppsala, Sweden), as described in prior work [50,56]. The li-8 brary was constructed using 10 of the 20 natural amino acids by selecting those capable of imparting the 9 full range of interaction modes (i.e., electrostatic, hydrophobic, polar, hydrogen bonding, etc.) to the 4-10 mer peptide ligands. The amination of Toyopearl HW-50F resin was performed by incubating 1 g of resin with 2 mmol of carbonyldiimidazole in 30 mL of acetone for 30 minutes at 25°C under mild agitation, 11 12 followed by rinsing in acetone and incubation with 2 mmol of tris(2-aminoethyl) amine under the same 13 conditions, to obtain an amine density of ~0.2 mmol per gram of resin. The peptide sequences RYWV-14 GSG, QEKK-GSG, VWHH-GSG, EWAK-GSG, RYWK-GSG, YHKH-GSG, RWYQ-GSG, and WYKK-GSG were con-15 jugated on Toyopearl AF-Amino-650M (TP650M) and aminated HW-50F resins (TP50F) following the same 16 protocol using an Alstra automated peptide synthesizer (Biotage, Uppsala, Sweden). Upon completing 17 chain elongation, the peptides were deprotected using TIPS deprotection cocktail (95% v/v TFA, 2.5% v/v 18 TIPS, 2.5% v/v water). Each sequence was conjugated to the Toyopearl beads at the loading of  $\sim$ 0.2 mmol 19 of peptide per gram of resin. The PichiaGuard-TP650M and PichiaGuard-TP50F adsorbents were finally 20 formulated by mixing the eight peptide-functionalized resins.

21

22 2.6. Protein labeling and formulation of a screening mix. Human IgG and the HCPs contained in the P. 23 pastoris (X-33 strain) cell culture fluid and were labeled using NHS ester AlexaFluor dyes AF488 (green) 24 and AF594 (red), respectively. Specifically, each dye was initially dissolved in dry DMSO at the concentra-25 tion of 10 mg/mL. Following diafiltration of the X-33 cell culture fluid into phosphate buffer saline at pH 26 7.4 using 3 kDa MWCO filters, 100 µL of K. phaffii HCPs at the concentration of 0.2 mg/mL were incubated 27 with 1 µL of AF594; in parallel, 100 µL of a solution of IgG at 5 mg/mL in PBS at pH 7.4 was incubated with 28 1 μL of AF594. The labeling reactions were allowed to proceed for 3 hours at 25°C under mild agitation. 29 Spin columns were used to remove the unreacted dyes. The labeled proteins were then mixed to obtain 30 a screening mix comprising AF488-IgG at 2 mg/mL and AF594-HCPs at 0.1 mg/mL. A volume of 20 µL of 31 OBOP library beads were then incubated with 1 mL of screening mix overnight at 4°C under mild agitation 32 and washed with PBS before proceeding with the screening protocol.

33

34 2.7. Library screening. The OBOP tetrameric library was screened against the screening mix (see Section 35 2.6) using a microfluidic device developed by our team [57]. Briefly, the microfluidic device comprises an 36 imaging chamber which hosts individual library beads in rapid sequence and is installed in a DMi8 micro-37 scope (Leica, Wetzlar, Germany) equipped with an ORCA-Flash4.0 V3 Digital CMOS camera with W View 38 Gemini image splitting optics (Hamamatsu, Shizuoka, Japan). A MATLAB® GUI developed to operate this 39 screening system was implemented to visualize the library beads under the green (human IgG; excita-40 tion/emission: 488/496 nm) and red (K. phaffii HCPs; excitation/emission: 594/617 nm) channels. Library 41 beads with green-only or green-and-red fluorescent emission were discarded, whereas beads with strong 42 red-only fluorescent emission were collected as positive leads and analyzed via Edman Degradation using 43 a PPSQ 33-A Protein Sequencer (Shimadzu, Kyoto, Japan) to identify the peptide sequences carried 44 thereon.

45

2.8. Diafiltration of feedstocks. Prior to chromatographic purification, the clarified null X-33 (non-produc-1 2 ing, for control studies) cell culture harvests were diafiltered to remove small media component, chiefly 3 glycerol remnants and unreacted protease inhibitors, using a HiPrep<sup>™</sup> 26/10 column (Cytiva, Marlbor-4 ough, MA) into two different buffers, namely 20 mM sodium acetate at pH 5.7- and 25-mM sodium phos-5 phate at pH 7.1. While these buffers were chosen to investigate the effects of pH on HCP capture (pH 5.7 6 to mimic the cell culture environment and pH 7.1 as a neutral control condition), the HCP titers were 7 varied respectively between 0.3 mg/mL and 0.5 mg/mL in both the acetate-conditioned and phosphate-8 conditioned feedstocks to study in detail the purification performance of PichiaGuard resins. 9

10 2.9. Static and dynamic HCP binding capacity measurements. Static binding experiments were conducted by incubating aliquots of 30 µL of settled PichiaGuard-TP650M or PichiaGuard-TP50F resin in a 12-well 11 12 plate (WP) with acetate-conditioned and phosphate-conditioned null X-33 feedstocks featuring HCP titers 13 of 0 - 1 mg/mL for 3.5 hours at 25°C under mild agitation (300 rpm). The plate was then spun down at 14 3000 rpm for 10 minutes and the supernatants were collected and analyzed to determine the residual 15 HCP concentration via BCA. Finally, the values of static binding capacity were calculated via mass balance. 16 The dynamic binding capacity was measured via breakthrough experiments (DBC<sub>20%</sub>) by loading ~6 mL of 17 acetate-conditioned or phosphate-conditioned X-33 feedstocks featuring a HCP titer of ~0.4 mg/mL onto 18 0.1 mL column packed with chromatographic resin – namely, PichiaGuard-TP650M and PichiaGuard-19 TP50F resin, Capto Q or CaptoAdhere (Cytiva), or CHO LigaGuard™ (LigaTrap Technologies, Raleigh, NC) – 20 pre-equilibrated with the respective buffers - namely, 20 mM sodium acetate at pH 5.7 and 25 mM so-21 dium phosphate at pH 7.1 – at a residence time (RT) of 1 minute until saturation was reached. The effluent 22 was continuously monitored via spectrophotometry at 280 nm and apportioned in 0.5 mL fractions that 23 were finally analyzed to determine the residual HCP titer via BCA.

24

25 2.10. Purification of ScFv and mAb via flow-through chromatography using PichiaGuard. PichiaGuard 26 resin was initially packed in a 0.1 mL column and equilibrated with 20 mM sodium acetate at pH 5.7 at the flowrate of 0.2 mL/min for 15 minutes. Two K. phaffi harvests were utilized as feedstock in this study -27 28 the first one featuring a ScFv titer of ~50 mg/L and an HCP titer of ~0.6 mg/mL; the second one featuring 29 a (spiked) mAb titer of ~0.4 mg/mL and an HCP titer of ~0.5 mg/mL – both conditioned in 20 mM sodium 30 acetate at pH 5.7. A volume of 4 mL of harvest was loaded on the column at the RT of 1 minute and flow-31 through fractions of 0.3 mL were collected durig the load and the final column wash for analytical charac-32 terization (Sections 2.2, 2.11, and 2.12). All purification studies were performed using an ÄKTA pure 33 (Cytiva, Chicago, IL, USA) while monitoring the effluents using UV spectroscopy at 280 nm.

34

35 2.11. Reverse phase (C18 RP-HPLC) analysis for the quantification of ScFv13R4 titer and purity. The titer 36 of ScFv13R4 in the collected chromatographic fractions was measured via C18 RP-HPLC using a Zorbax 37 C18-SB column (Agilent, Santa Clara, CA) installed on a Waters Alliance 2690 HPLC. The column was equil-38 ibrated by flowing 95% mobile phase B (MPB: 0.5% v/v TFA in acetonitrile) and 5% mobile phase A (MPA: 39 0.5% v/v TFA in water), at a flowrate of 0.5 mL/min. A 20-minutes gradient method (95 – 45% MPB) was 40 conducted at the constant flow rate of 0.5 mL/min to achieve separation. Pure ScFv13R4 expressed in E. 41 coli cells was utilized to identify the product retention time and calculate the ScFv13R4 purity in the chro-42 matographic fractions.

43 **2.12.** Other Analytical Assays. The total protein and *P. pastoris* HCP titers were measured respectively via

44 BCA assay (Thermo Fisher Scientific, Waltham, MA) and Pichia pastoris HCP G.2 ELISA kit (Cygnus Tech-

nologies, Southport, SC) following the manufacturer's protocol. Product purity was evaluated via BCA assay (reading wavelength: 660 nm) for histidine-rich proteins and SDS-PAGE analysis under non-reducing
conditions. The mAb titer was measured using analytical Protein A HPLC as described previously [50].

- 4
- 5 6

#### 3. Results

7

8 3.1. Discovery of K. phaffii HCP-targeting peptide ligands and classification of host cell proteins identi-9 fied via proteomics approaches. A sequence-based bioinformatics analysis of HCPs identified in the har-10 vest fluid via proteomics approaches (Figure S1A) was utilized to curate a list of HCPs/proteins (Table S2) with an assigned risk factor to each protein depending on their immunogenic propensity or similarity to 11 12 known problematic HCPs (Figure S2). The HCPs identified to possess significant sequential similarity to 13 known CHO HCPs with a registered detrimental effect on downstream processes has also been curated 14 and presented in **Table 1**. The bioinformatics workflow has been described in detail in the supplemental 15 information. Biochemically, these proteins presents a rather unique landscape of physicochemical prop-16 erties (Figure S1B; note: all values are calculated based on the amino acid sequence of the HCPs): (i) the 17 molecular weight distribution, centered at around 46.14 kDa, is significantly sharper (standard deviation, 18  $\sigma$  ~29.03 kDa) and a narrower (7 – 193 kDa) compared to its CHO counterpart ( $\sigma$  ~ 35.91, 5 – 776 kDa, 19 [50]); (ii) the values of isoelectric point feature a distinct bimodal distribution peaking at the values of pH 20 5 and 9.5, with 40% of the species being anionic and 60% cationic at the physiological culture pH, which 21 represents a second sharp difference compared to CHO HCPs (61% anionic and 39% cationic); (iii) the 22 values of grand average hydropathy (GRAVY), varying between -1.84 and 0.49 ( $\sigma \sim 0.37$ ), indicate that K. 23 *phaffii* HCPs are more hydrophilic than CHO HCPs (-1.90 to 0.97;  $\sigma \sim 0.26$ ); and *(iv)* an average polarity of 24 50.1% across all K. phaffii HCPs, with most species between a range of 42 - 62%, indicate a strong pro-25 pensity to form hydrogen-bond networks.

26 This analysis informed the design of a library whence peptides could be selected that target the whole 27 spectrum of the K. phaffii secretome: firstly, both positively and negatively charged amino acids (Glu, Lys, 28 His, Arg; Asp was excluded from the library due to the abundance of aspartyl proteases in K. phaffii har-29 vests) were included to address anionic and cationic ensembles of HCPs; furthermore, given the abun-30 dance of polar species, neutral amino acids that are hydrogen bond-forming (Gln and the spacer Ser) were 31 included to ensure the formation of hydrogen-bonding interactions; for the same reason, only one ali-32 phatic amino acid (Val; lle and Leu were excluded) and two aromatic amino acids capable of forming hy-33 drogen-bonds (Tyr and Trp; Phe was excluded) were included; finally, two amino acid linkers, the flexible 34 Gly and the semi-rigid Ala (the rigid Pro was excluded) to sample the broadest possible range of confor-35 mations. The multi-polar composition of the library was inspired by that of the ligands forming the CHO 36 LigaGuard™, among which multi-polar peptides hold a prominent role as key contributors to the capture 37 of high-risk and persistent CHO HCPs [56]. A short peptide length of 7 total residues in the format X1-X2-38 X3-X4-G-S-G, wherein the 4 residues on the N-terminus form the HCP-binding segment and are varied 39 within the library, while the Gly-Ser-Gly tripeptide on the C-terminus acts as a flexible spacer to enhance 40 ligand display. This format was inspired by prior studies [58], where we observed that most of the energy 41 of protein:peptide binding is contributed by the 4 residues on the N-terminus, and the recommendation 42 of using PichiaGuard as a disposable – and ideally single-use – adsorbent, which requires moderate cost 43 and thus short, easy-to-synthesize peptides.

Table 1: List of K. phaffii host cell proteins flagged as "high-risk" (HR-HCPs) due to their homology to known CHO HR-HCPs. While the full classification and list has been issued in 1 2

the supplemental information, this table summarizies protein entries that were found in the K. phaffii secretome with significant similarities to previously identified CHO HCPs and

3

their observed effects on downstream processing trains (DSP).

Name	Homologous to CHO	Effects on DSP	Name	Homologous to CHO	Effects on DSP	
Protein component of the large (60S) ribosomal subunit, nearly identical to Rpl4Ap (C4QV16)		Recurrent observation in downstream processes	Enolase I, a phosphopyruvate hydra- tase that catalyzes the conversion of 2- phosphoglycerate to phosphopyruvate (C4R3H8)	Beta-enolase isoform x3	Can lead to drug product modification via catalysis of dehydration reaction	
Heat shock protein that cooperates with Ydj1p (Hsp40) and Ssa1p (Hsp70) (C4QV89)		Sorting and degradation of proteins	Peptidase A1 domain-containing pro- tein (C4R3Q7)		Protease activity	
Ubiquitin-specific protease that deubiquitinates complexes(C4QV95)		Recurrent observation in downstream processes	ATPase involved in protein folding and the response to stress (C4R3X8)	Heat shock cognate protein	Recurrent observation in downstream processes	
Phosphoglycerate kinase (C4QY07)	Phosphoglycerate Ki- nase 1	Recurrent observation in downstream processes	Aspartic protease, attached to the plasma membrane via GPI (C4R458)		Protease activity	
Protein component of the large (60S) ribosomal subunit (C4QZL4)	60S Ribosomal Subu- nit Protein	Recurrent observation in downstream processes	Putative GPI-anchored aspartic prote- ase (C4R4F5)		Protease activity	
Endoplasmic reticulum chaperone BiP (C4QZS3)	ER Chaperone BiP Precursor	Increases aggregation pro- pensity	Tetrameric phosphoglycerate mutase (C4R5P4)	Phosphoglycerate mutase 1	Recurrent observation in downstream processes	
Proteasome endopeptidase complex (C4R080)	Alpha Enolase	Leads to drug product modification via catalysis of dehydration reaction	Triosephosphate isomerase (C4R626)		Recurrent observation in downstream processes	
Alpha 6 subunit of the 20S proteasome (C4R080)	proteasome subunit alpha type 7 isoform x1	Recurrent observation in downstream processes	Vacuolar aspartyl protease (Proteinase A) (C4R6G8)	Cathepsin E		
Kex2 proprotein convertase (C4R095)		Signal peptidase for secre- tion, often overexpressed with recombinant gene	Protein component of the large (60S) ribosomal subunit (C4R6P3)		Recurrent observation in downstream processes	
Glyceraldehyde-3-phosphate dehydro- genase, isozyme 3 (C4R0P1)	Glyceraldehyde-3 Phosphate Dehydro- genase	Recurrent observation in downstream processes	Lysophospholipase (C4R703)	Lysosomal Phospho- lipase A2 (LPLA2)	Leads to polysorbate deg- radation by cleaving acyl ester bonds of glycer- ophospholipids	
GTP-binding nuclear protein (C4R0Q2)		Recurrent observation in downstream processes	40S ribosomal protein S4 (C4R7A1)		Recurrent observation in downstream processes	
Acetyl-coenzyme A synthetase (C4R1P7)	Glutaredoxin-2	Recurrent observation in downstream processes	Lon protease homolog, mitochondrial (C4R7W9)		Recurrent observation in downstream processes	
Pyruvate Kinase (C4R1P9)	Pyruvate Kinase PKM isoform X1	Potential immunogenicity due to glycolytic action	Protein component of the small (40S) ribosomal subunit (C4R7Y4)		Recurrent observation in downstream processes	
FK506-binding protein (C4R2G3)	Peptidyl-prolyl cis- trans isomerase	Increases aggregation pro- pensity by influencing fold- ing and assembly	Aspartic protease, attached to the plasma membrane via (GPI) anchor (C4R8B8)		Protease activity	
Histone H3 (C4R2J7)		Recurrent observation in downstream processes	Cytoplasmic thioredoxin isoenzyme of the thioredoxin system (C4R8R1)	Periredoxin-1 family		
Catalase (C4R2S1)		Recurrent observation in downstream processes	Protein disulfide isomerase, multifunc- tional protein in the endoplasmic retic- ulum lumen (C4R938)	Protein Disulfide Iso- merase	Affects drug product qual- ity by reduction of protein disulfide bonds	



1 2

Figure 1. Identification of HCP-binding peptides. (A) A screening mix was formulated containing red-fluorescently labeled K. 3 phaffi HCPs at the titer of 0.1 mg/mL and green fluorescently labeled IgG at the titer of 2 mg/mL; (B) aliquots of 10 µL of 4-mer 4 peptide-ChemMatrix library beads equilibrated in PBS at pH 7.4 were incubated with the screening mix for overnight at room 5 temperature; (C) the beads were washed and fed to a microfluid bead-sorting device installed in a fluorescent micro-6 scope: (D) every bead that displays high red fluorescence emission and high red-to-green signal ratio was retained, while all other 7 beads were discarded; (E) the selected beads were washed with 30% acetonitrile in water (v/v), and individually analyzed by 8 Edman degradation using a PPSQ-33A protein sequencer (Shimadzu, Kyoto, Japan) to identify the peptide sequences.

9

10 These design criteria were implemented in the synthesis of a One-Bead-One-Peptide (OBOP) combi-11 natorial library on ChemMatrix<sup>™</sup> resin, whose translucent, porous, hydrophilic beads are ideal for library 12 selection in competitive mode [56]. To ensure identifying HCP-selective ligands capable of purifying cul-13 tures producing a variety of protein products – such as full mAbs or engineered fragments [59] in flow-14 through mode, competitive screening was ensured (HCPs and product). Our discovery workflow which is 15 based on fluorescence screening using a bead-imaging-and-sorting device [57], utilizes an automated bead-selection algorithm that processes images of the beads in real time, thus enabling the sampling of a 16 17 large portion of the library and utilizes multiple fluorescence emission wavelengths to select leads with 18 high target-binding strength and selectivity. To that end, a library screening feedstock mimicking industrial 19 harvests was prepared by combining K. phaffii HCPs and human IgG – including whole antibody, Fc/Fab 20 fragments, and nanobody fragments – at a titer of 0.2 mg/mL and 2 mg/mL respectively. The HCPs and 21 IgG species were collectively labeled with a red and a green-fluorescent dye, respectively. The library 22 beads displaying high-intensity red-only fluorescent emission (i.e., strong HCP-only capture behavior at 23 thermodynamic equilibrium) were selected and the peptides carried thereupon were sequenced via Ed-24 man degradation (Figure 1). The resulting sequences were analyzed to identify position-based homology 25 of residues (Figure S3A) and amino acid frequency in the identified peptides (Figure S3B).

26 The sequences present a strong enrichment in aromatic and cationic/H-bonding residues, and a de-27 pletion in anionic and aliphatic residues. The physicochemical properties of the candidate ligands are com-28 plementary to those of *K. phaffii* HCPs, thus providing confidence in the outcome of library screening. We 29 note that, while the enrichment of cationic residues was anticipated, given the abundance of anionic 30 HCPs, the presence of a population of cationic HCPs (Figure S1B) seemed to warrant some enrichment of 31 Asp, which was not registered. Accordingly, to ensure the broadest possible HCP-binding activity, eight 32 peptides were selected – namely, RYWV, QEKK, VWHH, EWAK, RYWK, YHKH, RWYQ, WYKK – that feature 33 a diverse amino acid arrangement and composition and are thus expected to encompass a broad spec-34 trum of binding modalities.

3.2. Capture of K. phaffii HCPs via flow-through affinity chromatography using PichiaGuard. The se-1 2 lected peptides were conjugated on Toyopearl beads, a polymethacrylate-based chromatographic resin 3 whose mechanical rigidity, particle size (65  $\mu$ m), and pore diameter (100 nm) and no non-specific binding 4 (Figure S7), are ideal for protein purification via flow-through affinity chromatography [50]. The K. phaffii 5 X-33 cell culture fluid (CCF) utilized to evaluate the HCP binding activity of PichiaGuard was initially diafil-6 tered to (i) remove residual glycerol and methanol that interfere with the assays for HCP quantification 7 (Figure S3C; note: while adopted in this study to ensure analytical accuracy, diafiltration may not be re-8 quired in a regular purification train); (ii) adjust the HCP titer to different concentrations, ranging from 9 0.01 to 1.5 mg/mL to study the effect of protein concentration on binding kinetics and capacity; and (iii) 10 test buffer compositions of either 20 mM sodium acetate at pH 5.7 (2.6 mS/cm) or 20 mM sodium phos-11 phate at pH 7.1 (7 mS/cm) to study the effect of ionic strength and pH on binding kinetics and capacity. 12 These buffers were adopted owing to their kosmotropic character that ensures a native, folded protein 13 state, while promoting affinity interactions; furthermore, the pH chosen to prepare the acetate buffer 14 matches that of K. phaffii culture conditions, while the neutral pH of the phosphate buffer matches that 15 utilized for the CHO HCP-targeting LigaGuard<sup>™</sup> and provides a control to interpret the HCP capture results. 16 Static binding studies were initially conducted by incubating the PichiaGuard resin with the con-17 ditioned harvests and the residual HCP titer in solution were measured to calculate the equilibrium bind-18 ing. The resulting adsorption points were fitted against Langmuir isotherm curves (Figures 2A and 2B), 19 from which the values of maximum binding capacity (Q<sub>max</sub>) and dissociation constant (K<sub>D</sub>) for the peptide 20 mixture were derived. The difference in the Q<sub>max</sub> of 25.4 and 18.3 mg HCP per mL resin respectively regis-21 tered in acetate and phosphate buffers can be attributed in part to the different ionic strength of the 22 buffers, where the higher conductivity of phosphate buffer shields Coulomb interactions; however, the 23 observation that cationic peptide ligands provide lower binding at pH 7, where the anionic HCPs should 24 be more accentuated, suggests that the electrostatic component of binding is one, but not the dominant, 25 component of the HCP:peptide interaction in this case. The difference in Q<sub>max</sub> can also be imputed to the 26 different positions of acetate and phosphate ions in the Hofmeister series, wherein acetate (less Kosmo-27 tropic) may be promoting the formation of ligand-accessible cavities/pockets on HCPs (*i.e.*, salting in), and 28 thus potentiate target sites for hydrophobic or aromatic and polar residues in the PichiaGuard peptides 29 to respectively form hydrophobic or  $\pi$ - $\pi$ /hydrogen bond interactions. On the other hand, the acetate and 30 phosphate ions are not expected to perturb hydrogen bonds and salt bridges between HCPs and peptides 31 [60–62] at their (low) concentrations in the buffers. The combination of these effects results in a strong 32 HCP-binding activity by the peptide-functionalized resin, and ultimately a high binding capacity.

The values of  $K_D - 17.5 \mu$ M and 30  $\mu$ M measured in acetate and phosphate buffer, respectively – indicate a moderate HCP:peptide affinity. These values, however, result from treating the *K. phaffii* HCPs as a single species, a characteristic trait of the evaluation of a chromatographic technology against fluids featuring varying HCP titers and profiles, and commonly found in the literature. On the other hand, HCP capture is driven by the values of  $K_D$  of the single HCP:peptide binding event, and since each HCP is present at very low concentration ( $\leq 1$  nM), the binding affinity of the selected ligands is actually high, as observed in prior work on LigaGuard<sup>TM</sup> ligands [63].

40

**3.3.** Clearance of K. phaffii HCPs in flow-through mode: PichiaGuard vs. ion exchange resins. The purification of biopharmaceuticals expressed by K. phaffii relies almost ubiquitously on ion exchange (IEX) and mixed-mode (MM) chromatography. In particular, the post-capture steps of product polishing are being increasingly conducted in flow-through mode, which combines speed of operation with lower capital and operational costs. In this context, the IEX and MM resins employed in flow-through operations often include a quaternary amine (Q) moiety, which, as a strong cationic group, provides high binding capacity

and strength of HCPs over a broad range of pH and conductivity values [36,64]. Owing to their high capac ity, IEX resins are often utilized at the capture step when affinity resins are not available.

3 On the other hand, these resins lack binding selectivity and must be combined in a series of or-4 thogonal chromatographic steps to ensure high product purity, which requires extensive process optimi-5 zation and is often achieved at the expense of yield [65]. The paradigm of flow-through affinity chroma-6 tography as a step dedicated to abating process-related impurities holds strong promise to overcome 7 these challenges and offers an ideal route to de-risk platform processes for established products as well 8 as products that do not benefit from a dedicated affinity resin. In prior work on mAb purification in flow-9 through mode, we observed that LigaGuard™ resin outperform commercial IEX and MM resins on clarified 10 and unconditioned CHO cell culture harvests, especially due to their lack of needing process optimization [56,63,66,67]. In this study, we resolved to conduct an analogous comparison between PichiaGuard, 11 12 LigaGuard™ [63], and a commercial anion exchanger (AEX, Capto Q) and cationic MM (CaptoAdhere) res-13 ins by tracking HCP capture upon continuous loading of a clarified K. phaffii harvest [68].

To study the effect of pore size, the PichiaGuard was conjugated on two different polymethacrylate-based resins, one with pore diameter of 100 nm (Toyopearl 650M, TP-650M) and one with pore diameter of 5 nm (Toyopearl HW-50F, TP50F). Engineered antibody fragments, such as the Fab and ScFv often produced in *K. phaffii* hosts [69], feature hydrodynamic radii ~2-4 nm [70,71].

18 Accordingly, we sought to explore the combination of size exclusion of the product (based on pore 19 size) and pseudo-affinity capture of HCPs by the PichiaGuard peptides as a means to improve product 20 yield and purity under optimal conditions of loading flowrate [72]. Accordingly, we resolved to operate 21 the flow-through process at the bioprocess-relevant value of residence time (RT) of 1 minute. Finally, 22 similar to static binding tests, the harvest was conditioned in either 20 mM phosphate buffer at pH 7.1 or 23 20 mM acetate buffer at pH 5.7 (whose pH closely mimics that of the K. phaffii harvest). The flow-through 24 effluent was continuously collected and apportioned in fractions at regular intervals to evaluate the tem-25 poral profiles of protein binding – both as global HCP titers and single species via proteomics analysis.

26 The results presented in Figure 2 provide a comparison of the HCP-clearing performance of the pep-27 tide-functionalized resin (PichiaGuard) vs. AEX, CHO LigaGuard<sup>™</sup> and MM resins as a function of load in 28 the two buffer systems. The PichiaGuard outperformed both LigaGuard™ and commercial resins when 29 operated in acetate buffer at the RT of 1 min, providing a cumulative HCP LRV > 2 (i.e., > 100-fold reduction 30 of HCPs) when loaded with up to 10 CVs, followed by LRVs of 1.5 - 2 (i.e., 30-100-fold reduction) for up to 31 20 CVs, 1.3 - 1.5 (i.e., 20-30-fold reduction) for up to 40 CVs, and plateauing at 1.3 for up to 100 CVs; 32 LigaGuard<sup>™</sup> resin afforded the second best HCP clearance profile, with LRVs of 1 - 1.5 for up to 10 CVs, 33 0.75 - 1 for up to 20 CVs, and plateaued to 0.45-0.55 CV when loaded up to 40 CVs. The reference AEX and 34 MM resins provided a rather poor HCP clearance (without keen buffer or process optimization), consist-35 ently below a 10-fold reduction and plateauing to a LRV ~ 0.5 (3-fold reduction) when loaded up to 40 36 CVs. A similar performance ranking was observed in phosphate buffer, with the sole exception of the AEX 37 resin, which performed on par with the PichiaGuard resin across the entire loading range (40 CVs). The 38 magnitude of HCP LRVs, however, decreased for all resins, thus mirroring the static binding results dis-39 cussed above.

A number of conclusions can be drawn from these results: (i) peptide ligands significantly outper formed simple MM or IEX ligands, likely due to their broader range of non-covalent interactions and con formations; and (ii) library selection targeted to K. phaffii HCPs proved successful in producing an ensem ble of bespoke ligands. The LigaGuard<sup>™</sup> peptides, selected for CHO HCP capture, delivered a higher HCP
 capture than commercial resins (Table 2).

45



Figure 1: HCP binding isotherms obtained by incubating PichiaGuard TP-650M resin with solutions of K. phaffii HCPs at concentrations ranging between 0 – 1 mg/mL in either (A) 20 mM sodium acetate at pH 5.7 or (B) 25 mM sodium phosphate at pH 7.1.
Values of cumulative logarithmic removal of K. phaffii HCPs afforded by CaptoAdhere, CaptoQ, CHO LigaGuard, and PichiaGuard TP-650M resins loaded (residence time: 1 minute) with X-33 feedstocks with HCP titer of ~0.4 mg/mL and conditioned in either (C) 20 mM sodium acetate at pH 5.7 (with and without protease inhibitors) or (D) 25 mM sodium phosphate at pH 7.1. N=2 for all.

8 In turn, the PichiaGuard peptides outperformed both LigaGuard<sup>™</sup> and the MM and IEX ligands, 9 demonstrating a biorecognition activity that is superior to that of conventional mixed-mode ligands when 10 evaluated under the same conditions and grants them the status of (pseudo-)affinity ligands. Further-11 more, (iii) the PichiaGuard peptides can be utilized under conditions that mimic the native environment of K. phaffii harvests, namely acidic pH. Finally, (iv) around the 30<sup>th</sup> CV of loading, we observed the cumu-12 13 lative LRV plateau at ~1 and the corresponding fractional LRV become lower than 0.5 (Figure S4E), indi-14 cating that the adsorbent had reached saturation. The 30<sup>th</sup> CV mark corresponds to a load ratio of ~18 g 15 of HCPs per L of resin, which corresponds to ~75% of Q<sub>max</sub> (see Section 3.3). This is a reasonable outcome 16 given the relationship between dynamic vs. static binding and can be improved by adjusting the residence 17 time as well as composition and pH of the running buffer. Nonetheless, contrary to affinity resins that 18 operate in bind-and-elute mode, the loading of 'Guard' resins is intended to exist between the values of 19 DBC<sub>10%</sub> and Q<sub>max</sub>.

20

**1 Table 2.** Number of K. phaffi HR-HCPs captured by PichiaGuard, CHO LigaGuard<sup>™</sup>, and CaptoQ resins loaded with X-33 feedstocks

2 (residence time: 1 minute) with HCP titer of ~0.4 mg/mL and conditioned in either 20 mM sodium acetate at pH 5.7- or 25-mM
 3 sodium phosphate at pH 7.1.

Acetate buffer at pH 5.7 сно PichiaGuard-PichiaGuard-CaptoQ LigaGuard™ TP50F TP650M Undetected 18 19 19 18 Not captured 15 19 5 10 Captured 16 10 11 20 % 40% 42% 80% 62% Phosphate buffer at pH 7.1 PichiaGuard-СНО PichiaGuard-CaptoQ LigaGuard™ TP50F TP650M 0 8 0 0 Undetected 20 17 8 Not captured 29 Captured 36 15 15 27 % 34% 41% 61% 82%

4

5 Further improvement of HCP clearance from K. phaffii harvests using PichiaGuard resin can be 6 achieved via process optimization. Conversely, achieving comparable purification under given conditions 7 from fluids that differ substantially in the profile of physicochemical properties of their HCPs is far from a 8 foregone conclusion. We therefore resolved to measure the global HCP removal from two fluids - pro-9 duced respectively with and without the addition of protease inhibitors to the cell culture medium -10 whose distribution of HCP molecular weights differed radically (Figures 2C and 2D). To that end, we im-11 plemented a design-of-experiments (DOE) algorithm to explore systematically the effects of both categor-12 ical (i.e., buffer and resin) and continuous (i.e., buffer conductivity and pH, load ratio, and RT) variables to 13 identify parameters that maximize HCP capture (Figure S4 A-D). The choice of resin, buffer, and interac-14 tion effects between these parameters were indeed found to be significant. The effect of RT appeared to 15 be the most pronounced when using phosphate buffer: the highest HCP reduction (LRV ~3) was obtained 16 at the RT of 1 min, which was therefore adopted for the remainder of this study. Notably, the combination 17 of phosphate buffer and a smaller pore diameter of the resin led to higher HCP clearance (LRV ~ 0.8 - 1) 18 than its acetate counterpart (LRV  $\sim 0.7 - 0.9$ ), possibly due to effects of phosphate counterions on protein 19 folding and increased ability to diffuse into smaller pores; however, limiting the total capture capacity. 20 PichiaGuard-TP650M, however, afforded a superlative HCP reduction (LRV ~ 3) and emerged as the resin 21 of choice. As observed above, performing flow-through purification in acetate buffer consistently 22 achieved a higher HCP clearance, thus providing conclusive evidence for the adoption of acetate buffer 23 for resin equilibration and, ideally, conditioning of the feedstock. 24

25 3.4. Tracking the removal of high-risk HCPs from K. phaffii cell culture harvests by PichiaGuard. In the 26 established biopharmaceutical practice, the validation of a batch of therapeutic proteins consists in certi-27 fying that the residual HCP and hcDNA titers are below an acceptable limit for monoclonal antibodies 28 (mAbs), these values are 100 ppm and 10 ppb, respectively [73]. The measurement of residual HCP titer 29 has been conducted for decades with ELISA kits. Like all assays that relies on polyclonal antibodies raised 30 against multiple antigens, most ELISAs do not provide a full coverage of HCPs, but only of those that are 31 abundant or more immunogenic to the animal host; furthermore, HCP aggregation and the formation of 32 HCP:product complexes – a widespread phenomenon in bioprocess fluids – can impair assay readout [74].

These elements, combined with the variability among assay lots and operator's performance, question 1 2 the accuracy of HCP quantification provided by ELISA assays. Furthermore, in recent years, the growth of 3 proteomics in biomanufacturing has shown that product batches with acceptable global level of impurities 4 can contain amounts of single high-risk HCPs (HR-HCPs) that pose a risk to patient health (e.g., are toxic 5 or immunogenic) or can degrade the mAb or its excipients during storage, resulting in reduced efficacy or 6 harmful products [74,75]. Recent studies have amply documented that commercial chromatographic res-7 ins may struggle to remove particular HR-HCPs: a number of these "persistent" HR-HCPs have been re-8 ported on both a process- and product-basis and have been linked to delays in clinical trials and process 9 approval or the recall of drug batches. 10 Under this premise, we resolved to complement the measurements of global residual HCPs presented

above with tracking the individual HCPs captured by PichiaGuard via proteomics analysis. To this end, the 11 12 effluents obtained in both acetate and phosphate buffers at the RT of 1 min were analyzed via LC-MS/MS 13 by implementing the proteomics workflow outlined above for top-speed data dependent acquisition 14 (DDA) [76]. Spectral interrogation using the Proteome Discoverer Suite was used to perform label-free 15 quantification (LFQ) of HCPs in the flow-through fractions vs. the corresponding feedstocks (note: due to 16 the lower number of proteins identified in K. phaffii harvests compared to CHO harvests, a 'match be-17 tween runs' was implemented to preserve information from low-intensity peptide hits that that may have 18 been masked by abundant contaminants like trypsin and any media components) [77]. Briefly, (i) data 19 processing prior to analysis of variance (ANOVA) involved the exclusion of contaminants such as trypsin 20 and keratin as well as the HCPs found in the effluents but not the load sample; (ii) the abundance of 21 individual proteins was used to calculate the HCPs removal values [78]. We noted that certain proteins 22 were identified in some flow-through fractions to have an abundance ratio > 1 (defined as the ratio of 23 HCP abundance in the effluent sample vs. the load sample), suggesting that their presence in the load was 24 masked by more abundant peptides with similar sequence or charge that were not excluded.

25 We adopted three main indicators to evaluate the HCP-capture performance of different adsorbents, 26 namely (a) global concentration and reduction; (b) between-group analysis, which presents the abun-27 dance trends of every identified HCP; and (c) clearance or persistence of HR-HCPs. These indicators cap-28 ture different variations within our study group and have been used collectively to make inferences. Fig-29 ure 3 presents the global landscape of the HCPs identified across all control experiments. The between-30 group significance analysis (group: individual resin tested at an assigned composition/pH) was conducted 31 using an iterative Tukey's HSD (honestly significant difference) test, known as Newman-Kuels method, to 32 construct a 'connected-color' plot that connects fractions with a similar normalized mean abundance us-33 ing the same color and ensures discontinuity with significant 'within group' difference (p-value > 0.01). 34 This test provides an at-a-glance presentation of changes in global HCP titer compared to the load and 35 other fractions (Table S3), while other statistics and visualization techniques have been employed to track

36 changes across individual species.





Figure 2: (A) Connected colors plot indicating significant differences in mean HCP abundance among flow-through fractions; groups connected with different colors feature significantly different HCP abundances calculated as an average value for the group; (B) Contour plots of K. phaffii HCP abundance ratios registered in the effluents obtained by loading CaptoQ, LigaGuard, and PichiaGuard resins loaded (residence time: 1 minute) with X-33 feedstocks with HCP titer of ~0.4 mg/mL and conditioned in either 20 mM sodium acetate at pH 5.7 or 25 mM sodium phosphate at pH 7.1: individual HCPs are aligned on the x-axis, while the number of collected flow-through fractions is on the y-axis; red to green indicates high to low ratio of individual HCP in the effluent vs. feedstock as detected by nanoLC-MS; (C) Flowchart describing the designation of removal status of 'high-risk' HCPs (HR-HCPs); (D) Number of individual HR-HCPs removed by CaptoQ, LigaGuard, and PichiaGuard resins grouped by risk category; the values outside the Venn diagrams indicate the number of HCPs that were not removed.

1 To visualize and compare HCPs in successive flow-through fractions within and across buffer groups – 2 (acetate, phosphate) and adsorbents (PichiaGuard-TP650M, PichiaGuard-TP50F resins, CHO LigaGuard™ 3 and Capto Q resin) – Figure 3B presents a reduced-dimension view of individual HCP abundance (Uniprot 4 Accession numbers, left-to-right) vs. volume of effluent (flow-through fraction, bottom-to-top). Specifi-5 cally, abundance ratios (fraction by load) are represented as contour plots wherein ratios  $\geq$  1 (in red) mark 6 HCPs that were not cleared or whose presence was masked in the feedstock by other species but detected 7 in a flow-through fraction, whereas ratios  $\leq 1$  (veering from yellow to green as the value decreases) mark 8 HCPs that were captured in the process.

9 These results illustrate the ability of PichiaGuard resins to capture most HCPs within both buffer sys-10 tems compared to reference commercial resins. The projected contour plots provide an effective visuali-11 zation method combining results from the chromatographic process and proteomics analyses, while per-12 mitting a clear comparison between resin subjects tested at similar experimental conditions. These result-13 ing plots provide a visual indication of individual HCPs that were not captured throughout a significant 14 part of the load due to insufficient HCP:ligand binding strength or possibly due to lack of charge/polarity 15 of HCPs at the pH condition of the experiment, based on their isoelectric points.

16 Notably, the proteomics analysis of the effluents allowed us to track the capture of K. phaffii HR-HCPs 17 by the various adsorbents (Figure 3C), specifically focusing on proteins that are (i) homologous to known 18 high-risk and persistent CHO HCPs; (ii) immunogenic, either reported or predicted based on their ability 19 to generate MHC-II binding peptides; and (iii) possess proteolytic or other enzymatic activity, as outlined 20 in Section 3.1. The number of bound HR-HCPs across groups were classified according to their risk sub-21 types/bins (Figure 3D). The corresponding logarithmic values of abundance variation of all identified HR-22 HCPs are presented in Figures S5A and S5B, whereas the number of captured HR-HCPs as a function of 23 loaded volume across all groups is reported in Figure S5C.

24 From these results, the following key observations were drawn. Firstly, the PichiaGuard resins (Figure 25 3B, top row) outperformed both LigaGuard<sup>™</sup> and AEX (Capto Q) resins in retaining a higher number of K. 26 phaffii HCPs, thus corroborating the results of total HCP quantification obtained from the total protein 27 assays (see Section 3.5). It is noteworthy that this behavior is observed across all flow-through fractions 28 for the HCPs, warranting the claim of pseudo-affinity capture. As shown in Figure S5C, of the 187 HCPs 29 identified in the acetate-conditioned feedstock, 180 were captured by PichiaGuard resins throughout the 30 entire feedstock loading, while 3 – 5 were not captured; similarly, of the 318 HCPs in phosphate-condi-31 tioned feedstock (for a total of 335 unique K. phaffii HCPs identified), 302 were captured, while 8 – 12 32 were not captured. Secondly, PichiaGuard-TP650M and PichiaGuard-TP50F resin performed similarly in 33 acetate buffer, whereas in phosphate buffer PichiaGuard-TP650M clearly outperformed its TP50F coun-34 terpart. The contour plots also show a greater extent of HCP dissociation from PichiaGuard resins in ace-35 tate media - especially from TP650M resins, likely a combined effect of pore size and residence time. 36 Thirdly, PichiaGuard demonstrated the ability to clear up to 82% of HR-HCPs – including various aspartic 37 proteases, ion protease homologs, etc. – whereas LigaGuard<sup>™</sup> and CaptoQ resins only bound up to 42% 38 of them. This corroborates our claim that ion-exchange resins, despite their satisfactory global HCP cap-39 ture under optimized conditions [79], can fail to clear a number of HR-HCPs and are therefore unlikely to 40 serve effectively as an HCP-scrubbing step prior to, or in lieu of, the affinity-based product capture step. 41 Conversely, PichiaGuard makes an excellent tool for orthogonal HCP removal, thus safeguarding the per-42 formance and lifetime of product-capture resins as well as promoting product quality and stability. 43 As is to be expected with any chromatographic adsorbent, process optimization in terms of loading 44 ratio, flow rate, and buffer composition is to some extent needed. In the context of this study, however,

45 these results demonstrate the potential of PichiaGuard resin to increase the performance and robustness

of current processes as well as provide a route towards platform processes with altogether novel design
 for isolating biopharmaceutics from *Pichia* harvests.

3

4 **3.5.** Purification of ScFv13R4 and mAb from K. phaffii harvests via flow-through affinity chromatography

5 using PichiaGuard. Purification processes fully operated in flow-through mode are the ideal embodiment 6 of continuous downstream biomanufacturing: their design minimizes process footprint and the number 7 of tanks for storing service buffers, simplifies the pipework and control systems, and accelerates opera-8 tions, thus increasing productivity. Straight-through processes combining column chromatography and 9 filtration unit operations have been proposed for the continuous manufacturing of mAbs from CHO cell 10 harvests [31,68]. These processes can be successfully designed based on the size of the target products (e.g., full mAbs), which are often larger than most of the process-related impurities. Conversely, small 11 12 proteins are significantly more challenging products (listed in **Table S1**), as they exclude filtration as an 13 orthogonal separation method, thus narrowing the purification toolbox to chromatography alone. In this 14 context, pseudo-continuous processes can be utilized by integrating periodic counter-current (PCC) or 15 simulated moving bed (SMB) modes [80] encompassing ion exchange chromatography, whose complexity 16 increases both capital and operation costs. An effective alternative is offered by leveraging resins specifically tasked with HCPs and hcDNA clearance, like LigaGuard<sup>™</sup> and PichiaGuard, which provide a product-17 18 agnostic route (from the standpoint of specificity to type of therapeutic product), towards fully chroma-19 tographic straight-through processes for the purification of both mAb and non-mAb drugs.

Under this premise, having demonstrated the broad HCP-capture activity of PichiaGuard, we moved to evaluate its ability to purify therapeutic proteins from *K. phaffii* harvests in flow-through mode. To this end, we adopted an antibody fragment (ScFv13R4, MW ~30 kDa and pl of 8.36) and a full antibody (referred to as 'mAb' hereon, MW ~150 kDa and pl of 7.56) as model targets. Most importantly, to demonstrate the robustness of the technology, the *K. phaffii* harvests were loaded on PichiaGuard until reaching a protein load of >15 grams per liter of resin. The analyses of the flow-through fractions – namely, global clearance of HCP and other impurities estimated and product yield – are reported in **Table 3**.

27 The purification of ScFv13R4 returned considerable values of impurity removal, with HCP clearance 28 ranging from LRV > 2.5 (~320-fold removal) to LRV > 1.5 (~32-fold removal) when loading respectively up 29 to 3.9 and 7.5 grams per liter of resin, finally reaching LRV ~ 1 upon loading 14.4 mg/mL resin, at which a 30 DNA LRV ~1.23 was also achieved. The electrophoretic and proteomics analyses of the flow-through frac-31 tions in Figure 4 provide at-a-glance presentation of impurity removal by PichiaGuard: particularly note-32 worthy is the comparison between the feedstock and effluents containing the ScFv13R4 (Figure 4C), which 33 demonstrate the removal of contaminants both heavier and lighter than the product. Notably, the prote-34 omics analysis of the effluents (Figure 4A, B) showed that, of the total number of HCPs identified in the 35 feedstock, the number of escaping species increased from 11 at the initial collection point of 4 g/L to 36 merely 34 at the final pool, with 10 not being captured throughout. This corresponds to > 84% of HCPs 37 completely removed – chiefly among them, 38 out of 44 HR-HCPs – while the remainder ~15% HCPs were 38 partially cleared. The reverse phase chromatograms show that the product-to-impurity ratio in the efflu-39 ent increases as the loading progresses up to the value of 14 mg/mL resin, translating in a growing product 40 enrichment. In earlier work, we reported a weak-partitioning effect with peptide ligands for CHO HCP 41 capture [66], where a higher load of culture leads to better separation of the mAb product from HCP 42 contaminants. However, this beneficial effect fades at the load value of > 14 mg/mL resin with PichiaGuard 43 and this ScFv culture, likely due to the relatively low product titer in the feed (~0.06 g of ScFv13R4 per liter 44 of K. phaffi harvest vs. >1 g of mAb per liter of CHO harvest) and the formation of HCP-HCP and HCP-45 product complexes observed in the effluent (Table 3 and Figure S6B).

1 2 

 Table 3. Results of ScFv13R4 and mAb purification obtained by loading PichiaGuard-TP650 resin loaded (residence time: 1

 minute) with X-33 K. phaffi feedstock with an ScFv13R4 titer of ~ 60 mg/L, and mAb at pH 5.7, HCP titer of ~0.4 mg/mL.

	Loading (mg proteins per mL resin)	Ratio of product <i>vs.</i> non-product peaks	HCP cLRV	DNA cLRV	cYield	HCP (ppm)
ScFv13R4	Load	0.19				
	2.10	n.d.	3.00			
	3.90	n.d.	2.55	1.23		
	5.10	2.06	1.92			
	6.30	2.83	1.55		75.3%	
	7.50	1.77	1.34			
	9.30	1.40	1.14			
	14.40	1.09	1.01			
		Product Titer (mg/mL)				
mAb	Load	0.41				69643.9
	16.0	0.40	1.14	1.40	80.1%	1206.3
	Wash	0.22	1.18			

# 3

10

The yield of ScFv13R4 reached ~75% at the end of the loading phase (estimated using 660 nm BCA for 6x-His tagged ScFv13R4) but was increased to 85% by 'chasing' the load with a high-conductivity buffer (**Figure S6A**). The relationship between HCP-capture using PichiaGuard resins as a function of ScFv13R4 product concentration will be studied as the next step. Similar results were obtained from the purification of a full mAb: overall HCP clearance of 1.15 LRV was obtained up to a load of 15 mg/mL HCP with a mAb product recovery of 80% estimated using an analytical Protein A HPLC assay.



11 12



1 Collectively, these results provide proof-of-concept of PichiaGuard as a specific, impurity-removal pu-2 rification technology for *K. phaffii* cultures in a product-agnostic manner. At the same time, improvement 3 in binding selectivity via optimizing ligand combinations or pretreatment of these cultures needs to be 4 explored, along with the development of additional analytics.

5

6 7 4. Discussion and Conclusions. The success of K. phaffii in pharmaceutical biomanufacturing is scripted in 8 its rapid growth in chemically defined media, low susceptibility to virus contamination [81], facile expres-9 sion inducibility, and ability to secrete human proteins with correct post-translational modifications at 10 high titer and purity [15,98]. These upstream-related benefits call for convergent efforts in the down-11 stream toolbox, especially in the context of continuous processing and isolation of products for which 12 dedicated affinity adsorbents are not available. In this context, new chromatographic tools are needed 13 that provide orthogonality in removing process-related impurities and/or complementary to commercial 14 adsorbents in clearing residual impurities across a spectrum of concentrations to ensure the safety of 15 patients treated with complex biological drugs [83,84]. Our group has contributed to this field by intro-16 ducing 'Guard' resins, whose HCP and hcDNA capture activity provides an orthogonal step to affinity-17 based resins for product capture or an alternative to IEX and MM resins for product polishing [63,66]. In 18 this context, we leveraged the heuristic power of designed peptide libraries coupled with high-throughput 19 dual-fluorescence screening to identify peptide ligands that can capture persistent and high-risk contam-20 inants; furthermore, we integrated advanced analytics, such as proteomics, to demonstrate the potential 21 of these ligands in biomanufacturing. Our initial efforts, conducted in the domain of full mAb and CHO 22 HCPs, were supported by an established portfolio of analytical technologies and a wealth of literature 23 highlighting target HCPs.

24 In this study, we extended the 'Guard' technology to K. phaffii HCPs by introducing PichiaGuard, the 25 first pseudo-affinity adsorbent dedicated to the clearance of process-related impurities present in Pichia 26 cell culture harvests. While the selection of K. phaffii HCP-binding peptides was streamlined by a robust 27 ligand identification technology, the evaluation of the resulting PichiaGuard adsorbent faced challenges 28 related to the quantification of product recovery and contaminant removal. In particular, the shortage of 29 relevant literature on the toxicological, immunological, bioprocess-relevant aspects of Pichia HCPs en-30 couraged us to formulate criteria for identifying species that pose – or are expected to pose – a risk to 31 patient's health (e.g., immunogenicity via B-cell activation [85]) or product stability, and can compromise 32 the efficiency of the purification process (e.g., co-elution with the product or persistence throughout the 33 downstream train [51]). Having established an analytical panel, we demonstrated that PichiaGuard affords 34 up to 99% reduction of HCPs and hcDNA concentration from a native K. phaffii cell culture supernatant 35 even when challenged with a harvest containing a therapeutic product – a full mAb or an ScFv fragment 36 - thus providing product enrichment through impurity removal. Most importantly, PichiaGuard's ability 37 to clear potentially immunogenic HCPs (i.e., species documented to trigger an immune response or ho-38 mologous to immunogenic species found in other organisms, such as 40S ribosomal proteins, Histone H3, 39 Thioredoxin peroxidase, Phosphatidylinositol transfer protein, Translation initiation factor eIF4G etc.) as 40 well as proteolytic or enzymatically active HCPs (e.g., Lon protease, Aspartic proteases – vacuolar, GPI-41 anchored and plasma membrane-attached forms, Lysophospholipase, Catalase, Peptidase A1, Ydj1p co-42 operating heat shock protein etc.) is key to process robustness and product safety. 43 These results warrant a path forward for PichiaGuard, chiefly its integration with existing purification pipelines, and into a straight-through process especially in combination with commercial resins for prod-44

- 45 uct isolation. A two-column process comprising polishing or scrubbing *K. phaffii* cultures followed by af-
- 46 finity capture or focusing of the target biologic can significantly de-risk and improve purification processes

by reducing protease-mediated product alteration, improvising formulation stability, and increasing the
 performance and lifetime of the subsequent chromatographic steps via early HCP removal.

Furthermore, the discovery of these peptides as the biorecognition elements also provide the oppor tunity of using them as membrane functionalized entities – therefore providing both separation and fil tration at the same time [86,87].

6 Accordingly, future studies on PichiaGuard will focus on (i) flexibility – whether it can be utilized in a 7 fully product-agnostic manner; (ii) robustness to variability in HCP titer, complexity and composition as 8 they result from the upstream process conditions, as discussed in Section 3.1; (iii) robustness to variability 9 in conductivity pH, and product titer in the feedstocks; (iv) ability to increase product recovery by adding 10 an optimized 'chasing' wash; and (v) lifetime – namely, how effectively it can be regenerated and re-used 11 safely over a high number of cycles. Furthermore, observations of commonalities in the HCP repertoire 12 among yeasts used in biomanufacturing [20] will provide both confidence and opportunity to evaluate the 13 use of PichiaGuard across these strains for their wider adaptability in purification processes. Setting the 14 stage for such work, this study presents a cogent message that the PichiaGuard technology holds potential 15 to de-risk platform processes, reduce the cost and time-to-market of established therapeutics, and bring 16 to fruition emerging medicines (e.g., cytokines, enzymes, engineered antibodies, VLP-based vaccines) 17 [3,21,88]. In so doing, the Guard peptide ligands are not anticipated to impose major financial burdens, 18 being manufactured affordably at large scale, or regulatory concerns, given their safety and ease of clear-19 ance (small size). This strengthens our confidence in their adaptability in next-generation, truly continuous 20 processes that can affordably and reliably meet the growing global demand [31,89,90]. 21

22

Acknowledgements. The authors wish to acknowledge the generous support by the Novo Nordisk Foundation (AIM-Bio Grant NNF19SA0035474) and the North Carolina Viral Vector Initiative in Research and Learning (NC-VVIRAL) at NC State University.

26 27

28 **Conflicts of Interest.** The authors declare no conflict of interest. The funders had no role in the design of 29 the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the

the study; in the collection, analydecision to publish the results.

#### References

- [1] Ergün, B. G.; Laçın, K.; Çaloğlu, B.; Binay, B. Second Generation Pichia Pastoris Strain and Bioprocess Designs. *Biotechnol. Biofuels Bioprod.* 2022, *15*, 150.
- [2] Guo, Y.; Liao, Y.; Wang, J.; Ma, C.; Qin, J.; Feng, J.; Li, Y.; Wang, X.; Chen, K. Methylotrophy of Pichia Pastoris: Current Advances, Applications, and Future Perspectives for Methanol-Based Biomanufacturing. ACS Sustain. Chem. Eng. 2022, 10, 1741.
- [3] Harzevili, F. D. Synthetic Biology of Yeasts; 2022.
- [4] Baghban, R.; Farajnia, S.; Rajabibazl, M.; Ghasemi, Y.; Mafi, A. A.; Hoseinpoor, R.; Rahbarnia, L.; Aria, M. Yeast Expression Systems: Overview and Recent Advances. *Mol. Biotechnol.* 2019, *61*, 365.
- [5] Vuree, S. Chapter 17 Pichia Pastoris Expression System: An Impending Candidate to Express Protein in Industrial and Biopharmaceutical Domains; Singh, J., Gehlot, P. B. T.-N. and F. D. in M. B. and B., Eds.; Elsevier, 2020; pp 223.
- [6] Datta, A.; Maryala, S.; John, R.; Nylen, A.; Chen, M.-T.; Montoliu-Gaya, L.; Esquerda-Canals, G.; Bronsoms, S.; Villegas, S.; Andrade, E. V. de; Albuquerque, F. C. de; Moraes, L. M. P. de; Brígido, M. M. de M.; Santos-Silva, M. A.; Tir, N.; Heistinger, L.; Grünwald-Gruber, C.; Jakob, L. A.; Dickgiesser, S.; Rasche, N.; Mattanovich, D.; Li, H.; Sethuraman, N.; Stadheim, T. A.; Zha, D.; Prinz, B.; Ballew, N.; Bobrowicz, P.; Choi, B.-K.; Cook, W. J.; Cukan, M.; Houston-Cummings, N. R.; Davidson, R.; Gong, B.; Hamilton, S. R.; Hoopes, J. P.; Jiang, Y.; Kim, N.; Mansfield, R.; Nett, J. H.; Rios, S.; Strawbridge, R.; Wildt, S.; Gerngross, T. U. Optimization of Humanized IgGs in Glycoengineered Pichia Pastoris. *Nat. Biotechnol.* 2000, *128*, 891.
- Peg, B.; Pharmaceuticals, E.; Valli, M.; Grillitsch, K.; Grünwald-Gruber, C.; Tatto, N. E.; Hrobath, B.;
   Klug, L.; Ivashov, V.; Hauzmayer, S.; Koller, M.; Tir, N.; Leisch, F.; Gasser, B.; Graf, A. B.; Altmann,
   F.; Daum, G.; Mattanovich, D. A Subcellular Proteome Atlas of the Yeast Komagataella Phaffii.
   *Bioconjugate Tech.* 2013, *20*, xv.
- [8] Tir, N.; Heistinger, L.; Grünwald-Gruber, C.; Jakob, L. A.; Dickgiesser, S.; Rasche, N.; Mattanovich,
   D. From Strain Engineering to Process Development: Monoclonal Antibody Production with an
   Unnatural Amino Acid in Pichia Pastoris. *Microb. Cell Fact.* 2022, *21*, 157.
- [9] Nylen, A.; Chen, M. T. Production of Full-Length Antibody by Pichia Pastoris. *Methods Mol. Biol.* 2018, *1674*, 37.
- [10] Andrade, E. V. de; Albuquerque, F. C. de; Moraes, L. M. P. de; Brígido, M. de M.; Santos-Silva, M.
   A. Single-Chain Fv with Fc Fragment of the Human IgG1 Tag: Construction, Pichia Pastoris Expression and Antigen Binding Characterization1. J. Biochem. 2000, 128, 891.
- [11] Montoliu-Gaya, L.; Esquerda-Canals, G.; Bronsoms, S.; Villegas, S. Production of an Anti-Aβ Antibody Fragment in Pichia Pastoris and in Vitro and in Vivo Validation of Its Therapeutic Effect. PLoS One 2017, 12, e0181480.
- Li, H.; Sethuraman, N.; Stadheim, T. A.; Zha, D.; Prinz, B.; Ballew, N.; Bobrowicz, P.; Choi, B.-K.;
   Cook, W. J.; Cukan, M.; Houston-Cummings, N. R.; Davidson, R.; Gong, B.; Hamilton, S. R.; Hoopes,
   J. P.; Jiang, Y.; Kim, N.; Mansfield, R.; Nett, J. H.; Rios, S.; Strawbridge, R.; Wildt, S.; Gerngross, T. U.
   Optimization of Humanized IgGs in Glycoengineered Pichia Pastoris. *Nat. Biotechnol.* 2006, *24*, 210.
- [13] Huang, C. J.; Damasceno, L. M.; Anderson, K. A.; Zhang, S.; Old, L. J.; Batt, C. A. A Proteomic Analysis of the Pichia Pastoris Secretome in Methanol-Induced Cultures. *Appl. Microbiol. Biotechnol.* 2011, 90, 235.
- [14] Hou, R.; Gao, L.; Liu, J.; Liang, Z.; Zhou, Y. J.; Zhang, L.; zhang, Y. Comparative Proteomics Analysis of Pichia Pastoris Cultivating in Glucose and Methanol. *Synth. Syst. Biotechnol.* 2022, *7*, 862.

- [15] Lin, X. Q.; Liang, S. L.; Han, S. Y.; Zheng, S. P.; Ye, Y. R.; Lin, Y. Quantitative ITRAQ LC-MS/MS Proteomics Reveals the Cellular Response to Heterologous Protein Overexpression and the Regulation of HAC1 in Pichia Pastoris. J. Proteomics 2013, 91, 58.
- [16] Zahrl, R. J.; Peña, D. A.; Mattanovich, D.; Gasser, B. Systems Biotechnology for Protein Production in Pichia Pastoris. *FEMS Yeast Res.* 2017, *17*, 1.
- [17] Nieto-Taype, M. A.; Garcia-Ortega, X.; Albiol, J.; Montesinos-Seguí, J. L.; Valero, F. Continuous Cultivation as a Tool Toward the Rational Bioprocess Development With Pichia Pastoris Cell Factory. *Front. Bioeng. Biotechnol.* 2020, *8*, 1.
- [18] Brady, J. R.; Whittaker, C. A.; Tan, M. C.; Kristensen 2nd, D. L.; Ma, D.; Dalvie, N. C.; Love, K. R.; Love, J. C. Comparative Genome-Scale Analysis of Pichia Pastoris Variants Informs Selection of an Optimal Base Strain. *Biotechnol. Bioeng.* 2020, *117*, 543.
- [19] Love, K. R.; Shah, K. A.; Whittaker, C. A.; Wu, J.; Bartlett, M. C.; Ma, D.; Leeson, R. L.; Priest, M.; Borowsky, J.; Young, S. K.; Love, J. C. Comparative Genomics and Transcriptomics of Pichia Pastoris. BMC Genomics 2016, 17, 1.
- [20] Mattanovich, D.; Graf, A.; Stadlmann, J.; Dragosits, M.; Redl, A.; Maurer, M.; Kleinheinz, M.; Sauer, M.; Altmann, F.; Gasser, B. Genome, Secretome and Glucose Transport Highlight Unique Features of the Protein Production Host Pichia Pastoris. *Microb. Cell Fact.* 2009, *8*, 29.
- [21] de Sá Magalhães, S.; Keshavarz-Moore, E. P. Pastoris (Komagataella Phaffii) as a Cost-effective Tool for Vaccine Production for Low- and Middle-income Countries (Lmics). *Bioengineering* 2021, *8*.
- [22] Walsh, G. Biopharmaceutical Benchmarks 2014. *Nat. Biotechnol.* 2014, *32*, 992.
- [23] Walsh, G.; Walsh, E. Biopharmaceutical Benchmarks 2022. Nat. Biotechnol. 2022, 40, 1722.
- [24] Love, K. R.; Politano, T. J.; Panagiotou, V.; Jiang, B.; Stadheim, T. A.; Love, J. C. Systematic Single-Cell Analysis of Pichia Pastoris Reveals Secretory Capacity Limits Productivity. *PLoS One* 2012, *7*, 1.
- [25] Staudacher, J.; Rebnegger, C.; Dohnal, T.; Landes, N.; Mattanovich, D.; Gasser, B. Going beyond the Limit: Increasing Global Translation Activity Leads to Increased Productivity of Recombinant Secreted Proteins in Pichia Pastoris. *Metab. Eng.* 2022, 70, 181.
- [26] Fischer, J. E.; Glieder, A. Current Advances in Engineering Tools for Pichia Pastoris. *Curr. Opin. Biotechnol.* 2019, *59*, 175.
- [27] Juturu, V.; Wu, J. C. Heterologous Protein Expression in Pichia Pastoris: Latest Research Progress and Applications. *ChemBioChem* 2018, *19*, 7.
- [28] Ahmad, M.; Winkler, C. M.; Kolmbauer, M.; Pichler, H.; Schwab, H.; Emmerstorfer-Augustin, A. Pichia Pastoris Protease-Deficient and Auxotrophic Strains Generated by a Novel, User-Friendly Vector Toolbox for Gene Deletion. *Yeast* 2019, *36*, 557.
- [29] Zhang, Y.; Liu, R.; Wu, X. The Proteolytic Systems and Heterologous Proteins Degradation in the Methylotrophic Yeast Pichia Pastoris. *Ann. Microbiol.* 2007, *57*, 553.
- [30] Molden, R.; Hu, M.; Yen E, S.; Saggese, D.; Reilly, J.; Mattila, J.; Qiu, H.; Chen, G.; Bak, H.; Li, N. Host Cell Protein Profiling of Commercial Therapeutic Protein Drugs as a Benchmark for Monoclonal Antibody-Based Therapeutic Protein Development. *MAbs* 2021, 13.
- [31] Crowell, L. E.; Lu, A. E.; Love, K. R.; Stockdale, A.; Timmick, S. M.; Wu, D.; Wang, Y. A.; Doherty, W.; Bonnyman, A.; Vecchiarello, N.; Goodwine, C.; Bradbury, L.; Brady, J. R.; Clark, J. J.; Colant, N. A.; Cvetkovic, A.; Dalvie, N. C.; Liu, D.; Liu, Y.; Mascarenhas, C. A.; Matthews, C. B.; Mozdzierz, N. J.; Shah, K. A.; Wu, S. L.; Hancock, W. S.; Braatz, R. D.; Cramer, S. M.; Love, J. C. On-Demand Manufacturing of Clinical-Quality Biopharmaceuticals. *Nat. Biotechnol.* 2018, *36*, 988.
- [32] De Groot, A. S. ISPRI EpiVax.

- [33] Duke, B. R.; Mitra-Kaushik, S. Current In Vitro Assays for Prediction of T Cell Mediated Immunogenicity of Biotherapeutics and Manufacturing Impurities. *J. Pharm. Innov.* 2019, *15*, 202.
- [34] Matthews, C. B.; Wright, C.; Kuo, A.; Colant, N.; Westoby, M.; Love, J. C. Reexamining Opportunities for Therapeutic Protein Production in Eukaryotic Microorganisms. *Biotechnol. Bioeng.* 2017, 114, 2432.
- [35] Yang, S.; Kuang, Y.; Li, H.; Liu, Y.; Hui, X.; Li, P.; Jiang, Z.; Zhou, Y.; Wang, Y.; Xu, A.; Li, S.; Liu, P.; Wu,
   D. Enhanced Production of Recombinant Secretory Proteins in Pichia Pastoris by Optimizing Kex2
   P1' Site. *PLoS One* 2013, *8*, e75347.
- [36] Heiss, S.; Maurer, M.; Hahn, R.; Mattanovich, D.; Gasser, B. Identification and Deletion of the Major Secreted Protein of Pichia Pastoris. *Appl. Microbiol. Biotechnol.* 2013, *97*, 1241.
- [37] Velez-Suberbie, M. L.; Morris, S. A.; Kaur, K.; Hickey, J. M.; Joshi, S. B.; Volkin, D. B.; Bracewell, D. G.; Mukhopadhyay, T. K. Holistic Process Development to Mitigate Proteolysis of a Subunit Rotavirus Vaccine Candidate Produced in Pichia Pastoris by Means of an Acid PH Pulse during Fed-Batch Fermentation. *Biotechnol. Prog.* 2020, *36*, 1.
- [38] Rajamanickam, V.; Krippl, M.; Herwig, C.; Spadiut, O. An Automated Data-Driven DSP Development Approach for Glycoproteins from Yeast. *Electrophoresis* 2017, *38*, 2886.
- [39] Crowell, L. E.; Goodwine, C.; Holt, C. S.; Rocha, L.; Vega, C.; Rodriguez, S. A.; Dalvie, N. C.; Tracey, M. K.; Puntel, M.; Wigdorovitz, A.; Parreño, V.; Love, K. R.; Cramer, S. M.; Love, J. C. Development of a Platform Process for the Production and Purification of Single-Domain Antibodies. *Biotechnol. Bioeng.* 2021, *118*, 3348.
- [40] Maurer, M. M.; Schillinger, H. Primary Recovery of Yeast Culture Supernatant for Recombinant Protein Purification. *Methods Mol. Biol.* 2019, *1923*, 335.
- [41] Motif-Foodworks. GRAS Notice for Myoglobin Preparation https://www.fda.gov/media/153921/download (accessed Apr 30, 2023).
- [42] Hudspeth, E. M.; Wang, Q.; Seid, C. A.; Hammond, M.; Wei, J.; Liu, Z.; Zhan, B.; Pollet, J.; Heffernan, M. J.; McAtee, C. P.; Engler, D. A.; Matsunami, R. K.; Strych, U.; Asojo, O. A.; Hotez, P. J.; Bottazzi, M. E. Expression and Purification of an Engineered, Yeast-Expressed Leishmania Donovani Nucleoside Hydrolase with Immunogenic Properties. *Hum. Vaccin. Immunother.* 2016, *12*, 1707.
- [43] Zhan, Y.; Wei, Y.; Chen, P.; Zhang, H.; Liu, D.; Zhang, J.; Liu, R.; Chen, R.; Zhang, J.; Mo, W.; Zhang, X. Expression, Purification and Biological Characterization of the Extracellular Domain of CD40 from Pichia Pastoris. *BMC Biotechnol.* 2016, *16*, 8.
- [44] Capela, E. V; Magnis, I.; Rufino, A. F. C. S.; Torres-Acosta, M. A.; Aires-Barros, M. R.; Coutinho, J. A.
   P.; Azevedo, A. M.; e Silva, F. A.; Freire, M. G. Using Three-Phase Partitioning for the Purification and Recovery of Antibodies from Biological Media. *Sep. Purif. Technol.* 2023, *316*, 123823.
- [45] Malpiedi, L. P.; Nerli, B. B.; Abdala, D. S. P.; Pessôa-Filho, P. de A.; Pessoa, A. Aqueous Micellar Systems Containing Triton X-114 and Pichia Pastoris Fermentation Supernatant: A Novel Alternative for Single Chain-Antibody Fragment Purification. Sep. Purif. Technol. 2014, 132, 295.
- [46] Zhang, C.; Fredericks, D.; Campi, E. M.; Florio, P.; Jespersgaard, C.; Schiødt, C. B.; Hearn, M. T. W. Purification of Monoclonal Antibodies by Chemical Affinity Mixed Mode Chromatography. *Sep. Purif. Technol.* 2015, *142*, 332.
- [47] Rosa, S. A. S. L.; dos Santos, R.; Aires-Barros, M. R.; Azevedo, A. M. Phenylboronic Acid Chromatography Provides a Rapid, Reproducible and Easy Scalable Multimodal Process for the Capture of Monoclonal Antibodies. *Sep. Purif. Technol.* 2016, *160*, 43.
- [48] Fan, X.; Zhou, Y.; Yu, J.; Li, F.; Chen, Q.; Song, H.; Feng, D.; Liu, W.; Nian, R.; Xian, M. Non-Affinity

Purification of a Nanobody by Void-Exclusion Anion Exchange Chromatography and Multimodal Weak Cation Exchange Chromatography. *Sep. Purif. Technol.* 2019, *225*, 88.

- [49] Burgard, J.; Grünwald-Gruber, C.; Altmann, F.; Zanghellini, J.; Valli, M.; Mattanovich, D.; Gasser, B. The Secretome of Pichia Pastoris in Fed-Batch Cultivations Is Largely Independent of the Carbon Source but Changes Quantitatively over Cultivation Time. *Microb. Biotechnol.* 2020, *13*, 479.
- [50] Sripada, S. A.; Chu, W.; Williams, T. I.; Teten, M. A.; Mosley, B. J.; Carbonell, R. G.; Lenhoff, A. M.; Cramer, S. M.; Bill, J.; Yigzaw, Y.; Roush, D. J.; Menegatti, S. Towards Continuous MAb Purification: Clearance of Host Cell Proteins from CHO Cell Culture Harvests via "Flow-through Affinity Chromatography" Using Peptide-Based Adsorbents. *Biotechnol. Bioeng.* 2022, *119*, 1873.
- Jones, M.; Palackal, N.; Wang, F.; Gaza-Bulseco, G.; Hurkmans, K.; Zhao, Y.; Chitikila, C.; Clavier, S.;
   Liu, S.; Menesale, E.; Schonenbach, N. S.; Sharma, S.; Valax, P.; Waerner, T.; Zhang, L.; Connolly, T.
   "High-risk" Host Cell Proteins (HCPs): A Multi-company Collaborative View. *Biotechnol. Bioeng.* 2021.
- [52] Disela, R.; Bussy, O. Le; Geldhof, G.; Pabst, M.; Ottens, M. Characterisation of the E. Coli HMS174 and BLR Host Cell Proteome to Guide Purification Process Development. *Biotechnol. J.* 2023, *n/a*, 2300068.
- [53] Wiśniewski, J. R. Filter Aided Sample Preparation A Tutorial. *Anal. Chim. Acta* 2019, *1090*, 23.
- [54] Martineau, P.; Jones, P.; Winter, G. Expression of an Antibody Fragment at High Levels in the Bacterial Cytoplasm11Edited by J. Karn. *J. Mol. Biol.* 1998, *280*, 117.
- [55] Omonde, A. K. Defined Media Design for High Cell Density Pichia Pastoris Fermentations in Shake Flasks, a 24 Well Microreactor and a 30L Bioreactor. *BTEC Masters Thesis* 2014.
- [56] Lavoie, R. A.; Fazio, A.; Williams, T. I.; Carbonell, R.; Menegatti, S. Targeted Capture of Chinese Hamster Ovary Host Cell Proteins: Peptide Ligand Binding by Proteomic Analysis. *Biotechnol. Bioeng.* 2020, *117*, 438.
- [57] Saberi-Bosari, S.; Omary, M.; Lavoie, A.; Prodromou, R.; Day, K.; Menegatti, S.; San-Miguel, A. Affordable Microfluidic Bead-Sorting Platform for Automated Selection of Porous Particles Functionalized with Bioactive Compounds. *Sci. Rep.* 2019, *9*, 7210.
- [58] Menegatti, S.; Bobay, B. G.; Ward, K.; Islam, T.; Kish, W. S.; Naik, A.; Carbonell, R. G. C. ". Design of Protease-Resistant Peptide Ligands for the Purification of Antibodies from Human Plasma. J. Chromatogr. A 2016, 1445, 93.
- [59] Gasser, B.; Prielhofer, R.; Marx, H.; Maurer, M.; Nocon, J.; Steiger, M.; Puxbaum, V.; Sauer, M.; Mattanovich, D. Pichia Pastoris : Protein Production Host and Model Organism for Biomedical Research. *Future Microbiol.* 2013, *8*, 191.
- [60] Zhang, Y.; Cremer, P. S. Interactions between Macromolecules and Ions: The Hofmeister Series. *Curr. Opin. Chem. Biol.* 2006, *10*, 658.
- [61] Omta, A. W.; Kropman, M. F.; Woutersen, S.; Bakker, H. J. Negligible Effect of Ions on the Hydrogen-Bond Structure in Liquid Water. *Science* 2003, *301*, 347.
- [62] Batchelor, J. D.; Olteanu, A.; Tripathy, A.; Pielak, G. J. Impact of Protein Denaturants and Stabilizers on Water Structure. *J. Am. Chem. Soc.* 2004, *126*, 1958.
- [63] Sripada, S. A.; Chu, W.; Williams, T. I.; Teten, M. A.; Mosley, B. J.; Carbonell, R. G.; Lenhoff, A. M.; Cramer, S. M.; Bill, J.; Yigzaw, Y.; Roush, D. J.; Menegatti, S. Towards Continuous MAb Purification: Clearance of Host Cell Proteins from CHO Cell Culture Harvests via "Flow-through Affinity Chromatography" Using Peptide-Based Adsorbents. *Biotechnol. Bioeng.* 2022, *119*, 1873.
- [64] Sánchez-Trasviña, C.; Flores-Gatica, M.; Enriquez-Ochoa, D.; Rito-Palomares, M.; Mayolo-Deloisa,

K. Purification of Modified Therapeutic Proteins Available on the Market: An Analysis of Chromatography-Based Strategies . *Frontiers in Bioengineering and Biotechnology* . 2021.

- [65] Castro, L. S.; Lobo, G. S.; Pereira, P.; Freire, M. G.; Neves, M. C.; Pedro, A. Q. Interferon-Based Biopharmaceuticals: Overview on the Production, Purification, and Formulation. *Vaccines* 2021, *9*.
- [66] Lavoie, R. A.; Chu, W.; Lavoie, J. H.; Hetzler, Z.; Williams, T. I.; Carbonell, R.; Menegatti, S. Removal of Host Cell Proteins from Cell Culture Fluids by Weak Partitioning Chromatography Using Peptide-Based Adsorbents. *Sep. Purif. Technol.* 2021, *257*, 117890.
- [67] Lavoie, R.; di Fazio, A.; Blackburn, R.; Goshe, M.; Carbonell, R.; Menegatti, S. Targeted Capture of Chinese Hamster Ovary Host Cell Proteins: Peptide Ligand Discovery. *Int. J. Mol. Sci.* 2019, *20*, 1729.
- [68] Ichihara, T.; Ito, T.; Kurisu, Y.; Galipeau, K.; Gillespie, C. Integrated Flow-through Purification for Therapeutic Monoclonal Antibodies Processing. *MAbs* 2018, *10*.
- [69] Tripathi, N. K.; Shrivastava, A. Recent Developments in Bioprocessing of Recombinant Proteins: Expression Hosts and Process Development. *Frontiers in Bioengineering and Biotechnology*. Frontiers Media S.A. December 20, 2019, p 420.
- [70] Thomas, G. D. Effect of Dose, Molecular Size, and Binding Affinity on Uptake of Antibodies BT -Drug Targeting: Strategies, Principles, and Applications; Francis, G. E., Delgado, C., Eds.; Humana Press: Totowa, NJ, 2000; pp 115.
- [71] Lüdel, F.; Bufe, S.; Bleymüller, W. M.; de Jonge, H.; Iamele, L.; Niemann, H. H.; Hellweg, T. Distinguishing Between Monomeric ScFv and Diabody in Solution Using Light and Small Angle X-Ray Scattering. *Antibodies*. 2019.
- [72] Noh, H.; Vogler, E. A. Volumetric Interpretation of Protein Adsorption: Competition from Mixtures and the Vroman Effect. *Biomaterials* 2007, *28*, 405.
- [73] Molden, R.; Hu, M.; Yen E, S.; Saggese, D.; Reilly, J.; Mattila, J.; Qiu, H.; Chen, G.; Bak, H.; Li, N. Host Cell Protein Profiling of Commercial Therapeutic Protein Drugs as a Benchmark for Monoclonal Antibody-Based Therapeutic Protein Development. *MAbs* 2021, 13.
- [74] Pilely, K.; Johansen, M. R.; Lund, R. R.; Kofoed, T.; Jørgensen, T. K.; Skriver, L.; Mørtz, E. Monitoring Process-Related Impurities in Biologics – Host Cell Protein Analysis. 2022, 747.
- [75] Bracewell, D. G.; Francis, R.; Smales, C. M. The Future of Host Cell Protein (HCP) Identification During Process Development and Manufacturing Linked to a Risk-Based Management for Their Control. *Biotechnol. Bioeng* 2015, *112*, 1727.
- [76] Wiśniewski, J. R.; Zougman, A.; Nagaraj, N.; Mann, M. Universal Sample Preparation Method for Proteome Analysis. *Nat. Methods* 2009, *6*, 359.
- [77] Tyanova, S.; Temu, T.; Cox, J. The MaxQuant Computational Platform for Mass Spectrometry-Based Shotgun Proteomics. *Nat. Protoc.* 2016, *11*, 2301.
- [78] Marcus, K. *Quantitative Methods in Proteomics*.
- [79] Singh, N.; Arunkumar, A.; Peck, M.; Voloshin, A. M.; Moreno, A. M.; Tan, Z.; Hester, J.; Borys, M. C.; Li, Z. J. Development of Adsorptive Hybrid Filters to Enable Two-Step Purification of Biologics. MAbs 2017, 9, 350.
- [80] Patil, R.; Walther, J. Continuous Manufacturing of Recombinant Therapeutic Proteins: Upstream and Downstream Technologies BT - New Bioprocessing Strategies: Development and Manufacturing of Recombinant Antibodies and Proteins; Kiss, B., Gottschalk, U., Pohlscheidt, M., Eds.; Springer International Publishing: Cham, 2018; pp 277.
- [81] Karbalaei, M.; Rezaee, S. A.; Farsiani, H. Pichia Pastoris: A Highly Successful Expression System for Optimal Synthesis of Heterologous Proteins. *J. Cell. Physiol.* 2020, *235*, 5867.

- [82] Heiss, S.; Puxbaum, V.; Gruber, C.; Altmann, F.; Mattanovich, D.; Gasser, B. Multistep Processing of the Secretion Leader of the Extracellular Protein Epx1 in Pichia Pastoris and Implications for Protein Localization. *Microbiology* 2015, *161*, 1356.
- [83] Jawa, V.; Hall, M.; Flynn, G. Evaluating Immunogenicity Risk Due to Host Cell Protein Impurities in Antibody-Based Biotherapeutics. *AAPS J.* 2016, *18*, 1439.
- [84] Jawa, V.; Terry, F.; Gokemeijer, J.; Mitra-Kaushik, S.; Roberts, B. J.; Tourdot, S.; De Groot, A. S. T-Cell Dependent Immunogenicity of Protein Therapeutics Pre-Clinical Assessment and Mitigation– Updated Consensus and Review 2020. *Front. Immunol.* 2020, *11*, 1.
- [85] FDA. Immunogenicity of Protein-based Therapeutics https://www.fda.gov/vaccines-bloodbiologics/biologics-research-projects/immunogenicity-protein-based-therapeutics.
- [86] Gupta, S. K.; Shukla, P. Sophisticated Cloning, Fermentation, and Purification Technologies for an Enhanced Therapeutic Protein Production: A Review. *Front. Pharmacol.* 2017, *8*, 1.
- [87] Gilgunn, S.; El-Sabbahy, H.; Albrecht, S.; Gaikwad, M.; Corrigan, K.; Deakin, L.; Jellum, G.; Bones, J. Identification and Tracking of Problematic Host Cell Proteins Removed by a Synthetic, Highly Functionalized Nonwoven Media in Downstream Bioprocessing of Monoclonal Antibodies. J. Chromatogr. A 2019, 1595, 28.
- [88] Shah, K. A.; Clark, J. J.; Goods, B. A.; Politano, T. J.; Mozdzierz, N. J.; Zimnisky, R. M.; Leeson, R. L.; Love, J. C.; Love, K. R. Automated Pipeline for Rapid Production and Screening of HIV-Specific Monoclonal Antibodies Using Pichia Pastoris. *Biotechnol. Bioeng.* 2015, *112*, 2624.
- [89] Goodrick, J. C.; Xu, M.; Finnegan, R.; Schilling, B. M.; Schiavi, S.; Hoppe, H.; Wan, N. C. High-Level Expression and Stabilization of Recombinant Human Chitinase Produced in a Continuous Constitutive Pichia Pastoris Expression System. *Biotechnol. Bioeng.* 2001, *74*, 492.
- [90] Brady, J. R.; Love, J. C. Alternative Hosts as the Missing Link for Equitable Therapeutic Protein Production. *Nat. Biotechnol.* 2021, *39*, 403.